



HAL
open science

Asymptotic behavior of the number of Eulerian orientations of graphs

Mikhail Isaev

► **To cite this version:**

Mikhail Isaev. Asymptotic behavior of the number of Eulerian orientations of graphs. 2011. hal-00631251v2

HAL Id: hal-00631251

<https://hal.science/hal-00631251v2>

Preprint submitted on 22 Dec 2011 (v2), last revised 10 Sep 2012 (v6)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Asymptotic behavior of the number of Eulerian orientations of graphs

M.I. Isaev

Moscow Institute of Physics and Technology, 141700 Dolgoprudny, Russia
Centre de Mathématiques Appliquées, Ecole Polytechnique, 91128 Palaiseau, France
e-mail: isaev.m.i@gmail.com

Abstract

We consider the class of simple graphs with large algebraic connectivity (the second-smallest eigenvalue of the Laplacian matrix). For this class of graphs we determine the asymptotic behavior of the number of Eulerian orientations. In addition, we establish some new properties of the Laplacian matrix, as well as an estimate of a conditionality of matrices with the asymptotic diagonal predominance

1. Introduction

The eulerian orientation of the graph G is the orientation of its edges such that for every vertex the number of incoming edges and outgoing edges are equal. We denote the number of Eulerian orientations $EO(G)$. It is easy clear that $EO(G) = 0$, If the degree of at least one vertex of G is odd. Eulerian orientations of the complete graph K_n are called regular tournaments.

In [6] it is shown that for even $n \rightarrow \infty$

$$EO(K_n) = \left(\frac{2^{n+1}}{\pi n} \right)^{(n-1)/2} n^{1/2} e^{-1/2} (1 + O(n^{-1/2+\varepsilon})) \quad (1.1)$$

for any $\varepsilon > 0$.

Undirected graphs without loops and multiple edges are called simple.

The problem of counting the number of the Eulerian orientations of an undirected simple graph is complete for the class $\#P$. Thus this problem is difficult in terms of complexity theory. The problem of counting Eulerian orientations can be reduced to counting perfect matching for a class of graphs for which it can be done approximately with high probability in polynomial time, see [4].

For the simple graph G we define $n \times n$ matrix Q by

$$Q_{jk} = \begin{cases} -1, & (v_j, v_k) \in EG, \\ d_j, & j = k, \\ 0, & \text{otherwise,} \end{cases} \quad (1.2)$$

where $n = |VG|$ and d_j denotes the degree of $v_j \in VG$. The matrix $Q = Q(G)$ is called the Laplacian matrix of the graph G . The eigenvalues $\lambda_0 \leq \lambda_1 \leq \dots \leq \lambda_{n-1}$ of the matrix Q are always non-negative real numbers and $\lambda_0 = 0$. The eigenvalue λ_1 is called the algebraic connectivity of the graph G . (For more information about the spectral properties of the Laplace matrix see, for example, [2] and [7].)

According to the Kirchhoff's Matrix-Tree-Theorem, see [3], we have that

$$t(G) = \frac{1}{n} \lambda_1 \lambda_2 \cdots \lambda_{n-1}, \quad (1.3)$$

where $t(G)$ denotes the number of spanning trees of the graph G .

In the present work we generalize approach of [6]. We determine the asymptotic behavior of the number of Eulerian orientations of simple graphs with large algebraic connectivity, see Theorem 2.1. In Section 2 we give conventions and notations and formulate the main result.

In Section 3 we prove some basic properties of the Laplacian matrix. Some statements seem to be of independent interest, for example, we obtain an estimate of a conditionality of matrices with the asymptotic diagonal predominance, see Lemma 3.2. Also, we prove the following property of simple graphs with large algebraic connectivity: when you remove the vertex the number of spanning trees decreases by no more than cn times, for some $c > 0$ depending only on λ_1/n , see. Corollary 3.3.

We prove the main result in Section 4. We express $EO(G)$ in terms of an n -dimensional integral using Cauchy's formula. The derivation of asymptotic estimation of this integral uses three lemmas (Lemma 4.1, Lemma 4.2, Lemma 4.3) from [1], whose proofs we omit in the text for reasons of volume.

2. Main result

Let $p \geq 1$ be a real number and $\mathbf{x} \in \mathbb{R}^n$. We use notation

$$\|\mathbf{x}\|_p = \left(\sum_{j=1}^n |x_j|^p \right)^{1/p}. \quad (2.1)$$

For $p = \infty$ we have the maximum norm

$$\|\mathbf{x}\|_\infty = \max_j |x_j|. \quad (2.2)$$

The matrix norm corresponding to the p -norm for vectors is

$$\|A\|_p = \sup_{\mathbf{x} \neq 0} \frac{\|A\mathbf{x}\|_p}{\|\mathbf{x}\|_p}. \quad (2.3)$$

One can show that for symmetric matrix A and $p \geq 1$

$$\|A\|_p \geq \|A\|_2. \quad (2.4)$$

For invertible matrices one can define the condition number.

$$\mu(A) \stackrel{def}{=} \|A\| \cdot \|A^{-1}\| \geq \|AA^{-1}\| = 1. \quad (2.5)$$

If f is bounded both above and below by g asymptotically, we use the notation

$$f(n) = \Theta_{k_1, k_2}(g(n)), \quad (2.6)$$

which implies as $n \rightarrow \infty$, eventually

$$k_1|g(n)| \leq |f(n)| \leq k_2|g(n)|. \quad (2.7)$$

When functions f and g depend not only on n , but also on other parameters ξ , we use notation (2.6) meaning that condition (2.7) holds uniformly for all possible values of ξ .

The main result of the present work is the following theorem.

Theorem 2.1. *Let G be simple graph with n vertices having even degree and algebraic connectivity $\lambda_1 \geq \sigma n$ for some $\sigma > 0$. Then as $n \rightarrow \infty$*

$$EO(G) = \Theta_{k_1, k_2} \left(2^{|EG| + \frac{n-1}{2}} \pi^{-\frac{n-1}{2}} / \sqrt{t(G)} \right), \quad (2.8)$$

where d_j denotes the degree of v_j , $t(G)$ denotes the number of spanning trees of the graph G and constants $k_1, k_2 > 0$ depend only on σ .

Remark 2.1. Taking into account (1.3), the value $t(G)$ can be represented as the principal minor of the Laplacian matrix Q .

Remark 2.2. For the complete graph $\lambda_1 = n$, $EK_n = \frac{n(n-1)}{2}$ and $t(K_n) = n^{n-2}$. The result of Theorem 2.1 for this case is in agreement with asymptotic formula (1.1).

Remark 2.3. There is the result on the asymptotic behavior of Eulerian circuits analogous to Theorem 2.1, see [1].

3. Some basic properties of the Laplacian matrix

In what follows we suppose that

$$G \text{ is a simple graph.} \quad (3.1)$$

The Laplacian matrix Q of the graph G has the eigenvector $[1, 1, \dots, 1]^T$, corresponding to the eigenvalue $\lambda_0 = 0$. We use notation $\hat{Q} = Q + J$, where J denotes the matrix with every entry 1. Note that Q and \hat{Q} have the same set of eigenvectors and eigenvalues, except for the eigenvalue corresponding to the eigenvector $[1, 1, \dots, 1]^T$, which equals 0 for Q and n for \hat{Q} .

Using (2.4), we get that

$$\lambda_{n-1} = \|Q\|_2 \leq \|\hat{Q}\|_2 \leq \|\hat{Q}\|_1 = \max_j \sum_{k=1}^n |\hat{Q}_{jk}| = n. \quad (3.2)$$

We denote by G_r the graph which arises from G by removing vertices v_1, v_2, \dots, v_r and all adjacent edges.

Lemma 3.1. *Let condition (3.1) holds for graph G with n vertices. Then*

$$\lambda_1(G) \leq \frac{n}{n-1} \min_j d_j, \quad (3.3)$$

$$\lambda_1(G_r) \geq \lambda_1(G) - r, \quad (3.4)$$

where $\lambda_1(G)$ is the algebraic connectivity of G and d_j is the degree of the vertex $v_j \in VG$.

The proof of Lemma 3.1 can be found in [2].

Lemma 3.2. *Let $a > 0$ and I be identity $n \times n$ matrix. Then for any n and $n \times n$ symmetric matrix X such that the matrix $I + X$ is nonsingular and $|X_{ij}| \leq a/n$,*

$$\mu_2(I + X) \leq \mu_\infty(I + X) \leq C\mu_2(I + X), \quad (3.5)$$

where C depends only on a . (does not depend on n)

Proof of Lemma 3.2. The left-hand side of (3.5) follows from (2.4). We order the eigenvalues of $I + X$ modulo

$$|\chi_1| \leq |\chi_2| \leq \dots \leq |\chi_n|. \quad (3.6)$$

Using (2.4), we get that

$$|\chi_n| = \|I + X\|_2 \leq \|I + X\|_\infty \leq \|I\|_\infty + \|X\|_\infty \leq 1 + a. \quad (3.7)$$

We consider $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$ such that $\|\mathbf{x}\|_\infty = 1$. For simplicity, we assume that $x_1 = \|\mathbf{x}\|_\infty = 1$. We denote by $\mathcal{J} = \left\{ j \mid x_j > \frac{1}{2a} \right\}$.

Case 1. $|\mathcal{J}| < \frac{n}{4a}$. Estimating the first coordinate of $(I + X)\mathbf{x}$, we get that

$$\|(I + X)\mathbf{x}\|_\infty \geq x_1 - \frac{a}{n} \left(\sum_{j \in \mathcal{J}} x_j + \sum_{j \notin \mathcal{J}} x_j \right) \geq 1 - \frac{a}{n} \left(\frac{n}{4a} \cdot 1 + n \cdot \frac{1}{2a} \right) = \frac{1}{4} \|\mathbf{x}\|_\infty. \quad (3.8)$$

Case 2. $|\mathcal{J}| \geq \frac{n}{4a}$. Note that

$$\sqrt{n} \|(I + X)\mathbf{x}\|_\infty^2 \geq \|(I + X)\mathbf{x}\|_2 \geq |\chi_1| \cdot \|\mathbf{x}\|_2 \geq |\chi_1| \cdot \sqrt{|\mathcal{J}| \cdot \frac{1}{4a^2}} \|\mathbf{x}\|_\infty. \quad (3.9)$$

Then

$$\|(I + X)\mathbf{x}\|_\infty \geq \frac{|\chi_1|}{4a^{3/2}} \|\mathbf{x}\|_\infty. \quad (3.10)$$

Combining (3.8) и (3.10), we get that at least one of the following inequalities holds.

$$\|(I + X)^{-1}\|_\infty \leq 4 \quad \text{or} \quad \|(I + X)^{-1}\|_\infty \leq \frac{4a^{3/2}}{|\chi_1|}. \quad (3.11)$$

From (3.7) we have that

$$\|I + X\|_\infty \leq 1 + a. \quad (3.12)$$

Taking into account $|\chi_n| \leq 1 + a$ and $\mu_2(I + X) = \frac{|\chi_n|}{|\chi_1|} \geq 1$, we obtain (3.5). \blacksquare

The following property of the Laplacian matrix is not used explicitly, but the proofs of Lemma 1, Lemma 2 and Lemma 3 are based on it (the proofs are given in [1]).

Corollary 3.1. *Let the assumptions of Theorem 2.1 hold. Then there is a constant $c_\infty > 0$ depending only on σ such that*

$$\|\hat{Q}^{-1}\|_1 = \|\hat{Q}^{-1}\|_\infty \leq \frac{c_\infty}{n}. \quad (3.13)$$

Proof of Corollary 3.1. Using (3.3), we get that

$$d_j \geq \lambda_1 \frac{n-1}{n} \geq \sigma(n-1) \geq \sigma n/2. \quad (3.14)$$

Taking into account (3.2), all eigenvalues of \hat{Q} are in the interval $[\sigma n; n]$. Inequality (3.13) follows easily from the assertion of Lemma 3.2 for the matrix $\Omega^T \hat{Q} \Omega$, where

$$\Omega_{jk} = \begin{cases} \frac{1}{\sqrt{d_j+1}}, & \text{if } j = k, \\ 0, & \text{otherwise.} \end{cases} \quad (3.15)$$

■

The following lemmas will be applied to estimate the determinant of a matrix close to the identity matrix I .

Lemma 3.3. *Let X be an $n \times n$ matrix such that $\|X\|_2 < 1$. Then for fixed $m \geq 2$*

$$\det(I + X) = \exp \left(\sum_{r=1}^{m-1} \frac{(-1)^{r+1}}{r} \operatorname{tr}(X^r) + E_m(X) \right), \quad (3.16)$$

where tr is the trace function and

$$|E_m(X)| \leq \frac{n}{m} \frac{\|X\|_2^m}{1 - \|X\|_2}. \quad (3.17)$$

Lemma 3.3 was also formulated and proved in [5].

Lemma 3.4. *Let G be a simple graph with n vertices and the algebraic connectivity $\lambda_1 \geq \sigma n$ for some $\sigma > 0$. Let G_1 be the graph which arises from G by removing vertex v_1 and all adjacent edges. Then there is a constant $c > 0$ depending only on σ such that*

$$\det \hat{Q}_1 \geq \frac{\det \hat{Q}}{cn}. \quad (3.18)$$

Proof of Lemma 3.4. Note that the matrix M_{11} that results from deleting the first row and the first column of \hat{Q} coincides with the matrix \hat{Q}_1 with the exception of the diagonal elements. Let Ω be a diagonal matrix such that

$$\Omega_{jj} = \begin{cases} 1, & \text{if } (v_1, v_j) \in EG, \\ 0, & \text{otherwise.} \end{cases} \quad (3.19)$$

Define the matrix X by

$$X_{jk} = \begin{cases} \frac{1}{d_1+1}, & \text{if } (v_1, v_j) \notin EG, (v_1, v_k) \notin EG \text{ and } j, k \neq 1 \\ 0, & \text{otherwise.} \end{cases} \quad (3.20)$$

After performing one step of the Gaussian elimination for $\hat{Q} + \Omega + X$, we obtain that

$$\det(\hat{Q} + \Omega + X) = (d_1 + 1) \det \hat{Q}_1, \quad (3.21)$$

Using (2.4), (3.14), we have that

$$\|\Omega + X\|_2 \leq \|\Omega\|_2 + \|X\|_2 \leq \|\Omega\|_2 + \|X\|_1 \leq 1 + \frac{n}{d_1 + 1} \leq \frac{3}{\sigma}. \quad (3.22)$$

Since the algebraic connectivity $\lambda_1 \geq \sigma n$

$$\|(\Omega + X)\hat{Q}^{-1}\|_2 \leq \|\Omega + X\|_2 \|\hat{Q}^{-1}\|_2 \leq \frac{3}{\sigma\lambda_1} \leq \frac{3}{\sigma^2 n} \quad (3.23)$$

Combining Lemma 3.3 and (3.23), we get that as $n \rightarrow \infty$

$$\begin{aligned} \det\left(I + (\Omega + X)\hat{Q}^{-1}\right) &= \exp\left(\operatorname{tr}\left((\Omega + X)\hat{Q}^{-1}\right) + E_2\left((\Omega + X)\hat{Q}^{-1}\right)\right) \geq \\ &\geq \exp\left(-n\frac{3}{\sigma^2 n} + O(n^{-1})\right). \end{aligned} \quad (3.24)$$

Using (3.21) and (3.24) we get that as $n \rightarrow \infty$

$$(d_1 + 1) \det \hat{Q}_1 = \det\left(I + (\Omega + X)\hat{Q}^{-1}\right) \det \hat{Q} \geq \det \hat{Q} \exp\left(-3/\sigma^2 + O(n^{-1})\right). \quad (3.25)$$

Since $d_1 + 1 \leq n$, we obtain (3.18). ■

Corollary 3.2. *Let the assumptions of Lemma 3.4 hold. Let G_r be the graph which arises from G by removing vertices v_1, v_2, \dots, v_r and all adjacent edges. Then there is a constant $c_1 > 0$ depending only on σ such that*

$$\det \hat{Q}(G_r) \geq \frac{\det \hat{Q}(G)}{(c_1 n)^r} \quad (3.26)$$

for any $r \leq \sigma n/2$.

Proof of Corollary 3.2. From (3.4) we have that

$$\lambda_1(G_r) \geq \sigma n - r \geq \sigma n/2. \quad (3.27)$$

Using r times the assertion of the Lemma 3.4, we obtain (3.29). ■

According to (1.3)

$$t(G) = \frac{1}{n} \lambda_1 \lambda_2 \cdots \lambda_{n-1} = \frac{\det \hat{Q}}{n^2}, \quad (3.28)$$

then the following proposition holds.

Corollary 3.3. *Let the assumptions of Lemma 3.4 hold. Then there is a constant $c > 0$ depending only on σ such that*

$$t(G_1) \geq \frac{t(G)}{cn}, \quad (3.29)$$

where $t(G)$ denotes the number of spanning trees of the graph G .

Lemma 3.5. *Let $a > 0$ and the assumptions of Lemma 3.4 hold. Then for any set $A \subset VG$ such, that $|A| \geq an$, there is a function $h : VG \rightarrow \mathbb{N}_0$, having following properties:*

$$h(v) = 0, \text{ if } v \in A, \quad h(v) \leq H, \text{ for any } v \in VG, \quad (3.30)$$

$$\left| \{w \in VG \mid (w, v) \in EG \text{ and } h(w) < h(v)\} \right| \geq \alpha n, \text{ if } v \notin A, \quad (3.31)$$

where constants $H, \alpha > 0$ depend only on a and σ .

Proof of Lemma 3.5. At first, we construct the set $A_1 = \{v \in VG \mid h(v) = 1\}$, having property (3.31).

If $|A| > n - \sigma n/4$, then let $A_1 = \{v \in VG \mid v \notin A\}$. Taking into account (3.14), we get that property (3.31) hold for $\alpha = \sigma/4$. In this case $H = 1$.

For $|A| \leq n - \sigma n/4$ define $\mathbf{x} \in \mathbb{R}^n$ such that

$$x_j = \begin{cases} 1 - |A|/n, & v_j \in A, \\ -|A|/n, & v_j \notin A. \end{cases} \quad (3.32)$$

Since $x_1 + x_2 + \dots + x_n = 0$

$$\mathbf{x}^T Q \mathbf{x} = \mathbf{x}^T \hat{Q} \mathbf{x} \geq \lambda_1 \|\mathbf{x}\|_2^2 \geq \lambda_1 |A| \left(\frac{n - |A|}{n} \right)^2 \geq \sigma n \alpha n (\sigma/4)^2 = \frac{a\sigma^3 n^2}{16}. \quad (3.33)$$

On the other hand,

$$\mathbf{x}^T Q \mathbf{x} = \sum_{(v_j, v_k) \in EG} (x_j - x_k)^2, \quad (3.34)$$

which is equal to the number of edges $(v, w) \in EG$, where $v \in A, w \notin A$. We denote A_1 the set of vertices $w \notin A$, having at least αn adjacent vertices in A , where $\alpha = \frac{1}{32} a\sigma^3$.

$$\mathbf{x}^T Q \mathbf{x} \leq n|A_1| + \alpha n|VG|. \quad (3.35)$$

Combining (3.33) and (3.35), we get that $|A_1| \geq \alpha n$.

We make further construction of the function h inductively, using for the k -th step the set $A^{(k)} = A \cup A_1 \cup \dots \cup A_k$. The number of steps does not exceed $1/\alpha$ as $|A_k| \geq \alpha n$ for each step, perhaps with the exception of the last one. ■

4. Proof of Theorem 2.1

Note that the function

$$\prod_{(v_j, v_k) \in EG} (x_j^{-1} x_k + x_k^{-1} x_j) \quad (4.1)$$

is the generating function the number of orientations of a graph G by the differences in the numbers of incoming and outgoing edges at each vertex. The value $EO(G)$ is the constant term, which we can extract via Cauchy's Theorem using the unit circle as a contour for each variable. Making the substitution $x_j = e^{i\theta_j}$ for each j , we find that

$$EO(G) = 2^{|EG|} \pi^{-n} S, \quad S = \int_{U_n(\pi/2)} \prod_{(v_j, v_k) \in EG} \cos \Delta_{jk} d\boldsymbol{\theta}, \quad (4.2)$$

where $\Delta_{jk} = \theta_j - \theta_k$,

$$U_n(R) = \{(x_1, x_2, \dots, x_n) \mid |x_j| \leq R \text{ для всех } j\}, \quad (4.3)$$

and using the fact that for graphs with vertices of even degree the integrand is unchanged by the substitutions $\theta_j \rightarrow \theta_j + \pi$.

Let's start the evaluation S from the part that makes a major contribution to the integral. We fix some sufficiently small constant $\varepsilon > 0$. Let

$$V_0 = \left\{ \boldsymbol{\theta} \in U_n(\pi/2) \mid |\theta_j - \bar{\theta}| \pmod{\pi} \leq n^{-1/2+\varepsilon}, \text{ где } \bar{\theta} = \frac{\theta_1 + \dots + \theta_n}{n} \right\}. \quad (4.4)$$

By Taylor's theorem we have that for $\boldsymbol{\theta} \in V_0$

$$\prod_{(v_j, v_k) \in EG} \cos \Delta_{jk} = \exp \left(-\frac{1}{2} \sum_{(v_j, v_k) \in EG} \Delta_{jk}^2 - \frac{1}{12} \sum_{(v_j, v_k) \in EG} \Delta_{jk}^4 + O(n^{-1+6\varepsilon}) \right). \quad (4.5)$$

We denote by S_0 the contribution to S in the integration over the region V_0 .

Lemma 4.1. *Let G be a simple graph with n vertices and the algebraic connectivity $\lambda_1 \geq \sigma n$ for some $\sigma > 0$. Then for any $a, b > 0$ as $n \rightarrow \infty$*

$$\int_{V_0} \exp \left(-a \sum_{(v_j, v_k) \in EG} \Delta_{jk}^2 - b \sum_{(v_j, v_k) \in EG} \Delta_{jk}^4 \right) d\boldsymbol{\theta} = \Theta_{k_1, k_2} \left(n \int_{\mathbb{R}^n} e^{-a\boldsymbol{\theta}^T \hat{Q} \boldsymbol{\theta}} d\boldsymbol{\theta} \right), \quad (4.6)$$

where constants $k_1, k_2 > 0$ depend only on a, b and σ .

Lemma 4.1 is also formulated and proved in [1] (see Lemma 8.3).

Combining (3.28), (4.2), (4.5), (4.6) and

$$\int_{\mathbb{R}^n} e^{-a\boldsymbol{\theta}^T \hat{Q} \boldsymbol{\theta}} d\boldsymbol{\theta} = \pi^{n/2} a^{-n/2} / \sqrt{\det \hat{Q}}, \quad (4.7)$$

we get that

$$S_0 = \Theta_{k_1, k_2} \left(2^{\frac{n-1}{2}} \pi^{\frac{n+1}{2}} / \sqrt{t(G)} \right), \quad (4.8)$$

where constants $k_1, k_2 > 0$ depend only on σ .

Thus it remains to show that the other parts are negligible. One can show that

$$|\cos(x)| \leq \exp\left(-\frac{1}{2}x^2\right) \text{ for } |x| \leq \frac{9}{16}\pi. \quad (4.9)$$

Divide the interval $[-\frac{1}{2}\pi, \frac{1}{2}\pi] \pmod{\pi}$ into 32 equal intervals H_0, \dots, H_{31} such that $H_0 = [-\frac{1}{64}\pi, \frac{1}{64}\pi]$. For each j , define the region $W_j \subseteq U_n(\pi/2)$ as the set of points $\boldsymbol{\theta} \in U_n(\pi/2)$, having at least $\frac{1}{32}n$ coordinates in H_j . Clearly, the W_j 's cover $U_n(\pi/2)$ and also each W_j can be mapped to W_0 by a uniform translation of the $\theta_j \pmod{\pi}$. This mapping preserves the integrand of (4.2), and also maps V_0 to itself, so we have that

$$\int_{U_n(\pi/2) - V_0} \prod_{(v_j, v_k) \in EG} \cos \Delta_{jk} d\boldsymbol{\theta} \leq 32Z, \quad (4.10)$$

where

$$Z = \int_{W_0 - V_0} \prod_{(v_j, v_k) \in EG} |\cos \Delta_{jk}| d\boldsymbol{\theta}. \quad (4.11)$$

We proceed by defining integrals S_1, S_2, S_3 in such a way that Z is obviously bounded by their sum. We then show that each of them is negligible. Let

$$\begin{aligned} V_1 &= \{\boldsymbol{\theta} \in W_0 \mid |\theta_j| \geq \frac{1}{32}\pi \text{ для менее чем } n^\varepsilon \text{ различных } j\}, \\ V_2 &= \{\boldsymbol{\theta} \in V_1 \mid |\theta_j| \geq \frac{1}{16}\pi \text{ для по крайней мере одного } j\}. \end{aligned} \quad (4.12)$$

Then our three integrals can be defined as

$$\begin{aligned} S_1 &= \int_{W_0 - V_1} \prod_{(v_j, v_k) \in EG} |\cos \Delta_{jk}| d\boldsymbol{\theta}, \\ S_2 &= \int_{V_2} \prod_{(v_j, v_k) \in EG} |\cos \Delta_{jk}| d\boldsymbol{\theta}, \\ S_3 &= \int_{V_1 - V_2 - V_0} \prod_{(v_j, v_k) \in EG} |\cos \Delta_{jk}| d\boldsymbol{\theta}. \end{aligned} \quad (4.13)$$

We begin with S_1 . Let h be the function from Lemma 3.5 for the set $A = \{v_j \mid |\theta_j| \leq \frac{1}{64}\pi\}$. We denote l_{min} such natural number that inequality

$$|\theta_j| \geq \frac{1}{64}\pi(1 + l/H) \quad (4.14)$$

holds for at least n^ε/H indices of the set $\{j \mid h(v_j) = l\}$. Existence of l_{min} follows from the definition of the region V_1 . If θ_j and θ_k are such that

$$|\theta_j| \geq \frac{1}{64}\pi(1 + l_{min}/H) \quad \text{and} \quad |\theta_k| \leq \frac{1}{64}\pi(1 + (l_{min} - 1)/H) \quad (4.15)$$

or vice versa, we have that $|\cos \Delta_{jk}| \leq \cos(\frac{1}{64}\pi/H)$. This includes at least $(\alpha n - n^\varepsilon)n^\varepsilon/H$ edges $(v_j, v_k) \in EG$. Using (3.2) and (3.28), we get that as $n \rightarrow \infty$

$$S_1 \leq \pi^n \left(\cos \frac{\pi}{64H} \right)^{(\alpha n - n^\varepsilon)n^\varepsilon/H} = O\left(\exp(-cn^{1+\varepsilon}) \right) 2^{\frac{n-1}{2}} \pi^{\frac{n+1}{2}} / \sqrt{t(G)} \quad (4.16)$$

for some constant $c > 0$, depending only on σ .

For $1 \leq r \leq n^\varepsilon$ let $S_2(r)$ denote the contribution to S_2 of those $\boldsymbol{\theta} \in V_2$ such that $|\theta_j| \geq \frac{1}{16}\pi$ for exactly r values of j . If $|\theta_j| \leq \frac{1}{32}\pi$ and $|\theta_k| \geq \frac{1}{16}\pi$ or vice versa, we have that

$$|\cos \Delta_{jk}| \leq \cos\left(\frac{1}{32}\pi\right) \quad (4.17)$$

This includes at least $r(\sigma n/2 - n^\varepsilon)$ edges $(v_j, v_k) \in EG$, because the degree of any vertex of the graph G is at least $\sigma n/2$, see (3.14). For pairs (j, k) such that $|\theta_j|, |\theta_k| \leq \frac{1}{16}\pi$, we use (4.9). We put $\boldsymbol{\theta}' = (\theta_1, \dots, \theta_{n-r})$.

$$S_2(r) \leq \pi^r \left(\cos \frac{\pi}{32} \right)^{r(\sigma n/2 - n^\varepsilon)} \sum_{G_r} \int_{U_{n-r}(\pi/2)} \exp\left(-\frac{1}{2} \sum_{(v_j, v_k) \in EG_r} \Delta_{jk}^2 \right) d\boldsymbol{\theta}', \quad (4.18)$$

where the first sum is over graphs, arises from G by removing all possible sets of r vertices.

Lemma 4.2. *Let the assumptions of Lemma 3.4 hold. Then*

$$\int_{U_n(\pi/2)} \exp\left(-\frac{1}{2} \sum_{(v_j, v_k) \in EG} \Delta_{jk}^2\right) d\boldsymbol{\theta} \leq \frac{2^{\frac{n-1}{2}} \pi^{\frac{n+1}{2}} n}{\sqrt{\det \hat{Q}}}. \quad (4.19)$$

Lemma 4.2 is also formulated and proved in [1] (see Lemma 6.1). Using Lemma 3.1 and Corollary 3.2, we get that

$$\lambda_1(G_r) \geq \sigma n - n^\varepsilon, \quad (4.20)$$

$$\det \hat{Q}(G_r) \geq \frac{\det \hat{Q}}{(c_1 n)^r}. \quad (4.21)$$

According to Lemma 4.2 we have that

$$\int_{U_{n-r}(\pi/2)} \exp\left(-\frac{1}{2} \sum_{(v_j, v_k) \in EG_r} \Delta_{jk}^2\right) d\boldsymbol{\theta}' \leq \frac{2^{\frac{n-r-1}{2}} \pi^{\frac{n-r+1}{2}} n}{\sqrt{\det \hat{Q}(G_r)}}. \quad (4.22)$$

Combining (4.18) with (4.21), (4.22) and allowing n^r for the choice of the set of r vertices for G_r , we get that

$$S_2(r) \leq 2^{\frac{n-r-1}{2}} \pi^{\frac{n-r+1}{2}} n^{r+1} \left(\cos \frac{\pi}{32}\right)^{r(\sigma n/2 - n^\varepsilon)} \frac{(c_1 n)^{r/2}}{\sqrt{\det \hat{Q}}}. \quad (4.23)$$

Using (3.2) and summing over $0 \leq r \leq n^\varepsilon$, we find that

$$S_2 = \sum_{r=1}^{n^\varepsilon} S_2(r) = O(c^{-n}) 2^{\frac{n-1}{2}} \pi^{\frac{n+1}{2}} / \sqrt{t(G)} \quad (4.24)$$

for some constant $c > 1$, depending only on σ .

Note that $\Delta_{jk} \leq \frac{1}{8}\pi$ for $\boldsymbol{\theta} \in V_1 - V_2$. Thus

$$V_1 - V_2 \subset V_3 = \{\boldsymbol{\theta} \in U_n(\pi/2) \mid |\theta_j - \bar{\theta}| \pmod{\pi} \leq \pi/8\}. \quad (4.25)$$

Since the integrand is invariant under uniform translation of all the θ_j 's mod π as well as V_0 and V_3 are mapped into itself, we can fix $\bar{\theta} = 0$ and multiply it by the ratio of its range π to the length $n^{-1/2}$ of the vector $\frac{1}{n}[1, 1, \dots, 1]^T$. Thus we get that

$$S_3 \leq \pi n^{1/2} \int_{L \cap U_n(\pi/8) \setminus V_0} \prod_{(v_j, v_k) \in EG} |\cos \Delta_{jk}| dL, \quad (4.26)$$

where L denotes the orthogonal complement to the vector $[1, 1, \dots, 1]^T$. In a similar way as (4.18) we find that

$$S_3 \leq \pi n^{1/2} \int_{L \cap U_n(\pi/8) \setminus V_0} \exp\left(-\frac{1}{2} \sum_{(v_j, v_k) \in EG} \Delta_{jk}^2\right) dL. \quad (4.27)$$

Lemma 4.3. *Let the assumptions of Lemma 3.4 hold. Then as $n \rightarrow \infty$*

$$\int_{L \setminus U_n(n^{-1/2+\varepsilon})} \exp \left(-\frac{1}{2} \sum_{(v_j, v_k) \in EG} \Delta_{jk}^2 \right) dL = O \left(\exp(-cn^{2\varepsilon}) \frac{2^{\frac{n-1}{2}} \pi^{\frac{n-1}{2}} n^{1/2}}{\sqrt{\det \hat{Q}}} \right) \quad (4.28)$$

for some $c > 0$, depending only on σ .

Lemma 4.3 is also formulated and proved in [1] (see Lemma 6.2). Combining (3.28), (4.28) and (4.27), we get that as $n \rightarrow \infty$

$$S_3 = O \left(\exp(-cn^{2\varepsilon}) \frac{2^{\frac{n-1}{2}} \pi^{\frac{n+1}{2}} n}{\sqrt{\det \hat{Q}}} \right) = O \left(\exp(-cn^{2\varepsilon}) \frac{2^{\frac{n-1}{2}} \pi^{\frac{n+1}{2}}}{\sqrt{t(G)}} \right) \quad (4.29)$$

for some $c > 0$, depending only on σ . Combining (4.8), (4.16), (4.24) and (4.29), we obtain (2.8) as well as the following lemma.

Lemma 4.4. *Let the assumptions of Lemma 3.4 hold. Then as $n \rightarrow \infty$*

$$S = \left(1 + O \left(\exp(-cn^{2\varepsilon}) \right) \right) S_0 \quad (4.30)$$

for some $c > 0$, depending only on σ .

5. Final remarks

In fact, the estimation of the number of the Eulerian circuits is reduced (see the first part of the proof of Theorem 2.1 and Lemma 4.4) to estimating the integral

$$\int_{V_0} \exp \left(-\frac{1}{2} \sum_{(v_j, v_k) \in EG} \Delta_{jk}^2 - \frac{1}{12} \sum_{(v_j, v_k) \in EG} \Delta_{jk}^4 \right) d\theta. \quad (5.1)$$

Integral (5.1) can be evaluated more precisely for specific classes of graphs in order to get asymptotic formulas for $EO(G)$ similar to (1.1). For example, for even $n \rightarrow \infty$

$$EO(K_{n,n}) = e^{-1} \frac{2^{n^2+n-\frac{1}{2}}}{\pi^{n-\frac{1}{2}} n^{n-1}} \left(1 + O(n^{-1/2+\varepsilon}) \right) \quad (5.2)$$

for any $\varepsilon > 0$, where $K_{n,n}$ denotes the complete bipartite graph with n vertices in each part.

Acknowledgements

This work was carried out under the supervision of S.P. Tarasov and supported in part by RFBR grant no 11-01-00398a.

References

- [1] M. Isaev, *Asymptotic behaviour of the number of Eulerian circuits*, Electronic Journal of Combinatorics, 2011, V.18(1), P.219, e-print arXiv: 1104.3046.
- [2] M. Fiedler, *Algebraic connectivity of graphs*, Czech. Math. J. 23 (98) (1973), 298-305.
- [3] G. Kirchhoff, *Über die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Verteilung galvanischer Ströme geführt wird*, Ann. Phys. Chem. 72 (1847), 497-508. Translated by J. B. O'Toole in I.R.E. Trans. Circuit Theory, CT-5 (1958) 4.
- [4] M. Mihail and P. Winkler, *On the number of Eulerian orientations of a graph*, Algorithmica 16 (1996), 402–414.
- [5] B. D. McKay, R. W. Robinson, *Asymptotic enumeration of eulerian circuits in the complete graph*. Combinatorica, 7(4), December 1998.
- [6] B. D. McKay, *The asymptotic numbers of regular tournaments, eulirian digraphs and eulirian oriented graphs*. Combinatorica 10 (1990), no. 4, 367–377.
- [7] B. Mohar, *The Laplacian spectrum of graphs*, Graph Theory, Combinatorics, and Applications, Vol. 2, Ed. Y. Alavi, G. Chartrand, O. R. Oellermann, A. J. Schwenk, Wiley, 1991, pp. 871-898.