



HAL
open science

The Godunov scheme for scalar conservation laws with discontinuous bell-shaped flux functions

Boris Andreianov, Clément Cancès

► **To cite this version:**

Boris Andreianov, Clément Cancès. The Godunov scheme for scalar conservation laws with discontinuous bell-shaped flux functions. 2011. hal-00631586v1

HAL Id: hal-00631586

<https://hal.science/hal-00631586v1>

Preprint submitted on 12 Oct 2011 (v1), last revised 7 Mar 2012 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Godunov scheme for scalar conservation laws with discontinuous bell-shaped flux functions

Boris Andreianov*

Clément Cancès†

Abstract

We consider hyperbolic scalar conservation laws with discontinuous flux function of type $\partial_t u + \partial_x f(x, u) = 0$, with $f(x, u) = f_L(u)\mathbb{1}_{\mathbb{R}^-}(x) + f_R(u)\mathbb{1}_{\mathbb{R}^+}(x)$. Here $f_{L,R}$ are compatible bell-shaped flux functions as appear in numerous applications. In [2] and [7], it was shown that several notions of solution make sense, according to a choice of the so-called (A, B) -connection. In this note, we remark that every choice of connection (A, B) corresponds to a limitation of the flux under the form $f(u)|_{x=0} \leq \bar{F}$, first introduced in [13]. Hence we derive a very simple and cheap to compute explicit formula for the the Godunov numerical flux across the interface $\{x = 0\}$, for each choice of the connection. This gives a simple to use numerical scheme governed only by the parameter \bar{F} .

1 Introduction

Since it arises in several real life applications like traffic flow modeling [8], multiphase flows in porous media [3, 9, 10, 16] or water treatment [14], the Cauchy problem of the type

$$\partial_t u + \partial_x f(x, u) = 0, \quad u(\cdot, 0) = u_0, \quad (1)$$

where the flux function f is discontinuous w.r.t. the space variable have been widely studied during the last 20 years. A particular attention has been paid to the most simple case, i.e.

$$f(x, u) = f_L(u)\mathbb{1}_{\mathbb{R}^-}(x) + f_R(u)\mathbb{1}_{\mathbb{R}^+}(x). \quad (2)$$

In the sequel, we assume that the flux functions $f_{L,R}$ are *compatible* and *bell-shaped*, i.e.,

(A1) the functions $f_{L,R}$ are Lipschitz continuous and such that $f_L(0) = f_R(0)$, $f_L(1) = f_R(1)$;

(A2) there exists $b_{L,R} \in [0, 1]$ such that $f'_{L,R}(u)(u - b_{L,R}) < 0$ for a.e. $u \in [0, 1]$.

We also require the following condition on the initial data u_0 :

(A3) u_0 is a measurable function satisfying $0 \leq u_0(x) \leq 1$ for a.e. $x \in \mathbb{R}$.

For such a problem, it is natural to consider entropy solutions in the sense of Kružkov [17] away from the flux discontinuity at $x = 0$, i.e. functions $u \in L^\infty(\mathbb{R} \times \mathbb{R}_+; [0, 1])$ such that (3) holds with $q_{L,R}(u, \kappa) := \text{sign}(u - \kappa)(f_{L,R}(u) - f_{L,R}(\kappa))$. It has been pointed out in [2] that prescribing the balance of the fluxes at the interface is not sufficient to ensure uniqueness for the solution of the problem (1). Namely, some entropy criterion has to be fulfilled by the solution at the interface, and different physical contexts lead to different interface coupling criteria and thus to different notions of solution. In Section 2, we give a short introduction to the problem by following the theory introduced in [7] and extensively developed in [5]. We re-interpret the “ (A, B) -connections” of [2, 7] in terms of interface flux constraints “ $f(u)|_{x=0} \leq \bar{F}_{(A,B)}$ ” introduced in [13]. Due to this idea of flux limitation at

*Laboratoire de Mathématiques de Besançon, Université de Franche-Comté, 16 route de Gray, 25030 Besançon, France, boris.andreianov@univ-fcomte.fr

†Laboratoire Jacques-Louis Lions, UPMC Paris 06, 4 place Jussieu, 75005 Paris, France, cances@ann.jussieu.fr

the interface, in Section 3, we establish an explicit formula for the flux at the interface corresponding to any Riemann problem. This yields the flux for the Godunov scheme for approximation of solutions to problem (1) for any choice of interface coupling (i.e., for any choice of a connection (A, B) or of an interface flux constraint $\bar{F} = \bar{F}_{(A,B)}$).

2 Connections, flux limitation and L^1 dissipative germs

Definition 2.1 (Connections and L^1 dissipative germs; see [2, 7] and [5]) *For $f_{L,R}$ satisfying **(A1)**, **(A2)**, a couple $(A, B) \in [0, 1]^2$ is said to be a connection if $A \in [b_L, 1]$, $B \in [0, b_R]$ and $f_L(A) = f_R(B)$. We define the corresponding L^1 dissipative germ $\mathcal{G}_{(A,B)}$ (cf. [5]) to be the singleton $\{(A, B)\}$, and we set*

$$\mathcal{G}_{(A,B)}^* = \left\{ (c_L, c_R) \in [0, 1]^2 \text{ s.t. } f_L(c_L) = f_R(c_R) \text{ and } q_R(c_R, B) - q_L(c_L, A) \leq 0 \right\}.$$

We denote by $\mathcal{U} \subset [0, 1]^2$ the set of all the connections corresponding to the flux functions f_L, f_R . Finally, we define the optimal connection $(A^{\text{opt}}, B^{\text{opt}})$ by

$$(A^{\text{opt}}, B^{\text{opt}}) \in \mathcal{U}, \text{ with either } A^{\text{opt}} = b_L \text{ or } B^{\text{opt}} = b_R.$$

As it was shown in [18], under Assumption **(A2)**, a function $u \in L^\infty(\mathbb{R}^* \times \mathbb{R}_+; [0, 1])$ satisfying (3) admits one-sided traces $\gamma_{L,R}(u) \in L^\infty(\mathbb{R}_+)$ achieved in a strong sense. This permits to give the next definition.

Definition 2.2 ($\mathcal{G}_{(A,B)}$ -entropy solution) *A function $u \in L^\infty(\mathbb{R}^* \times \mathbb{R}_+; [0, 1])$ is said to be a $\mathcal{G}_{(A,B)}$ -entropy solution of (1),(2) if it satisfies*

$$\forall \kappa \in [0, 1] \quad \partial_t |u - \kappa| - \partial_x q_{L,R}(u, \kappa) \leq 0 \quad \text{in } \mathcal{D}'(\Omega_{L,R}), \quad (3)$$

and for a.e. $t > 0$, one has $(\gamma_L(u)(t), \gamma_R(u)(t)) \in \mathcal{G}_{(A,B)}^*$.

The theory developed in [5] shows that for all $(A, B) \in \mathcal{U}$, there exists a unique $\mathcal{G}_{(A,B)}$ -entropy solution to problem (1) in the sense of Definition 2.2. Equivalent characterizations of the $\mathcal{G}_{(A,B)}$ -entropy solutions in terms of up-to-the-interface entropy inequalities were used in [5, 7]. In this paper, we will rather benefit from the point of view developed in [13] and then in [4, 12]; to this end, we establish the link between connections and flux limitation at the interface. We need more notations (see Fig. 1). For $(A, B) \in \mathcal{U}$,

$$\text{set } \bar{F}_{(A,B)} := f_L(A) = f_R(B); \text{ notice that } \bar{F}_{(A^{\text{opt}}, B^{\text{opt}})} = \bar{F}^{\text{opt}} := \max_{(A,B) \in \mathcal{U}} \bar{F}_{(A,B)}.$$

The set \mathcal{U} of connections can be parametrized by \bar{F} (we write $\bar{F}_{(A,B)}$ or $(A_{\bar{F}}, B_{\bar{F}})$ to stress this link) which takes values in $[\bar{F}^{\text{barr}}, \bar{F}^{\text{opt}}] := [\max(f_{L,R}(0), f_{L,R}(1)), \min(f_L(b_L), f_R(b_R))]$.

Set $\mathcal{O} := \mathcal{G}_{(A^{\text{opt}}, B^{\text{opt}})}^*$. Then $\mathcal{O} \setminus \{(A^{\text{opt}}, B^{\text{opt}})\}$ is the set of all couples $(a, b) \in [0, 1]^2 \setminus \mathcal{U}$ such that $f_L(a) = f_R(b)$. In contrast to *under-compressive states* $(A, B) \in \mathcal{U}$, every couple $(a, b) \in \mathcal{O}$ will be called an *over-compressive state* (note that $(A^{\text{opt}}, B^{\text{opt}}) \in \mathcal{U} \cap \mathcal{O}$ is both under- and over-compressive). We have

$$\mathcal{G}_{(A,B)}^* = \{(A, B)\} \cup \mathcal{O}_{\bar{F}_{(A,B)}}, \text{ where } \mathcal{O}_{\bar{F}_{(A,B)}} := \{(c_L, c_R) \in \mathcal{O} \text{ s.t. } f_L(c_L) = f_R(c_R) \leq \bar{F}_{(A,B)}\} \quad (4)$$

is a restriction of \mathcal{O} . The connection (A, B) is the only under-compressive state belonging to $\mathcal{G}_{(A,B)}^*$. From (4), we readily see that $\mathcal{O}_{\bar{F}}$ depends in a monotone way on $\bar{F} \in [\bar{F}^{\text{barr}}, \bar{F}^{\text{opt}}]$.

In [13] (see also [4, 12]), L^1 -contractive semigroups of solutions were constructed even for the classical case $f_L = f_R$, by imposing an interface flux constraint of the form $f_{L,R}(\gamma_{L,R}(u)) \leq \bar{F}$ at $\{x = 0\}$. In the case $f_L \neq f_R$, the situation is exactly similar. Namely, each connection (A, B) makes appear a set of trace couples $\mathcal{G}_{(A,B)}^*$ satisfying (4), so that the different $\mathcal{G}_{(A,B)}$ -entropy solutions for (1),(2) for different $(A, B) \in \mathcal{U}$ correspond to different levels $\bar{F}_{(A,B)}$ of interface flux constraint. Kruřkov solutions (in the case $f_L = f_R$) and optimal entropy solutions (in the general case) shall be seen as the unconstrained ones.




Figure 1: On the left hand side, the two flux functions $f_{L,R}$ have been plotted together. A choice $(A, B) \in \mathcal{U}$ of connection is drawn, as well as the particular values A^* and B^* such that $f_L(A^*) = f_R(B^*) = \bar{F}_{(A,B)}$. As it plays a particularly important role, the connection $(A^{\text{opt}}, B^{\text{opt}})$ is also represented. On the right hand side, we have drawn the sets \mathcal{U} (red dashed line), $\mathcal{O} \equiv \mathcal{O}_{(A^{\text{opt}}, B^{\text{opt}})}$ (solid line) and its subset $\mathcal{O}_{(A,B)}$ (green solid line, outside of the grey rectangle). The grey rectangle represents the open set of $(u_L, u_R) \in [0, 1]^2$ that fail to satisfy $[f_L(u_L) \leq \bar{F}_{(A,B)}] \& [f_R(u_R) \leq \bar{F}_{(A,B)}]$.

3 The Godunov scheme

Consider the Riemann problem (1),(2) with initial datum $u_0 = u_L \mathbb{1}_{\mathbb{R}^-} + u_R \mathbb{1}_{\mathbb{R}^+}$. Let us compute the flux across the interface $\{x = 0\}$ of the $\mathcal{G}_{(A,B)}$ -entropy solution u of the Riemann problem in order to be able to build the Godunov scheme (see [15]). Note that such scheme is proved to be convergent in [5].

In the case $f_L = f_R$, the numerical scheme proposed in [4] used the flux $\min\{\bar{F}, \mathcal{F}(u_L, u_R)\}$, i.e., a given interface numerical flux $\mathcal{F}(\cdot, \cdot)$ for the unconstrained problem was limited to a given maximal value \bar{F} . Moreover, in the particular case where $\mathcal{F}(\cdot, \cdot)$ is the Godunov flux for the unconstrained problem, it is shown in [12] that the resulting scheme for the constrained problem is also the Godunov one. Here, we show that the same property holds for general $f_{L,R}$, namely, the Godunov flux through the interface $\{x = 0\}$ corresponding to the $\mathcal{G}_{(A,B)}$ -entropy solution is the Godunov flux corresponding to the optimal entropy solution on which we apply a constraint afterwards. Notice that, in addition, an explicit formula for the Godunov flux for the optimal entropy solution is well known since [1].

Theorem 3.1 (Main result) *The Godunov flux for $\mathcal{G}_{(A,B)}$ -entropy solutions at the interface $x = 0$ is given by*

$$\mathcal{F}(u_L, u_R) = \min(\bar{F}_{(A,B)}, f_L(\min(u_L, b_L)), f_R(\max(u_R, b_R))). \quad (5)$$

Moreover, whenever $\mathcal{F}^{\text{opt}}(u_L, u_R) > \bar{F}_{(A,B)}$, i.e., the constraint is active, one has

$$\gamma_L(u) = A, \quad \gamma_R(u) = B.$$

Proof: As it has been explicitly stated in [1], it follows from the *bell-shaped* behavior of the flux functions (see Assumption **(A2)**) that the flux of the $(A^{\text{opt}}, B^{\text{opt}})$ -entropy solution of the above Riemann problem across the discontinuity $\{x = 0\}$ is given by

$$\mathcal{F}^{\text{opt}}(u_L, u_R) = \min(f_L(\min(u_L, b_L)), f_R(\max(u_R, b_R))). \quad (6)$$

We have two possibilities. First, assume that $\mathcal{F}^{\text{opt}}(u_L, u_R) \leq \bar{F}_{(A,B)}$; we see from Fig. 1 that the traces couple $(\gamma_L(u^{\text{opt}}), \gamma_R(u^{\text{opt}}))$ belongs to $\mathcal{O} \cap \mathcal{G}_{(A,B)}^* = \mathcal{O}_{\bar{F}_{(A,B)}}$. Therefore, in this case the $\mathcal{G}_{(A^{\text{opt}}, B^{\text{opt}})}$ -entropy solution of the Riemann problem coincides with the $\mathcal{G}_{(A,B)}$ -entropy solution. Therefore, in the case under consideration the flux across the interface, which is given by formula (6), is also given by formula (5).

Second, assume that $\mathcal{F}^{\text{opt}}(u_L, u_R) > \bar{F}_{(A,B)}$, so that $(A, B) \neq (A^{\text{opt}}, B^{\text{opt}})$. In this case, one has

$$f_L(A) < f_L(\gamma_L(u^{\text{opt}})) = \min(f_L(\min(u_L, b_L)), f_L(\max(b_L, \gamma_L(u^{\text{opt}}))). \quad (7)$$

Denoting by $A^* \in [0, b_L]$ and $B^* \in [b_R, 0]$ the values such $f_L(A^*) = f_L(A) = \bar{F}_{(A,B)} = f_R(B^*) = f_R(B)$ (see Fig. 1), one deduces from (7) that $u_L > A^*$ and $u_R < B^*$. Therefore, using **(A2)**, one obtains that

$$f_L(A) = \bar{F}_{(A,B)} = \min(f_L(\min(u_L, b_L)), f_L(\max(b_L, A))).$$

Similarly, one obtains that $f_R(B) = \bar{F}_{(A,B)} = \min(f_R(\min(B, b_R)), f_R(\max(b_R, u_R)))$. These two relations imply that the boundary $\{x = 0\}$ is characteristic for each of the Cauchy-Dirichlet problems

$$\begin{cases} \partial_t u + \partial_x f_L(u) = 0 \text{ in } \mathcal{D}'(\mathbb{R}_-^* \times \mathbb{R}_+^*), \\ u(x, 0) = u_L \text{ for } x < 0, \\ u(0, t) = A \text{ for } t > 0; \end{cases} \quad \begin{cases} \partial_t u + \partial_x f_R(u) = 0 \text{ in } \mathcal{D}'(\mathbb{R}_-^* \times \mathbb{R}_+^*), \\ u(x, 0) = u_R \text{ for } x > 0, \\ u(0, t) = B \text{ for } t > 0. \end{cases} \quad (8)$$

This ensures that the boundary conditions prescribed in (8) are fulfilled in a strong sense by the function u (see [6]). Defining u as the juxtaposition of the entropy solutions of problems (8) in the sense of [6], we see that u satisfies (3) and it takes the initial datum $u_L \mathbb{1}_{\mathbb{R}^-} + u_R \mathbb{1}_{\mathbb{R}^+}$. Moreover, we have $(\gamma_L(u)(t), \gamma_R(u)(t)) = (A, B) \in \mathcal{G}_{(A,B)}^*$ for all $t > 0$, ensuring that u is the unique $\mathcal{G}_{(A,B)}$ -entropy solution (see [7, 5]) to the Riemann problem under study. Thus the Godunov flux for this Riemann problem is the flux of u across the interface. The latter is given by $\mathcal{F}(u_L, u_R) = \bar{F}_{(A,B)}$, so that formula (5) is true also in this case. \square

As a conclusion, remark that the numerical fluxes of the Godunov scheme given by formula (5) are cheap to compute. In particular, no integration is needed to compute the solution of the Godunov scheme, in contrast, e.g., to the Engquist-Osher-type scheme proposed in [7].

Moreover, the scheme based on (5) readily adapts to any level \bar{F} of interface flux constraint. We refer to [3] (see also [9, 10]) for an example of determination of the level of constraint in the setting of Buckley-Leverett equations for two-phase flow in a two-rocks' medium. Indeed, in this model different values $\bar{F} \in [\bar{F}^{\text{barr}}, \bar{F}^{\text{opt}}]$ can appear as physically motivated ones, depending to the capillary pressure profiles.

References

- [1] Adimurthi, J. Jaffré, and G. D. Veerappa Gowda. Godunov-type methods for conservation laws with a flux function discontinuous in space. *SIAM J. Numer. Anal.*, 42(1):179–208, 2004.
- [2] Adimurthi, S. Mishra, and G. D. Veerappa Gowda. Optimal entropy solutions for conservation laws with discontinuous flux-functions. *J. Hyperbolic Differ. Equ.*, 2(4):783–837, 2005.
- [3] B. Andreianov and C. Cancès. Vanishing capillarity solutions of Buckley-Leverett equation with gravity in two-rocks' medium. In preparation.
- [4] B. Andreianov, P. Goatin, and N. Seguin. Finite volume schemes for locally constrained conservation laws. *Numer. Math.*, 115(4):609–645, 2010. With supplementary material available online.
- [5] B. Andreianov, K. Karlsen, and N. Risebro. A theory of L^1 -dissipative solvers for scalar conservation laws with discontinuous flux. *Arch. Ration. Mech. Anal.* 201(1):27–86, 2011.
- [6] C. Bardos, A. Y. le Roux, and J.-C. Nédélec. First order quasilinear equations with boundary conditions. *Comm. Partial Differential Equations*, 4(9):1017–1034, 1979.
- [7] R. Bürger, K. H. Karlsen, and J. D. Towers. An Engquist-Osher-type scheme for conservation laws with discontinuous flux adapted to flux connections. *SIAM J. Numer. Anal.*, 47(3):1684–1712, 2009.
- [8] R. Bürger, A. García, K. H. Karlsen, and J. D. Towers. Difference schemes, entropy solutions, and speedup impulse for an inhomogeneous kinematic traffic flow model. *Netw. Heterog. Media*, 3(1):1–41, 2008.

- [9] C. Cancès. Asymptotic behavior of two-phase flows in heterogeneous porous media for capillarity depending only on space. I. Convergence to the optimal entropy solution. *SIAM J. Math. Anal.*, 42(2):946–971, 2010.
- [10] C. Cancès. Asymptotic behavior of two-phase flows in heterogeneous porous media for capillarity depending only on space. II. Nonclassical shocks to model oil-trapping. *SIAM J. Math. Anal.*, 42(2):972–995, 2010.
- [11] C. Cancès and T. Gallouët. On the time continuity of entropy solutions. *J. Evol. Equ.*, 11(1):43–55, 2011.
- [12] C. Cancès and N. Seguin. Error estimate for godunov approximation of locally constrained conservation laws. *HAL : hal-00599581*, 2011 (submitted).
- [13] R. M. Colombo and P. Goatin. A well posed conservation law with a variable unilateral constraint. *J. Differential Equations*, 234(2):654–675, 2007.
- [14] S. Diehl. A conservation law with point source and discontinuous flux function modelling continuous sedimentation. *SIAM J. Appl. Math.*, 56(2):388–419, 1996.
- [15] S. K. Godunov. A difference method for numerical calculation of discontinuous solutions of the equations of hydrodynamics. *Mat. Sb. (N.S.)*, 47 (89):271–306, 1959.
- [16] E. F. Kaasschieter. Solving the Buckley-Leverett equation with gravity in a heterogeneous porous medium. *Comput. Geosci.*, 3(1):23–48, 1999.
- [17] S. N. Kružkov. First order quasilinear equations with several independent variables. *Mat. Sb.* 81 (123):228–255, 1970.
- [18] E. Yu. Panov. Existence of strong traces for quasi-solutions of multidimensional conservation laws. *J. Hyperbolic Differ. Equ.*, 4(4):729–770, 2007.