



HAL
open science

Efficient estimation of conditional covariance matrices for dimension reduction

Jean-Michel Loubes, Clément Marteau, Michael Solis, Sébastien da Veiga

► **To cite this version:**

Jean-Michel Loubes, Clément Marteau, Michael Solis, Sébastien da Veiga. Efficient estimation of conditional covariance matrices for dimension reduction. 2011. hal-00632576

HAL Id: hal-00632576

<https://hal.science/hal-00632576>

Preprint submitted on 14 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Efficient estimation of conditional covariance matrices for dimension reduction

JEAN-MICHEL LOUBES

Institut de Mathématiques de Toulouse

loubes@math.univ-toulouse.fr

CLÉMENT MARTEAU

Institut de Mathématiques de Toulouse

clement.marteau@insa-toulouse.fr

MICHAEL SOLÍS CHACÓN

Institut de Mathématiques de Toulouse

msolisch@math.univ-toulouse.fr

SEBASTIEN DA VEIGA

Institut Français du Pétrole

sebastien.da-veiga@ifp.fr

Abstract

We consider the problem of estimating a conditional covariance matrix in an inverse regression setting. We show that this estimation can be achieved by estimating a quadratic functional extending the results of Da Veiga & Gamboa (2008). We prove that this method provides a new efficient estimator whose asymptotic properties are studied.

1 Introduction

Consider the nonparametric regression

$$Y = \varphi(X) + \epsilon,$$

where $X \in \mathbb{R}^p$, $Y \in \mathbb{R}$ and $\mathbb{E}[\epsilon] = 0$. The main difficulty with any regression method is that, as the dimension of X becomes larger, the number of observations needed for a good estimator increases exponentially. This phenomena is usually called the *curse of dimensionality*. All the “classical” methods could break down, as the dimension p increases, unless we have at hand a very huge sample.

For this reason, there have been along the past decades a very large number of methods to cope with this issue. Their aim is to reduce the dimensionality of the problem, using just to name a few, the generalized linear model in Brillinger (1983), the additive models in Hastie & Tibshirani (1990), sparsity constraint models as Li (2007) and references therein.

Alternatively, Li (1991a) proposed the procedure of Sliced Inverse Regression (SIR) considering the following semiparametric model,

$$Y = \phi(v_1^\top X, \dots, v_K^\top X, \epsilon)$$

where the v 's are unknown vectors in \mathbb{R}^p , ϵ is independent of X and ϕ is an arbitrary function in \mathbb{R}^{K+1} . This model can gather all the relevant information about the variable Y , with only the projection of X onto the $K \ll p$ dimensional subspace $(v_1^\top X, \dots, v_K^\top X)$. In the case when K is small, it is possible to reduce the dimension by estimating the v 's efficiently. This method is also used to search nonlinear structures in data and to estimate the projection directions v 's. For a review on SIR methods, we refer to Li (1991a,b); Duan & Li (1991); Hardle & Tsybakov (1991) and references therein. The v 's define the effective dimension reduction (e.d.r) direction and the eigenvectors of $\mathbb{E}[\text{Cov}(X|Y)]$ are the e.d.r directions. Many estimators have been proposed in order to study the e.d.r directions in many different cases. For example, Zhu & Fang (1996) and Ferré & Yao (2005, 2003) use kernel estimators, Hsing (1999) combines nearest neighbor and SIR, Bura & Cook (2001) assume that $\mathbb{E}[X|Y]$ has some parametric form, Setodji & Cook (2004) use k-means and Cook & Ni (2005) transform SIR to least square form.

In this paper, we propose an alternate estimation of the matrix

$$\text{Cov}(\mathbb{E}[X|Y]) = \mathbb{E}[\mathbb{E}[X|Y]\mathbb{E}[X|Y]^\top] - \mathbb{E}[X]\mathbb{E}[X]^\top,$$

using ideas developed by Da Veiga & Gamboa (2008), inspired by the prior work of Laurent (1996). More precisely since $\mathbb{E}[X]\mathbb{E}[X]^\top$ can be easily estimated with many usual methods, we will focus on finding an estimator of $\mathbb{E}[\mathbb{E}[X|Y]\mathbb{E}[X|Y]^\top]$. For this we will show that this estimation implies an estimation of a quadratic functional rather than plugging non parametric estimate into this form as commonly used. This method has the advantage of getting an efficient estimator in a semi-parametric framework.

This paper is organized as follows. Section 2 is intended to motivate our investigation of $\text{Cov}(\mathbb{E}[X|Y])$ using a Taylor approximation. In Section 3.1 we set up notation and hypothesis. Section 3.2 is devoted to demonstrate that each coordinate of $\text{Cov}(\mathbb{E}[X|Y])$ converge efficiently. Also we find the normality asymptotic for the whole matrix. An asymptotic bound of the variance for the quadratic part for the Taylor's expansion of $\text{Cov}(\mathbb{E}[X|Y])$ is found in Section 4. All technical Lemmas and their proofs are postponed to Sections 6 and 5 respectively.

2 Methodology

Our aim is to estimate $\text{Cov}(\mathbb{E}[X|Y])$ efficiently when observing $X \in \mathbb{R}^p$, for $p \geq 1$, and $Y \in \mathbb{R}$. For this, write the matrix

$$\text{Cov}(\mathbb{E}[X|Y]) = \mathbb{E}[\mathbb{E}[X|Y]\mathbb{E}[X|Y]^\top] - \mathbb{E}[X]\mathbb{E}[X]^\top,$$

where A^\top means the transpose of A . If $\mathbb{E}[X]$ can be easily estimated by classical methods, the remainder term

$$\mathbb{E}[\mathbb{E}[X|Y]\mathbb{E}[X|Y]^\top] = (T_{ij}^*)_{i,j} \quad i, j = 1, \dots, p;$$

is a non linear term whose estimation is the main topic of this paper. Each term of this matrix can be written as

$$T_{ij}^* = \int \left(\frac{\int x_i f(x_i, x_j, y) dx_i dx_j}{\int f(x_i, x_j, y) dx_i dx_j} \right) \left(\frac{\int x_j f(x_i, x_j, y) dx_i dx_j}{\int f(x_i, x_j, y) dx_i dx_j} \right) f(x_i, x_j, y) dx_i dx_j dy, \quad (1)$$

where $f(x_i, x_j, y)$ for i and j fixed, is the joint density of (X_i, X_j, Y) $i, j = 1, \dots, p$.

Hence, we focus on the efficient estimation of the corresponding non linear functional for $f \in \mathbb{L}(dx_i, dx_j, dy)$

$$f \mapsto T_{ij}(f) = \int \left(\frac{\int x_i f(x_i, x_j, y) dx_i dx_j}{\int f(x_i, x_j, y) dx_i dx_j} \right) \left(\frac{\int x_j f(x_i, x_j, y) dx_i dx_j}{\int f(x_i, x_j, y) dx_i dx_j} \right) f(x_i, x_j, y) dx_i dx_j dy. \quad (2)$$

In the case $i = j$, this estimation has been considered in Da Veiga & Gamboa (2008); Laurent (1996). Here we extend their methodology to this case. Assume we have at hand an i.i.d sample $(X_i^{(k)}, X_j^{(k)}, Y^{(k)})$, $k = 1, \dots, n$ such that it is possible to build a preliminary estimator \hat{f} of f with a subsample of size $n_1 < n$. Now, the main idea is to make a Taylor's expansion of $T_{ij}(f)$ in a neighborhood of \hat{f} which will play the role of a suitable approximation of f . More precisely, define an auxiliary function $F : [0, 1] \rightarrow \mathbb{R}$;

$$F(u) = T_{ij}(uf + (1-u)\hat{f})$$

with $u \in [0, 1]$. The Taylor's expansion of F between 0 and 1 up to the third order is

$$F(1) = F(0) + F'(0) + \frac{1}{2}F''(0) + \frac{1}{6}F'''(\xi)(1-\xi)^3 \quad (3)$$

for some $\xi \in [0, 1]$. Moreover, we have

$$F(1) = T_{ij}(f)$$

$$F(0) = T_{ij}(\hat{f}) = \int \left(\frac{\int x_i \hat{f}(x_i, x_j, y) dx_i dx_j}{\int \hat{f}(x_i, x_j, y) dx_i dx_j} \right) \left(\frac{\int x_j \hat{f}(x_i, x_j, y) dx_i dx_j}{\int \hat{f}(x_i, x_j, y) dx_i dx_j} \right) \hat{f}(x_i, x_j, y) dx_i dx_j dy.$$

To simplify the notations, let

$$m_i(f_u, y) = \frac{\int x_i f_u(x_i, x_j, y) dx_i dx_j}{\int f_u(x_i, x_j, y) dx_i dx_j}$$

$$m_i(f_0, y) = m_i(\hat{f}, y) = \frac{\int x_i \hat{f}(x_i, x_j, y) dx_i dx_j}{\int \hat{f}(x_i, x_j, y) dx_i dx_j},$$

where $f_u = uf + (1 - u)\hat{f}$, $\forall u \in [0, 1]$. Then, we can rewrite $F(u)$ as

$$F(u) = \int m_i(f_u, y)m_j(f_u, y)f_u(x_i, x_j, y)dx_idx_jdy.$$

The Taylor's expansion of $T_{ij}(f)$ is given in the next Proposition.

Proposition 1 (Linearization of the operator T). *For the functional $T_{ij}(f)$ defined in (2), the following decomposition holds*

$$T_{ij}(f) = \int H_1(\hat{f}, x_i, x_j, y)f(x_i, x_j, y)dx_idx_jdy + \int H_2(\hat{f}, x_{i1}, x_{j2}, y)f(x_{i1}, x_{j1}, y)f(x_{i2}, x_{j2}, y)dx_{i1}dx_{j1}dx_{i2}dx_{j2}dy + \Gamma_n \quad (4)$$

where

$$H_1(\hat{f}, x_i, x_j, y) = x_im_j(\hat{f}, y) + x_jm_i(\hat{f}, y) - m_i(\hat{f}, y)m_j(\hat{f}, y) \quad (5)$$

$$H_2(\hat{f}, x_{i1}, x_{j2}, y) = \frac{1}{\int \hat{f}(x_i, x_j, y)dx_idx_j} (x_{i1} - m_i(\hat{f}, y))(x_{j2} - m_j(\hat{f}, y)) \quad (6)$$

$$\Gamma_n = \frac{1}{6}F'''(\xi)(1 - \xi)^3, \quad (7)$$

for some $\xi \in]0, 1[$.

This decomposition has the main advantage of separating the terms to be estimated into a linear functional of f , which can be easily estimated and a second part which is a quadratic functional of f . In this case, Section 4 will be dedicated to estimate this kind of functionals and specifically to control its variance. This will enable to provide an efficient estimator of $T_{ij}(f)$ using the decomposition of Proposition 1.

3 Main Results

In this section we build a procedure to estimate $T_{ij}(f)$ efficiently. Since we used $n_1 < n$ to build a preliminary approximation \hat{f} , we will use a sample of size $n_2 = n - n_1$ to estimate (5) and (6). Since (5) is a linear functional of the density f , it can be estimated by its empirical counterpart

$$\frac{1}{n_2} \sum_{k=1}^{n_2} H_1(\hat{f}, X_i^{(k)}, X_j^{(k)}, Y^{(k)}). \quad (8)$$

Since (6) is a nonlinear functional of f , the estimation is harder. Its estimation will be a direct consequence of the technical results presented in Section 4, where we build an estimator for the general functional

$$\theta(f) = \int \eta(x_{i1}, x_{j2}, y)f(x_{i1}, x_{j1}, y)f(x_{i2}, x_{j2}, y)dx_{i1}dx_{j1}dx_{i2}dx_{j2}dy$$

where $\eta : \mathbb{R}^3 \rightarrow \mathbb{R}$ is a bounded function. The estimator $\hat{\theta}_n$ of $\theta(f)$ is an extension of the method developed in Da Veiga & Gamboa (2008).

3.1 Hypothesis and Assumptions

The following notations will be used throughout the paper. Let d_s and b_s for $s = 1, 2, 3$ be real numbers where $d_s < b_s$. Let, for i and j fixed, $\mathbb{L}^2(dx_i dx_j dy)$ be the squared integrable functions in the cube $[d_1, b_1] \times [d_2, b_2] \times [d_3, b_3]$. Moreover, let $(p_l(x_i, x_j, y))_{l \in D}$ be an orthonormal basis of $\mathbb{L}^2(dx_i dx_j dy)$, where D is a countable set. Let $a_l = \int p_l f$ denote the scalar product of f with p_l .

Furthermore, denote by $\mathbb{L}^2(dx_i dx_j)$ (resp. $\mathbb{L}^2(dy)$) the set of squared integrable functions in $[d_1, b_1] \times [d_2, b_2]$ (resp. $[d_3, b_3]$). If $(\alpha_{l_\alpha}(x_i, x_j))_{l_\alpha \in D_1}$ (resp. $(\beta_{l_\beta}(y))_{l_\beta \in D_2}$) is an orthonormal basis of $\mathbb{L}^2(dx_i dx_j)$ (resp. $\mathbb{L}^2(dy)$) then $p_l(x_i, x_j, y) = \alpha_{l_\alpha}(x_i, x_j) \beta_{l_\beta}(y)$ with $l = (l_\alpha, l_\beta) \in D_1 \times D_2$.

We also use the following subset of $\mathbb{L}^2(dx_i dx_j dy)$

$$\mathcal{E} = \left\{ \sum_{l \in D} e_l p_l : (e_l)_{l \in D} \text{ is such that } \sum_{l \in D} \left| \frac{e_l}{c_l} \right|^2 < 1 \right\}$$

where $(c_l)_{l \in D}$ is a given fixed sequence.

Moreover assume that (X_i, X_j, Y) have a bounded joint density f on $[d_1, b_1] \times [d_2, b_2] \times [d_3, b_3]$ which lies in the ellipsoid \mathcal{E} .

In what follows, $X_n \xrightarrow{\mathcal{D}} X$ (resp. $X_n \xrightarrow{\mathcal{P}} X$) denotes the convergence in distribution or weak convergence (resp. convergence in probability) of X_n to X . Additionally, the support of f will be denoted by $\text{supp } f$.

Let $(M_n)_{n \geq 1}$ denote a sequence of subsets D . For each n there exists M_n such that $M_n \subset D$. Let us denote by $|M_n|$ the cardinal of M_n .

We shall make three main assumptions:

Assumption 1. For all $n \geq 1$ there is a subset $M_n \subset D$ such that $(\sup_{l \notin M_n} |c_l|^2)^2 \approx |M_n|/n^2$ ($A_n \approx B$ means $\lambda_1 \leq A_n/B \leq \lambda_2$ for some positives constants λ_1 and λ_2). Moreover, $\forall f \in \mathbb{L}^2(dx dy dz)$, $\int (S_{M_n} f - f)^2 dx dy dz \rightarrow 0$ when $n \rightarrow 0$, where $S_{M_n} f = \sum_{l \in M_n} a_l p_l$

Assumption 2. $\text{supp } f \subset [d_1, b_1] \times [d_2, b_2] \times [d_3, b_3]$ and $\forall (x, y, z) \in \text{supp } f$, $0 < \alpha \leq f(x, y, z) \leq \beta$ with $\alpha, \beta \in \mathbb{R}$.

Assumption 3. It is possible to find an estimator \hat{f} of f built with $n_1 \approx n/\log(n)$ observations, such that for $\epsilon > 0$,

$$\forall (x, y, z) \in \text{supp } f, 0 < \alpha - \epsilon \leq \hat{f}(x, y, z) \leq \beta + \epsilon$$

and,

$$\forall 2 \leq q \leq +\infty, \forall l \in \mathbb{N}^*, \mathbb{E}_f \| \hat{f} - f \|_q^l \leq C(q, l) n_1^{-l\lambda}$$

for some $\lambda > 1/6$ and some constant $C(q, l)$ not depending on f belonging to the ellipsoid \mathcal{E} .

Assumption 1 is necessary to bound the bias and variance of $\hat{\theta}_n$. Assumption 2 and 3 allow to establish that the remainder term in the Taylor expansion is negligible, i.e $\Gamma_n = O(1/n)$. Assumption 3 depends on the regularity of the density function. For instance for $x \in \mathbb{R}^p$, $s > 0$ and $L > 0$, consider the class

$\mathcal{H}_q(s, L)$ of Nikol'skii of functions $f \in \mathbb{L}^q(dx)$ with partials derivatives up to order $r = \lfloor s \rfloor$ inclusive, and for each of these derivatives $g^{(r)}$

$$\|f^{(r)}(\cdot + h) - f^{(r)}(\cdot)\|_q \leq L |h|^{s-r} \quad \forall h \in \mathbb{R}.$$

Then, Assumption 3 is satisfied for $f \in \mathcal{H}_q(s, L)$ with $s > \frac{p}{4}$.

3.2 Efficient Estimation of $T_{ij}(f)$

As seen in Section 2, $T_{ij}(f)$ can be decomposed as (4). Hence, using (8) and (14) we consider the following estimate

$$\begin{aligned} \widehat{T}_{ij}^{(n)} &= \frac{1}{n_2} \sum_{k=1}^{n_2} H_1(\hat{f}, X_i^{(k)}, X_j^{(k)}, Y^{(k)}) \\ &+ \frac{1}{n_2(n_2 - 1)} \sum_{l \in M} \sum_{k \neq k'=1}^{n_2} p_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) \int p_l(x_i, x_j, Y^{(k')}) H_3(\hat{f}, x_i, x_j, X_i^{(k')}, X_j^{(k')}, Y^{(k')}) dx_i dx_j \\ &- \frac{1}{n_2(n_2 - 1)} \sum_{l, l' \in M} \sum_{k \neq k'=1}^{n_2} p_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) p_{l'}(X_i^{(k')}, X_j^{(k')}, Y^{(k')}) \\ &\quad \int p_l(x_{i1}, x_{j1}, y) p_{l'}(x_{i2}, x_{j2}, y) H_2(\hat{f}, x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy. \end{aligned}$$

where $H_3(f, x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) = H_2(f, x_{i1}, x_{j2}, y) + H_2(f, x_{i2}, x_{j1}, y)$ and $n_2 = n - n_1$. The remainder Γ_n does not appear because we will prove that it is negligible when compared to the other error terms.

The asymptotic behavior of $\widehat{T}_{ij}^{(n)}$ for i and j fixed is given in the next Theorem.

Theorem 1. *Let Assumptions 1-3 hold and $|M_n|/n \rightarrow 0$ when $n \rightarrow \infty$. Then:*

$$\sqrt{n}(\widehat{T}_{ij}^{(n)} - T_{ij}(f)) \xrightarrow{\mathcal{D}} \mathcal{N}(0, C_{ij}(f)), \quad (9)$$

and

$$\lim_{n \rightarrow \infty} n \mathbb{E}[\widehat{T}_{ij}^{(n)} - T_{ij}(f)]^2 = C_{ij}(f), \quad (10)$$

where

$$C_{ij}(f) = \text{Var}(H_1(f, X_i, X_j, Y))$$

Note that, in Theorem 1, it appears that the asymptotic variance of $T_{ij}(f)$ depends only on $H_1(f, X_i, X_j, Y)$. Hence the asymptotic variance of $\widehat{T}_{ij}^{(n)}$ is explained only by the linear part of (4). This will entail that the estimator is naturally efficient as proved in the following.

Indeed, the semi-parametric Cramér-Rao bound is given in the next theorem.

Theorem 2 (Semi-parametric Cramér-Rao bound.). *Consider the estimation of*

$$T_{ij}(f) = \int \left(\frac{\int x_i f(x_i, x_j, y) dx_i dx_j}{\int f(x_i, x_j, y) dx_i dx_j} \right) \left(\frac{\int x_j f(x_i, x_j, y) dx_i dx_j}{\int f(x_i, x_j, y) dx_i dx_j} \right) f(x_i, x_j, y) dx_i dx_j dy$$

for a random vector (X_i, X_j, Y) with joint density $f \in \mathcal{E}$. Let $f_0 \in \mathcal{E}$ be a density verifying the assumptions of Theorem 1. Then, for all estimator $\widehat{T}_{ij}^{(n)}$ of $T_{ij}(f)$ and every family $\{\mathcal{V}_r(f_0)\}_{r>0}$ of neighborhoods of f_0 we have

$$\inf_{\{\mathcal{V}_r(f_0)\}_{r>0}} \liminf_{n \rightarrow \infty} \sup_{f \in \mathcal{V}_r(f_0)} n\mathbb{E}[\widehat{T}_{ij}^{(n)} - T_{ij}(f_0)]^2 \geq C_{ij}(f_0)$$

where $\mathcal{V}_r(f_0) = \{f : \|f - f_0\|_2 < r\}$ for $r > 0$.

Consequently, the estimator $\widehat{T}_{ij}^{(n)}$ is efficient.

In the case of our estimate, its variance is $C_{ij}(f)$, which proves its asymptotically efficiency.

Remark that Theorem 1 proves asymptotic normality entry by entry of the matrix $\mathbf{T}(f) = (T_{ij}(f))_{p \times p}$. To extend the result for the whole matrix it is necessary to introduce the half-vectorization operator vech . This operator, stacks only the columns from the principal diagonal of a square matrix downwards in a column vector, that is, for an $p \times p$ matrix $\mathbf{A} = (a_{ij})$,

$$\text{vech}(\mathbf{A}) = [a_{11}, \dots, a_{p1}, a_{22}, \dots, a_{p2}, \dots, a_{33}, \dots, a_{pp}]^\top.$$

Let define the estimator matrix $\widehat{\mathbf{T}}^{(n)} = (\widehat{T}_{ij}^{(n)})$ and $\mathbf{H}_1(f)$ denote the matrix with entries $(H_1(f, x_i, x_j, y))_{i,j}$. Now we are able to state the following

Corollary 1. *Let Assumptions 1-3 hold and $|M_n|/n \rightarrow 0$ when $n \rightarrow \infty$. Then $\widehat{\mathbf{T}}^{(n)}$ has the following properties:*

$$\sqrt{n} \text{vech}(\widehat{\mathbf{T}}^{(n)} - \mathbf{T}(f)) \xrightarrow{\mathcal{D}} \mathcal{N}(0, \mathbf{C}(f)), \quad (11)$$

$$\lim_{n \rightarrow \infty} n\mathbb{E}[\text{vech}(\widehat{\mathbf{T}}^{(n)} - \mathbf{T}(f)) \text{vech}(\widehat{\mathbf{T}}^{(n)} - \mathbf{T}(f))^\top] = \mathbf{C}(f) \quad (12)$$

where

$$\mathbf{C}(f) = \text{Cov}(\text{vech}(\mathbf{H}_1(f)))$$

Previous results depend on the accurate estimation of the quadratic part of the estimator of $T_{ij}^{(n)}$, which is the issue of the following section.

4 Estimation of quadratic functionals

As pointed out in Section 2 the decomposition (4) has a quadratic part (6) that we want to estimate. To achieve this we will construct a general estimator of the form:

$$\theta = \int \eta(x_{i1}, x_{j2}, y) f(x_{i1}, x_{j1}, y) f(x_{i2}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy,$$

for $f \in \mathcal{E}$ and $\eta : \mathbb{R}^3 \rightarrow \mathbb{R}$ a bounded function.

Given M_n a subset of D , consider the estimator

$$\begin{aligned}\hat{\theta}_n &= \frac{1}{n(n-1)} \sum_{l \in M} \sum_{k \neq k'=1}^n p_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) \\ &\quad \int p_l(x_i, x_j, Y^{(k')}) \left(\eta(x_i, X_j^{(k')}, Y^{(k')}) + \eta(X_i^{(k')}, x_j, Y^{(k')}) \right) dx_i dx_j \\ &\quad - \frac{1}{n(n-1)} \sum_{l, l' \in M} \sum_{k \neq k'=1}^n p_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) p_{l'}(X_i^{(k')}, X_j^{(k')}, Y^{(k')}) \\ &\quad \int p_l(x_{i1}, x_{j1}, y) p_{l'}(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy. \quad (13)\end{aligned}$$

In order to simplify the presentation of the main Theorem, let $\psi(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) = \eta(x_{i1}, x_{j2}, y) + \eta(x_{i2}, x_{j1}, y)$ verifying

$$\int \psi(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy = \int \psi(x_{i2}, x_{j2}, x_{i1}, x_{j1}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy.$$

With this notation we can simplify (13) in

$$\begin{aligned}\hat{\theta}_n &= \frac{1}{n(n-1)} \sum_{l \in M} \sum_{k \neq k'=1}^n p_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) \int p_l(x_i, x_j, Y^{(k')}) \psi(x_i, x_j, X_i^{(k')}, X_j^{(k')}, Y^{(k')}) dx_i dx_j \\ &\quad - \frac{1}{n(n-1)} \sum_{l, l' \in M} \sum_{k \neq k'=1}^n p_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) p_{l'}(X_i^{(k')}, X_j^{(k')}, Y^{(k')}) \\ &\quad \int p_l(x_{i1}, x_{j1}, y) p_{l'}(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy. \quad (14)\end{aligned}$$

Using simple algebra, it is possible to prove that this estimator has bias equal to

$$\begin{aligned}- \int (S_M f(x_{i1}, x_{j1}, y) - f(x_{i1}, x_{j1}, y)) (S_M f(x_{i2}, x_{j2}, y) - f(x_{i2}, x_{j2}, y)) \\ \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \quad (15)\end{aligned}$$

The following Theorem gives an explicit bound for the variance of $\hat{\theta}_n$.

Theorem 3. *Let Assumption 1 hold. Then if $|M_n|/n \rightarrow 0$ when $n \rightarrow 0$, then $\hat{\theta}_n$ has the following property*

$$\left| n\mathbb{E}[(\hat{\theta}_n - \theta)^2] - \Lambda(f, \eta) \right| \leq \gamma \left[\frac{|M_n|}{n} + \|S_{M_n} f - f\|_2 + \|S_{M_n} g - g\|_2 \right], \quad (16)$$

where $g(x_i, x_j, y) = \int f(x_{i2}, x_{j2}, y) \psi(x_i, x_j, x_{i2}, x_{j2}, y) dx_{i2} dx_{j2}$ and

$$\Lambda(f, \eta) = \int g(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy - \left(\int g(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2,$$

where γ is constant depending only on $\|f\|_\infty, \|\eta\|_\infty$, and $\Delta_{x_i x_j} = (b_1 - a_1) \times (b_2 - a_2)$. Moreover, this constant is an increasing function of these quantities.

Note that equation (16) implies that

$$\lim_{n \rightarrow \infty} n\mathbb{E}[\hat{\theta}_n - \theta]^2 = \Lambda(f, \eta).$$

These results will be stated in order to control the term

$$Q = \int H_2(\hat{f}, x_{i1}, x_{j2}, y) f(x_{i1}, x_{j1}, y) f(x_{i2}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy$$

which has the form of the quadratic functional θ with the particular choice $\eta(x_{i1}, x_{j2}, y) = H_2(\hat{f}, x_{i1}, x_{j2}, y)$. We point out that we also show that in this particular frame, we get $\Lambda(f, \eta) = 0$. This the reason why the asymptotic variance of the estimate $\hat{T}_{ij}^{(n)}$ built in the previous section, is only governed by its linear part, yielding asymptotic efficiency.

5 Proofs

Proof of Proposition 1.

We need to calculate the three first derivatives of $F(u)$. In order to facilitate the calculation, we are going to differentiate $m_i(f_u, y)$:

$$\begin{aligned} \frac{d}{du} (m_i(f_u, y)) &= \frac{d}{du} \left(\frac{\int x_i f_u(x_i, x_j, y) dx_i dx_j}{\int f_u(x_i, x_j, y) dx_i dx_j} \right) \\ &= \frac{\int x_i (f(x_i, x_j, y) - \hat{f}(x_i, x_j, y)) dx_i dx_j}{\int f_u(x_i, x_j, y) dx_i dx_j} \\ &\quad - \frac{\int x_i f_u(x_i, x_j, y) dx_i dx_j \int (f(x_i, x_j, y) - \hat{f}(x_i, x_j, y)) dx_i dx_j}{\left(\int f_u(x_i, x_j, y) dx_i dx_j \right)^2}, \\ &= \frac{\int x_i (f(x_i, x_j, y) - \hat{f}(x_i, x_j, y)) dx_i dx_j}{\int f_u(x_i, x_j, y) dx_i dx_j} \\ &\quad - \frac{m_i(f_u, y) \int (f(x_i, x_j, y) - \hat{f}(x_i, x_j, y)) dx_i dx_j}{\int f_u(x_i, x_j, y) dx_i dx_j}, \\ &= \frac{\int (x_i - m_i(f_u, y)) (f(x_i, x_j, y) - \hat{f}(x_i, x_j, y)) dx_i dx_j}{\int f_u(x_i, x_j, y) dx_i dx_j}. \end{aligned} \quad (17)$$

Now, using (17) we first compute $F'(u)$,

$$\begin{aligned} &\int \frac{d}{du} (m_i(f_u, y)) m_j(f_u, y) f_u(x_i, x_j, y) + m_i(f_u, y) \frac{d}{du} (m_j(f_u, y)) f_u(x_i, x_j, y) \\ &+ m_i(f_u, y) m_j(f_u, y) \frac{d}{du} (f_u(x_i, x_j, y)) dx_i dx_j dy, \\ &= \int [x_i m_j(f_u, y) + x_j m_i(f_u, y) - m_i(f_u, y) m_j(f_u, y)] (f(x_i, x_j, y) - \hat{f}(x_i, x_j, y)) dx_i dx_j dy. \end{aligned}$$

Taking $u = 0$ we have

$$F'(0) = \int [x_i m_j(\hat{f}, y) + x_j m_i(\hat{f}, y) - m_i(\hat{f}, y) m_j(\hat{f}, y)] (f(x_i, x_j, y) - \hat{f}(x_i, x_j, y)) dx_i dx_j dy. \quad (18)$$

We derive now $m_i(f_u, y)m_j(f_u, y)$ to obtain

$$\begin{aligned}
\frac{d}{du} (m_i(f_u, y)m_j(f_u, y)) &= \frac{d}{du} (m_i(f_u, y)) m_j(f_u, y) + m_i(f_u, y) \frac{d}{du} (m_j(f_u, y)) \\
&= m_j(f_u, y) \frac{\int (x_i - m_i(f_u, y))(f(x_i, x_j, y) - \hat{f}(x_i, x_j, y))dx_i dx_j}{\int f_u(x_i, x_j, y)dx_i dx_j} \\
&\quad + m_i(f_u, y) \frac{\int (x_j - m_j(f_u, y))(f(x_i, x_j, y) - \hat{f}(x_i, x_j, y))dx_i dx_j}{\int f_u(x_i, x_j, y)dx_i dx_j}.
\end{aligned} \tag{19}$$

Following with $F''(u)$ and using (17) and (19) we get,

$$\begin{aligned}
F''(u) &= \int \left[x_{i1} \frac{\int (x_{j2} - m_j(f_u, y))(f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y))dx_{i2} dx_{j2}}{\int f_u(x_i, x_j, y)dx_i dx_j} \right. \\
&\quad + x_{j1} \frac{\int (x_{i2} - m_i(f_u, y))(f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y))dx_{i2} dx_{j2}}{\int f_u(x_i, x_j, y)dx_i dx_j} \\
&\quad - m_j(f_u, y) \frac{\int (x_{i2} - m_i(f_u, y))(f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y))dx_i dx_j}{\int f_u(x_i, x_j, y)dx_i dx_j} \\
&\quad \left. - m_i(f_u, y) \frac{\int (x_{j2} - m_j(f_u, y))(f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y))dx_{i2} dx_{j2}}{\int f_u(x_i, x_j, y)dx_i dx_j} \right] \\
&\quad (f(x_{i1}, x_{j1}, y) - \hat{f}(x_{i1}, x_{j1}, y))dx_{i1} dx_{j1} dy.
\end{aligned}$$

Simplifying the last expression we obtain

$$\begin{aligned}
F''(u) &= \\
&\int \frac{1}{\int f_u(x_i, x_j, y)dx_i dx_j} \left\{ (x_{i1} - m_i(f_u, y))(x_{j2} - m_j(f_u, y)) + (x_{i2} - m_i(f_u, y))(x_{j1} - m_j(f_u, y)) \right\} \\
&\quad (f(x_{i1}, x_{j1}, y) - \hat{f}(x_{i1}, x_{j1}, y))(f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y))dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy.
\end{aligned}$$

Besides, when $u = 0$

$$\begin{aligned}
F''(0) &= \tag{20} \\
&\int \frac{1}{\int \hat{f}(x_i, x_j, y)dx_i dx_j} \left\{ (x_{i1} - m_i(\hat{f}, y))(x_{j2} - m_j(\hat{f}, y)) + (x_{i2} - m_i(\hat{f}, y))(x_{j1} - m_j(\hat{f}, y)) \right\} \\
&\quad (f(x_{i1}, x_{j1}, y) - \hat{f}(x_{i1}, x_{j1}, y))(f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y))dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\
&= \int \frac{2}{\int \hat{f}(x_i, x_j, y)dx_i dx_j} (x_{i1} - m_i(\hat{f}, y))(x_{j2} - m_j(\hat{f}, y)) \\
&\quad (f(x_{i1}, x_{j1}, y) - \hat{f}(x_{i1}, x_{j1}, y))(f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y))dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy. \tag{21}
\end{aligned}$$

Using the previous arguments we can finally find $F'''(u)$:

$$F'''(u) = \int \frac{-6}{\int f_u(x_i, x_j, y) dx_i dx_j} (x_{i1} - m_j(f_u, y))(x_{j2} - m_j(f_u, y)) (f(x_{i1}, x_{j1}, y) - \hat{f}(x_{i1}, x_{j1}, y))(f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y)) (f(x_{i3}, x_{j3}, y) - \hat{f}(x_{i3}, x_{j3}, y)) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dx_{i3} dx_{j3} dy \quad (22)$$

Replacing (18), (21) and (22) into (3) we get the desired decomposition. \square

Proof of Theorem 1.

We will first control the remaining term (7),

$$\Gamma_n = \frac{1}{6} F'''(\xi)(1 - \xi)^3.$$

Remember that

$$F'''(\xi) = -6 \int \frac{(x_{i1} - m_i(f_\xi, y))(x_{j2} - m_j(f_\xi, y))}{(\int f_\xi(x_i, x_j, y) dx_i dx_j)^2} \left(f(x_{i1}, x_{j1}, y) - \hat{f}(x_{i1}, x_{j1}, y) \right) \left(f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y) \right) \left(f(x_{i3}, x_{j3}, y) - \hat{f}(x_{i3}, x_{j3}, y) \right) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dx_{i3} dx_{j3} dy,$$

Assumptions 1 and 2 ensure that the first part of the integrand is bounded by a constant μ . Furthermore,

$$\begin{aligned} |\Gamma_n| &\leq \mu \int \left| f(x_{i1}, x_{j1}, y) - \hat{f}(x_{i1}, x_{j1}, y) \right| \left| f(x_{i2}, x_{j2}, y) - \hat{f}(x_{i2}, x_{j2}, y) \right| \\ &\quad \left| f(x_{i3}, x_{j3}, y) - \hat{f}(x_{i3}, x_{j3}, y) \right| dx_{i1} dx_{j1} dx_{i2} dx_{j2} dx_{i3} dx_{j3} dy \\ &= \mu \int \left(\int \left| f(x_i, x_j, y) - \hat{f}(x_i, x_j, y) \right| dx_i dx_j \right)^3 dy \\ &\leq \mu \Delta_{x_i x_j}^3 \int \left| f(x_{i1}, x_{j1}, y) - \hat{f}(x_{i1}, x_{j1}, y) \right|^3 dx_i dx_j dy \end{aligned}$$

by the Hölder inequality. Then $\mathbb{E}[\Gamma_n^2] = O(\mathbb{E}(f |f - \hat{f}|^3)^2) = O(\mathbb{E}\|f - \hat{f}\|_3^6)$. Since \hat{f} verifies Assumption 3, this quantity is of order $O(n_1^{-6\lambda})$. Since we also assume $n_1 \approx n/\log(n)$ and $\lambda > 1/6$, then $n_1^{-6\lambda} = o(\frac{1}{n})$. Therefore, we get $\mathbb{E}[\Gamma_n^2] = o(1/n)$ which implies that the remaining term Γ_n is negligible.

To prove the asymptotic normality of $\hat{T}_{ij}^{(n)}$, we shall show that $\sqrt{n} \left(\hat{T}_{ij}^{(n)} - T_{ij}(f) \right)$ and define

$$Z_{ij}^{(n)} = \frac{1}{n_2} \sum_{k=1}^{n_2} H_1(f, X_i^{(k)}, X_j^{(k)}, Y^{(k)}) - \int H_1(f, x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \quad (23)$$

have the same asymptotic behavior. We can get for $Z_{ij}^{(n)}$ a classic central limit theorem with variance

$$\begin{aligned} C_{ij}(f) &= \text{Var}(H_1(f, x_i, x_j, y)) \\ &= \int H_1(f, x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy - \left(\int H_1(f, x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2 \end{aligned}$$

which implies (9) and (10). In order to establish our claim, we will show that

$$R_{ij}^{(n)} = \sqrt{n} \left[\widehat{T}_{ij}^{(n)} - T_{ij}(f) - Z_{ij}^{(n)} \right] \quad (24)$$

has second-order moment converging to 0.

Define $\widehat{Z}_{ij}^{(n)}$ as $Z_{ij}^{(n)}$ with f replaced by \hat{f} . Let us note that $R_{ij}^{(n)} = R_1 + R_2$ where

$$\begin{aligned} R_1 &= \sqrt{n} \left[\widehat{T}_{ij}^{(n)} - T_{ij}(f) - \widehat{Z}_{ij}^{(n)} \right] \\ R_2 &= \sqrt{n} \left[\widehat{Z}_{ij}^{(n)} - Z_{ij}^{(n)} \right]. \end{aligned}$$

It only remains to state that $\mathbb{E}[R_1^2]$ and $\mathbb{E}[R_2^2]$ converges to 0. We can rewrite R_1 as

$$R_1 = -\sqrt{n} \left[\widehat{Q} - Q + \Gamma_n \right]$$

where we note that

$$\begin{aligned} Q &= \int H_2(\hat{f}, x_{i1}, x_{j2}, y) f(x_{i1}, x_{j1}, y) f(x_{i2}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\ H_2(\hat{f}, x_{i1}, x_{j2}, y) &= \frac{1}{\int \hat{f}(x_i, x_j, y) dx_i dx_j} \left(x_{i1} - m_i(\hat{f}, y) \right) \left(x_{j2} - m_j(\hat{f}, y) \right) \end{aligned}$$

has the form of a quadratic functional studied in Section 4 with $\eta(x_{i1}, x_{j2}, y) = H_2(\hat{f}, x_{i1}, x_{j2}, y)$. Hence such functional can be estimated as done in Section 4 and let \widehat{Q} be its corresponding estimator. Since $\mathbb{E}[\Gamma_n^2] = o(1/n)$, we only have to control the term $\sqrt{n}(\widehat{Q} - Q)$ which is such that $\lim_{n \rightarrow \infty} n \mathbb{E}[\widehat{Q} - Q]^2 = 0$ by Lemma 7. This Lemma implies that $\mathbb{E}[R_1^2] \rightarrow 0$ as $n \rightarrow \infty$. For R_2 we have

$$\begin{aligned} \mathbb{E}[R_2^2] &= \frac{n}{n_2} \left[\int \left(H_1(f, x_i, x_j, y) - H_1(\hat{f}, x_i, x_j, y) \right)^2 f(x_i, x_j, y) dx_i dx_j dy \right] \\ &\quad - \frac{n}{n_2} \left[\int H_1(f, x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy - \int H_1(\hat{f}, x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy \right]^2. \end{aligned}$$

The same arguments as the ones of Lemma 7 (mean value and Assumptions 2 and 3) show that $\mathbb{E}[R_2^2] \rightarrow 0$. \square

Proof of Theorem 2. To prove the inequality we will use the usual framework described in Ibragimov & Khas'minskii (1991). The first step is to calculate the Fréchet derivative of $T_{ij}(f)$ at some point $f_0 \in \mathcal{E}$. Assumptions 2 and 3 and equation (4), imply that

$$\begin{aligned} T_{ij}(f) - T_{ij}(f_0) &= \int \left(x_i m_j(f_0, y) + x_j m_i(f_0, y) - m_i(f_0, y) m_j(f_0, y) \right) \\ &\quad \left(f(x_i, x_j, y) - f_0(x_i, x_j, y) \right) dx_i dx_j dy + O \left(\int (f - f_0)^2 \right) \end{aligned}$$

where $m_i(f_0, y) = \int x_i f_0(x_i, x_j, y) dx_i dx_j dy / \int f_0(x_i, x_j, y) dx_i dx_j dy$. Therefore, the Fréchet derivative of $T_{ij}(f)$ at f_0 is $T'_{ij}(f_0) \cdot h = \langle H_1(f_0, \cdot), h \rangle$ with

$$H_1(f_0, x_i, x_j, y) = x_i m_j(f_0, y) + x_j m_i(f_0, y) - m_i(f_0, y) m_j(f_0, y).$$

Using the results of Ibragimov & Khas'minskii (1991), denote $H(f_0) = \{u \in \mathbb{L}^2(dx_i dx_j dy), \int u(x_i, x_j, y) \sqrt{f_0(x_i, x_j, y)} dx_i dx_j dy = 0\}$ the set of functions in $\mathbb{L}^2(dx_i dx_j dy)$ orthogonal to $\sqrt{f_0}$, $\text{Pr}_{H(f_0)}$ the projection onto $H(f_0)$, $A_n(t) = (\sqrt{f_0})t/\sqrt{n}$ and $P_{f_0}^{(n)}$ the joint distribution of $(X_i^{(k)}, X_j^{(k)})$ $k = 1, \dots, n$ under f_0 . Since $(X_i^{(k)}, X_j^{(k)})$ $k = 1, \dots, n$ are i.i.d., the family $\{P_{f_0}^{(n)}, f \in \mathcal{E}\}$ is differentiable in quadratic mean at f_0 and therefore locally asymptotically normal at all points $f_0 \in \mathcal{E}$ in the direction $H(f_0)$ with normalizing factor $A_n(f_0)$ (see the details in Van der Vaart (2000)). Then, by the results of Ibragimov & Khas'minskii (1991) say that under these conditions, denoting $K_n = B_n \theta'(f_0) A_n \text{Pr}_{H(f_0)}$ with $B_n = \sqrt{n}u$, if $K_n \xrightarrow{\mathcal{D}} K$ and if $K(u) = \langle t, u \rangle$, then for every estimator $\widehat{T}_{ij}^{(n)}$ of $T_{ij}(f)$ and every family $\mathcal{V}(f_0)$ of vicinities of f_0 , we have

$$\inf_{\{\mathcal{V}(f_0)\}} \liminf_{n \rightarrow \infty} \sup_{f \in \mathcal{V}(f_0)} n \mathbb{E} [\widehat{T}_{ij}^{(n)} - T_{ij}(f_0)]^2 \geq \|t\|_{\mathbb{L}^2(dx_i dx_j dy)}^2.$$

Here,

$$K_n(u) = \sqrt{n} T'(f_0) \cdot \frac{\sqrt{f_0}}{\sqrt{n}} \text{Pr}_{H(f_0)}(u) = T'(f_0) \left(\sqrt{f_0} \left(u - \sqrt{f_0} \int u \sqrt{f_0} \right) \right),$$

since for any $u \in \mathbb{L}^2(dx_i dx_j dy)$ we can write it as $u = \sqrt{f_0} \langle \sqrt{f_0}, u \rangle + \text{Pr}_{H(f_0)}(u)$. In this case $K_n(u)$ does not depend on n and

$$\begin{aligned} K(h) &= T'(f_0) \cdot \left(\sqrt{f_0} \left(u - \sqrt{f_0} \int h \sqrt{f_0} \right) \right) \\ &= \int H_1(f_0, \cdot) \sqrt{f_0} u - \int H_1(f_0, \cdot) \sqrt{f_0} \int u \sqrt{f_0} \\ &= \langle t, u \rangle \end{aligned}$$

with

$$t(x_i, x_j, y) = H_1(f_0, x_i, x_j, y) \sqrt{f_0} - \left(\int H_1(f_0, x_i, x_j, y) f_0 \right) \sqrt{f_0}.$$

The semi-parametric Cramér-Rao bound for this problem is thus

$$\|t\|_{\mathbb{L}^2(dx_i, dx_j, dy)}^2 = \int H_1(f_0, x_i, x_j, y)^2 f_0 dx_i dx_j dy - \left(\int H_1(f_0, x_i, x_j, y) f_0 dx_i dx_j dy \right)^2 = C_{ij}(f_0)$$

and we recognize the expression $C_{ij}(f_0)$ found in Theorem 1. \square

Proof of Corollary 1. The proof is based in the following observation. Employing equation (24) we have

$$\widehat{\mathbf{T}}^{(n)} - \mathbf{T}(f) = \mathbf{Z}^{(n)}(f) + \frac{\mathbf{R}^{(n)}}{\sqrt{n}}$$

where $\mathbf{Z}^{(n)}(f)$ and $\mathbf{R}^{(n)}$ are matrices with elements $Z_{ij}^{(n)}$ and $R_{ij}^{(n)}$, defined in (23) and (24), respectively.

Hence we have,

$$n\mathbb{E}[\|\text{vech}\left(\widehat{\mathbf{T}}^{(n)} - \mathbf{T}(f) - \mathbf{Z}^{(n)}(f)\right)\|^2] = \mathbb{E}[\|\text{vech}\left(\mathbf{R}^{(n)}\right)\|^2] = \sum_{i \leq j} \mathbb{E}[\left(R_{ij}^{(n)}\right)^2].$$

We see by Lemma 7 that $\mathbb{E}[R_{ij}^2] \rightarrow 0$ as $n \rightarrow 0$. It follows that

$$n\mathbb{E}[\|\text{vech}\left(\widehat{\mathbf{T}}^{(n)} - \mathbf{T}(f) - \mathbf{Z}^{(n)}(f)\right)\|^2] \rightarrow 0 \text{ as } n \rightarrow 0.$$

We know that if X_n , X and Y_n are random variables, then if $X_n \xrightarrow{\mathcal{D}} X$ and $(X_n - Y_n) \xrightarrow{\mathcal{P}} 0$, follows that $Y_n \xrightarrow{\mathcal{D}} X$.

Remember also that convergence in \mathbb{L}^2 implies convergence in probability, therefore

$$\sqrt{n} \text{vech}\left(\widehat{\mathbf{T}}^{(n)} - \mathbf{T}(f) - \mathbf{Z}^{(n)}(f)\right) \xrightarrow{\mathcal{P}} 0.$$

By the multivariate central limit theorem we have that $\sqrt{n} \text{vech}\left(\mathbf{Z}^{(n)}(f)\right) \xrightarrow{\mathcal{D}} \mathcal{N}(0, \mathbf{C}(f))$. Therefore, $\sqrt{n} \text{vech}\left(\widehat{\mathbf{T}}^{(n)} - \mathbf{T}(f)\right) \xrightarrow{\mathcal{D}} \mathcal{N}(0, \mathbf{C}(f))$. \square

Proof of Theorem 3. For abbreviation, we write M instead of M_n and set $m = |M_n|$. We first compute the mean squared error of $\hat{\theta}_n$ as

$$\mathbb{E}[\hat{\theta}_n - \theta]^2 = \text{Bias}^2(\hat{\theta}_n) + \text{Var}(\hat{\theta}_n)$$

where $\text{Bias}(\hat{\theta}_n) = \mathbb{E}[\hat{\theta}_n] - \theta$.

We begin the proof by bounding $\text{Var}(\hat{\theta}_n)$. Let A and B be $m \times 1$ vectors with components

$$\begin{aligned} a_l &= \int p_l(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \quad l = 1, \dots, m, \\ b_l &= \int p_l(x_{i1}, x_{j1}, y) f(x_{i2}, x_{j2}, y) \psi(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\ &= \int p_l(x_i, x_j, y) g(x_i, x_j, y) dx_i dx_j dy \quad l = 1, \dots, m \end{aligned}$$

where $g(x_i, x_j, y) = \int f(x_{i2}, x_{j2}, y) \psi(x_i, x_j, x_{i2}, x_{j2}, y) dx_{i2} dx_{j2}$. Let Q and R be $m \times 1$ vectors of centered functions

$$\begin{aligned} q_l(x_i, x_j, y) &= p_l(x_i, x_j, y) - a_l \\ r_l(x_i, x_j, y) &= \int p_l(x_{i2}, x_{j2}, y) \psi(x_i, x_j, x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} - b_l \end{aligned}$$

for $l = 1, \dots, m$. Let C a $m \times m$ matrix of constants

$$c_{ll'} = \int p_l(x_{i1}, x_{j1}, y) p_{l'}(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \quad l, l' = 1, \dots, m.$$

Let us denote by U_n the process

$$U_n h = \frac{1}{n(n-1)} \sum_{k \neq k'=1}^n h(X_i^{(k)}, X_j^{(k)}, Y^{(k)}, X_i^{(k')}, X_j^{(k')}, Y^{(k')})$$

and P_n the empirical measure

$$P_n h = \frac{1}{n} \sum_{k=1}^n h(X_i^{(k)}, X_j^{(k)}, Y^{(k)})$$

for some h in $\mathbb{L}^2(dx_i, dx_j, dy)$. With these notations, $\hat{\theta}_n$ has the Hoeffding's decomposition

$$\begin{aligned} \hat{\theta}_n &= \frac{1}{n(n-1)} \sum_{l \in M} \sum_{k \neq k'=1}^n (q_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) + a_l) (r_l(X_i^{(k')}, X_j^{(k')}, Y^{(k')}) + b_l) \\ &\quad - \frac{1}{n(n-1)} \sum_{l, l' \in M} \sum_{k \neq k'=1}^n (q_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) + a_l) (q_{l'}(X_i^{(k')}, X_j^{(k')}, Y^{(k')}) + a_{l'}) c_{ll'} \\ &= U_n K + P_n L + A^\top B - A^\top C A \end{aligned}$$

where

$$\begin{aligned} K(x_{i1}, x_{j1}, y_1, x_{i2}, x_{j2}, y_2) &= Q^\top(x_{i1}, x_{j1}, y_1) R(x_{i2}, x_{j2}, y_2) - Q^\top(x_{i1}, x_{j1}, y_1) C Q(x_{i2}, x_{j2}, y_2) \\ L(x_i, x_j, y) &= A^\top R(x_i, x_j, y) + B Q(x_i, x_j, y) - 2A^\top C Q(x_i, x_j, y). \end{aligned}$$

Therefore $\text{Var}(\hat{\theta}_n) = \text{Var}(U_n K) + \text{Var}(P_n L) - 2 \text{Cov}(U_n K, P_n L)$. These three terms are bounded in Lemmas 2 - 4, which gives

$$\text{Var}(\hat{\theta}_n) \leq \frac{20}{n(n-1)} \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2 (m+1) + \frac{12}{n} \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2.$$

For n enough large and a constant $\gamma \in \mathbb{R}$,

$$\text{Var}(\hat{\theta}_n) \leq \gamma \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2 \left(\frac{m}{n^2} + \frac{1}{n} \right).$$

The term Bias $(\hat{\theta}_n)$ is easily computed, as proven in Lemma 5, is equal to

$$\begin{aligned} & - \int (S_M f(x_{i1}, x_{j1}, y) - f(x_{i1}, x_{j1}, y)) (S_M f(x_{i2}, x_{j2}, y) - f(x_{i2}, x_{j2}, y)) \\ & \quad \eta(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy. \end{aligned}$$

From Lemma 5, the bias of $\hat{\theta}_n$ is bounded by

$$\left| \text{Bias}(\hat{\theta}_n) \right| \leq \Delta_{x_i x_j} \|\eta\|_\infty \sup_{l \notin M} |c_l|^2.$$

The assumption of $\left(\sup_{l \notin M} |c_l|^2 \right)^2 \approx m/n^2$ and since $m/n \rightarrow 0$, we deduce that $\mathbb{E}[\hat{\theta}_n - \theta]^2$ has a parametric rate of convergence $O(1/n)$.

Finally to prove (16), note that

$$\begin{aligned} n\mathbb{E}[\hat{\theta}_n - \theta]^2 &= n \text{Bias}^2(\hat{\theta}_n) + n \text{Var}(\hat{\theta}_n) \\ &= n \text{Bias}^2(\hat{\theta}_n) + n \text{Var}(U_n K) + n \text{Var}(P_n L). \end{aligned}$$

We previously proved that for some $\lambda_1, \lambda_2 \in \mathbb{R}$

$$\begin{aligned} n \text{Bias}^2(\hat{\theta}_n) &\leq \lambda_1 \Delta_{x_i x_j}^2 \|\eta\|_\infty^2 \frac{m}{n} \\ n \text{Var}(U_n K) &\leq \lambda_2 \Delta_{x_i x_j}^2 \|f\|_\infty^2 \|\eta\|_\infty^2 \frac{m}{n}. \end{aligned}$$

Thus, Lemma 6 implies

$$|n \text{Var}(P_n L) - \Lambda(f, \eta)| \leq \lambda [\|S_M f - f\|_2 + \|S_M g - g\|_2],$$

where λ is an increasing function of $\|f\|_\infty^2$, $\|\eta\|_\infty^2$ and $\Delta_{x_i x_j}$. From all this we deduce (16) which ends the proof of Theorem 3. \square

6 Technical Results

Lemma 1 (Bias of $\hat{\theta}_n$). *The estimator $\hat{\theta}_n$ defined in (14) estimates θ with bias equal to*

$$-\int (S_M f(x_{i1}, x_{j1}, y) - f(x_{i1}, x_{j1}, y)) (S_M f(x_{i2}, x_{j2}, y) - f(x_{i2}, x_{j2}, y)) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy.$$

Proof. Let $\hat{\theta}_n = \hat{\theta}_n^1 - \hat{\theta}_n^2$ where

$$\begin{aligned} \hat{\theta}_n^1 &= \frac{1}{n(n-1)} \sum_{l \in M} \sum_{k \neq k'=1} p_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) \int p_l(x_i, x_j, Y^{(k')}) \psi(x_i, x_j, X_i^{(k')}, X_j^{(k')}, Y^{(k')}) dx_i dx_j \\ \hat{\theta}_n^2 &= -\frac{1}{n(n-1)} \sum_{l, l' \in M} \sum_{k \neq k'=1}^n p_l(X_i^{(k)}, X_j^{(k)}, Y^{(k)}) p_{l'}(X_i^{(k')}, X_j^{(k')}, Y^{(k')}) \\ &\quad \int p_l(x_{i1}, x_{j1}, y) p_{l'}(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy. \end{aligned}$$

Let us first compute $\mathbb{E}[\hat{\theta}_n^1]$.

$$\begin{aligned}
\mathbb{E}[\hat{\theta}_n^1] &= \sum_{l \in M} \int p_l(x_{i1}, x_{j1}, y) f(x_{i1}, x_{j1}, y) dx_{i1} dx_{j1} dy \\
&\quad \int p_l(x_{i1}, x_{j1}, y) \psi(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) f(x_{i2}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\
&= \sum_{l \in M} a_l \int p_l(x_{i1}, x_{j1}, y) \psi(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) f(x_{i2}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\
&= \int \left(\sum_{l \in M} a_l p_l(x_{i2}, x_{j2}, y) \right) \psi(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) f(x_{i2}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\
&= \int S_M f(x_{i1}, x_{j1}, y) f(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\
&\quad + \int S_M f(x_{i2}, x_{j2}, y) f(x_{i1}, x_{j1}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy
\end{aligned}$$

Now for $\hat{\theta}_n^2$, we get

$$\begin{aligned}
\mathbb{E}[\hat{\theta}_n^2] &= \sum_{l, l' \in M} \int p_l(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \int p_{l'}(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \\
&\quad \int p_l(x_{i1}, x_{j1}, y) p_{l'}(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\
&= \sum_{l, l' \in M} a_l a_{l'} \int p_l(x_{i1}, x_{j1}, y) p_{l'}(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\
&= \int \left(\sum_{l \in M} a_l p_l(x_{i1}, x_{j1}, y) \right) \left(\sum_{l' \in M} a_{l'} p_{l'}(x_{i2}, x_{j2}, y) \right) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\
&= \int S_M f(x_{i1}, x_{j1}, y) S_M f(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy.
\end{aligned}$$

Arranging these terms and using

$$\text{Bias}(\hat{\theta}_n) = \mathbb{E}[\hat{\theta}_n] - \theta = \mathbb{E}[\hat{\theta}_n^1] - \mathbb{E}[\hat{\theta}_n^2] - \theta$$

we obtain the desire bias. \square

Lemma 2 (Bound of $\text{Var}(U_n K)$). *Under the assumptions of Theorem 3, we have*

$$\text{Var}(U_n K) \leq \frac{20}{n(n-1)} \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2 (m+1)$$

Proof. Note that $U_n K$ is centered because Q and R are centered and $(X_i^{(k)}, X_j^{(k)}, Y^{(k)})$, $k =$

$1, \dots, n$ is an independent sample. So $\text{Var}(U_n K)$ is equal to

$$\begin{aligned} \mathbb{E}[U_n K]^2 &= \mathbb{E}\left(\frac{1}{(n(n-1))^2} \sum_{k_1 \neq k'_1=1}^n \sum_{k_2 \neq k'_2=1}^n K(X_i^{(k_1)}, X_j^{(k_1)}, Y^{(k_1)}, X_i^{(k'_1)}, X_j^{(k'_1)}, Y^{(k'_1)}) \right. \\ &\quad \left. K(X_i^{(k_2)}, X_j^{(k_2)}, Y^{(k_2)}, X_i^{(k'_2)}, X_j^{(k'_2)}, Y^{(k'_2)})\right) \\ &= \frac{1}{n(n-1)} \mathbb{E}\left(K^2(X_i^{(1)}, X_j^{(1)}, Y^{(1)}, X_i^{(2)}, X_j^{(2)}, Y^{(2)}) \right. \\ &\quad \left. + K(X_i^{(1)}, X_j^{(1)}, Y^{(1)}, X_i^{(2)}, X_j^{(2)}, Y^{(2)})K(X_i^{(2)}, X_j^{(2)}, Y^{(2)}, X_i^{(1)}, X_j^{(1)}, Y^{(1)})\right) \end{aligned}$$

By the Cauchy-Schwarz inequality, we get

$$\text{Var}(U_n K) \leq \frac{2}{n(n-1)} \mathbb{E}[K^2(X_i^{(1)}, X_j^{(1)}, Y^{(1)}, X_i^{(2)}, X_j^{(2)}, Y^{(2)})].$$

Moreover, using the fact that $2|\mathbb{E}[XY]| \leq \mathbb{E}[X^2] + \mathbb{E}[Y^2]$, we obtain

$$\begin{aligned} \mathbb{E}[K^2(X_i^{(1)}, X_j^{(1)}, Y^{(1)}, X_i^{(2)}, X_j^{(2)}, Y^{(2)})] &\leq 2 \left[\mathbb{E}[(Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)})R(X_i^{(2)}, X_j^{(2)}, Y^{(2)}))^2] \right. \\ &\quad \left. + \mathbb{E}[(Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)})CQ(X_i^{(2)}, X_j^{(2)}, Y^{(2)}))^2] \right]. \end{aligned}$$

We will bound these two terms. The first one is

$$\begin{aligned} &\mathbb{E}[(Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)})R(X_i^{(2)}, X_j^{(2)}, Y^{(2)}))^2] \\ &= \sum_{l, l' \in M} \left(\int p_l(x_i, x_j, y)p_{l'}(x_i, x_j, y)f(x_i, x_j, y)dx_i dx_j dy - a_l a_{l'} \right) \\ &\quad \left(\int p_l(x_{i2}, x_{j2}, y)p_{l'}(x_{i3}, x_{j3}, y)\psi(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) \right. \\ &\quad \left. \psi(x_{i1}, x_{j1}, x_{i3}, x_{j3}, y)f(x_{i1}, x_{j1}, y)dx_{i1} dx_{j1} dx_{i2} dx_{j2} dx_{i3} dx_{j3} dy - b_l b_{l'} \right) \\ &= W_1 - W_2 - W_3 + W_4 \end{aligned}$$

where

$$\begin{aligned}
W_1 &= \int \sum_{l,l' \in M} p_l(x_{i_1}, x_{j_1}, y) p_{l'}(x_{i_1}, x_{j_1}, y) p_l(x_{i_2}, x_{j_2}, y') p_{l'}(x_{i_3}, x_{j_3}, y') \psi(x_{i_4}, x_{j_4}, x_{i_2}, x_{j_2}, y') \\
&\quad \psi(x_{i_4}, x_{j_4}, x_{i_3}, x_{j_3}, y') f(x_{i_1}, x_{j_1}, y) f(x_{i_4}, x_{j_4}, y') dx_{i_1} dx_{j_1} dx_{i_2} dx_{j_2} dx_{i_3} dx_{j_3} dx_{i_4} dx_{j_4} dy dy' \\
W_2 &= \int \sum_{l,l' \in M} b_l b_{l'} p_l(x_{i_1}, x_{j_1}, y) p_{l'}(x_{i_1}, x_{j_1}, y) f(x_{i_1}, x_{j_1}, y) dx_{i_1} dx_{j_1} dy \\
W_3 &= \int \sum_{l,l' \in M} a_l a_{l'} p_l(x_{i_2}, x_{j_2}, y') p_{l'}(x_{i_3}, x_{j_3}, y') \\
&\quad \psi(x_{i_4}, x_{j_4}, x_{i_2}, x_{j_2}, y') \psi(x_{i_4}, x_{j_4}, x_{i_3}, x_{j_3}, y') f(x_{i_4}, x_{j_4}, y') dx_{i_2} dx_{j_2} dx_{i_3} dx_{j_3} dx_{i_4} dx_{j_4} dy' \\
W_4 &= \sum_{l,l' \in M} a_l a_{l'} b_l b_{l'}.
\end{aligned}$$

W_2 and W_3 are positive, hence

$$\mathbb{E} \left[\left(2Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) R(X_i^{(2)}, X_j^{(2)}, Y^{(2)}) \right)^2 \right] \leq W_1 + W_4.$$

$$\begin{aligned}
W_1 &= \int \sum_{l,l' \in M} p_l(x_{i_1}, x_{j_1}, y) p_{l'}(x_{i_1}, x_{j_1}, y) \left(\int p_l(x_{i_2}, x_{j_2}, y') \psi(x_{i_4}, x_{j_4}, x_{i_2}, x_{j_2}, y') dx_{i_2} dx_{j_2} \right) \\
&\quad \left(\int p_{l'}(x_{i_3}, x_{j_3}, y') \psi(x_{i_4}, x_{j_4}, x_{i_3}, x_{j_3}, y') dx_{i_3} dx_{j_3} \right) f(x_{i_1}, x_{j_1}, y) f(x_{i_4}, x_{j_4}, y') dx_{i_1} dx_{j_1} dx_{i_4} dx_{j_4} dy dy' \\
&\leq \|f\|_\infty^2 \sum_{l,l' \in M} \int p_l(x_{i_1}, x_{j_1}, y) p_{l'}(x_{i_1}, x_{j_1}, y) dx_{i_1} dx_{j_1} dy \\
&\quad \int \left(\int p_l(x_{i_2}, x_{j_2}, y') \psi(x_{i_4}, x_{j_4}, x_{i_2}, x_{j_2}, y') dx_{i_2} dx_{j_2} \right) \\
&\quad \left(\int p_{l'}(x_{i_3}, x_{j_3}, y') \psi(x_{i_4}, x_{j_4}, x_{i_3}, x_{j_3}, y') dx_{i_3} dx_{j_3} \right) dx_{i_2} dx_{j_2} dx_{i_4} dx_{j_4} dy'
\end{aligned}$$

Since p_l 's are orthonormal we have

$$W_1 \leq \|f\|_\infty^2 \sum_{l \in M} \int \left(\int p_l(x_{i_2}, x_{j_2}, y') \psi(x_{i_4}, x_{j_4}, x_{i_2}, x_{j_2}, y') dx_{i_2} dx_{j_2} \right)^2 dx_{i_4} dx_{j_4} dy'.$$

Moreover by the Cauchy-Schwarz inequality and $\|\psi\|_\infty \leq 2\|\eta\|_\infty$

$$\begin{aligned}
\left(\int p_l(x_{i_2}, x_{j_2}, y') \psi(x_{i_4}, x_{j_4}, x_{i_2}, x_{j_2}, y') dx_{i_2} dx_{j_2} \right)^2 &\leq \int p_l(x_{i_2}, x_{j_2}, y')^2 dx_{i_2} dx_{j_2} \\
&\quad \int \psi(x_{i_4}, x_{j_4}, x_{i_2}, x_{j_2}, y')^2 dx_{i_2} dx_{j_2} \\
&\leq \|\psi\|_\infty^2 \Delta_{x_i x_j} \int p_l(x_{i_2}, x_{j_2}, y')^2 dx_{i_2} dx_{j_2} \\
&\leq 4\|\eta\|_\infty^2 \Delta_{x_i x_j} \int p_l(x_{i_2}, x_{j_2}, y')^2 dx_{i_2} dx_{j_2},
\end{aligned}$$

and then

$$\begin{aligned}
& \int \left(\int p_l(x_{i2}, x_{j2}, y') \psi(x_{i4}, x_{j4}, x_{i2}, x_{j2}, y') dx_{i2} dx_{j2} \right)^2 dx_{i4} dx_{j4} dy' \\
& \leq 4 \|\eta\|_\infty^2 \Delta_{x_i x_j}^2 \int p_l(x_{i2}, x_{j2}, y')^2 dx_{i2} dx_{j2} dy' \\
& = 4 \|\eta\|_\infty^2 \Delta_{x_i x_j}^2.
\end{aligned}$$

Finally,

$$W_1 \leq 4 \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2 m.$$

For the term W_4 using the facts that $S_M f$ and $S_M g$ are projection and that $\int f = 1$, we have

$$W_4 = \left(\sum_{l \in M} a_l b_l \right)^2 \leq \sum_{l \in M} a_l^2 \sum_{l \in M} b_l^2 \leq \|f\|_2^2 \|g\|_2^2 \leq \|f\|_\infty \|g\|_2^2.$$

By the Cauchy-Schwartz inequality we have $\|g\|_2^2 \leq 4 \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2$ and then

$$W_4 \leq 4 \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2$$

which leads to

$$\mathbb{E}[(Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)})R(X_i^{(2)}, X_j^{(2)}, Y^{(2)}))^2] \leq 4 \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2 (m+1). \quad (25)$$

The second term $\mathbb{E}[(Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)})CQ(X_i^{(2)}, X_j^{(2)}, Y^{(2)}))] = W_5 - 2W_6 + W_7$ where

$$\begin{aligned}
W_5 &= \int \sum_{l_1, l'_1} \sum_{l_2, l'_2} c_{l_1 l'_1} c_{l_2 l'_2} p_{l_1}(x_{i1}, x_{j1}, y) p_{l_2}(x_{i1}, x_{j1}, y) p_{l'_1}(x_{i2}, x_{j2}, y') p_{l'_2}(x_{i2}, x_{j2}, y') \\
& \quad f(x_{i1}, x_{j1}, y) f(x_{i2}, x_{j2}, y') dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy' dy \\
W_6 &= \int \sum_{l_1, l'_1} \sum_{l_2, l'_2} c_{l_1 l'_1} c_{l_2 l'_2} a_{l_1} a_{l_2} p_{l'_1}(x_i, x_j, y) p_{l'_2}(x_i, x_j, y) dx_i dx_j dy \\
W_7 &= \sum_{l_1, l'_1} \sum_{l_2, l'_2} c_{l_1 l'_1} c_{l_2 l'_2} a_{l_1} a_{l'_1} a_{l_2} a_{l'_2}.
\end{aligned}$$

Using the previous manipulation, we show that $W_6 \geq 0$. Thus

$$\mathbb{E}[(Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)})CQ(X_i^{(2)}, X_j^{(2)}, Y^{(2)}))] \leq W_5 + W_7.$$

First, observe that

$$\begin{aligned}
W_5 &= \sum_{l_1, l'_1} \sum_{l_2, l'_2} c_{l_1 l'_1} c_{l_2 l'_2} \left(\int p_{l_1}(x_{i1}, x_{j1}, y) p_{l_2}(x_{i1}, x_{j1}, y) f(x_{i1}, x_{j1}, y) dx_{i1} dx_{j1} dy \right) \\
&\quad \left(\int p_{l'_1}(x_{i2}, x_{j2}, y') p_{l'_2}(x_{i2}, x_{j2}, y') f(x_{i2}, x_{j2}, y') dx_{i2} dx_{j2} dy' \right) \\
&\leq \|f\|_\infty^2 \sum_{l_1, l'_1} \sum_{l_2, l'_2} c_{l_1 l'_1} c_{l_2 l'_2} \left(\int p_{l_1}(x_{i1}, x_{j1}, y) p_{l_2}(x_{i1}, x_{j1}, y) dx_{i1} dx_{j1} dy \right) \\
&\quad \left(\int p_{l'_1}(x_{i2}, x_{j2}, y') p_{l'_2}(x_{i2}, x_{j2}, y') dx_{i2} dx_{j2} dy' \right) \\
&= \|f\|_\infty^2 \sum_{l, l'} c_{ll'}^2
\end{aligned}$$

again using the orthonormality of the the p_l 's. Besides given the decomposition $p_l(x_i, x_j, y) = \alpha_{l_\alpha}(x_i, x_j) \beta_{l_\beta}(y)$,

$$\begin{aligned}
\sum_{l, l'} c_{ll'}^2 &= \int \sum_{l_\beta, l'_\beta} \beta_{l_\beta}(y) \beta_{l'_\beta}(y) \beta_{l_\beta}(y') \beta_{l'_\beta}(y') \\
&\quad \sum_{l_\alpha, l'_\alpha} \left(\int \alpha_{l_\alpha}(x_{i1}, x_{j1}) \alpha_{l'_\alpha}(x_{i2}, x_{j2}) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} \right) \\
&\quad \left(\int \alpha_{l_\alpha}(x_{i3}, x_{j3}) \alpha_{l'_\alpha}(x_{i4}, x_{j4}) \eta(x_{i3}, x_{j4}, y') dx_{i3} dx_{j3} dx_{i4} dx_{j4} \right) dy dy'
\end{aligned}$$

But

$$\begin{aligned}
&\sum_{l_\alpha, l'_\alpha} \left(\int \alpha_{l_\alpha}(x_{i1}, x_{j1}) \alpha_{l'_\alpha}(x_{i2}, x_{j2}) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} \right) \\
&\quad \left(\int \alpha_{l_\alpha}(x_{i3}, x_{j3}) \alpha_{l'_\alpha}(x_{i4}, x_{j4}) \eta(x_{i3}, x_{j4}, y') dx_{i3} dx_{j3} dx_{i4} dx_{j4} \right) \\
&= \sum_{l_\alpha, l'_\alpha} \int \alpha_{l_\alpha}(x_{i1}, x_{j1}) \alpha_{l'_\alpha}(x_{i2}, x_{j2}) \eta(x_{i1}, x_{j2}, y) \alpha_{l_\alpha}(x_{i3}, x_{j3}) \\
&\quad \alpha_{l'_\alpha}(x_{i4}, x_{j4}) \eta(x_{i3}, x_{j4}, y') dx_{i1} dx_{j1} dx_{i2} dx_{j2} dx_{i3} dx_{j3} dx_{i4} dx_{j4} \\
&= \int \sum_{l_\alpha} \left(\int \alpha_{l_\alpha}(x_{i1}, x_{j1}) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} \right) \alpha_{l_\alpha}(x_{i3}, x_{j3}) \\
&\quad \sum_{l'_\alpha} \left(\int \alpha_{l'_\alpha}(x_{i4}, x_{j4}) \eta(x_{i3}, x_{j4}, y') dx_{i4} dx_{j4} \right) \alpha_{l'_\alpha}(x_{i2}, x_{j2}) dx_{i2} dx_{j2} dx_{i3} dx_{j3} \\
&\leq \int \eta(x_{i3}, x_{j3}, x_{i2}, x_{j2}, y) \eta(x_{i3}, x_{j2}, y') dx_{i2} dx_{j2} dx_{i3} dx_{j3} \\
&\leq \Delta_{x_i x_j}^2 \|\eta\|_\infty^2
\end{aligned}$$

using the orthonormality of the basis α_{l_α} . Then we get

$$\begin{aligned}
\sum_{l,l'} c_{ll'}^2 &\leq \Delta_{x_i x_j}^2 \|\eta\|_\infty^2 \left(\int \sum_{l_\beta, l'_\beta} \beta_{l_\beta}(y) \beta_{l'_\beta}(y) \beta_{l_\beta}(y') \beta_{l'_\beta}(y') dy dy' \right) \\
&= \Delta_{x_i x_j}^2 \|\eta\|_\infty^2 \sum_{l_\beta, l'_\beta} \left(\int \beta_{l_\beta}(y) \beta_{l'_\beta}(y) dy \right)^2 \\
&\leq \Delta_{x_i x_j}^2 \|\eta\|_\infty^2 \sum_{l_\beta} \left(\int \beta_{l_\beta}^2(y) dy \right)^2 \\
&\leq \Delta_{x_i x_j}^2 \|\eta\|_\infty^2 m
\end{aligned}$$

since the β_{l_β} are orthonormal. Finally

$$W_5 \leq \|f\|_\infty^2 \|\eta\|_\infty^2 \Delta_{x_i x_j}^2 m.$$

Now for W_7 we first will bound,

$$\begin{aligned}
\left| \sum_{l,l'} c_{ll'} a_l a_{l'} \right| &= \left| \int \sum_{l,l' \in M} a_l a_{l'} p_{l_2}(x_{i1}, x_{j1}, y) p_{l'_1}(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \right| \\
&\leq \int |S_M(x_{i1}, x_{j1}, y) S_M(x_{i2}, x_{j2}, y) \eta(x_{i1}, x_{j2}, y)| dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\
&\leq \|\eta\|_\infty \int \left(\int |S_M(x_{i1}, x_{j1}, y) S_M(x_{i2}, x_{j2}, y)| dy \right) dx_{i1} dx_{j1} dx_{i2} dx_{j2}.
\end{aligned}$$

Taking squares in both sides and using the Cauchy-Schwartz inequality twice, we get

$$\begin{aligned}
\left(\sum_{l,l'} c_{ll'} a_l a_{l'} \right)^2 &= \|\eta\|_\infty^2 \left(\int \left(\int |S_M(x_{i1}, x_{j1}, y) S_M(x_{i2}, x_{j2}, y)| dy \right) dx_{i1} dx_{j1} dx_{i2} dx_{j2} \right)^2 \\
&\leq \|\eta\|_\infty^2 \Delta_{x_i x_j}^2 \int \left(\int |S_M(x_{i1}, x_{j1}, y) S_M(x_{i2}, x_{j2}, y)| dy \right)^2 dx_{i1} dx_{j1} dx_{i2} dx_{j2} \\
&\leq \|\eta\|_\infty^2 \Delta_{x_i x_j}^2 \int \left(\int S_M(x_{i1}, x_{j1}, y)^2 dy \right) \left(\int S_M(x_{i2}, x_{j2}, y')^2 dy' \right) dx_{i1} dx_{j1} dx_{i2} dx_{j2} \\
&= \|\eta\|_\infty^2 \Delta_{x_i x_j}^2 \int S_M(x_{i1}, x_{j1}, y)^2 S_M(x_{i1}, x_{j1}, y')^2 dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy dy' \\
&= \|\eta\|_\infty^2 \Delta_{x_i x_j}^2 \left(\int S_M(x_i, x_j, y)^2 dx_i dx_j dy \right) \\
&\leq \|\eta\|_\infty^2 \Delta_{x_i x_j}^2 \|f\|_\infty^2.
\end{aligned}$$

Finally,

$$\mathbb{E}[(Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)})CQ(X_i^{(2)}, X_j^{(2)}, Y^{(2)}))^2] \leq \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2 (m+1). \quad (26)$$

Collecting (25) and (26), we obtain

$$\text{Var}(U_n K) \leq \frac{20}{n(n-1)} \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2 (m+1)$$

which concludes the proof of Lemma 2. \square

Lemma 3 (Bound for $\text{Var}(P_n L)$). *Under the assumptions of Theorem 3, we have*

$$\text{Var}(P_n L) \leq \frac{12}{n} \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2.$$

Proof. First note that given the independence of $(X_i^{(k)}, X_j^{(k)}, Y^{(k)})$ $k = 1, \dots, n$ we have

$$\text{Var}(P_n L) = \frac{1}{n} \text{Var}(L(X_i^{(1)}, X_j^{(1)}, Y^{(1)}))$$

we can write $L(X_i^{(1)}, X_j^{(1)}, Y^{(1)})$ as

$$\begin{aligned} & A^\top R \left(X_i^{(1)}, X_j^{(1)}, Y^{(1)} \right) + B^\top Q \left(X_i^{(1)}, X_j^{(1)}, Y^{(1)} \right) - 2A^\top C Q \left(X_i^{(1)}, X_j^{(1)}, Y^{(1)} \right) \\ &= \sum_{l \in M} a_l \left(\int p_l(x_i, x_j, Y^{(1)}) \psi(x_i, x_j, X_i^{(1)}, X_j^{(1)}, Y^{(1)}) dx_i dx_j - b_l \right) \\ & \quad + \sum_{l \in M} b_l \left(p_l(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) - a_l \right) - 2 \sum_{l, l' \in M} c_{ll'} a_{l'} \left(p_l(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) - a_l \right) \\ &= \int \sum_{l \in M} a_l p_l(x_i, x_j, Y^{(1)}) \psi(x_i, x_j, X_i^{(1)}, X_j^{(1)}, Y^{(1)}) dx_i dx_j \\ & \quad + \sum_{l \in M} b_l p_l(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) - 2 \sum_{l, l' \in M} c_{ll'} a_{l'} p_l(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) - 2A^\top B - 2A^\top C A. \\ &= \int S_M f(x_i, x_j, Y^{(1)}) \psi(x_i, x_j, X_i^{(1)}, X_j^{(1)}, Y^{(1)}) dx_i dx_j + S_M g(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) \\ & \quad - 2 \sum_{l, l' \in M} c_{ll'} a_{l'} p_l(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) - 2A^\top B - 2A^\top C A. \end{aligned}$$

Let $h(x_i, x_j, y) = \int S_M f(x_{i2}, x_{j2}, y) \psi(x_i, x_j, x_{i2}, x_{j2}, y) dx_{i2} dx_{j2}$, we have

$$\begin{aligned} & S_M h(x_i, x_j, y) \\ &= \sum_{l \in M} \left(\int h(x_{i2}, x_{j2}, y) p_l(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} dy \right) p_l(x_i, x_j, y) \\ &= \sum_{l \in M} \left(\int S_M f(x_{i3}, x_{j3}, y) \psi(x_{i2}, x_{j2}, x_{i3}, x_{j3}, y) p_l(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} dx_{i3} dx_{j3} dy \right) p_l(x_i, x_j, y) \\ &= \sum_{l, l' \in M} \left(\int a_{l'} p_{l'}(x_{i3}, x_{j3}, y) \psi(x_{i2}, x_{j2}, x_{i3}, x_{j3}, y) p_l(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} dx_{i3} dx_{j3} dy \right) p_l(x_i, x_j, y) \\ &= 2 \sum_{l, l' \in M} \left(\int a_{l'} p_{l'}(x_{i3}, x_{j3}, y) \eta(x_{i2}, x_{j3}, y) p_l(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} dx_{i3} dx_{j3} dy \right) p_l(x_i, x_j, y) \\ &= 2 \sum_{l, l' \in M} a_{l'} c_{ll'} p_l(x_i, x_j, y) \end{aligned}$$

and we can write

$$\begin{aligned} L(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) &= h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) + S_M g(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) \\ &\quad - S_M h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) - 2A^\top B - 2A^\top C A. \end{aligned}$$

Thus,

$$\begin{aligned} &\text{Var}(L(X_i^{(1)}, X_j^{(1)}, Y^{(1)})) \\ &= \text{Var}(h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) + S_M g(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) + S_M h(X_i^{(1)}, X_j^{(1)}, Y^{(1)})) \\ &\leq \mathbb{E}[(h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) + S_M g(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) + S_M h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}))^2] \\ &\leq \mathbb{E}[(h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}))^2 + (S_M g(X_i^{(1)}, X_j^{(1)}, Y^{(1)}))^2 + (S_M h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}))^2]. \end{aligned}$$

Each of these terms can be bounded

$$\begin{aligned} &\mathbb{E}[(h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}))^2] \\ &= \int \left(\int S_M f(x_{i2}, x_{j2}, y) \psi(x_{i1} x_{j2}, x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} \right)^2 f(x_{i1}, x_{j1}, y) dx_{i1} dx_{j1} dy \\ &\leq \Delta_{x_i x_j} \int S_M f(x_{i2}, x_{j2}, y)^2 \psi(x_{i1} x_{j2}, x_{i2}, x_{j2}, y)^2 f(x_{i1}, x_{j1}, y) dx_{i1} dx_{j1} dx_{i2} dx_{j2} dy \\ &\leq 4\Delta_{x_i x_j}^2 \|f\|_\infty \|\eta\|_\infty^2 \int S_M f(x_i, x_j, y)^2 dx_i dx_j dy \\ &= 4\Delta_{x_i x_j}^2 \|f\|_\infty \|\eta\|_\infty^2 \|S_M f\|_2^2 \\ &\leq 4\Delta_{x_i x_j}^2 \|f\|_\infty \|\eta\|_\infty^2 \|f\|_2^2 \\ &\leq 4\Delta_{x_i x_j}^2 \|f\|_\infty^2 \|\eta\|_\infty^2 \end{aligned}$$

and similar calculations are valid for the others two terms,

$$\begin{aligned} \mathbb{E}[(S_M g(X_i^{(1)}, X_j^{(1)}, Y^{(1)}))^2] &\leq \|f\|_\infty \|S_M g\|_2^2 \leq \|f\|_\infty \|g\|_2^2 \leq 4\Delta_{x_i x_j}^2 \|f\|_\infty \|\eta\|_\infty^2 \\ \mathbb{E}[(S_M h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}))^2] &\leq \|f\|_\infty \|S_M h\|_2^2 \leq \|f\|_\infty \|h\|_2^2 \leq 4\Delta_{x_i x_j}^2 \|f\|_\infty \|\eta\|_\infty^2. \end{aligned}$$

Finally we get,

$$\text{Var}(P_n L) \leq \frac{12}{n} \|\eta\|_\infty^2 \|f\|_\infty^2 \Delta_{x_i x_j}^2.$$

□

Lemma 4 (Computation of $\text{Cov}(U_n K, P_n L)$). *Under the assumptions of Theorem 3, we have*

$$\text{Cov}(U_n K, P_n L) = 0.$$

Proof of Lemma 4. Since $U_n K$ and $P_n L$ are centered, we have

$$\begin{aligned}
& \text{Cov}(U_n K, P_n L) \\
&= \mathbb{E}[U_n K P_n L] \\
&= \mathbb{E}\left[\frac{1}{n^2(n-1)} \sum_{k \neq k'=1}^n K(X_i^{(k)}, X_j^{(k)}, Y^{(k)}, X_i^{(k')}, X_j^{(k')}, Y^{(k')}) \sum_{k=1}^n L(X_i^{(k)}, X_j^{(k)}, Y^{(k)})\right] \\
&= \frac{1}{n} \mathbb{E}[K(X_i^{(1)}, X_j^{(1)}, Y^{(1)}, X_i^{(2)}, X_j^{(2)}, Y^{(2)}) (L(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) + L(X_i^{(2)}, X_j^{(2)}, Y^{(2)}))] \\
&= \frac{1}{n} \mathbb{E}[(Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)})R(X_i^{(2)}, X_j^{(2)}, Y^{(2)}) - Q^\top(X_i^{(1)}, X_j^{(1)}, Y^{(1)})CQ(X_i^{(2)}, X_j^{(2)}, Y^{(2)})) \\
&\quad (A^\top R(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) + B^\top Q(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) - 2A^\top CQ(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) \\
&\quad + A^\top R(X_i^{(2)}, X_j^{(2)}, Y^{(2)}) + B^\top Q(X_i^{(2)}, X_j^{(2)}, Y^{(2)}) - 2A^\top CQ(X_i^{(2)}, X_j^{(2)}, Y^{(2)}))] \\
&= 0.
\end{aligned}$$

Since K , L , Q and R are centered. □

Lemma 5 (Bound of Bias $(\hat{\theta}_n)$). *Under the assumptions of Theorem 3, we have*

$$|\text{Bias}(\hat{\theta}_n)| \leq \Delta_{x_i x_j} \|\eta\|_\infty \sup_{l \notin M} |c_l|^2.$$

Proof.

$$\begin{aligned}
|\text{Bias}(\hat{\theta}_n)| &\leq \|\eta\|_\infty \int \left(\int |S_M f(x_{i1}, x_{j1}, y) - f(x_{i1}, x_{j1}, y)| dx_{i1} dx_{j1} \right) \\
&\quad \left(\int |S_M f(x_{i2}, x_{j2}, y) - f(x_{i2}, x_{j2}, y)| dx_{i2} dx_{j2} \right) dy \\
&= \|\eta\|_\infty \int \left(\int |S_M f(x_i, x_j, y) - f(x_i, x_j, y)| dx_i dx_j \right)^2 dy \\
&\leq \Delta_{x_i x_j} \|\eta\|_\infty \int (S_M f(x_i, x_j, y) - f(x_i, x_j, y))^2 dx_i dx_j dy \\
&= \Delta_{x_i x_j} \|\eta\|_\infty \sum_{l, l' \notin M} a_l a_{l'} \int p_l(x_i, x_j, y) p_{l'}(x_i, x_j, y) dx_i dx_j dy \\
&= \Delta_{x_i x_j} \|\eta\|_\infty \sum_{l \notin M} |a_l|^2 \leq \Delta_{x_i x_j} \|\eta\|_\infty \sup_{l \notin M} |c_l|^2.
\end{aligned}$$

We use the Hölder's inequality and the fact that $f \in \mathcal{E}$ then $\sum_{l \notin M} |a_l|^2 \leq \sup_{l \notin M} |c_l|^2$. □

Lemma 6 (Asymptotic variance of $\sqrt{n}(P_n L)$). *Under the assumptions of Theorem 3, we have*

$$n \text{Var}(P_n L) \rightarrow \Lambda(f, \eta)$$

where

$$\Lambda(f, \eta) = \int g(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy - \left(\int g(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2.$$

Proof. We proved in Lemma 3 that

$$\begin{aligned}
& \text{Var}(L(X_i^{(1)}, X_j^{(1)}, Y^{(1)})) \\
&= \text{Var}(h(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) + S_M g(X_i^{(1)}, X_j^{(1)}, Y^{(1)}) + S_M h(X_i^{(1)}, X_j^{(1)}, Y^{(1)})) \\
&= \text{Var}(A_1 + A_2 + A_3) \\
&= \sum_{k,l=1}^3 \text{Cov}(A_k, A_l).
\end{aligned}$$

We claim that $\forall k, l \in \{1, 2, 3\}^2$, we have

$$\left| \text{Cov}(A_k, A_l) - \epsilon_{kl} \left[\int g(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy - \left(\int g(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2 \right] \right| \leq \lambda [\|S_M f - f\|_2 + \|S_M g - g\|_2] \quad (27)$$

where

$$\epsilon_{kl} = \begin{cases} -1 & \text{if } k = 3 \text{ or } l = 3 \text{ and } k \neq l \\ 1 & \text{otherwise} \end{cases},$$

and where λ depends only on $\|f\|_\infty$, $\|\eta\|_\infty$ and $\Delta_{x_i x_j}$. We will do the details only for the case $k = l = 3$ since the calculations are similar for others configurations.

$$\text{Var}(A_3) = \int S_M^2 h(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy - \left(\int S_M h(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2.$$

The computation will be done in two steps. We first bound the quantity by the Cauchy-Schwartz inequality

$$\begin{aligned}
& \left| \int S_M^2 h(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy - \int g(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy \right| \\
&\leq \int |S_M^2 h(x_i, x_j, y) f(x_i, x_j, y) - S_M^2 g(x_i, x_j, y) f(x_i, x_j, y)| dx_i dx_j dy \\
&\quad + \int |S_M^2 g(x_i, x_j, y) f(x_i, x_j, y) - g(x_i, x_j, y)^2 f(x_i, x_j, y)| dx_i dx_j dy \\
&\leq \|f\|_\infty \|S_M h + S_M g\|_2 \|S_M h - S_M g\|_2 + \|f\|_\infty \|S_M g + g\|_2 \|S_M g - g\|_2.
\end{aligned}$$

Using several times the fact that since S_M is a projection, $\|S_M g\|_2 \leq \|g\|_2$, the sum is bounded by

$$\begin{aligned}
& \|f\|_\infty \|h + g\|_2 \|h - g\|_2 + 2\|f\|_\infty \|g\|_2 \|S_M g - g\|_2 \\
&\leq \|f\|_\infty (\|h\|_2 + \|g\|_2) \|h - g\|_2 + 2\|f\|_\infty \|g\|_2 \|S_M g - g\|_2.
\end{aligned}$$

We saw previously that $\|g\|_2 \leq 2\Delta_{x_i x_j} \|f\|_\infty^{1/2} \|\eta\|_\infty$ and $\|h\|_2 \leq 2\Delta_{x_i x_j} \|f\|_\infty^{1/2} \|\eta\|_\infty$. The sum is then bound by

$$4\Delta_{x_i x_j} \|f\|_\infty^{3/2} \|\eta\|_\infty \|h - g\|_2 + 4\Delta_{x_i x_j} \|f\|_\infty^{3/2} \|\eta\|_\infty \|S_M g - g\|_2.$$

We now have to deal with $\|h - g\|_2$:

$$\begin{aligned}
& \|h - g\|_2^2 \\
&= \int \left(\int (S_M f(x_{i2}, x_{j2}, y) - f(x_{i2}, x_{j2}, y)) \psi(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} \right)^2 dx_{i1} dx_{j1} dy \\
&\leq \int \left(\int (S_M f(x_{i2}, x_{j2}, y) - f(x_{i2}, x_{j2}, y))^2 dx_{i2} dx_{j2} \right) \left(\int \psi^2(x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} \right) dx_{i1} dx_{j1} dy \\
&\leq 4\Delta_{x_i x_j}^2 \|\eta\|_\infty^2 \|S_M f - f\|_2^2.
\end{aligned}$$

Finally this first part is bounded by

$$\begin{aligned}
& \left| \int S_M^2 h(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy - \int g(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy \right| \\
&\leq 4\Delta_{x_i x_j} \|f\|_\infty^{3/2} \|\eta\|_\infty \left(2\Delta_{x_i x_j} \|\eta\|_\infty \|S_M f - f\|_2 + \|S_M g - g\|_2 \right).
\end{aligned}$$

Following with the second quantity

$$\begin{aligned}
& \left| \left(\int S_M h(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2 - \left(\int g(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2 \right| \\
&= \left| \left(\int (S_M h(x_i, x_j, y) - g(x_i, x_j, y)) f(x_i, x_j, y) dx_i dx_j dy \right) \right. \\
&\quad \left. \left(\int (S_M h(x_i, x_j, y) + g(x_i, x_j, y)) f(x_i, x_j, y) dx_i dx_j dy \right) \right|.
\end{aligned}$$

By using the Cauchy-Schwartz inequality, it is bounded by

$$\begin{aligned}
& \|f\|_2 \|S_M h - g\|_2 \|f\|_2 \|S_M h + g\|_2 \\
&\leq \|f\|_2^2 (\|h\|_2 + \|g\|_2) (\|S_M h - S_M g\|_2 + \|S_M g - g\|_2) \\
&\leq 4\Delta_{x_i x_j} \|f\|_\infty^{3/2} \|\eta\|_\infty (\|h - g\|_2 + \|S_M g - g\|_2) \\
&\leq 4\Delta_{x_i x_j} \|f\|_\infty^{3/2} \|\eta\|_\infty \left(2\Delta_{x_i x_j} \|\eta\|_\infty \|S_M f - f\|_2 + \|S_M g - g\|_2 \right)
\end{aligned}$$

using the previous calculations. Collecting the two inequalities gives (27) for $k = l = 3$. Finally, since by assumption $\forall t \in \mathbb{L}^2(d\mu)$, $\|S_M t - t\|_2 \rightarrow 0$ when $n \rightarrow \infty$ a direct consequence of (27) is

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \text{Var}(L(X_i^{(1)}, X_j^{(1)}, Y^{(1)})) \\
&= \int g^2(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy - \left(\int g(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2 \\
&= \Lambda(f, \eta).
\end{aligned}$$

We conclude by noting that $\text{Var}(\sqrt{n}(P_n L)) = \text{Var}(L(X_i^{(1)}, X_j^{(1)}, Y^{(1)}))$. \square

Lemma 7 (Asymptotics for $\sqrt{n}(\hat{Q} - Q)$). *Under the assumptions of Theorem 1, we have*

$$\lim_{n \rightarrow \infty} n\mathbb{E}[\hat{Q} - Q]^2 = 0.$$

Proof. The bound given in (16) states that if $|M_n|/n \rightarrow 0$ we have

$$\begin{aligned} & \left| n\mathbb{E}[(\hat{Q} - Q)^2 | \hat{f}] - \left[\int \hat{g}(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy - \left(\int \hat{g}(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2 \right] \right| \\ & \leq \gamma(\|f\|_\infty, \|\eta\|_\infty, \Delta_{x_i x_j}) \left[\frac{|M_n|}{n} + \|S_M f - f\|_2 + \|S_M \hat{g} - \hat{g}\|_2 \right] \end{aligned}$$

where $\hat{g}(x_i, x_j, y) = \int H_3(\hat{f}, x_i, x_j, x_{i2}, x_{j2}, y) f(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2}$, where we recall that $H_3(f, x_{i1}, x_{j1}, x_{i2}, x_{j2}, y) = H_2(f, x_{i1}, x_{j2}, y) + H_2(f, x_{i2}, x_{j1}, y)$ with $H_2(\hat{f}, x_{i1}, x_{j2}, y) = \frac{1}{\int \hat{f}(x_i, x_j, y) dx_i dx_j} (x_{i1} - m_i(\hat{f}, y))(x_{j2} - m_j(\hat{f}, y))$. By deconditioning we get

$$\begin{aligned} & \left| n\mathbb{E}[(\hat{Q} - Q)^2] - \mathbb{E} \left[\int \hat{g}(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy - \left(\int \hat{g}(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2 \right] \right| \\ & \leq \gamma(\|f\|_\infty, \|\eta\|_\infty, \Delta_{x_i x_j}) \left[\frac{|M_n|}{n} + \|S_M f - f\|_2 + \mathbb{E}[\|S_M \hat{g} - \hat{g}\|_2] \right] \end{aligned}$$

Note that

$$\begin{aligned} \mathbb{E}[\|S_{M_n} \hat{g} - \hat{g}\|_2] & \leq \mathbb{E}[\|S_M \hat{g} - S_M g\|_2] + \mathbb{E}[\|\hat{g} - g\|_2] + \mathbb{E}[\|S_{M_n} g - g\|_2] \\ & \leq 2\mathbb{E}[\|\hat{g} - g\|_2] + \mathbb{E}[\|S_{M_n} g - g\|_2] \end{aligned}$$

where $g(x_i, x_j, y) = \int H_3(f, x_i, x_j, x_{i2}, x_{j2}, y) f(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2}$. The second term converges to 0 since $g \in \mathbb{L}^2(dx dy dz)$ and $\forall t \in \mathbb{L}^2(dx dy dz)$, $\int (S_M t - t)^2 dx dy dz \rightarrow 0$. Moreover

$$\begin{aligned} \|\hat{g} - g\|_2^2 & = \int [\hat{g}(x_i, x_j, y) - g(x_i, x_j, y)]^2 dx_i dx_j dy \\ & = \int \left[\int \left(H_3(\hat{f}, x_i, x_j, x_{i2}, x_{j2}, y) - H_3(f, x_i, x_j, x_{i2}, x_{j2}, y) \right) f(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} \right]^2 dx_i dx_j dy \\ & \leq \int \left[\int \left(H_3(\hat{f}, x_i, x_j, x_{i2}, x_{j2}, y) - H_3(f, x_i, x_j, x_{i2}, x_{j2}, y) \right)^2 dx_{i2} dx_{j2} \right] \\ & \quad \left[\int f(x_{i2}, x_{j2}, y)^2 dx_{i2} dx_{j2} \right] dx_i dx_j dy \\ & \leq \Delta_{x_i x_j} \|f\|_\infty^2 \int \left(H_2(\hat{f}, x_i, x_j, x_{i2}, x_{j2}, y) - H_2(f, x_i, x_j, x_{i2}, x_{j2}, y) \right)^2 dx_i dx_j dx_{i2} dx_{j2} dy \\ & \leq \delta \Delta_{x_i x_j}^2 \|f\|_\infty^2 \int \left(\hat{f}(x_i, x_j, y) - f(x_i, x_j, y) \right)^2 dx_i dx_j dy \end{aligned}$$

for some constant δ that comes out of applying the mean value theorem to $H_3(\hat{f}, x_i, x_j, x_{i2}, x_{j2}, y) - H_3(f, x_i, x_j, x_{i2}, x_{j2}, y)$. The constant δ was taken under Assumptions 1-3. Since $\mathbb{E}[\|f - \hat{f}\|_2] \rightarrow 0$ then $\mathbb{E}[\|g - \hat{g}\|_2] \rightarrow 0$. Now show that the expectation of

$$\int \hat{g}(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy - \left(\int \hat{g}(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j dy \right)^2$$

converges to 0. We develop the proof for only the first term. We get

$$\begin{aligned}
& \left| \int \hat{g}(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy - \int g(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j dy \right| \\
& \leq \int |\hat{g}(x_i, x_j, y)^2 - g(x_i, x_j, y)^2| f(x_i, x_j, y) dx_i dx_j dy \\
& \leq \lambda \int (\hat{g}(x_i, x_j, y) - g(x_i, x_j, y))^2 dx_i dx_j dy \\
& = \lambda \|\hat{g} - g\|_2^2
\end{aligned}$$

for some constant λ . By taking the expectation of both sides, we see it is enough to show that $\mathbb{E}[\|\hat{g} - g\|_2^2] \rightarrow 0$. Besides, we can verify

$$\begin{aligned}
g(x_i, x_j, y) &= \int H_3(f, x_i, x_j, x_{i2}, x_{j2}, y) f(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} \\
&= \frac{2}{\int f(x_i, x_j, y) dx_i dx_j} (x_i - \hat{m}_i(y)) \\
&\quad \left(\int x_{j2} f(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} - \hat{m}_j(y) \int f(x_{i2}, x_{j2}, y) dx_{i2} dx_{j2} \right) \\
&= 0
\end{aligned}$$

which proves that the expectation of $\int \hat{g}(x_i, x_j, y)^2 f(x_i, x_j, y) dx_i dx_j$ converges to 0. Similar computations shows that the expectation of $(\int \hat{g}(x_i, x_j, y) f(x_i, x_j, y) dx_i dx_j)^2$ also converges to 0. Finally we have

$$\lim_{n \rightarrow \infty} n \mathbb{E}[\hat{Q} - Q]^2 = 0.$$

□

References

- Brillinger, D. R. (1983). A generalized linear model with " gaussian" regressor variables. *A Festschrift for Erich L. Lehmann in Honor of His Sixty-Fifth Birthday*, (pp. 97–114).
- Bura, E. & Cook, R. D. (2001). Estimating the structural dimension of regressions via parametric inverse regression. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 63(2), 393–410.
- Cook, R. D. & Ni, L. (2005). Sufficient dimension reduction via inverse regression. *Journal of the American Statistical Association*, 100(470), 410–428.
- Da Veiga, S. & Gamboa, F. (2008). Efficient estimation of nonlinear conditional functionals of a density. *Submitted in The Annals of Statistics*. 42 pages 62G20, 62G06, 62G07, 62P30.

- Duan, N. & Li, K. C. (1991). Slicing regression: a link-free regression method. *The Annals of Statistics*, (pp. 505–530).
- Ferré, L. & Yao, A. F. (2003). Functional sliced inverse regression analysis. *Statistics*, 37(6), 475–488.
- Ferré, L. & Yao, A.-F. (2005). Smoothed functional inverse regression. *Statist. Sinica*, 15(3), 665–683.
- Hardle, W. & Tsybakov, A. B. (1991). Sliced inverse regression for dimension reduction: Comment. *Journal of the American Statistical Association*, 86(414), pp. 333–335.
- Hastie, T. J. & Tibshirani, R. J. (1990). *Generalized Additive Models*. Monographs on Statistics and Applied Probability ; 43. Chapman & Hall/CRC, 1st ed. edition.
- Hsing, T. (1999). Nearest neighbor inverse regression. *The Annals of Statistics*, 27(2), 697–731.
- Ibragimov, I. A. & Khas'minskii, R. Z. (1991). Asymptotically normal families of distributions and efficient estimation. *The Annals of Statistics*, (pp. 1681–1724).
- Laurent, B. (1996). Efficient estimation of integral functionals of a density. *The Annals of Statistics*, 24(2), 659–681.
- Li, K. C. (1991a). Sliced inverse regression for dimension reduction. *Journal of the American Statistical Association*, 86(414), 316–327.
- Li, K. C. (1991b). Sliced inverse regression for dimension reduction: Rejoinder. *Journal of the American Statistical Association*, 86(414), pp. 337–342.
- Li, L. (2007). Sparse sufficient dimension reduction. *Biometrika*, 94(3), 603.
- Setodji, C. M. & Cook, R. D. (2004). K-means inverse regression. *Technometrics*, 46(4), 421–429.
- Van der Vaart, A. W. (2000). *Asymptotic Statistics*. Cambridge Series on Statistical and Probabilistic Mathematics. Cambridge University Press.
- Zhu, L. X. & Fang, K. T. (1996). Asymptotics for kernel estimate of sliced inverse regression. *The Annals of Statistics*, 24(3), 1053–1068.