



HAL
open science

Ab initio DFT - the seamless connection between WFT and DFT

Ireneusz Grabowski, So Hirata, Victor F Lotrich

► **To cite this version:**

Ireneusz Grabowski, So Hirata, Victor F Lotrich. Ab initio DFT - the seamless connection between WFT and DFT. *Molecular Physics*, 2010, 108 (21), pp.3313. 10.1080/00268976.2010.523441 . hal-00632718

HAL Id: hal-00632718

<https://hal.science/hal-00632718>

Submitted on 15 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Ab initio DFT - the seamless connection between WFT and DFT

Journal:	<i>Molecular Physics</i>
Manuscript ID:	TMPH-2010-0212.R1
Manuscript Type:	Special Issue Paper - Electrons, Molecules, Solids and Biosystems: Fifty Years of the Quantum Theory Project
Date Submitted by the Author:	14-Aug-2010
Complete List of Authors:	Grabowski, Ireneusz; Institute of Physics, Nicolaus Copernicus University Hirata, So; University of Florida, Department of Chemistry Lotrich, Victor; University of Florida, Chemistry
Keywords:	optimized effective potential method, electron correlation, density functional theory
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
source files.zip	

SCHOLARONE™
Manuscripts

1
2
3 *Molecular Physics*

4 Vol. 00, No. 00, Month 2010, 1–20
5
6
7
8
9

10 **ARTICLE**

11
12 ***Ab initio* DFT – the seamless connection between WFT and DFT**

13
14 Ireneusz Grabowski^{a*}, Victor Lotrich^b and So Hirata^b

15
16
17 ^a*Institute of Physics, Nicolaus Copernicus University, 87-100 Toruń, Poland;*

18 ^b*Quantum Theory Project, Department of Chemistry, University of Florida, Gainesville,*
19 *FL, USA*

20
21 ()
22

23
24 Orbital-dependent exchange-correlation functionals and potentials play an increasingly impor-
25 tant role in Density Functional Theory (DFT). Methods which use explicit orbital-dependent
26 functionals can be viewed as a natural extension to the standard Kohn-Sham (KS) procedure
27 in DFT, that traditionally have used functionals with explicit density-dependence but only
28 implicit orbital-dependence. *Ab initio* DFT, invented at the Quantum Theory Project, is the
29 method which could define rigorous orbital-dependent exchange-correlation functionals and
30 potentials in the context of KS DFT theory. The local and multiplicative exchange-correlation
31 potentials are derived from a general theoretical framework based on the density condition
32 in KS theory and from coupled-cluster theory and many-body perturbation theory. *Ab initio*
33 DFT guarantees to converge to the right answer in the correlation and basis set limit, just as
34 does *ab initio* Wave Function Theory (WFT) and solves in a rigorous way most of the short-
35 comings of standard density-dependent KS DFT. It is also the route toward understanding
36 the relationships between traditional *ab initio* WFT and DFT.

37 The Optimized Effective Potential "journey" on the borderline of WFT and DFT was
38 inspired and possible only because of the Quantum Theory Project where we stay as a postdocs
39 in the 1999-2001. It seems to us then we were in the right place and the right time, and
40 certainly with the right people. The QTP scientific melting pot and Sanibel's meetings gave us
41 an excellent possibility to work together, learn and hopefully solve many important scientific
42 problems.

43
44 **Keywords:** optimized effective potential method, electron correlation, density functional
45 theory, *ab initio* DFT

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
1. Introduction

The most powerful and popular contemporary many-electron theory methods used
for calculating the electronic structure of atoms, molecules, and solids, originate
from two main categories of quantum mechanical theories: the traditional *ab initio*
theories directly grounded on the concept of the wave function (Wave Function
Theories – WFT) and the density functional theory (DFT) essentially based on
the concept that the electron density is fundamental. WFT provides methods in
which we are able to control and predict the accuracy of the results. There is a the

*Corresponding author. Email: ig@physics.ufl.edu or <http://mc.manuscriptcentral.com/tandf/tmph>

August 14, 2010 17:48 Molecular Physics sanib2010

2

Ireneusz Grabowski, Victor Lotrich and So Hirata

well known hierarchy of converging, size extensive methods MBPT(2) < CCD < CCSD < CCSD(T) < CCSDT < ... < FCC. = FCI establishes the standards in *ab initio* WFT methods. MBPT(2) means second order many-body perturbation theory, CC - coupled cluster methods with S- single, D-double, T-triple excitations. Taking into account n -fold excitations for n electrons defines the Full CC method which is equal to the FCI (CI -configuration interaction) method and is the best possible solution for the given basis set used in the calculations. Unfortunately as the inclusion of electron correlation effects is systematically improved the WFT methods become extremely expensive for larger systems, so, we still are limited to the systems with only few dozen of atoms. The computationally much cheaper DFT methods are the most widely used electronic structure methods and are routinely used for calculating ground state and dynamical properties of many-electron systems. Currently DFT seems to be the only practical method which can perform calculations on systems of hundreds and thousands of electrons. The huge reduction of the computational cost is due to the single-particle effective Hamiltonian in the Kohn-Sham [1, 2](KS) implementation of DFT, compared to the two-particle Hamiltonian in most WFT methods. In this method all unknown information is contained in the exchange-correlation (xc) functional E_{xc} , which must be approximated in any practical implementation. Unfortunately DFT establishes the existence of E_{xc} but does not provide the energy functional or even a prescription how to approximate it. Existing standard density-dependent approximations to the xc functionals, e.g. local-density LDA [2], generalized gradient approximations GGA [3–5], or hybrid functionals [6], despite their great popularity and success, suffer for many fundamental problems like lack of strict cancellation of self-interaction energies between Coulomb and exchange energies, incorrect asymptotic behaviour of the KS xc-potentials [7], missing the derivative discontinuity of the energy, qualitatively incorrect correlation potentials [8], and lack of the possibility of systematic improvement of existing functionals. KS-DFT cannot describe van der Waals interactions, there are problems with computation of response properties of molecular

systems (polarization, excitation, etc.) , also activation barrier heights for a variety of chemical reactions are predicted incorrectly [9].

On one side we have excellent but unfortunately very expensive methods (WFT) which cannot routinely be applied to very large systems. On the other side KS DFT methods can be, but we have little confidence in the reliability of the results. The main problems are not fundamental rather "technical" problems - approximations of the xc functionals and potentials in KS DFT. Since the beginning of KS DFT an enormous amount of effort has been invested in the search for realistic representations of the unknown exchange and correlation functionals (see e.g. the discussion in Ref. [10]), adding new rungs to the Jacob's Ladder [11], and adding more and more empirical adjustable parameters in new xc functionals. The new xc functionals usually only correct existing ones and only partially solve some of their problems but do not really improve the reliability of the DFT methods.

So, the important questions are: can we make DFT methods as predictive as e.g. coupled cluster theory? Can we find a method capable of rigorously defining an xc functional and potential in the context of KS theory? And how can we effectively use our experience in WFT to help DFT to be more reliable?

One of the possible solutions to aforementioned problems is by exploiting orbital-dependent functionals and potentials in KS DFT. This step, from explicitly density-dependent xc functionals, to orbital-dependent ones can be viewed in some sense as analogous to the transition from the Thomas-Fermi equation to the KS equations. Therefore the KS DFT with orbital-dependent xc functionals can be called a third generation of DFT [12]. By using orbitals to define the xc orbital-dependent functionals we have much more flexibility in the construction of the functional. It presents the possibility for using, almost directly, orbital-dependent expressions for the xc energy functionals from well known and established WFT methods. This is particularly visible e.g. in the case of the exchange energy functional E_x , which is naturally formulated in terms of orbitals, and has the form of the usual

1
2 4 *Ireneusz Grabowski, Victor Lotrich and So Hirata*

3
4 Hartree-Fock (HF) exchange energy functional

$$E_x^{\text{OEP}}[\{\phi_p\}] = -\frac{1}{2} \sum_{i,j}^N (ij|ji), \quad (1)$$

5
6
7
8
9
10 where two-electron integrals are defined as:

$$(pq|rs) = \int \phi_p^*(\mathbf{r}) \phi_q(\mathbf{r}) \frac{1}{|\mathbf{r} - \mathbf{r}'|} \phi_r^*(\mathbf{r}') \phi_s(\mathbf{r}') d\mathbf{r} d\mathbf{r}'. \quad (2)$$

11
12
13
14
15
16
17 and $\phi_p(\mathbf{r})$ are the KS orbitals, which are the self-consistent solutions of the stan-
18
19 dard KS equation. Here and in the following we use the convention that i,j,k,l label
20
21 occupied orbitals, a,b,c,d label virtual orbitals, and p,q,r,s label either. With this
22
23 choice for the exchange functional there is no self interaction error in KS DFT,
24
25 and we have precisely defined the dominant part of the xc energy functional. Un-
26
27 fortunately the corresponding KS exchange potential, which is by definition the
28
29 functional derivative of E_x with respect to the electron density, cannot be obtained
30
31 directly as in the case of density-dependent functionals. Nonetheless the orbital-
32
33 dependent exchange-only (x-only) approximation combined with the KS principles
34
35 most naturally leads to the optimized effective potential (OEP) method of Talman
36
37 and Shadwick [13] following the original suggestion of Sharp and Horton [14]. In
38
39 the KS DFT context, OEP is a rigorous exchange-only method in DFT akin to the
40
41 HF method in *ab initio* WFT and it has been implemented using various modern
42
43 algorithms for the ground state [15–22], and even for excited states [23].

44
45 Extensions of the x-only OEP to the correlation problem are not straightforward
46
47 and have been the subject of substantial interest since the development of the x-
48
49 only OEP method [18, 24–27]. The first fully self-consistent inclusion of correlation
50
51 effects in the DFT OEP procedure used the second-order many-body perturbation
52
53 theory (MBPT(2)) energy functional (OEP2-KS method [28]), where numerical
54
55 results on a few atoms supported the methodology. Numerical results for a set of
56
57 atoms and molecules, was presented by us in 2002 [28]. The proposed correlation
58
59 potential has the correct overall shape as compared to quantum Monte-Carlo re-
60

sults, especially for He-isoelectronic series, and is much better than the potentials obtained from the standard DFT calculations, which are frequently qualitatively wrong. However, we have also observed that, for larger systems, the depth of the correlation potentials may be significantly overestimated and correlation energies too negative. We have also found much slower convergence, or even divergence, of the iterative solutions of the self-consistent KS-OEP equation for the orbitals and potential [29] of the Be atom, where the quasi-degeneracy of the $(1s)^2 (2s)^2$ and $(1s)^2 (2p)^2$ configurations is present [30]. This baffling behaviour of the OEP2-KS method caused much confusion. Misunderstandings and failures to describe and solve this problem (see Refs. [31–33]), ending with the very pessimistic, but incorrect conclusion [31] ” This failure basically excludes PT based orbital functionals for practical applications in DFT”. Since then we have found [29] that the cause of this aforementioned problem of OEP2-KS is due to the choice of the zeroth-order Hamiltonian H_0 , which, in this method, is the sum of eigenvalues of the KS equation, and which is not optimal although it is also made in Görling–Levy perturbation theory (GLPT)[25, 34], and it appears to be a natural choice from the KS DFT point of view. A solution of this and other problems has been proposed by us [29] and consists of making a different choice of the unperturbed Hamiltonian to define the second-order OEP correlation functionals and potentials. Also in that paper [29], we have rederived the correlated OEP equation from a more general perspective, starting with the density condition [35] in conjunction with the CC energy functional. Defined this way the OEP exchange-only functional and OEP2-KS, OEP2-sc, OEP-ccpt2 [36] etc., correlation functionals constitute *ab initio DFT* a hierarchy of systematic approximations [37–41] converging to the exact solution of the Schrödinger equation in the correlation and basis set limit, just as does *ab initio* WFT. Independently other interesting algorithms which include correlation effects in the OEP method were proposed [42–46] which extend the applicability, and theoretical foundations of orbital-dependent functionals and potentials in DFT.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

6 *Ireneusz Grabowski, Victor Lotrich and So Hirata*

This article is dedicated to a survey of the work done at QTP mainly on correlated OEP focusing on challenges, problems and applications. In the following we present a short theory of correlated OEP, a few important results and conclusions.

2. Orbital-dependent functionals in KS DFT, Optimized Effective Potential Method and *ab initio* DFT - theory and numerical results

2.1. Theory and implementation

In the KS implementation of DFT, the computed electron density, $\rho(\mathbf{r}) = \sum_{i=1}^N |\phi_i(\mathbf{r})|^2$, is formed from the occupied KS orbitals $\phi_i(\mathbf{r})$ which are the self-consistent solutions of the KS equation [2]:

$$\left\{ -\frac{1}{2}\nabla^2 + v_s[\rho](\mathbf{r}) \right\} \phi_p(\mathbf{r}) = \varepsilon_p \phi_p(\mathbf{r}), \quad (3)$$

with the local effective KS potential

$$v_s[\rho](\mathbf{r}) = v(\mathbf{r}) + \int d\mathbf{r}' \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + V_{xc}[\rho](\mathbf{r}), \quad (4)$$

where

$$V_{xc}[\rho](\mathbf{r}) = \frac{\delta E_{xc}[\rho]}{\delta \rho(\mathbf{r})} \quad (5)$$

is the local xc potential formally defined as the functional derivative of the xc energy with respect to the electron density $\rho(\mathbf{r})$. In the case of orbital-dependent xc functionals in KS DFT $E_{xc}[\{\phi_p\}]$ the common route to the obtain orbital-dependent form of xc OEP potentials is by using the chain rule for functional derivatives and first order perturbation theory [18, 19, 28, 35]. But this seems not to be a very effective route to obtain explicit expressions for xc potentials, especially in the case when correlation is included and causes problems with extending correlation to higher orders effects.

In the following we will utilize the ideas of Bartlett *et al.* [29], where the density condition together with the CC methodology is employed to derive orbital-dependent multiplicative exchange-correlation potentials in KS OEP, and define

correlated OEP methods in an effective way. Thus, for the sake of completeness, we only briefly recall the basic ideas and notation employed, skipping the detailed diagrammatic and algebraic derivation [29, 36, 37, 39] and focusing on the main steps and basics ideas of *ab initio* DFT.

By construction, the KS density is an exact density. This means that any corrections to the converged KS density introduced by changes in $\phi_i(\mathbf{r})$ have to vanish [29, 35, 42];

$$\rho(\mathbf{r}) = \rho^{KS}(\mathbf{r}) + \delta\rho^{KS}(\mathbf{r}), \quad (6)$$

and

$$\delta\rho^{KS}(\mathbf{r}) = 0. \quad (7)$$

We use this condition to define a local exchange-correlation potential, instead of following the alternative derivative definition of the exchange-correlation potential \widehat{V}_{xc} Eq.(5). The density correction $\delta\rho^{KS}(\mathbf{r})$ can be written using density matrix correction $\Delta\gamma_{qp}$ from the CC theory [47]

$$\delta\rho^{KS}(\mathbf{r}) = \sum_{p,q} \varphi_q(\mathbf{r}) \Delta\gamma_{qp} \varphi_p^*(\mathbf{r}), \quad (8)$$

where

$$\Delta\gamma_{qp} = \langle 0 | e^{T^\dagger} \{p^\dagger q\} e^T | 0 \rangle / \langle 0 | e^{T^\dagger} e^T | 0 \rangle = \langle 0 | [e^{T^\dagger} \{p^\dagger q\} e^T]_C | 0 \rangle, \quad (9)$$

and T is the cluster operator [47–49].

We separate the total Hamiltonian

$$H = \sum_i \widehat{h}(i) + \sum_{i<j} 1/r_{ij} \quad (10)$$

into the zeroth-order H_0 and a perturbation V

$$H = H_0 + V, \quad (11)$$

1
2 8 *Ireneusz Grabowski, Victor Lotrich and So Hirata*

3
4 where

$$5 H_0 = \sum_i [\hat{h}(i) + \hat{u}(i)], \quad (12)$$

$$6 V = \sum_{i<j} 1/r_{ij} - \sum_i \hat{u}(i), \quad (13)$$

7
8
9
10
11
12 and $\hat{u} = \hat{J} + \hat{V}_{xc}$ is the one-particle potential which we determine in our approach. In
13 the above, $\hat{h}(i)$ is the sum of the kinetic energy operator and the external potential
14 and \hat{J} is the Coulomb potential. Then we can expand the density corrections,
15 Eq.(8), from CC theory in orders of the perturbation V .
16
17
18
19

20 The density condition requirement at first order, $\delta\rho^{(1)}(\mathbf{r}) = 0$, leads to the x-only
21 OEP (OEPx) equation which defines local exchange potential \hat{V}_x [13, 29] by
22
23

$$24 \sum_{a,i} \varphi_a(\mathbf{r}_1) \varphi_i^*(\mathbf{r}_1) [\langle a | \hat{K} + \hat{V}_x | i \rangle / (\epsilon_i - \epsilon_a)] = 0, \quad (14)$$

25
26
27
28
29 and \hat{K} is the non-local Hartree-Fock (HF) exchange potential. This OEP exchange
30 potential \hat{V}_x^{OEP} corresponds to the x-only OEP functional E_x^{OEP} , which has the
31 form of usual HF exchange energy functional Eq.(1) in terms of KS orbitals . In
32 matrix form, as shown elsewhere[29, 50] the x-only OEP equation which would
33 define V_X for a particular iteration can be written,
34
35
36
37
38

$$39 \mathbf{XV}_X^{(1)} = \mathbf{Y}^{(1)} \quad (15)$$

40
41
42
43 where \mathbf{X} is the matrix (auxiliary basis) representation of the density response
44 function [19, 28]
45

46
47 To include correlation effects in *ab initio* DFT and introduce the correlation
48 potential, we have to go to higher orders in perturbation theory. At the next –
49 second-order, the density condition requires that the second-order correction to the
50 KS density must vanish: $\delta\rho^{(2)}(\mathbf{r}) = 0$. This directly results in the orbital-dependent
51 OEP equations for the second-order correlation potential - the detailed derivation
52 and the explicit form of the correlated OEP equations can be found elsewhere.
53
54
55
56
57
58
59
60

[29, 37, 39] In a compact matrix form the relevant OEP equations become

$$\mathbf{XV}_{XC}^{(1+2)} = \mathbf{Y}^{(1)} + \mathbf{Y}^{(2)} \quad (16)$$

where $\mathbf{Y}^{(2)}$ arises from the correlation functional, MBPT2, and we understand $\mathbf{V}_{XC}^{(1+2)}$ to mean the combined potential through second order, which would then be used in an iterative, self-consistent scheme.

As was indicated elsewhere [29, 36, 40], the partitioning of the total Hamiltonian plays a prime role in determining the performance of the second-order correlated OEP functional. The zeroth-order Hamiltonian can be chosen in different ways. The natural choice of H_0 in OEP DFT would seem to be a simple sum of KS one-particle Hamiltonians, $H_0^{KS} = \sum_i [h(i) + v_s(i)] = \sum_p \varepsilon_p^{KS} \{p^\dagger p\}$. This is the choice made by Görling–Levy [18], by Jiang and Engel [32], by Mori-Sanchez, *et al.* [31], and by Rohr, *et al.* [33] and also by us: Grabowski *et al.* [28] in our first attempt to solve the OEP correlation problem (OEP2-KS method). However, as indicated above, this choice is a very poor one, which causes many problems in OEP2-KS calculations, and is *not* the one used to define *ab initio* DFT. [29, 36, 37, 41, 46]

By removing the large diagonal contribution $\langle p | \hat{K} + \hat{V}_{xc} | p \rangle$ from the perturbation and include it in the new H_0 .

$$H = \underbrace{\sum_p [\varepsilon_p^{KS} - \langle p | \hat{K} + \hat{V}_{xc} | p \rangle] \{p^\dagger p\}}_{H_0 = \sum_p f_{pp} \{p^\dagger p\}} - \sum_{p \neq q} \langle p | \hat{K} + \hat{V}_{xc} | q \rangle \{p^\dagger q\} + \frac{1}{4} \sum_{p,q,r,s} \langle pq || rs \rangle \{p^\dagger q^\dagger sr\} + \langle 0 | H | 0 \rangle, \quad (17)$$

and following the idea of generalized many body perturbation theory (GMBPT) [47], to reinstate orbital invariance of the MBPT(2) energy for rotations which mix occupied or virtual orbitals among themselves, we immediately define a more optimal partitioning of the Hamiltonian [29], the semi-canonical one. In this case the semi-canonical rotation of the orbitals additionally eliminates off-diagonal f_{ij}

1
2 10 *Ireneusz Grabowski, Victor Lotrich and So Hirata*

3
4 and f_{ab} terms from the perturbation

$$5
6
7 V = \sum_{ai} f_{ai} \{a^\dagger i + i^\dagger a\} + W, \quad (18)$$

8
9
10 where f_{pq} are the usual Fock matrix elements defined in terms of KS OEP spinor-
11
12 bitals

$$13
14
15 f_{pq} = \varepsilon_p^{KS} \delta_{pq} - \langle p | \widehat{K} + \widehat{V}_{xc} | q \rangle. \quad (19)$$

16
17
18 Then H_0 becomes

$$19
20
21 H_0 = \sum_p f_{pp} \{p^\dagger p\} + \sum_{i \neq j} f_{ij} \{i^\dagger j\} + \sum_{a \neq b} f_{ab} \{a^\dagger b\}. \quad (20)$$

22
23
24 Defined in this way, the OEP2-sc method, has been found to be the most effective
25
26 and stable second-order *ab initio* DFT method [29, 37, 40, 41], in which all of the
27
28 aforementioned problems of OEP2-KS were solved. A further generalization of *ab*
29
30 *initio* DFT including higher than second-order terms was made by a change of H_0 in
31
32 the OEP2-sc case to a form that has selected double excitation contributions along
33
34 with single excitations [36, 51]. The two-particle terms are those that correspond to
35
36 operators that do not change the particle number of the diagrams that constitute
37
38 the OEP functional (CCPT2). The new H_0 which defines OEP-ccpt2 method can
39
40 be written as

$$41
42
43 H_0^{CCPT} = \sum_p f_{pp} \{p^\dagger p\} + \frac{1}{4} \sum_{a,b,c,d} \langle ab || cd \rangle \{a^\dagger b^\dagger dc\} + \frac{1}{4} \sum_{i,j,k,l} \langle ij || kl \rangle \{i^\dagger j^\dagger lk\} \quad (21)$$

$$44
45
46 + \sum_{a,b,i,j} \langle aj || bi \rangle \{a^\dagger j^\dagger ib\} + \langle \Phi_{KS} | H | \Phi_{KS} \rangle \quad (22)$$

47
48
49 The OEP equations Eq. (16) for V_{xc} functional differs only in the form of $\mathbf{Y}^{(2)}$
50
51 which arises from the correlation functional, and certainly depends on the choice
52
53 of H_0 in our derivation.

54
55 We have to stress here that it is not necessary to solve the CC equations in
56
57 our implementation of *ab initio* DFT. In our methodology, we use the CC theory

to properly define orbital-dependent exchange and correlation potentials in the context of KS DFT. All correlated OEP equations (OEP2-KS, OEP2-sc and OEP-ccpt2) for the potentials are solved fully self-consistently together with KS-DFT equations until final (usually 10^{-9}) convergence is achieved.

Our implementations of *ab initio* DFT are based on the linear combination of atomic orbital (LCAO) OEP method [19, 20], which permits OEP calculations of atoms and molecules on an equal footing to the conventional molecular orbital or DFT calculations, i.e., with conventional Gaussian-type basis sets, but without any further approximations. The finite basis set implementation of OEP involves a projection method [19, 50] for solving the required integral equation, and by construction all potentials are expanded in terms of auxiliary Gaussian functions. With these LCAO OEP methods, it is possible to extract local exchange and correlation potentials and other properties from the OEP calculations of atoms and molecules. The local exchange-correlation potentials generated from the OEP method, can be plotted and may be used in developing accurate exchange and correlation potentials and functionals which can be systematically improved. [21, 28, 29, 36].

Some computational difficulties in the LCAO OEP method have been found in routine calculations with exchange-only calculations [21, 50], and with correlation included [36, 39], were the manifestation of the well-known instability associated with the numerical solutions of Fredholm integral equation of the first kind. To obtain numerical solutions of reasonable accuracy for this class of equations, an appropriate choice of basis functions is crucial. Common numerical problems which appear are the slow convergence of the projected exchange-correlation potential with respect to the size of the expansion basis set for the orbitals. The basis set incompleteness has a much more important effect on the finite basis set OEP implementation than on other SCF procedures i.e., HF method. Certainly by making a judicious choice of the orbital basis set, we can obtain reasonably accurate exchange and correlation potentials for atoms and molecules. Unfortunately, large uncontracted basis sets must be employed for orbitals, and additionally we must

August 14, 2010 17:48 Molecular Physics sanib2010

12 *Ireneusz Grabowski, Victor Lotrich and So Hirata*

make an adequate basis set choice to include correlation effects in correlated OEP calculations.

2.2. Numerical results

We have implemented various OEP methods defined above in the ACES II package [52] developed in Bartlett's group in QTP. Many testing calculations have been performed using all variants of OEP methods showing their properties, successes and failings (see the Refs. [28, 29, 36, 37, 40, 41] and also the Conclusions).

Here we present another example, the comparison of fundamental DFT quantities - electron density and KS correlation potentials obtained in different WFT and DFT methods. This analysis helps us to get more insight into the links between the coverage of dynamic electron correlation effects defined in *ab initio* WFT, standard DFT and *ab initio* DFT.

We have performed similar kind of analysis before [53–55], and here we present only few selected results obtained for the Neon atom. The Neon atom in its ground state is a perfect subject for this kind of analysis, because it has an almost ideal closed-shell structure and its correlation effects are classified as almost entirely dynamic. This allows us to avoid confusions at time of analysis with the presence of the non-dynamical correlation effects, which is often used as an explanation of failings of some methods. All calculations were performed with the uncontracted ROOS-ATZP basis set.[56] In order to make such a comparison of describing correlation effects by a different classes of various methods we have made the calculations using the main representative members of *ab initio* WFT methods: MP2, CCSD and CCSD(T), standard DFT methods SVWN5[57, 58] from LDA class of functionals, and gradient corrected BLYP [6], and *ab initio* DFT methods OEP2-KS, OEP2-sc and OEP-ccpt2.

In Fig.1. we compare the KS correlation potentials of the Neon atom calculated by standard density-dependent DFT methods, *ab initio* DFT ones and from MP2, CCSD and CCSD(T) *ab initio* WFT methods with the exact correlation KS poten-

tial [8]. To calculate the xc potentials from the MP2, CCSD and CCSD(T) densities, we employ the direct optimization technique of Wu and Yang [44]. Details of this procedure adopted to calculate xc KS potentials can be found elsewhere. [55] The resulting xc potential can be written

$$v_{xc}([\rho^A]; \mathbf{r}) = \sum_t b_t g_t(\mathbf{r}) + v_0([\rho^A]; \mathbf{r}) - J([\rho^A]; \mathbf{r}) \quad (23)$$

where $J = \int \frac{\rho^A(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|} d\mathbf{r}'$ is the Coulomb potential, $v_0(\mathbf{r})$ is the Fermi-Amaldi potential [59] used to ensure the correct $-1/r$ asymptotic behaviour of the resulting xc potential. The first term is an expansion in an auxiliary basis set of Gaussian functions $g(\mathbf{r})$ and the coefficients $\{b_t\}$ are the only unknown parameters and are determined in our procedure [44, 55]. In the following we perform calculations of v_{xc} for a variety of *ab initio* WFT densities ρ^A , where A = HF, MP2, CCSD or CCSD(T). In order to determine an approximate correlation potential, we calculate it by simply taking difference of the xc potential calculated in MP2, CCSD or CCSD(T) and x-only potential obtained in the same procedure from HF density, e.g. for the MP2 we have

$$v_c([\rho^{\text{MP2}}]; \mathbf{r}) = v_{xc}([\rho^{\text{MP2}}]; \mathbf{r}) - v_{xc}([\rho^{\text{HF}}]; \mathbf{r}) \quad (24)$$

Calculated this way correlation potentials are plotted together with the exact one and other KS potentials. As we can see in the Fig.1., the *ab initio* WFT potentials are very close to the exact one and can be used as a reference correlation potentials in a comparison with DFT results. The standard density-dependent correlation potentials VWN and LYP are qualitatively incorrect, they have the wrong shape, magnitude and even the opposite sign to the exact results almost everywhere in the space. This fact is well known [8, 28], but for some reason ignored by the DFT community. On the other side, the orbital-dependent correlation potentials calculated from *initio* DFT methods are very close to the exact results. Only the OEP2-KS correlation potential, as we indicated before, significantly overestimates

1
2 14 *Ireneusz Grabowski, Victor Lotrich and So Hirata*

3
4 the exact correlation potential, which is the result of the wrong choice of the H_0
5
6 in the definition of perturbation problem. But as we have shown elsewhere [29, 36,
7
8 37, 40] it can be fixed by proper splitting of the total Hamiltonian. The improved
9
10 OEP2-sc and OEP-ccpt2 are very close not only to the exact correlation potential,
11
12 but they are also close to their *ab initio* WFT counterparts - correlation potentials
13
14 calculated from MP2 and CCSD densities respectively. This shows the systematic
15
16 improvement of the *ab initio* DFT methods mentioned above, which is reflected
17
18 not only by e.g. the correlation and total energies but also by the KS correlation
19
20 potentials, which are the very subtle correlation effect in the total xc potential.

21 In the following we include in our analysis of the behaviour of the correlated OEP
22
23 methods also the electron density. The studies of the impact of electron correlation
24
25 effects on the electron density are based on an analysis of graphical representations
26
27 of difference radial-density distributions [53, 54] calculated between radial densities
28
29 $D(r)$ obtained from correlated method $D^{XC}(r)$ and x-only method $D^X(r)$,

$$\delta^{XC/X}(r) = D^{XC}(r) - D^X(r), \quad (25)$$

30
31
32
33
34 where the radial charge densities are defined as

$$D^A(r) = 4\pi r^2 \rho^A(r), \quad (26)$$

35
36
37
38
39
40
41 and $\rho^A(r)$ denotes the electron density at the distance r from the nucleus calcu-
42
43 lated by means of the method A. In the Fig. 2 we present a comparison of the plots
44
45 which represent the response of the density on a correlation effects. We plot curves
46
47 representing *ab initio* WFT difference radial density distributions: $\delta^{MP2/HF}(r)$
48
49 $\delta^{CCSD/HF}(r)$ and $\delta^{CCSD(T)/HF}(r)$, together with the curves which represent im-
50
51 pact on the density $\delta^{XC/X}(r)$ of the correlation functionals C (C=VWN5,LYP,
52
53 OEP2-KS, OEP2-sc, OEP-ccpt2) on the exchange functionals X (X=S, B88, and
54
55 OEPX), all generated for the Ne atom. The curves $\delta^{MP2/HF}(r)$, $\delta^{CCSD/HF}(r)$ and
56
57 $\delta^{CCSD(T)/HF}(r)$ represent *ab initio* WFT correlation effects on the density, and
58
59
60

are our reference response density distribution curves. Going to the standard DFT results, we plot the response curves for $\delta^{SVWN5/S}(r)$, $\delta^{LYP/B88}(r)$, representing the impact of inclusion of the correlation functionals on the density distributions obtained at the X-only level. One can see in this figure that the plot obtained for the VWN5 and the LYP correlation functionals, do not show similarity with the $\delta^{MP2/HF}(r)$ and $\delta^{CCSD/HF}(r)$ ones. This disagreement may be explained by the fact that the LYP and VWN5 correlation functionals do not represent any considerable dynamic correlation effects.[53] A completely different situation takes place for the curves corresponding to the OEP2-KS, OEP2-sc and OEP2-ccpt2 functionals. One can see in this figure that the plots obtained for the orbital-dependent OEP correlation functional very well, except OEP2-KS, resemble the MP2 and CCSD distributions. The OEP2-KS correlation response density curve significantly overestimates the reference WFT results, but still have qualitative good shape. It confirms the previous findings on the energy and correlation potential level and additionally supports the previous conclusions about problems with OEP2-KS method.[29, 36] Using of the OEP2-sc and OEP-ccpt2 methods significantly improves the results. OEP2-sc and OEP-ccpt2 curves are very close to the *ab initio* WFT reference ones. The shape and magnitude are almost the same, and additionally, like in correlation potential case (see. Fig. 1.), the OEP2-sc curve is closer to the MP2 one, and curve generated from OEP-ccpt2 method approaches the CCSD result. This is very systematic behaviour of the correlated OEP results, visible on the density level, which confirms findings and conclusions from the energy- and correlation KS potential-analysis. Improvement of the correlation KS-OEP functional, not only improves correlation energies, KS-potentials and other properties, but is also reflected in a systematic way on the electron density.

3. Conclusions

The use of systematically improvable, orbital-dependent approximations to the exact energy in DFT allows us to circumvent virtually all of the aforementioned shortcomings of the density-dependent functionals in KS DFT. There is no problem with the self interaction error, correlation potentials calculated in *ab initio* DFT provide the correct shape compared to those from reference quantum Monte Carlo calculations [28, 29, 36]. Exchange-only and exchange-correlation potentials are reasonable, and have the correct long-range behavior [21, 28]. As a consequence, other calculated quantities are also reasonable, e.g. total and correlation energies calculated in correlated OEP methods are almost of CC accuracy [29, 36]. Moreover *ab initio* DFT provides good ionization potentials and excitations energies [23, 37, 38]. We also have shown that the OEP2 method is capable of consistently recovering dispersion effects in weakly bounded dimers [41], which extends the applicability of *ab initio* DFT to areas which were inaccessible by standard DFT methods.

The different correlated OEP methods could be easily derived thanks to the *ab initio* DFT formalism of Bartlett *et al.*[29] based on the density condition and the CC functional. Our general approach also has important technical features: It eliminates the necessity of directly taking the cumbersome functional derivatives to arrive at the working equations to be implemented, giving us greater flexibility in defining new correlation potentials, even at infinite order. Indeed, it had been considered impossible to improve the approximations to exchange and/or correlation functionals in a systematic fashion until the concept and practical realisation of *ab initio* DFT was put forward capitalizing on the systematic orbital-dependent approximations to the exact energy in *ab initio* wave function theory (WFT). So, we are able to define a hierarchy of DFT methods (with orbital-dependent functionals) $\text{OEP}_x < \text{OEP2-KS} < \dots < \text{OEP2-sc} < \text{OEP-ccpt2} < \dots$ which converge to the right answer in the infinite and complete basis set limit as the correlation level is increased. This systematic improvement of the *ab initio* DFT methods is

reflected not only in total and correlation energies, ionization potentials, and excitation energies, but also in fundamental DFT quantities like electron density or correlation potentials.[53, 55]

The derivation of *ab initio* DFT from CC theory and the KS density condition along with the numerical results, directly compared with their WFT counterparts (MP2 – OEP2-sc, CCSD – OEP-ccpt2) shows very systematic behaviour and clearly indicates the interconnections between these two independent types of theories DFT and WFT.

But certainly, "nothing is for free". We have to mention some negative aspects of using orbital-dependent functionals in DFT. First of all, xc functionals and potentials are orbital-dependent, so in the KS procedure we have to solve additional OEP equations, which are numerically demanding and in the LCAO-OEP implementation large uncontracted basis sets must be used in calculations to provide high quality of the solutions. Additionally our equations for correlated OEP are complicated (even if the *ab initio* DFT procedure helps significantly in their derivation) and numerically more demanding the standard density-dependent DFT - they scale as $N_{it}n_{occ}^2n_{virt}^3$ (at the OEP2 level). Also, the extension of our procedure to TD-DFT is possible but is extremely complicated especially at the correlated level [60].

But the price seems to be acceptable, because right now, even in DFT calculations, we are able to control and predict the accuracy level of the results, like in *ab initio* WFT, which makes DFT method much more reliable than before and brings a hope to the DFT world that there is a way to do DFT correctly, not only theoretically but also in practice.

3.1. Acknowledgments

This work was supported by the Polish Committee for Scientific Research MNiSW under Grant no. N N204 560839. S. Hirata was supported by the U.S. National Science Foundation (CHE-0844448), the U.S. Department of Energy (DE-FG02-

04ER15621), and the Donors of the American Chemical Society Petroleum Research Fund (48440-AC6). S. Hirata is a Camille Dreyfus Teacher-Scholar. We appreciate many informative discussions with Dr. Ajith Perera and Dr. Igor Schweigert. And we would like to thank Rod Bartlett for his ideas, stimulation, "black board discussions", and for giving us opportunity to work together in Quantum Theory Project.

References

- [1] P. Hohenberg and W. Kohn, *Phys. Rev.* **136**, B864 (1964).
- [2] W. Kohn and L.J. Sham, *Phys. Rev.* **140**, A1133–A1138 (1965).
- [3] A. Becke, *Phys. Rev. A* **38**, 3098 (1988).
- [4] C. Lee, W. Yang and R.G. Parr, *Phys. Rev. B* **37**, 785 (1988).
- [5] J.P. Perdew, K. Burke and M. Ernzerhof, *Phys. Rev. Lett.* **77**, 865 (1996).
- [6] A. Becke, *J. Chem. Phys.* **98**, 5648 (1993).
- [7] E.J. Baerends and O.V. Gritsenko, *J. Phys. Chem. A* **101**, 5383 (1997).
- [8] C.J. Umrigar and X. Gonze, *Phys. Rev. A* **50**, 3827 (1994).
- [9] K.D. Dobbs and D.A. Dixon, *J. Phys. Chem.* **98**, 12584 (1994).
- [10] J.P. Perdew, A.R.J. Tao, V.N. Staroverov, G.E. Scuseria and G.I. Csonka, Prescription for the design and selection of density functional approximations: More constraint satisfaction with fewer fits. *J. Chem. Phys.* **123**, 062201–9 (2005).
- [11] J. Perdew and K. Schmidt, Jacob's Ladder of Density Functional Approximations for the Exchange-Correlation Energy, in *Density Functional Theory and Its Applications to Materials*, edited by V.E. Van Doren, K. Van Alsenoy and P. Geerlings (, , 2001), pp. 1–20.
- [12] E. Engel, Orbital-dependent functional for the Exchange-Correlation Energy: A Third Generation of DFT, in *A Primer in Density Functional Theory*, edited by Carlos Fiolhais, Fernando Nogueira and Miguel A.L. Marques (, , 2003).
- [13] J.D. Talman and W.F. Shadwick, *Phys. Rev. A* **14**, 36 (1976).
- [14] R.T. Sharp and G.K. Horton, *Phys. Rev.* **90**, 317 (1953).
- [15] J.B. Krieger, Y. Li and G.J. Iafrate, *Phys. Lett. A* **146**, 256 (1990).
- [16] J.B. Krieger, Y. Li and G.J. Iafrate, *Phys. Rev. A* **45**, 101 (1992).
- [17] E. Engel and S.H. Vosko, *Phys. Rev. A* **47**, 2800 (1993).
- [18] A. Görling and M. Levy, *Phys. Rev. A* **50**, 196 (1994).
- [19] S. Ivanov, S. Hirata and R.J. Bartlett, *Phys. Rev. Lett.* **83**, 5455 (1999).
- [20] A. Görling, *Phys. Rev. Lett.* **83**, 5459 (1999).
- [21] S. Hirata, S. Ivanov, I. Grabowski, R.J. Bartlett, K. Burke and J.D. Talman, *J. Chem. Phys.* **115**, 1635 (2001).
- [22] S. Kümmel and J.P. Perdew, *Phys. Rev. B* **68**, 035103 (2003).
- [23] S. Hirata, S. Ivanov, I. Grabowski and R.J. Bartlett, *J. Chem. Phys.* **116**, 6468 (2002).
- [24] T. Grabo and E.K.U. Gross, *Int. J. Quantum Chem.* **64**, 95 (1997).
- [25] A. Görling and M. Levy, *Phys. Rev. B* **47**, 13105 (1993).
- [26] S. Ivanov, S. Hirata and R.J. Bartlett, *Chem. Phys. Lett.* **308**, 449 (1999).
- [27] A.F. Bonetti, E. Engel, R.H. Schmidt and E.M. Dreizler, *Phys. Rev. Lett.* **86**, 2241 (2001).
- [28] I. Grabowski, S. Hirata, S. Ivanov and R.J. Bartlett, *J. Chem. Phys.* **116**, 4415 (2002).
- [29] R.J. Bartlett, I. Grabowski, S. Hirata and S. Ivanov, *J. Chem. Phys.* **122**, 034104 (2005).
- [30] B.G. Adams and K. Jankowski, *Int. J. Quantum. Chem.* **17**, 297 (1983).
- [31] P. Mori-Sánchez, Q. Wu and W. Yang, *J. Chem. Phys.* **123**, 062204 (2005).
- [32] H. Jiang and E. Engel, *J. Chem. Phys.* **123**, 224102 (2005).
- [33] D. Rohr, O. Gritsenko and E.J. Baerends, *Chem. Phys. Lett.* **432**, 336 (2006).
- [34] S. Ivanov, S. Hirata, I. Grabowski and R.J. Bartlett, *J. Chem. Phys.* **118**, 461 (2003).
- [35] A. Görling and M. Levy, *Int. J. Quantum Chem. Symp.* **29**, 93 (1995).
- [36] I. Grabowski, V. Lotrich and R.J. Bartlett, *J. Chem. Phys.* **127**, 154111 (2007).
- [37] R.J. Bartlett, V. Lotrich and I. Schweigert, *J. Chem. Phys.* **123**, 62205 (2005).
- [38] V.F.L. I. V. Schweigert and R.J. Bartlett, *J. Chem. Phys.* **125**, 104108 (2006).
- [39] I. Grabowski and V. Lotrich, *Mol. Phys.* **103**, 2087 (2005).
- [40] I. Grabowski, *Int. J. Quantum Chem.* **108**, 2076 (2008).
- [41] V. Lotrich, R.J. Bartlett and I. Grabowski, *Chem. Phys. Lett.* **405**, 33 (2005).
- [42] S. Ivanov and M. Levy, *J. Chem. Phys.* **116**, 6924 (2002).
- [43] Y.M. Niquet, M. Fuchs and X. Gonze, *J. Phys. Chem.* **118**, 9504 (2003).

- [44] Q. Wu and W. Yang, *J. Chem. Phys.* **118**, 2498 (2003).
- [45] A. Hesselmann, *J. Chem. Phys.* **122**, 224108 (2005).
- [46] D. Bokhan and R.J. Bartlett, *Chem. Phys. Lett.* **427**, 466 (2006).
- [47] R.J. Bartlett, Coupled-cluster Theory: An Overview of Recent Developments, in *Modern Electronic Structure Theory, Part II*, edited by David R. Yarkony (, , 1995), pp. 1047–1131.
- [48] R.J. Bartlett, Coupled-Cluster Approach to Molecular Structure and Spectra: A Step Toward Predictive Quantum Chemistry *J. Phys. Chem.* **93**, 1697–1708 (1989).
- [49] J. Paldus and J. Čížek, Coupled-cluster Theory: An Overview of Recent Developments, in *Energy Structure and Reactivity*, edited by D. Smith and W. B. McRae (, , 1973), pp. 198–212.
- [50] S. Ivanov, S. Hirata and R.J. Bartlett, *J. Chem. Phys.* **116**, 1269 (2002).
- [51] V. Lotrich, M. Musiał and R.J. Bartlett, To be published. (????).
- [52] J.S. Stanton, J. Gauss, J.D. Watts, M. Nooijen, N.O. andn S. A. Perera, P.G. Szalay, W.J. Lauderdale, S.A. Kucharski, S.R. Gwaltney, S. Beck, A. Balkov, D.E. Bernholdt, K.K. Baeck, P. Rozyczko, H. Sekino, C. Hober and R.J. Bartlett, ACES 2 (Quantum Theory Project, University of Florida); VMOL (J. Almlöf and P. R. Taylor); VPROPS (P. Taylor); ABACUS; (T. Helgaker, H. J. Aa. Jensen, P. Jorgensen, J. Olsen, and P. R. Taylor) .
- [53] K. Jankowski, K. Nowakowski, I. Grabowski and J. Wasilewski, *J. Chem. Phys.* **130**, 164102 (2009).
- [54] K. Jankowski, K. Nowakowski, I. Grabowski and J. Wasilewski, *Theoret. Chem. Acc.* **125**, 433 (2010).
- [55] I. Grabowski, A. Teale, S. Śmiga, K. Jankowski and R.J. Bartlett, to be published (2010).
- [56] P.O. Widmark, P.A. Malmqvist and B. Roos, *Theor. Chim. Acta* **77**, 291 (1990).
- [57] S.H. Vosko, L. Wilk and M. Nusair, *Can. J. Phys.* **55**, 1200–XXX (1980).
- [58] J.C. Slater, *Quantum Theory of Molecules and Solids*, Vol. 4: The Self-Consistent Field for Molecules and Solids (McGraw-Hill, New York, 1974).
- [59] E. Fermi and E. Amaldi, *Mem. Accad. Italia* **6**, 117 (1934).
- [60] D. Bokhan and R.J. Bartlett, *Phys. Rev. A* **73**, 022502 (2006).

Figure captions

Figure 1. Comparison of correlation potentials of the Neon atom obtained from the OEP2-KS, OEP2-sc, OEP-ccpt2 with the correlation potentials obtained from MP2, CCSD and CCSD(T), and from standard DFT: VWN5 and LYP and the exact correlation potential Ref. ([8]).

Figure 2. Comparison of difference radial density distributions, $\delta^{MP2/HF}(r)$, $\delta^{CCSD/HF}(r)$ and $\delta^{CCSD(T)/HF}(r)$ with the impact on the density $\delta^{XC/X}(r)$ of the correlation functionals C (C=VWN5, LYP, OEP2-KS, OEP2-sc, OEP-ccpt2) on the exchange functionals X (X=S, B88, and OEPX) generated for the Ne atom.




Figure 1. Grabowski, Lotrich, Hirata




Figure 2. Grabowski, Lotrich, Hirata