



HAL
open science

Specific targeting of the GABA-A receptor $\alpha 5$ subtype by a selective inverse agonist restores cognitive deficits in Down syndrome mice.

Jérôme Braudeau, B. Delatour, Arnaud Duchon, Philippe Lopes Pereira, Luce Dauphinot, Fabrice de Chaumont, Jean-Christophe Olivo-Marin, Robert H. Dodd, Y. Hérault, Marie-Claude Potier

► To cite this version:

Jérôme Braudeau, B. Delatour, Arnaud Duchon, Philippe Lopes Pereira, Luce Dauphinot, et al.. Specific targeting of the GABA-A receptor $\alpha 5$ subtype by a selective inverse agonist restores cognitive deficits in Down syndrome mice.. *Journal of Psychopharmacology*, 2011, 25 (8), pp.1030-42. 10.1177/02698811111405366 . hal-00634143

HAL Id: hal-00634143

<https://hal.science/hal-00634143v1>

Submitted on 25 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Specific targeting of the GABA-A receptor $\alpha 5$ subtype by a selective inverse agonist restores cognitive deficits in Down syndrome mice

J Braudeau^{1,2,*}, B Delatour^{1,2,*}, A Duchon^{3,4}, P Lopes Pereira³,
 L Dauphinot¹, F de Chaumont⁵, J-C Olivo-Marin⁵, RH Dodd⁶,
 Y Hérault^{3,4} and M-C Potier¹

Abstract

An imbalance between inhibitory and excitatory neurotransmission has been proposed to contribute to altered brain function in individuals with Down syndrome (DS). Gamma-aminobutyric acid (GABA) is the major inhibitory neurotransmitter in the central nervous system and accordingly treatment with GABA-A antagonists can efficiently restore cognitive functions of Ts65Dn mice, a genetic model for DS. However, GABA-A antagonists are also convulsant which preclude their use for therapeutic intervention in DS individuals. Here, we have evaluated safer strategies to release GABAergic inhibition using a GABA-A-benzodiazepine receptor inverse agonist selective for the $\alpha 5$ -subtype ($\alpha 5$ IA). We demonstrate that $\alpha 5$ IA restores learning and memory functions of Ts65Dn mice in the novel-object recognition and in the Morris water maze tasks. Furthermore, we show that following behavioural stimulation, $\alpha 5$ IA enhances learning-evoked immediate early gene products in specific brain regions involved in cognition. Importantly, acute and chronic treatments with $\alpha 5$ IA do not induce any convulsant or anxiogenic effects that are associated with GABA-A antagonists or non-selective inverse agonists of the GABA-A-benzodiazepine receptors. Finally, chronic treatment with $\alpha 5$ IA did not induce histological alterations in the brain, liver and kidney of mice. Our results suggest that non-convulsant $\alpha 5$ -selective GABA-A inverse agonists could improve learning and memory deficits in DS individuals.

Keywords

Down syndrome, GABA-A, inverse agonist, learning, memory, therapy

Introduction

Down syndrome (DS) is the consequence of trisomy 21, the most common genetic cause of mental retardation (1/800 live births), and is characterized by varying degrees of cognitive impairments (Sherman et al., 2007). Advances in teaching methods and educational mainstreaming have proven to be beneficial to people with DS, but are clearly not sufficient to counteract all cognitive deficits (Wishart et al., 2007). Since these individuals now have a life expectancy of 55 years and often survive their parents, treatments aimed at enhancing cognitive skills to provide higher autonomy are long-awaited. Unfortunately, attempts with off-label use of various drugs have not been successful (Reeves and Garner, 2007; Wiseman et al., 2009).

Recent data strongly suggest that changes associated with learning and memory dysfunction in DS might result, in part, from defects in the hippocampus associated with increased inhibition (GABAergic activity) in the brain, opening new avenues for pharmacological intervention (Best et al., 2007; Kleschevnikov et al., 2004). As a consequence, treatment of DS mouse models with non-competitive GABA-A

¹Centre de Recherche de l'Institut du Cerveau et de Moelle Epinière, CNRS UMR7225, INSERM UMR5 975, UPMC, Paris, France.

²CNRS, Lab NAMC, UMR8620, Université Paris Sud, Orsay, France.

³Institut de Génétique et de Biologie Moléculaire et Cellulaire (IGBMC), Institut National de Santé et de Recherche Médicale (INSERM) U964/ Centre National de Recherche Scientifique (CNRS) UMR 1704/Université de Strasbourg, 67404 Illkirch, France.

⁴Institut Clinique de la Souris (ICS), Groupe d'Intérêt Economique Centre Européen de Recherche en Biologie et en Médecine (GIE-CERBM), INSERM, CNRS, Université de Strasbourg, Illkirch, France.

⁵Institut Pasteur, Quantitative Image Analysis Unit, CNRS URA 2582, 75015 Paris, France.

⁶Institut de Chimie des Substances Naturelles - CNRS UPR 2301, Gif-sur-Yvette, France.

*First co-authors.

Corresponding author:

M-C Potier, Centre de Recherche de l'Institut du Cerveau et de Moelle Epinière, CNRS UMR7225, INSERM UMR5 975, UPMC, 75013 Paris, France
 Email: marie-claude.potier@upmc.fr

antagonists, such as picrotoxin or pentylenetetrazol, can restore impaired phenotypes in DS mice (Fernandez et al., 2007; Rueda et al., 2008). However, these drugs are convulsant at high doses, precluding their use as cognition enhancers in humans, particularly considering that DS patients are more prone to convulsions (Menendez, 2005). Frequency of seizures has been reported to reach 6–17% in DS people (Veall, 1974) with a triphasic distribution of seizure onset depending on age (infancy, early adulthood and late onset) (Pueschel et al., 1991).

As an alternative to GABA-A antagonists, we searched among ligands of the GABA-A-benzodiazepine receptors that could decrease GABAergic transmission without inducing convulsant activity. This selective pharmacological profile can be obtained using molecules that are active at the $\alpha 5$ subunit-containing GABA-A-benzodiazepine receptors (Sur et al., 1999). These receptors are largely expressed in the hippocampus (Wisden et al., 1992), an area integral to learning and memory. Molecules that specifically decrease GABAergic transmission through these receptors, such as $\alpha 5$ -selective inverse agonists, have been shown to enhance cognition and synaptic plasticity without having any adverse convulsant/pro-convulsant or anxiogenic effects (Ballard et al., 2009; Collinson et al., 2006; Dawson et al., 2006). To the best of the authors' knowledge, these compounds have not yet been evaluated for the treatment of cognitive impairments associated with brain dysfunction.

The goal of the present work was to assess the therapeutic potential of an $\alpha 5$ -selective inverse agonist, the orally active 3-(5-methylisoxazol-3-yl)-6-[(1-methyl-1, 2, 3-triazol-4-yl)methoxy]-1,2,4-triazolo[3,4-a]phthalazine (Sternfeld et al., 2004), referred to herein as compound $\alpha 5IA$, in cognitively impaired mouse models of DS. We used Ts65Dn mice, which are trisomic for orthologues of about half of the genes on human chromosome 21 (Reeves et al., 1995). These mice demonstrate learning and memory defects, as well as synaptic plasticity abnormalities and are widely used for preclinical research on DS (Escorihuela et al., 1995; Kleschevnikov et al., 2004; Reeves et al., 1995).

Materials and methods

Animals

Male mice were produced at the Intragene resource centre (TAAM, CNRS UPS44 Orléans, France) and bred on a mixed genetic background B6C3, derived from C57BL/6J (B6) and a congenic inbred line C3H/HeH for the BALB/c wild-type *Pde6b* allele (Hoelter et al., 2008), thus avoiding retinal degeneration and impaired visual acuity. On this background, Ts65Dn mice show similar behavioural phenotypes when compared with the original Ts65Dn line (AD and YH, personal communication; see also Costa et al., 2010). Mice were acclimated in our animal facility for at least 2 weeks before initiating behavioural testing. For each experiment, different batches of mice (3 months old) were used (total number of animals used: Ts65Dn mice, $n = 90$; euploid littermates, $n = 122$).

All experiments were conducted in accordance with the ethical standards of French and European regulations

(European Communities Council Directive of 24 November 1986). The supervisor of *in vivo* studies (B Delatour) received official authorization from the French Ministry of Agriculture to carry out research and experiments on animals (authorization number 91-282).

Real-time quantitative PCR of *Gabra-5*

Total RNA was extracted from dissected hippocampi of nine euploid and seven Ts65Dn mice and treated with DNase using the Nucleospin RNA II kit (Macherey-Nagel, France). RNAs (500 ng) were individually reverse-transcribed into cDNAs overnight at 37°C using the Verso cDNA kit (ThermoFisher Scientific, Waltham, USA) according to the manufacturer's instructions. qPCR assays were performed in a Lightcycler[®] 480 System (Roche), in the presence of 200nM of each primer (*Gabra5* 5'gacgactcttgatggcta3'_forward and 5'acctgcgtgattcgtct3'_reverse; *pPib* 5'tcttcataaccacagt-caagacc3'_forward and 5'acctccgtaccacatccat3'_reverse for normalization), 100nM of specific hydrolysis probe and 1X Lightcycler[®] 480 Probes Master mix (Roche, France), and normalized using the Lightcycler[®] 480 SW 1.5 software.

$\alpha 5IA$ synthesis and formulation

The drug used was 3-(5-methylisoxazol-3-yl)-6-[(1-methyl-1,2,3-triazol-4-yl)methoxy]-1, 2, 4-triazolo[3, 4-a]phthalazine ($\alpha 5IA$). It was synthesized by Orga-Link SARL (Magny-les-Hameaux, France), according to Sternfeld et al. (2004). The hydrochloride salt was prepared by dissolving the base in hot ethanol and adding a solution of 5% hydrochloric acid in ethanol until the solution was slightly acidic. Upon cooling, a precipitate formed which was collected by filtration, washed with cold ethanol and dried.

The HCl salt of $\alpha 5IA$ was solubilized in a mixture of DMSO, Cremophor El (BASF, Ludwigshafen, Germany) and hypotonic water (ProAmp[®]) (10:15:75). $\alpha 5IA$ or vehicle (solubilization solution) was injected intraperitoneally (i.p.) at different doses ranging from 1 to 50 mg/kg.

Morris water maze

Experiments were performed in a 150-cm diameter Morris water maze filled with opacified water kept at 19°C and equipped with a 9 cm diameter platform submerged 1 cm under the water surface.

In a first pilot experiment, a total of 27 C57BL/6 mice were used to study dose–response cognitive-enhancement effects of $\alpha 5IA$ (vehicle, 1 mg/kg, 5 mg/kg, $n = 9$ in each group) in a delayed matching-to-place task (DMTP) (see Figure 1(a)) as described previously (Collinson et al., 2002).

In a second experiment, 16 Ts65Dn (vehicle $n = 8$, $\alpha 5IA$ 5 mg/kg $n = 8$) mice and 16 euploid littermates (vehicle $n = 8$, $\alpha 5IA$ 5 mg/kg $n = 8$) were trained during 6 days in the standard Morris water maze task (MWM) (see Figure 2(a)). Training consisted of daily sessions (two trials per session). Start positions varied pseudo-randomly among the four cardinal points. Mean inter-trial interval was 2 hours. During the habituation and spatial training phases each trial ended when the animal reached the platform. A 90-second cut-off




Figure 1. Dose-response effect of $\alpha 5$ IA. The optimal $\alpha 5$ IA promnesic dose was determined in euploid mice trained in the DMTP task. (a) Schematic representation of the DMTP protocol. Training was performed during 7 consecutive days. Each day, mice underwent one acquisition trial (T1) and three retention trials (T2-T4); inter-trial interval was 60 seconds. The position of the platform was changed every day, but remained constant within each session. (b) Performance (distance to platform; mean \pm SEM) of the mice between acquisition and retention trials. Data from the seven training days have been pooled. All mice showed a significant increase in behavioural accuracy within each session. While vehicle-treated mice and mice receiving 1 mg/kg of $\alpha 5$ IA showed similar retention, mice that were treated with the 5 mg/kg dose of $\alpha 5$ IA displayed a significantly higher retention performance in comparison to other groups (* $p < 0.05$, ANOVA with Fisher's *post hoc* comparisons).

was used, after which mice were manually guided to the platform. Once on the platform, animals were given a 20-second rest before being returned to their cage. Twenty four hours after the last training trial, retention was assessed during a probe trial in which the platform was no longer available. During the four subsequent sessions visual ability of mice was controlled: platform location was cued by a white styrene ball placed 12 cm above water surface and access to external indices was prevented by a black curtain surrounding the pool.

In all navigation tasks (DMTP, MWM) mice were injected daily with vehicle or $\alpha 5$ IA 30 min before each first (T1) trial of each daily session. Animals were monitored with the Any-Maze (DMTP task; Stoelting, Wood Dale, USA) or the VideoTrack (MWM task; Viewpoint, Lyon, France) video analysis systems.

Novel-object recognition

The apparatus consisted of a square open field (50 cm \times 50 cm) placed in a room with weak controlled luminosity (4-6 lux) and constant 60 dB white noise.

The first day, all animals (16 euploid and 16 Ts65Dn mice) were handled by the experimenter. On day 2, mice were habituated for 20 min to the empty arena. On day 3, four identical objects were placed symmetrically 14 cm away from the arena corners. Mice were free to explore the objects for 20 min. On the test day (day 4), mice were injected i.p. with either vehicle or $\alpha 5$ IA 5 mg/kg (eight euploid and eight Ts65Dn mice in each group). Thirty minutes after injections, mice were placed in the arena containing two identical objects, and allowed to explore them for 10 min. Mice then returned to their home cage for a 10-min retention interval. To test short-term recognition memory, one familiar object and one novel object were placed in the apparatus, and mice were free to explore for a 10-min period. Between each trial the arena and objects were cleaned with 70% ethanol to reduce olfactory cues.

During all sessions mice were monitored using the Any-Maze video-tracking software. Object exploration was manually scored with an ethological keyboard and defined as the orientation of the nose to the object at a distance < 4 cm. The amount of time exploring familiar vs. novel objects was calculated to assess memory performance.

Measure of cerebral Fos immunoreactivity

Euploid ($n = 13$) and Ts65Dn ($n = 6$) mice were pseudo-trained in the object recognition task using the same protocol as described in the novel-object recognition (NOR) task, but with no retention phase. Thirty minutes before acquisition, six euploid and three Ts65Dn mice, and seven euploid and three Ts65Dn were injected i.p. with $\alpha 5$ IA (5 mg/kg) or vehicle, respectively. Following the acquisition session, mice returned to their home cage. Ninety minutes following behavioural stimulation, mice were perfused transcardially with phosphate buffered saline (PBS), their brains fixed in 10% formalin, cryoprotected and sectioned on a freezing microtome. Fos immunoreactivity (polyclonal AB-5, Calbiochem-VWR, France; dilution 1:10,000) was detected using the ABC




Figure 2. $\alpha 5$ IA restores spatial learning in Ts65Dn mice. (a) Schematic representation of the MWM protocol (see the text for explanations; PT: probe trial). (b) Data on learning performance have been pooled into two blocks of 3 days. Vehicle-treated Ts65Dn mice demonstrated decreased learning index in comparison with the three other trained groups. This deficit was corrected by treatment with $\alpha 5$ IA. (c) A hit was defined as reaching the platform before 90 seconds. Vehicle-treated Ts65Dn mice showed a clear delay in conditioning that was rescued after treatment with $\alpha 5$ IA. (d) Only euploid mice showed a spatial bias for the platform target quadrant during probe trial. Impaired retention of the platform location in Ts65Dn mice was not rescued after drug treatment. For (b) and (d), horizontal dotted lines at 25% correspond to random performance. For (d), percentage of time spent in other quadrants ('others') calculated as the mean time spent in these three non-target quadrants. * $p < 0.05$: ANOVA with repeated measures and contrast analysis. # $p < 0.05$: paired Student's t -tests.

method with nickel-enhanced diaminobenzidine as final chromogen. Immunoreactivity was quantified using QUAA software (see <http://www.bioimageanalysis.org>) that automatically calculated the proportion of stained tissue ($p = \text{stained area}/\text{total area}$), providing unbiased stereological measurements. Four regions of interest (ROIs) were analysed: posterior cingulate cortex, perirhinal cortex, dentate gyrus and CA1 field of the hippocampus on several serial sections. Results were then averaged to give a reliable quantitative evaluation of local Fos immunostaining.

Convulsant and pro-convulsant effects

The convulsant effects were evaluated after a single i.p. injection of $\alpha 5$ IA at high dosage (50 mg/kg) or vehicle in Ts65Dn or euploid littermates. For testing the pro-convulsant effects

of $\alpha 5$ IA, a sub-convulsant dose of pentylenetetrazole (45 mg/kg i.p.) was injected i.p. 20 min after injection of $\alpha 5$ IA or vehicle. Six or seven mice were used for each condition. Mice were observed for 20 min (convulsant effects) or 30 min (pro-convulsant effects): the occurrence of tonic convulsions and latency to the first myoclonic jerk episode were recorded.

Locomotor activity

Locomotor activity was evaluated in a total of 33 mice 30 min after i.p. injections (vehicle: 8 euploid and 7 Ts65Dn mice; $\alpha 5$ IA 5 mg/kg: 10 euploid and 8 Ts65Dn mice). Locomotion was measured in a square open field (50 cm \times 50 cm; luminosity: 30 lux) with black walls 30 cm high. Each animal was allowed to freely explore the arena for 10 min. Horizontal

activity was monitored using the Any-Maze software. Time spent in the 10-cm wide peripheral zone and in the complementary 30 cm × 30 cm central zone was recorded to evaluate anxiety.

Anxiety-related behavioural testing

Modulation of anxiety-related behaviours by $\alpha 5IA$ was assessed using an elevated plus maze, in a total of 42 mice, 30 min after i.p. injections (vehicle: 11 euploid and 7 Ts65Dn mice; $\alpha 5IA$ 15 mg/kg: 14 euploid and 10 Ts65Dn mice). The maze was constructed of black Perspex (length, 28 cm; width, 5 cm; height from floor, 40 cm; overall luminosity in open arms: 70 lux) with two opposing open arms, and two enclosed arms equipped with three 16-cm high walls. Mice were placed in the central region of the maze and behaviour was recorded for a 5-min period using the Any-Maze software.

To explore the potential adversity of chronic injections of $\alpha 5IA$, another group of euploid mice was treated for 2 weeks (5 mg/kg, five injections/week; five $\alpha 5IA$ -treated mice; five vehicle-treated mice) and then evaluated in the elevated plus maze as described previously.

Anatomopathology after chronic treatment with $\alpha 5IA$

Mice treated for 2 weeks with $\alpha 5IA$ 5 mg/kg and tested in the elevated plus maze (see the previous section) were further treated for another 3 weeks. On the last day of treatment, urine samples were collected 2 hours after $\alpha 5IA$ or vehicle i.p. administration. The next day, mice were sacrificed. For anatomic-pathological examination, three additional euploid non-injected mice were also sacrificed. Liver, kidney, brain and spleen were dissected and fixed in a 10% formalin solution. Tissues were then paraffin-embedded, cut and processed for routine histopathological evaluation (haematoxylin–eosin and periodic acid–Schiff stainings).

Statistical analysis

In most cases, data were analysed using an analysis of variance (ANOVA) with Fisher's *post hoc* comparisons. ANOVA with repeated measures or within-subjects designs and contrast analysis were carried out when required by the experimental plan to assess complementary statistical effects. Also in some designs, statistical analysis was performed using Student's *t*-tests. For all analysis statistical significance was set to a *p*-value <0.05. All analyses were performed using Statistica v6 (StatSoft, Inc., Tulsa, OK, USA) or GraphPad Prism (GraphPad Software, La Jolla, CA, USA) software.

Results

$\alpha 5IA$ acts as a cognition enhancer and alleviates learning and memory deficits in Ts65Dn mice

Synthesis of the $\alpha 5IA$ and determination of the pharmacologically active dose. As a prerequisite we checked that the level of expression of the *Gabra5* gene encoding the $\alpha 5$ GABA-A subunit was unchanged in the hippocampus of Ts65Dn mice as compared with euploids ($t_{14} = 0.40$, $p = 0.69$;

data not shown) confirming the presence of the pharmacological target in Ts65Dn mice. We concurrently synthesized $\alpha 5IA$ and showed that the spectral characteristics and binding affinity of the compound conformed with published data (see Supplementary Figure S1) (Sternfeld et al., 2004). We then determined the optimal dose of $\alpha 5IA$ that induced clear promnesic effects in mice trained in the DMTP version of the MWM task (Figure 1(a)). As illustrated in Figure 1(b), a large decrease in distance travelled was observed between acquisition and retention trials underlying memory of the goal location ($F_{1,24} = 66.39$, $p < 0.0001$). The three groups (vehicle, $\alpha 5IA$ 1 mg/kg and $\alpha 5IA$ 5 mg/kg) showed similar performances during acquisition ($F < 1$) but a group effect was observed during retention trial ($F_{2,24} = 4.5$, $p < 0.05$). Indeed while the vehicle and the $\alpha 5IA$ 1 mg/kg groups demonstrated comparable retention performance ($F < 1$), mice treated with $\alpha 5IA$ 5 mg/kg displayed a clear improvement of performance (comparison with vehicle mice: $F_{1,24} = 8.5$, $p < 0.01$). We therefore selected the dose of 5 mg/kg to be used in subsequent behavioural tests in DS models.

Effects of $\alpha 5IA$ on reference memory in Ts65Dn mice using the Morris water maze task.

To evaluate the rescuing potential of $\alpha 5IA$ in behaviourally impaired Ts65Dn mice, we first assessed the effect of the drug on spatial reference memory in the standard MWM task, in which mice have to swim in their environment to locate a hidden platform at a constant location (Figures 2–4). Out of 16 Ts65Dn and 16 euploid mice, 1 Ts65Dn mouse was discarded from statistical analysis because it displayed abnormal floating behaviour and decreased swim speed in the maze. During the probe trial one euploid mouse was removed from the analysis for the same reason.

We first analysed the acquisition of place location (Figure 2(b) and 2(c)). ANOVA on swim speeds revealed an effect of group factor ($F_{3,26} = 2.99$, $p < 0.05$). Owing to variations in swim speeds between conditions that may impact non-specifically on performances, we calculated an unbiased index of spatial learning that is the percentage of the path length spent by mice in the target quadrant (Faure et al., 2009; Janus et al., 2004) (Figure 2(b)). ANOVA (main factors: group and block of sessions) on this learning index indicated significant effect of group ($F_{3,27} = 4.77$, $p < 0.01$) and block ($F_{1,27} = 16.63$, $p < 0.001$) factors with no significant interactions between these main factors ($F_{3,27} < 1$). Vehicle-treated Ts65Dn mice displayed a low learning index when compared with mice from the three other groups (all $F_{1,27} > 6.53$, $p < 0.05$). ANOVA on the percentage of trials performed within the cut-off limit (that is, percentage of hits, Figure 2(c)), a complementary measure of learning proficiency, indicated significant effect of group ($F_{3,27} = 3.44$, $p < 0.05$) and session ($F_{5,135} = 4.32$, $p < 0.002$) factors with no significant interactions between these main factors ($F_{5,135} = 0.91$, $p > 0.55$). Vehicle-treated Ts65Dn mice were once again severely impaired in terms of hits performed when compared with mice from the three other groups (percentage of hits: all $F_{1,27} > 6.10$, $p < 0.05$, Figure 2(c)). Finally, ANOVA indicated that $\alpha 5IA$ significantly potentiated the acquisition proficiency of Ts65Dn mice ($F_{1,27} > 6.10$, $p < 0.025$ for the learning index




Figure 3. $\alpha 5IA$ relieves the use of inadequate behavioural navigating strategies in the Morris water maze. A robust effect of $\alpha 5IA$ was observed on thigmotaxis (percentage of time spent performing thigmotaxis has been pooled over the six training sessions, mean \pm SEM). This inadequate strategy to locate the platform in the water maze was strongly decreased following treatment with $\alpha 5IA$ but more particularly in Ts65Dn mice and to a lesser (non-significant) extent in euploid mice, likely due to some ceiling effects as Ts65Dn mice displayed an overall increased basal level of thigmotaxis in comparison to euploid mice. * $p < 0.05$; ** $p < 0.001$; *** $p < 0.0001$; ANOVA with repeated measures and contrast analysis.

and hit measures) allowing them to regain normal levels of performance. This promnesic effect of the treatment was not observed in euploid mice that performed equally well in this test with or without $\alpha 5IA$ ($F < 1$ for the learning index and percentage of hits measures).

We then analysed navigation strategies of mice (Figure 3). Indeed, in association with an impaired learning capacity, Ts65Dn mice also displayed high levels of thigmotaxis. We measured this ‘wall-seeking behaviour’ as the time spent by mice in the 10-cm-wide peripheral annulus of the pool. As shown in Figure 3 thigmotaxis of Ts65Dn mice appeared to be strongly decreased after $\alpha 5IA$ treatment. ANOVA on the time spent performing thigmotaxis confirmed significant effects of group ($F_{3,28} = 13.70$; $p < 0.0001$) and day ($F_{5,140} = 7.48$; $p < 0.0001$), underlining that thigmotaxis decreased across training sessions (Figure 3). The group \times day interaction was non-significant ($F_{15,140} = 1.37$, $p = ns$). *Post-hoc* analysis showed that Ts65Dn mice were more thigmotactic in comparison to euploid mice in vehicle condition ($F_{1,28} = 20.75$, $p < 0.0001$). While $\alpha 5IA$ strongly reduced thigmotaxis in Ts65Dn mice ($F_{1,28} = 15.74$; $p < 0.001$) these mice still displayed increased thigmotaxis after $\alpha 5IA$ treatment (comparison with euploids: $F_{1,28} = 4.81$, $p < 0.05$). In addition, in euploid mice, the thigmotaxis-reducing effect of $\alpha 5IA$, although observed (see Figure 3), did not reach statistical significance ($F_{1,28} = 2.57$, $p = ns$).

Retention of place location was evaluated during a single probe trial (PT) (no platform available, see Figure 2(a)). Examination of each group separately




Figure 4. Spatial impairments in Ts65Dn mice are not due to visual deficits. Following evaluation of spatial memory in the Morris water maze (MWM), mice were trained in a visually guided navigation task (cued visible platform). Performance was assessed using an unbiased learning index (mean \pm SEM, same as in Figure 2). Analysis indicated that behavioural accuracy to locate the visible platform increased across sessions with no effect of genotype or treatment. The horizontal dotted line at 25% represents level of performance due to random navigation in the pool. As illustrated, all trained groups performed largely above this level.

showed that euploid mice clearly located the target quadrant as demonstrated by their biased exploration (comparison between target vs. non-target quadrants, paired *t*-test: $t_{12} > 2.91$, $p < 0.025$ for vehicle and $\alpha 5IA$ conditions; see Figure 2(d)). In contrast, Ts65Dn mice, even after $\alpha 5IA$ treatment, did not show exploratory preference for the target quadrant during probe test ($t < 1.7$, $p = ns$ for all treatment conditions), indicating that they could not efficiently remember the goal location.

Finally, although mice produced for this study carried a functional allele of Pd6b avoiding retinal degeneration (see the Materials and methods section), their visual ability was controlled using a non-spatial training procedure (Figure 4). ANOVA showed no effects of the group factor ($F_{3,28} < 1$, $p = ns$). The repetition of training trials (day factor) had a significant impact on performance ($F_{3,84} = 3.11$, $p < 0.05$) and there was no group \times day interaction ($F < 1$) thus indicating that all groups gradually increased their performance in the visual discrimination task and performed equally, whatever the genotype or treatment.

In summary it can be concluded that $\alpha 5IA$ treatment rescued the MWM spatial learning deficits present in Ts65Dn mice and mitigated their use of inadequate navigation strategies.

Effects of $\alpha 5IA$ on short-term memory in Ts65Dn mice using the novel object recognition task. We then evaluated $\alpha 5IA$ treatment effects on non-spatial memory




Figure 5. $\alpha 5$ IA alleviates recognition memory deficits in Ts65Dn mice and potentiates neuronal activity (a) Upper part: general protocol of the novel-object recognition (NOR) (see the text for explanations). Lower part: Learning index (see Table 1 for raw data). Under vehicle, Ts65Dn mice were found to be impaired. Following i.p. injection of $\alpha 5$ IA (5 mg/kg), both euploid and Ts65Dn mice improved their NOR performance and the deficit of Ts65Dn mice was abolished. * $p < 0.05$; ** $p < 0.001$; *** $p < 0.0001$; ANOVA with Fisher's *post hoc* comparisons. (b) Upper part: general protocol for assessing the levels of Fos after behavioural stimulation (see the text for explanations). Lower part: histograms depict the relative increase of Fos immunoreactivity in $\alpha 5$ IA-treated mice normalized against values obtained for vehicle-treated littermates. In all brain regions sampled, except the dentate gyrus, a significant increase of Fos was observed after $\alpha 5$ IA injection. # $p < 0.05$; ### $p < 0.001$; two way ANOVA with repeated measures and contrast analysis. No differences between genotypes were observed.

Table 1. $\alpha 5$ IA modulates the time spent by mice exploring familiar versus novel objects

Genotype	Treatment	New object mean \pm SEM	Familiar object mean \pm SEM
Euploids	Vehicle	44.91 \pm 6.43	32.43 \pm 3.47*
	$\alpha 5$ IA (50 mg/kg)	44.40 \pm 6.83	28.48 \pm 4.16***
Ts65Dn	Vehicle	44.91 \pm 5.79	48.35 \pm 5.09
	$\alpha 5$ IA (50 mg/kg)	49.28 \pm 7.27	21.05 \pm 2.16***

In contrast to Ts65Dn vehicle-treated mice, vehicle-treated euploids and $\alpha 5$ IA treated mice (Ts65Dn and euploids) discriminated between familiar and novel objects. Comparison between objects: * $p < 0.05$, *** $p < 0.001$, paired *t*-test.

using the NOR paradigm assessing short-term recognition memory (Figure 5(a)).

Out of 16 Ts65Dn and 16 euploid mice, 2 Ts65Dn and 1 euploid were removed from statistical analysis because they displayed abnormally low levels of object exploration ($t < 7$ s) during retention test, hence precluding analysis of their memory performance. The remaining mice spent a large amount of time exploring objects ($t = 77 \pm 4.9$ s).

A preliminary analysis of global levels of object exploration was carried out during the acquisition and retention phases of the object recognition task (data not shown). ANOVA did not show any effects of the group ($F_{3,25} = 2.40$, $p = ns$) and testing phase ($F_{1,25} = 3.67$, $p = ns$) nor of the interaction between these factors ($F_{3,25} < 1$, $p = ns$). These results

demonstrate that whatever their genotype and treatment, mice displayed the same overall levels of exploration directed towards objects.

Object recognition memory performance was then specifically evaluated during the retention phase by analysing the time spent by mice exploring familiar versus novel objects (Table 1). Unpaired *t*-tests showed that euploid mice, treated or not with $\alpha 5$ IA, were able to discriminate between the two objects (vehicle condition: $t_6 = 2.49$, $p < 0.05$; $\alpha 5$ IA condition: $t_7 = 6.3$, $p < 0.001$) indicating normal recognition memory. In contrast, vehicle-treated Ts65Dn did not show any significant exploratory preference towards the novel object ($t_7 < 1$) underscoring impaired recognition memory. However, Ts65Dn mice treated with $\alpha 5$ IA were able to clearly

differentiate between the two objects, indicating that $\alpha 5IA$ treatment was able to restore normal recognition memory ($t_5 = 4.85$, $p < 0.005$).

In order to better clarify the effects of genotype and $\alpha 5IA$ treatment on recognition memory we calculated a learning index (I) according to the following formula:

$$I = \left[\frac{\left(\frac{\text{Novel Object}}{\text{Exploration Time}} \right) - \left(\frac{\text{Familiar Object}}{\text{Exploration Time}} \right)}{\text{Total Exploration Time}} \right]$$

ANOVA on this learning index indicated significant effect of group ($F_{3,25} = 12.52$, $p < 0.001$). *Post-hoc* analysis indicated a significant effect of $\alpha 5IA$ treatment which largely potentiated recognition memory (Figure 5(a)). The effect was observed in both euploid (comparison vehicle vs. drug conditions: $F_{1,25} = 8.42$, $p < 0.01$) and Ts65Dn mice ($F_{1,25} = 24.34$, $p < 0.0001$). The analysis also showed that vehicle-treated Ts65Dn mice had a lower learning index as compared with euploid mice ($F_{1,25} = 4.97$, $p < 0.05$). However, following $\alpha 5IA$ treatment, the learning index of Ts65Dn and euploid mice were found to be similar ($F < 1$), underlining recovery of performance following treatment in this genotype.

In summary, Ts65Dn mice under vehicle condition presented impaired recognition memory in the NOR task that was recovered, after $\alpha 5IA$ treatment.

$\alpha 5IA$ potentiates evoked-neuronal activity

In order to determine how $\alpha 5IA$ modulated behaviour-evoked neuronal activity in euploid and Ts65Dn mice, we performed a brain mapping analysis of an immediate early gene product (Fos protein). Animals were trained as described previously in the NOR task until completion of the acquisition phase (Figure 5(b)). We first confirmed that all groups displayed the same level of object exploration, with no effect of genotype, treatment and of their interactions (all $F < 1$; data not shown). In addition, the distance travelled by mice did not vary significantly with genotype ($F < 1$) and treatment ($F_{1,29} = 2.29$, $p = \text{ns}$) (data not shown). It was then concluded that all mice received the same sensorimotor stimulation during the acquisition phase of the NOR task. Ninety minutes after completion of behaviour, mice were sacrificed and their brains processed for quantitative assessment of the neuronal activity marker Fos (Figure 5(b)). The proportion of brain tissue immunolabelled against Fos was quantified and analysed using ANOVA. This analysis revealed a significant effect of Treatment as immunoreactivity was found to be significantly increased in $\alpha 5IA$ treated mice ($F_{1,13} = 6.376$, $p < 0.025$). There was, however, no effect of the genotype or of the interactions between genotype and treatment ($F < 1$), suggesting that euploid and Ts65Dn mice displayed the same overall levels of Fos immunoreactivity and underwent similar effects after $\alpha 5IA$ treatment. Complementary analysis showed that the effect of $\alpha 5IA$ was not the same throughout brain regions ($F_{3,39} = 85.93$, $p < 0.0001$; Figure 5(b)) illustrating that NOR-evoked neuronal activity was restricted to some brain areas (CA1, perirhinal and posterior cingulated

cortices). The interaction between region and treatment was found to be significant ($F_{3,39} = 5.612$, $p < 0.005$), likely due to the lack of $\alpha 5IA$ -induced increase of Fos immunoreactivity in one of the four regions analysed, the dentate gyrus (effect of treatment: posterior cingulate cortex $F_{1,15} = 35.59$, $p < 0.0001$; perirhinal cortex $F_{1,15} = 6.37$; $p < 0.025$; CA1 $F_{1,15} = 5.30$, $p < 0.05$; dentate gyrus $F < 1$).

We thus concluded that following behavioural stimulation, $\alpha 5IA$ enhanced evoked immediate early gene products in specific brain regions such as hippocampus, perirhinal and posterior cingulate cortices.

$\alpha 5IA$ treatment does not induce side effects in Ts65Dn and euploid mice

Convulsant and pro-convulsant effects. The $\alpha 5IA$ molecule was demonstrated previously to be neither convulsant nor anxiogenic in wild-type mice and rats (Dawson et al., 2006); however, this characteristic had never been tested in DS mouse models. We tested the putative convulsant effect of $\alpha 5IA$ after a single injection of 50 mg/kg (10 \times the dose producing promnesic effects). Neither euploid nor Ts65Dn mice displayed any convulsions after injection (Table 2). We then tested the pro-convulsant effect of $\alpha 5IA$ by injecting it (50 mg/kg) 20 min before a sub-convulsant dose of pentylenetetrazol (45 mg/kg) that induces myoclonic convulsions in about 50% of mice. Injection of $\alpha 5IA$ did not potentiate convulsant activity of pentylenetetrazol in either euploid or Ts65Dn mice (ANOVA on the latency of myoclonic jerks: effect of group $F_{3,8} = 1.43$, $p = \text{ns}$).

Locomotor activity. In the open field task, ANOVA on travelled distances (Figure 6(a)) did not show any effect of group ($F_{3,29} = 2.64$, $p = \text{ns}$). To evaluate anxiety during the open field test, a periphery-to-centre exploration ratio was measured. ANOVA on this measurement did not reveal any effect of group ($F < 1$; Figure 6(b)).

Putative anxiogenic effects. In order to better assess the level of anxiety in euploid and Ts65Dn mice treated or not with $\alpha 5IA$, we used the elevated plus maze task. Time spent in the open arms of the elevated plus maze was taken as a measure of anxiety levels (the greater the time spent, the less anxious). ANOVA of this measure did not show any significant effect of group ($F_{3,40} = 2.59$, $p = 0.06$). We nevertheless observed that vehicle-treated Ts65Dn mice had an increased propensity to stay in open arms as compared to euploid mice ($F_{1,40} = 3.68$, $p = 0.062$) and hence displayed some trends for hypo-anxiety traits (for similar findings see Demas et al., 1996). In addition, as illustrated in Figure 7, it appears that $\alpha 5IA$ slightly decreased time spent in the open arms. This tendency was significant in Ts65Dn mice ($F_{1,40} = 4.56$, $p < 0.05$) but not in euploid mice ($F < 1$). We therefore propose that the weak 'anxiogenic-like' effects of $\alpha 5IA$ in Ts65Dn mice are mainly due, in our experimental design, to a normalization of behaviour, from low to normal levels of anxiety, in these mice.

Table 2. Lack of convulsant and pro-convulsant activities of $\alpha 5IA$

Genotype	Treatment	Latency of myoclonic jerks mean \pm SEM	Rate of convulsant mice
Convulsant effects			
Euploids	Vehicle	/	0/6
	$\alpha 5IA$ (50 mg/kg)	/	0/6
Ts65Dn	Vehicle	/	0/6
	$\alpha 5IA$ (50 mg/kg)	/	0/7
Pro-convulsant effects after pentylenetetrazol (45 mg/kg)			
Euploids	Vehicle	448 \pm 145	4/6
	$\alpha 5IA$ (50 mg/kg)	330 \pm 90	3/6
Ts65Dn	Vehicle	507 \pm 39	3/6
	$\alpha 5IA$ (50 mg/kg)	796 \pm 354	4/7

Data indicate that $\alpha 5IA$ (50 mg/kg) did not induce any convulsant effects in either euploid or Ts65Dn mice. The drug also did not promote the convulsant action of pentylenetetrazol (45mg/kg) in the two genotypes.

Effects of chronic treatment with $\alpha 5IA$. Euploid mice treated with $\alpha 5IA$ (5 mg/kg) for 2 weeks did not show any change in their gross behaviour. Body weights were comparable between vehicle and $\alpha 5IA$ mice ($F < 1$) and both groups showed normal progressive growth (ANOVA on body weights: $F_{8,64} = 22.45$, $p < 0.0001$, data not shown).

More importantly, mice treated chronically with $\alpha 5IA$ 5 mg/kg showed similar levels of anxiety as vehicle-treated mice (unpaired t -test on the time spent in open arms: $t_8 = 1.04$, $p = ns$; Figure 7, right panel), suggesting that $\alpha 5IA$ chronic treatment did not alter anxiety-related behaviours.

Following 5 weeks of chronic treatment with $\alpha 5IA$, various organs were collected and processed for routine histopathological examination. Haematoxylin–eosin (Figure 8) and periodic acid-Schiff stained sections (not shown) did not reveal any significant macroscopic nor microscopic tissue alterations in any of the three experimental groups (non-injected, vehicle-injected or $\alpha 5IA$ -treated mice). In particular, examination of brain, hepatic and renal tissues under polarized light did not show the occurrence of abnormal crystals in mice that did receive injections of $\alpha 5IA$.

In summary, it appears that treatment with $\alpha 5IA$ did not promote any significant liabilities as it did not induce convulsant/pro-convulsant activity nor affected locomotion and anxiety-related behaviours.

Discussion

$\alpha 5IA$ restores cognitive dysfunction in Ts65Dn mice

In this study we demonstrated that treatment with $\alpha 5IA$ largely alleviates the cognitive deficits of Ts65Dn mice. Indeed Ts65Dn mice receiving a single administration of $\alpha 5IA$ increased their memory performance in the NOR task and behaved as $\alpha 5IA$ -treated euploid littermates. Furthermore, repeated $\alpha 5IA$ treatment across training sessions in the MWM task allowed Ts65Dn mice to decrease their anomalous foraging behaviours, and to learn a fixed goal location with the same efficiency as euploid mice. Rescue of learning deficits in Ts65Dn mice by $\alpha 5IA$ appeared to be specific since

sensory functions in the MWM test or motivation to explore objects in the NOR task remained unchanged in this genotype and were not affected by $\alpha 5IA$ treatment. These exciting findings provide, for the first time, important preclinical evidence for the hypothesis that release of GABAergic inhibition by $\alpha 5$ GABA-A benzodiazepine inverse agonists may improve cognitive function in DS individuals.

Treatment with GABA_A antagonists (e.g. pentylenetetrazol) was previously shown to rescue memory performances in Ts65Dn mice trained in the NOR task (Fernandez et al., 2007) and in the MWM task (Rueda et al., 2008). However, the use of GABA antagonists as well as of non-specific GABA-A benzodiazepine inverse agonists as therapeutic molecules has serious limitations because of their known adverse effects: convulsant, pro-convulsant and anxiogenic effects. The $\alpha 5$ GABA-A benzodiazepine inverse agonists, thanks to their unique pharmacological profile, are devoid of such liabilities (for a review see Atack, 2009). In the present study, we further show that Ts65Dn mice treated with $\alpha 5IA$ did not display any alteration in their locomotor behaviour. More importantly and as opposed to treatments with pentylenetetrazol, Ts65Dn did not develop significant alterations of anxiety-related behaviours nor any convulsant or pro-convulsant activity. A putative renal toxicity of $\alpha 5IA$ has been claimed in some reports (Atack, 2008; Merschman et al., 2005) because of the *in vivo* formation and crystallization of insoluble metabolites at extremely high dosages (240 mg/kg/day for 5 weeks). However, we did not find evidence of any anatomopathological lesions in mice chronically treated with $\alpha 5IA$ at 5 mg/kg, the pharmacologically active dose (see Supplemental Text T1 for additional discussion).

From these observations it can be concluded that $\alpha 5IA$ has a better therapeutic profile than GABA antagonists. Indeed the first successful use of $\alpha 5IA$ as a cognitive enhancer candidate in human subjects has been recently published (Nutt et al., 2007) affirming its good safety and tolerability.

$\alpha 5IA$ effects on acquisition and retrieval of memories

In addition to its therapeutic effects in Ts65Dn mice, $\alpha 5IA$ displayed some promnesic action in euploid mice trained in




Figure 6. $\alpha 5$ IA does not alter locomotor activity and anxiety of Ts65Dn and euploid mice in the open field. Effects of $\alpha 5$ IA (5mg/kg) were evaluated on locomotion and anxiety in the open field. (a) Analysis of horizontal activity (travelled distances; mean \pm SEM) did not show any effect of treatment, underscoring that a single $\alpha 5$ IA injection did not modify the gross locomotor activity of both euploid and Ts65Dn mice. (b) To assess anxiety during the open field session, a periphery-to-centre exploration ratio was measured (P/C ratio; mean \pm SEM). Analysis of this measure did not reveal any effects of Genotype or Treatment.

short-term memory tasks using the NOR or DMTP paradigms. Most studies investigating cognitive-enhancing properties of $\alpha 5$ -specific GABA-A inverse agonists were indeed conducted in rodents trained in the DMTP test (Collinson et al., 2002; Dawson et al., 2006). However, in a spatial reference memory task requiring gradual memorization of an invariant goal location throughout trials and days (MWM task) we showed that $\alpha 5$ IA largely facilitated the performance of Ts65Dn mice but not those of euploid mice. This underscores that $\alpha 5$ IA, under non-pathological conditions, might have positive outcomes but only in specific (short-term memory) training conditions.




Figure 7. $\alpha 5$ IA does not induce anxiety-related behaviours. Anxiety was assessed in the standard elevated plus maze task, in both euploid and Ts65Dn mice under vehicle or $\alpha 5$ IA (one single 15 mg/kg i.p. injection; left panel of the figure). Under vehicle condition, Ts65Dn mice showed a trend for hypoanxiety (increased time in open arms) in comparison to euploid mice. Acute treatment with $\alpha 5$ IA did not modify the behaviour of euploid mice, but significantly reduced the time spent in open arms by Ts65Dn mice. This effect can be ascribed to a normalization of behaviour in the Ts65Dn mice. Semi-chronic injections of $\alpha 5$ IA in euploid mice (5 mg/kg five times a week for 2 weeks; right panel of the figure) did not alter the anxiety levels. Horizontal dotted line indicates the baseline performance of mice acutely treated with vehicle. * $p < 0.05$, ANOVA with Fisher's *post hoc* comparisons.

When evaluating the effects of $\alpha 5$ IA on the retrieval of long-term (24 hours) spatial memory during the probe test of the MWM task, we showed that $\alpha 5$ IA did not actually increase retention performance in either euploid or Ts65Dn mice. This indicates that $\alpha 5$ IA mainly exerts its nootropic action during the acquisition of information but might be less potent in stimulating accurate retrieval of the previously formed memories. Collinson, Attack and colleagues suggested that GABA-A $\alpha 5$ inverse agonists could, under some circumstances, improve both the acquisition and the retrieval of spatial memories. However, they used memory paradigms based on short-intermediate retention intervals (15–180 min) that do not fully assess long-term recall (at least 24 hours post-acquisition) as usually performed during probe tests in spatial navigation tasks (Attack et al., 2006; Collinson et al., 2006).

Altogether these studies suggest that $\alpha 5$ IA stimulates short-term memories in normal and cognitively impaired mice, likely through a modulation of the attentional-working memory process. In addition, gradual learning across training sessions, as evaluated in the MWM task, can also be potentiated by $\alpha 5$ IA in Ts65Dn mice but the effects are less pronounced in euploid mice displaying high learning proficiencies in this task. Finally, the stabilization and late recall of




Figure 8. $\alpha 5$ IA does not induce any histological lesions after chronic treatment. Following chronic treatment with $\alpha 5$ IA (5 mg/kg; five injections/week for 5 weeks), different organs were ablated and processed for routine histopathological examination. As illustrated, haematoxylin-eosin staining did not reveal any significant macroscopic or microscopic tissue alterations in liver or kidney in any of the three experimental groups (non-injected, vehicle-injected or $\alpha 5$ IA-treated mice). The same negative findings were observed following periodic acid-Schiff staining of the tissues (not illustrated). Examination of brain, hepatic and renal tissues under polarized light revealed the lack of abnormal crystals in mice receiving injections of $\alpha 5$ IA. The size and distribution of urine crystals (not illustrated) appeared to be very similar in the different groups. Scale bar = 100 μ m.

reference memories do not appear to be impacted by $\alpha 5$ IA treatment.

Putative mechanisms of action of $\alpha 5$ IA in normal and diseased brain

In close association with an enhancement of cognitive proficiency, we showed that treatment with $\alpha 5$ IA also increased immediate early gene products (Fos protein levels) following a behavioural stimulation that mimics a learning episode (encoding of new information). Increased Fos immunoreactivity was observed in all of the sampled brain areas involved in recognition memory (posterior cingulate and perirhinal cortices, pyramidal cell layer of the hippocampus) but not at the level of the dentate gyrus. This latter observation was expected as the dentate gyrus is a sector of the hippocampus that displays only low concentrations of GABA_A $\alpha 5$ receptors (Pirker et al., 2000; Sperk et al., 1997). Paucity of targets might hence explain the local lack of drug-induced increased neuronal activity. Importantly we did not find any differences between euploid and Ts65Dn mice in terms of Fos immunoreactivity levels. The absence of a genotype effect under vehicle conditions underscores that Ts65Dn mice did not sustain an overall pattern of reduced neuronal activity, at least during the exploration–memorization of a new environment. Following drug administration, both genotypes displayed

significant (and comparable) increases in the levels of neuronal activity markers. This potentiation of brain activity during acquisition of new information might therefore be the substratum of the ‘general’ promnesic effects of $\alpha 5$ IA that should be independent of the disease status.

While we showed that Ts65Dn mice displayed similar levels of brain activity as euploid mice, it is known from the literature that these mice concurrently develop synaptic plasticity anomalies as exemplified by impaired LTP (Siarey et al., 1997). Reduction of synaptic plasticity in Ts65Dn mice is observed in the absence of any notable changes in the general properties of excitatory synaptic transmission (Kleschevnikov et al., 2004). Importantly these LTP deficits can be rescued following release of the GABAergic inhibitory transmission by means of picrotoxin (Kleschevnikov et al., 2004). In parallel it has been shown recently that $\alpha 5$ GABA-A inverse agonists, including the drug used in the present study, potentiate LTP in mouse hippocampal slices (Ballard et al., 2009; Dawson et al., 2006) and it can be postulated that these drugs likely have the potential to reverse LTP deficits and concomitantly to improve cognition in Ts65Dn mice.

In conclusion, we have demonstrated that an $\alpha 5$ -selective GABA-A inverse agonist can restore cognitive function (short-term recognition memory and spatial learning) in a mouse model of DS. Our results strengthen the hypothesis that modifying the GABAergic-mediated balance between excitatory and inhibitory neurotransmission can efficiently

alleviate cognitive impairments in preclinical models of DS. The exact mechanism of action of $\alpha 5$ IA remains to be clarified, but might involve potentiation of neuronal activity and of synaptic plasticity of neural networks.

$\alpha 5$ IA, because of its lack of convulsant or anxiogenic effects, has a more favourable therapeutic profile than other GABAergic drugs such as pentylentetrazol. Also we did not detect any toxicity of $\alpha 5$ IA following repeated injections. The first successful use of $\alpha 5$ IA as a cognitive enhancer for blocking alcohol's amnesic activity in human subjects has indeed been published, confirming it as safe and well tolerated (Nutt et al., 2007). The excellent safety profile of $\alpha 5$ IA and of similar recently developed compounds will undoubtedly facilitate their clinical investigation in individuals with DS.

Acknowledgements

We are grateful to Professor JJ Hauw, Professor C Duyckaerts and technicians from the Escourolle Neuropathology Laboratory (Paris) for assistance/expertise in histological studies, and to zootechnicians from CNRS UMR 8195 (Orsay) and UPS 44 TAAM (Orléans) for providing animal care.

Funding

The authors wish to thank the Foundation AMIPI Bernard Vendre and particularly Mr and Mrs Vendre and Mr Jean-Marc Richard, for their funding and support. We are also grateful for support from the EEC AnEUploidy program (grant number LSHG-CT-2006-037627), and from the Fondation Jérôme Lejeune. JB, BD, YH, RHD and MCP received funding from the AMIPI – Bernard Vendre foundation to carry out this research (including salary for post-doctoral fellowships to JB).

Conflict of interest statement

The authors are not aware of any biases that might be perceived as affecting the objectivity of the present research work.

References

Atack JR (2008) GABA(A) receptor subtype-selective efficacy: TPA023, an alpha2/alpha3 selective non-sedating anxiolytic and alpha5IA, an alpha5 selective cognition enhancer. *CNS Neurosci Ther* 14: 25–35.

Atack JR (2009) Preclinical and clinical pharmacology of the GABA(A) receptor alpha5 subtype-selective inverse agonist alpha5IA. *Pharmacol Ther*, in press.

Atack JR, Bayley PJ, Seabrook GR, et al. (2006) L-655,708 enhances cognition in rats but is not proconvulsant at a dose selective for alpha5-containing GABAA receptors. *Neuropharmacology* 51: 1023–1029.

Ballard TM, Knoflach F, Prinssen E, et al. (2009) RO4938581, a novel cognitive enhancer acting at GABA(A) alpha5 subunit-containing receptors. *Psychopharmacology (Berl)* 202: 207–223.

Best TK, Siarey RJ and Galdzicki Z (2007) Ts65Dn, a mouse model of Down syndrome, exhibits increased GABAB-induced potassium current. *J Neurophysiol* 97: 892–900.

Collinson N, Atack JR, Laughton P, et al. (2006) An inverse agonist selective for alpha5 subunit-containing GABAA receptors improves encoding and recall but not consolidation in the Morris water maze. *Psychopharmacology (Berl)* 188: 619–628.

Collinson N, Kuenzi FM, Jarolimek W, et al. (2002) Enhanced learning and memory and altered GABAergic synaptic transmission in mice lacking the alpha 5 subunit of the GABAA receptor. *J Neurosci* 22: 5572–5580.

Costa AC, Stasko MR, Schmidt C, et al. (2010) Behavioral validation of the Ts65Dn mouse model for Down syndrome of a genetic background free of the retinal degeneration mutation Pde6b(rd1). *Behav Brain Res* 206: 52–62.

Dawson GR, Maubach KA, Collinson N, et al. (2006) An inverse agonist selective for alpha5 subunit-containing GABAA receptors enhances cognition. *J Pharmacol Exp Ther* 316: 1335–1345.

Demas GE, Nelson RJ, Krueger BK, et al. (1996) Spatial memory deficits in segmental trisomic Ts65Dn mice. *Behav Brain Res* 82: 85–92.

Escorihuela RM, Fernandez-Teruel A, Vallina IF, et al. (1995) A behavioral assessment of Ts65Dn mice: a putative Down syndrome model. *Neurosci Lett* 199: 143–146.

Faure A, Verret L, Bozon B, et al. (2009) Impaired neurogenesis, neuronal loss, and brain functional deficits in the APPxPS1-Ki mouse model of Alzheimer's disease. *Neurobiol Aging*, in press.

Fernandez F, Morishita W, Zuniga E, et al. (2007) Pharmacotherapy for cognitive impairment in a mouse model of Down syndrome. *Nat Neurosci* 10: 411–413.

Hoelter SM, Dalke C, Kallnik M, et al. (2008) "Sighted C3H" mice—a tool for analysing the influence of vision on mouse behaviour? *Front Biosci* 13: 5810–5823.

Janus C, Welzl H, Hanna A, et al. (2004) Impaired conditioned taste aversion learning in APP transgenic mice. *Neurobiol Aging* 25: 1213–1219.

Kleschevnikov AM, Belichenko PV, Villar AJ, et al. (2004) Hippocampal long-term potentiation suppressed by increased inhibition in the Ts65Dn mouse, a genetic model of Down syndrome. *J Neurosci* 24: 8153–8160.

Menendez M (2005) Down syndrome, Alzheimer's disease and seizures. *Brain Dev* 27: 246–252.

Merschman SA, Rose MJ, Pearce GE, et al. (2005) Characterization of the solubility of a poorly soluble hydroxylated metabolite in human urine and its implications for potential renal toxicity. *Pharmazie* 60: 359–363.

Nutt DJ, Besson M, Wilson SJ, et al. (2007) Blockade of alcohol's amnesic activity in humans by an alpha5 subtype benzodiazepine receptor inverse agonist. *Neuropharmacology* 53: 810–820.

Pirker S, Schwarzer C, Wieselthaler A, et al. (2000) GABA(A) receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* 101: 815–850.

Pueschel SM, Louis S and McKnight P (1991) Seizure disorders in Down syndrome. *Arch Neurol* 48: 318–320.

Reeves RH and Garner CC (2007) A year of unprecedented progress in Down syndrome basic research. *Ment Retard Dev Disabil Res Rev* 13: 215–220.

Reeves RH, Irving NG, Moran TH, et al. (1995) A mouse model for Down syndrome exhibits learning and behaviour deficits. *Nat Genet* 11: 177–184.

Rueda N, Florez J and Martinez-Cue C (2008) Chronic pentylentetrazole but not donepezil treatment rescues spatial cognition in Ts65Dn mice, a model for Down syndrome. *Neurosci Lett*.

Sherman SL, Allen EG, Bean LH, et al. (2007) Epidemiology of Down syndrome. *Ment Retard Dev Disabil Res Rev* 13: 221–227.

Siarey RJ, Stoll J, Rapoport SI, et al. (1997) Altered long-term potentiation in the young and old Ts65Dn mouse, a model for Down syndrome. *Neuropharmacology* 36: 1549–1554.

Sperk G, Schwarzer C, Tsunashima K, et al. (1997) GABA(A) receptor subunits in the rat hippocampus I: immunocytochemical distribution of 13 subunits. *Neuroscience* 80: 987–1000.

Sternfeld F, Carling RW, Jelley RA, et al. (2004) Selective, orally active gamma-aminobutyric acid alpha5 receptor inverse agonists as cognition enhancers. *J Med Chem* 47: 2176–2179.

- Sur C, Fresu L, Howell O, et al. (1999) Autoradiographic localization of alpha5 subunit-containing GABAA receptors in rat brain. *Brain Res* 822: 265–270.
- Veall RM (1974) The prevalence of epilepsy among mongols related to age. *J Ment Defic Res* 18: 99–106.
- Wisden W, Laurie DJ, Monyer H, et al. (1992) The distribution of 13 GABAA receptor subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon. *J Neurosci* 12: 1040–1062.
- Wiseman FK, Alford KA, Tybulewicz VL, et al. (2009) Down syndrome—recent progress and future prospects. *Hum Mol Genet* 18: R75–R83.
- Wishart JG, Willis DS, Cebula KR, et al. (2007) Collaborative learning: comparison of outcomes for typically developing children and children with intellectual disabilities. *Am J Ment Retard* 112: 361–374.

Copyright of Journal of Psychopharmacology is the property of Sage Publications, Ltd. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.