



HAL
open science

Volunteer effect and compromised randomization in the Mayo Project of screening for lung cancer

Lorenzo Dominioni, Albino Poli, William Mantovani, Nicola Rotolo, Andrea Imperatori

► **To cite this version:**

Lorenzo Dominioni, Albino Poli, William Mantovani, Nicola Rotolo, Andrea Imperatori. Volunteer effect and compromised randomization in the Mayo Project of screening for lung cancer. *European Journal of Epidemiology*, 2010, 26 (1), pp.79-80. 10.1007/s10654-010-9519-4 . hal-00634766

HAL Id: hal-00634766

<https://hal.science/hal-00634766>

Submitted on 23 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Volunteer effect and compromised randomization in the Mayo Project of screening for lung cancer

Lorenzo Dominioni¹, Albino Poli², William Mantovani², Nicola Rotolo¹, Andrea Imperatori¹

¹ *Center for Thoracic Surgery, University of Insubria - Varese, Italy*

² *Department of Public Health, University of Verona, Italy*

Full address for correspondence:

Lorenzo Dominioni, MD

Center for Thoracic Surgery, University of Insubria

Ospedale di Circolo

Viale Borri 57,

21100 Varese, Italy

e-mail: lorenzo.dominioni@uninsubria.it

phone: +39 0332 278868

fax: +39 0332 260260

Abstract

It has been confirmed recently that the volunteer effect in lung cancer screening is characterized by higher lung cancer mortality risk in self-selected screening participants. The Mayo Lung Project, the most influential trial of screening for lung cancer ever completed, was conducted in *nonvolunteer* Mayo Clinic outpatients, with a peculiar study design that rendered the randomization vulnerable to the volunteer effect.

Of all nonvolunteers randomized in the Mayo Lung Project, only those allocated in the screened group were asked consent to participate in the trial. The final Mayo Lung Project report stated that 655 randomized nonvolunteers refused screening and were excluded from the study, thus documenting violation of the rule that no selection should occur after randomization. The long-term follow-up of the Mayo Lung Project showed an enigmatic result which has never been explained: the lung cancer mortality was 13% higher in the screening intervention group than in the control group [4.4 (95% CI 3.9-4.9) vs. 3.9 (95% CI 3.5-4.4) per 1,000 person-years; $p=0.09$]. Such overrepresented mortality is consistent with the volunteer effect and supports the concept that the Mayo Lung Project randomization was compromised by the post-randomization self-selection of participant nonvolunteers.

Key words:

Compromised randomization, lung cancer screening, Mayo Lung Project, volunteer effect

Abbreviations:

Chest X-rays: CXR

Confidence interval: CI

Lung cancer: LC

Johns Hopkins Lung Project: JHLP

Mayo Lung Project: MLP

LETTER TO THE EDITOR

Dear Editor,

The volunteer effect has been studied extensively in screening trials for cancer of the breast, colon-rectum and prostate [1,2,3], but it has been scarcely investigated in the context of screening for lung cancer (LC). The Johns Hopkins Lung Project (JHLP) first documented that the LC mortality of volunteers screened by chest X-rays (CXR) was significantly higher than expected on the basis of two large population studies, namely the Veterans Study [observed/expected LC mortality ratio 1.72; 95% confidence interval (CI) 1.28-2.18] and the American Cancer Society Study (observed/expected LC mortality ratio 1.46; 95% CI 1.09-1.83) [4]. Recently, in a population-based study of smokers offered CXR screening we confirmed that the volunteer effect is characterized by significantly higher LC mortality risk in self-selected participants compared to nonparticipants (risk ratio 1.40; 95% CI, 1.03-1.91) [5]. The volunteer effect may bias the results of comparative LC screening studies of *nonvolunteers*, if the latter are not randomized adequately. The Mayo Lung Project (MLP), the most influential trial of CXR screening for LC [6,7], was conducted in *nonvolunteer* Mayo Clinic outpatients with a peculiar design that rendered the randomization vulnerable.

Indeed the details of the MLP randomization, described on page 1375 in the preliminary report [8], specify that: *“the random assignment to the close-surveillance or the control group is made at the interview. Up to this point, the Mayo Lung Project has not been mentioned, and, if the assignment is to the control group, nothing is said of it and the interview ends. If the assignment is to the close-surveillance group, however, the interviewer explains that program to the candidate and obtains his informed consent and agreement to take part in it”*.

Accordingly, the MLP candidates were first randomized and subsequently only the nonvolunteers allocated in the screened group were asked consent to participate in the trial; because of that study design we hypothesize that the MLP randomization was likely compromised by the volunteer bias.

The final MLP report stated that 655 randomized nonvolunteers refused screening and were excluded from the study [9], thus documenting violation of the rule that no selection should occur after randomization. After such selection and exclusion of candidates, an important methodological weakness, there is no guarantee that the screening and control groups at baseline were well matched for all known and unknown LC risks [10]. Notably, the extended follow-up of the MLP showed two puzzling results: 1) in 16 years after the end of screening (1983-1999) the total number of newly diagnosed LCs was 11% greater in the screened group than in controls (379 vs. 340) [11], an excess certainly not attributable to screening overdiagnosis; 2) the LC mortality was 13% higher in the screening intervention group than in the control group [4.4 (95% CI 3.9-4.9) vs. 3.9 (95% CI 3.5-4.4) per 1,000 person-years; $p=0.09$]. [12]. These enigmatic results are consistent with the volunteer effect in the LC screening group; moreover, they support the concept that the documented post-randomization selection of nonvolunteers compromised the MLP randomization.

References

1. Richardson A, Wells JE. Breast cancer screening: the effect of self selection for screening on comparisons of randomized controlled trials. *J Med Screen* 1997; 4: 16-18.
2. Blom J, Yin L, Lidén A, Dolk A, Jeppson B, Pahlman L et al. A 9-year follow-up study of participants and nonparticipants in sigmoidoscopy screening: importance of self-selection. *Cancer Epidemiol Biomarkers Prev* 2008; 17: 1163-1168.
3. Otto SJ, Schroeder FH, De Koning HJ. Low all-cause mortality in the volunteer-based Rotterdam section of the European randomised study of screening for prostate cancer: self-selection bias? *J Med Screen* 2004; 11:89-92.
4. Levin ML, Tockman MS, Frost JK, Ball WCJr. Lung cancer mortality in males screened by chest X-ray and cytologic sputum examination: a preliminary report. *Recent Results Cancer Res* 1982; 82: 138-146.
5. Dominioni L, Rotolo N, Poli A, Paolucci M, Sessa F, D'Ambrosio V et al. Self-selection effects in smokers attending lung cancer screening. A 9.5-year population-based cohort study in Varese, Italy. *J Thorac Oncol* 2010; 5: 428-435.
6. Fontana RS, Sanderson DR, Woolner LB, Taylor WF, Miller WE, Muhm JR et al. Screening for lung cancer: a critique of the Mayo Lung Project. *Cancer* 1991; 67(Suppl): 1155-1164.

7. National Cancer Institute. Available from:
<http://www.cancer.gov/cancertopics/pdq/screening/lung/HealthProfessional/page4>.
Accessed Aug. 1, 2010
8. Fontana RS, Sanderson DR, Miller WE, Woolner LB, Taylor WF, Uhlenhopp MA.
The Mayo Lung Project: preliminary report of “early cancer detection” phase.
Cancer 1972; 30: 1373-1382.
9. Woolner LB, Fontana RS, Sanderson DR, Miller WE, Muhm JR, Taylor WF et al.
Mayo Lung Project. Evaluation of lung cancer screening through December 1979.
Mayo Clin Proc 1981; 56: 544-555.
10. Manser R, Irving LB, Stone C, Byrnes G, Abramson MJ, Campbell D. Screening for
lung cancer. *Cochrane Database of Systematic Reviews* 2004, Issue 1. Art. No.:
CD001991. DOI: 10.1002/14651858.CD001991.pub2.
11. Marcus PM, Bergstralh EJ, Zweig MH, Harris A, Offord KP, Fontana RS. Extended
lung cancer incidence follow-up in the Mayo Lung Project and overdiagnosis. *J Natl
Cancer Inst* 2006; 98: 748-756.
12. Marcus PM, Bergstralh EJ, Fagerstrom RM, Williams DE, Fontana RS, Taylor WF
et al. Lung cancer mortality in the Mayo Lung Project: impact of extended follow-
up. *J Natl Cancer Inst* 2000; 92: 1308-1316.