



HAL
open science

Nature or nurture: Let food be your epigenetic medicine in chronic inflammatory disorders

Katarzyna Szarc Vel Szic, 'Matladi Ndlovu N, Guy Haegeman, Wim Vanden
Berghe

► **To cite this version:**

Katarzyna Szarc Vel Szic, 'Matladi Ndlovu N, Guy Haegeman, Wim Vanden Berghe. Nature or nurture: Let food be your epigenetic medicine in chronic inflammatory disorders. *Biochemical Pharmacology*, 2010, 80 (12), pp.1816. 10.1016/j.bcp.2010.07.029 . hal-00637154

HAL Id: hal-00637154

<https://hal.science/hal-00637154>

Submitted on 31 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Nature or nurture: Let food be your epigenetic medicine in chronic inflammatory disorders

Authors: Katarzyna Szarc vel Szic, 'Matladi Ndlovu N, Guy Haegeman, Wim Vanden Berghe




PII: S0006-2952(10)00567-8
DOI: doi:10.1016/j.bcp.2010.07.029
Reference: BCP 10662

To appear in: *BCP*

Received date: 26-5-2010
Revised date: 19-7-2010
Accepted date: 23-7-2010

Please cite this article as: Szic KS, N MN, Haegeman G, Berghe WV, Nature or nurture: Let food be your epigenetic medicine in chronic inflammatory disorders, *Biochemical Pharmacology* (2010), doi:10.1016/j.bcp.2010.07.029

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1
2
3 **Nature or nurture: Let food be your epigenetic medicine in chronic inflammatory**
4 **disorders**
5
6

7
8
9 Katarzyna Szarc vel Szi^{1,2}, Matladi Ndlovu N.^{1,3}, Guy Haegeman¹, Wim Vanden Berghe^{1,2}
10

11
12
13
14 ¹*Laboratory of Eukaryotic Gene Expression and Signal Transduction (LEGEST), Department*
15 *of Physiology, Ghent University, K.L.Ledeganckstraat 35, Gent, Belgium*

16
17
18 ²*Present address: Lab Protein Science, Proteomics and Epigenetic Signaling, Department of*
19 *Biomedical Sciences, University Antwerp, Campus Drie Eiken, Universiteitsplein 1, Wilrijk,*
20 *Belgium*

21
22
23 ³*Present address: Laboratory of Cancer Epigenetics, Free University of Brussels, Faculty of*
24 *Medicine, 808 route de Lennik, 1070 Brussels, Belgium*
25
26

27
28
29 *Corresponding author: wim.vandenbergh@ua.ac.be*
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract:

1 Numerous clinical, physiopathological and epidemiological studies have underlined the
2 detrimental or beneficial role of nutritional factors in complex inflammation related disorders
3 such as allergy, asthma, obesity, type 2 diabetes, cardiovascular disease, rheumatoid arthritis
4 and cancer. Today, nutritional research has shifted from alleviating nutrient deficiencies to
5 chronic disease prevention. It is known that lifestyle, environmental conditions and nutritional
6 compounds influence gene expression. Gene expression states are set by transcriptional
7 activators and repressors and are often locked in by cell-heritable chromatin states. Only
8 recently, it has been observed that the environmental conditions and daily diet can affect
9 transgenerational gene expression via “reversible” heritable epigenetic mechanisms.
10 Epigenetic changes in DNA methylation patterns at CpG sites (epimutations) or corrupt
11 chromatin states of key inflammatory genes and noncoding RNAs, recently emerged as major
12 governing factors in cancer, chronic inflammatory and metabolic disorders. Reciprocally,
13 inflammation, metabolic stress and diet composition can also change activities of the
14 epigenetic machinery and indirectly or directly change chromatin marks. This has recently
15 launched re-exploration of anti-inflammatory bioactive food components for characterization
16 of their effects on epigenome modifying enzymatic activities (acetylation, methylation,
17 phosphorylation, ribosylation, oxidation, ubiquitination, sumoylation). This may allow to
18 improve healthy aging by reversing disease prone epimutations involved in chronic
19 inflammatory and metabolic disorders.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

1 A large body of epidemiological and experimental data has demonstrated a direct link
2 between chronic inflammation and complex diseases such as obesity, allergy, asthma,
3 cardiovascular disease, type 2 diabetes, rheumatoid arthritis, bowel disease and several types
4 of cancer. Nutrition, as the main aspect of the environment, is said to play a key role in some
5 of this disparities, however, the molecular mechanisms involved still remain to be unraveled
6 [1-3]. Genome-wide association studies have identified hundreds of genetic variants
7 associated with complex human diseases and traits, and have provided valuable insights into
8 their genetic architecture. However, most variants identified so far confer relatively small
9 increments in risk, leaving many questions how the remaining 'missing' heritability can be
10 explained, although polygenic disease traits may account for some of this limitations [4, 5].
11 Furthermore, genetics *per se* cannot explain the vast diversity of phenotypes. A substantial
12 number of studies have shown that constraint in the early life environment is associated with
13 increased risk of cardiometabolic disease, affective disorders and cognitive decline,
14 osteoporosis, allergy, inflammation and specific cancers. Furthermore, genetics *per se* cannot
15 explain the vast diversity of phenotypes. The differences between identical twins and their
16 different susceptibility to most diseases has recently been attributed to epigenetic changes
17 which accumulate during life following exposure to different environmental conditions [6-8].
18 The contribution of epigenetic changes (epimutations) to human disease is probably
19 underestimated. Epigenetics encompasses several extra-genetic processes such as DNA
20 methylation (methylation of cytosines within CpG dinucleotides), histone tail modifications
21 (including acetylation, phosphorylation, methylation, sumoylation, ribosylation and
22 ubiquitination), noncoding RNA functions, regulation of polycomb group proteins and the
23 epigenetic cofactor modifiers, all of which may alter gene expression but do not involve
24 changes in the DNA sequence itself [9-13]. The combinatorial nature of DNA methylation
25 and histone modifications significantly extends the information potential of the genetic code.
26 In a large scale comparative study, authors concluded a lack of genetic factor in chronic
27 obstructive pulmonary disorder (COPD), cardiovascular disease (CVD), rheumatoid arthritis
28 and Crohn's disease, all typical examples of diseases with an important inflammatory
29 component [14]. A recent survey of the global incidence of cancer shows that the age-adjusted
30 cancer incidence in the Western world is above 300 cases per 100,000 population, whereas
31 that in Asian countries is less than 100 cases per 100,000. Sedentary lifestyle, diet, obesity
32 and metabolic syndrome have been named as the major contributors to this phenomena, which
33 is further emphasized by the increase in cancer cases among immigrants from Asian to

1 Western countries [15, 16]. As such, a reasonable good fraction of cancer deaths maybe
2 prevented by modifying the diet composition (i.e. content of fiber, fruit, vegetable, fat/oil,
3 protein, spices, cereals, xenobiotics) and regular physical exercise [3, 17]. Abnormalities in
4 DNA methylation, histone modifications, chromatin remodelling and microRNA (miR)
5 patterns [12, 18, 19] are important hallmarks of inflammatory disease states and cancer [10,
6 12, 13, 20-22].
7
8
9

10 The most studied epigenetic lesion, which is DNA hypermethylation at the promoter region
11 of many genes [20, 23], is proved to be responsible for silencing of more than 600 cancer-
12 related genes and this number is still rising. They include tumour suppressor genes, as well as
13 genes involved in the cell-cycle regulation, DNA repair, angiogenesis, metastasis and
14 apoptosis [24]. On the other hand, global hypomethylation of the DNA is said to activate
15 endoparasitic sequences and causes the global chromosome instability leading to various
16 mutations and cancer progression [21]. Now, it is becoming clear that epigenetic changes
17 complement genetic mutations and importantly drive the development and progression of
18 various diseases ranging from allergy, asthma, rheumatoid arthritis, type 2 diabetes, obesity,
19 bowel disease, cardiovascular disease and cancer [12, 13, 20, 21, 25-29]. Recent successes of
20 therapeutic intervention in chronic inflammatory diseases using epigenetic modifiers such as
21 histone deacetylase (HDAC) and DNA methyltransferase (DNMT) inhibitors has fuelled
22 interest in exploring epigenomic maps of inflammatory disease states [30, 31]. Interestingly,
23 emerging data demonstrate the direct influence of certain anti-inflammatory dietary factors
24 (for example polyphenols, isothiocyanates, epicatechins) and micronutrients (for example
25 folic acid, selenium) on heritable gene expression and DNA methylation or chromatin
26 remodelling [32-40]. Because epigenetic changes are reversible, developing drugs that control
27 epigenetic regulation now attract substantial research investment, including the development
28 of functional foods or supplements as nutrition based epigenetic modulators which will be
29 discussed in more detail below [2, 30, 41]
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 **Extragenomic epigenetic information**

50 The definition of ‘epigenetics’ evolved during the past 50 years into the current and common
51 definition of ‘heritable changes in gene function that cannot be explained by changes in DNA
52 sequence, although some definitions have excluded heritability [42]. While “inheritance” and
53 “heritability” strictly spoken, refer to intergenerational organismal phenomena, they are also
54 widely used today for describing both molecular and phenotypic characteristics at the cellular
55 level that are transmitted between mitotic cell division. Expression of our DNA genotype can
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

be twiddled by various volume knobs. Considerable cross-talk occurs between three main players in silencing: DNA methylation, histone modifications and noncoding RNAs. The chromatin template therefore serves as the major platform for epigenetic regulation in the form of DNA methylation, histone modifications and histone variants. More recently, an epigenetic role for RNA has also been demonstrated in normal situations as well as in disease [12]. Before most activators of a gene access their DNA-binding sites, a transition from a condensed ("solenoid-like fiber") to a decondensed ("beads on a string") chromatin structure appears to take place. Conversely, the acquisition of a more condensed chromatin structure is often associated with gene silencing [11]. This structural restriction of chromatin on gene expression can be overcome by chromatin remodeling cofactor complexes, which reversibly modify (acetylation, phosphorylation, ubiquitylation, glycosylation, sumoylation) on lysine, arginistine, serine or threonine residues of amino-terminal histone tails. In general, DNA is wrapped around nucleosomes, which are arranged as regularly spaced beads (146 bp DNA/nucleosome) along the DNA. Typically, nucleosomes consist of a histone octamer of histones (H)2A/B, H3 and H4. The DNA bridging two adjacent nucleosomes is normally bound by the linker histone H1 and is termed linker DNA. While the core histones are bound relatively tightly to DNA, chromatin is largely maintained by the dynamic association with its architectural proteins. Since the discovery of histone modifying enzymes, N-terminal histone tails protruding from nucleosomes were found to be 'velcro patches' for (de)acetylases (HDAC/HAT), (de)methylases (HMT/HDMT), ubiquitin ligases, small ubiquitin-related modifier (SUMO) ligases, kinases, phosphatases, ribosylases, which together establish specific histone modification patterns involved in transcription [11]. Specific sets of histone modifications and/or variants are associated with genes that are actively transcribed or are repressed, a phenomenon defined as the "histone code" [11]. Since the modifications do not involve underlying changes in nucleotide or amino acid sequences, they are "above the genome" and as such termed "epigenomic". However, to establish specificity of epigenetic marks, histone modifying complexes have to be recruited to relevant genomic locations by DNA-binding proteins, RNAs or protein-RNA complexes that bind to their specific DNA sites as a consequence of their own binding specificities and cellular concentrations [43-46]. It cannot come from the enzymatic activities per se as neither DNMTs, nor enzymes which modify histones know which part of the genome needs to be tagged. Furthermore, there is now a large body of evidence showing that modifications of the histone tails provide signals ("binary switches") that are recognized by specific binding proteins, such as chromo-, bromo- or tudor-domains which in turn can influence gene expression and other chromatin functions

[47-49]. The spatial and time-dependent combinations of histone modifications further increase the complexity of information contained in chromatin [11, 48, 49].

DNA methylation is the best-known epigenetic mark [20, 50]. It is catalyzed by two types of DNMTs: DNMT1 is a maintenance methyltransferase, whereas both DNMT3A and DNMT3B are *de novo* methyltransferases [51, 52]. The role of DNMT2 in DNA methylation is minor, its enzymology being largely directed to tRNA. DNA methylation is normally associated with gene inactivation and it usually occurs in CpG dinucleotides. Alternatively, DNA methylation of transcription factor binding sites which prevents binding of repressor proteins, may paradoxically induce gene activation. CpGs are normally methylated when scattered throughout the genome, but are mostly unmethylated when they are clustered as CpG islands at 5' ends of many genes. Hypermethylation of CpG-rich promoters triggers local histone code modifications resulting in a cellular camouflage mechanism that sequesters gene promoters away from transcription factors and results in stable silencing of gene expression. DNA methylation at CpG dinucleotides occurs upon transfer of S-adenosylmethionine (SAM) on cytosine by DNMTs. Whereas DNMT3A/B are responsible for DNA methylation during development (differentiation), DNMT1 is in charge of maintaining DNA methylation patterns in DNA replication during cell division. In mammalian cells, the fidelity of maintenance of methylation is 97–99.9% per mitosis, whereas *de novo* methylation is as high as 3–5% per mitosis, thus creating possibilities for epigenetic changes. DNA methylation also regulates genomic imprinting [53], X-chromosome inactivation [54] and silencing of repetitive sequences [55]. Although in most cases DNA methylation is a stable epigenetic mark, reduced levels of methylation can also be observed during development. This net loss of methylation can either occur passively by replication in the absence of cunitional maintenance methylation pathways, or actively, by removing methylated cytosines. In plants active demethylation is achieved in plants by DNA glycosylase activity, probably in combination with the base excision repair pathway. In mammals, coupling of 5-methylcytosine deaminase and thymine DNA glycosylase activities maybe responsible for DNA demethylation. Alternatively, a role for the 5-hydroxymethylcytosine modification in mammalian DNA demethylation has also been proposed as an intermediate in an active DNA demethylation pathway involving DNA repair and 5-hydroxymethylcytosine-specific DNA glycosylase activity [52].

Although DNA methylation is the best-known epigenetic mark [51, 56, 57], DNA methylation does not act alone. It occurs in the context of nucleosome positioning and histone modifications [9, 58, 59]. For example, high resolution DNA methylation analysis has

1 revealed 10-base periodicities (i.e one helical turn) in the DNA methylation status of
2 nucleosome-bound DNA and found that nucleosomal DNA was more highly methylated than
3 flanking DNA [59]. These data revealed that nucleosome positioning influences DNA
4 methylation patterning of promoters and intron-exon boundaries throughout the genome and
5 that DNA methyltransferases preferentially target nucleosome-bound DNA. Whether
6 nucleosome strings provide a combinatorial histone code is a matter of debate [9, 11, 12, 48,
7 60-62], but in any event, histone modifications influence gene activity and regulation. For
8 example, acetylation of lysines is generally associated with transcriptional activation whereas
9 lysine methylation can dictate either activation (e.g. H3K4, H3K36, H3K79) or suppression
10 (e.g. H3K9, H3K27 or H4K20). Specific histone modifications have been shown to be
11 associated with DNA hypermethylation of CpG islands, including deacetylation of histones
12 H3 and H4, loss of H3K4me, and gain of H3K9me3 and H3K27me3 [63, 64]. More
13 importantly, specific histone modifications have been implicated in several diseases, such as
14 cancer and CVD, while recently, evidence for modifications in allergic asthma and COPD is
15 also emerging [21, 25, 65-68]. DNA methylation marks are recognized by DNA methyl-
16 binding proteins (MBD) which can interact with corepressor-associated enzymes (i.e.
17 HDACs, enhancer of zeste homologue (EZH)2, ...), thus further linking DNA methylation and
18 chromatin regulation [43, 69]. Altogether, "histone code" may only become biologically
19 meaningful at the level of the chromatin fiber ("chromatin regulatory code") which, upon
20 integration of conformations of multiple nucleosomes, translates allosteric changes into
21 specific gene (cluster) activities, in order to establish specific regulatory programs at the
22 genome level [11, 70-72]. In analogy to allosteric control of enzymes, specific gene activity
23 may be determined by the spatial organization (compartmentalization in discrete territories)
24 and structural landscape (three-dimensional structure) of a gene locus, by altering the higher
25 order structure of chromatin (cis mechanism) or by generating a binding platform for effector
26 proteins (trans mechanisms) [70, 71, 73, 74].

27 There is good evidence that also noncoding RNAs regulate chromatin architecture [12, 75-
28 80]. The term noncoding RNA (ncRNA) is commonly employed for RNA that does not
29 encode a protein. Although it has been generally assumed that most genetic information is
30 transacted by proteins, recent evidence suggests that the majority of the genomes of mammals
31 and other complex organisms is in fact transcribed into ncRNAs, many of which are
32 alternatively spliced and/or processed into smaller products. Besides tRNA and rRNA, these
33 ncRNAs include long-noncodingRNAs (lncRNAs), microRNAs (miRNAs) and tinyRNAs
34 (tiRNAs) as well as several other classes of, sometimes yet-to-be-discovered, small regulatory
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

RNAs such as snoRNAs [75-77]. These RNAs (including those derived from introns) appear to comprise a hidden layer of internal signals that control various levels of gene expression in physiology and development, including chromatin architecture/epigenetic memory, transcription (enhancer function), RNA splicing, editing, translation and turnover [81]. RNA regulatory networks may determine most of our complex characteristics and play a significant role in disease [81]. For example, miRNAs can change expression levels of the epigenetic machinery (DNMT, HDAC, sirtuin (SIRT), polycomb (Pc) proteins, etc.) by posttranscriptional gene regulation involving base pairing with 3'untranslated (UTR) regions in their target mRNAs resulting in mRNA degradation or inhibition of translation [12, 18, 19, 82]. Alternatively, long ncRNAs and tiRNAs can regulate gene expression and/or DNA methylation by promoter association [75, 76, 81]. DNA-methylation can thus also be RNA-directed [12, 83].

Inflammatory and metabolic stress fuel epigenetic plasticity

Inflammation is a fundamental adaptation to the loss of cellular and tissue homeostasis covering many important processes, including host defense, tissue remodeling and repair, and the regulation of metabolism. The complexity of the inflammatory response requires that its many functional programs are controlled coordinately in some situations but independently in others [84, 85]. This is achieved through multiple mechanisms that operate at different levels, including alterations in the composition of immune cells in tissues, changes in cell responsiveness to inflammatory stimuli, regulation of signaling pathways and epigenetic control of gene expression.

Cell-specific mechanisms operate at the level of different cell types, and include regulation of their recruitment and activation. The cellular component involves the movement of leukocytes from blood vessels into the inflamed tissue, while leukocytes are involved in the initiation and maintenance of inflammation [84, 86]. Macrophages participate in host defense, immunity and inflammatory responses, where they are potently activated resulting in the production of pro-inflammatory cytokines, oxygen and nitrogen species and eicosanoids. Acute inflammation is mediated by granulocytes or polymorphonuclear leucocytes, while chronic inflammation is mediated by mononuclear cells such as monocytes and macrophages which can be further stimulated to maintain inflammation through the action of an adaptive cascade involving dendritic cells, T- and B-lymphocytes and antibodies [84, 87-92]. Imbalances in control of haematopoiesis and lineage differentiation of Th1, Th2 and Treg cells have a major impact on prevalence of sensitization to allergens and allergic diseases and has

1 reached epidemic proportions in Western societies [93-96]. Furthermore, tumors are typically
2 infiltrated with immune cells and inflammation which impacts on most, if not all, stages of
3 tumorigenesis [97-100]. Moreover, most cancers contain an inflammatory infiltrate that is
4 hijacked by tumor cells to promote angiogenesis, tissue invasion and cell proliferation. What
5 is more, overnutrition and obesity activate the immune system which at long-term switches to
6 chronic inflammatory condition that is as fertile soil for cancer development [3, 101-106].

7
8
9
10 Signal-specific mechanisms operate at the level of pathways which for example activate the
11 key transcription factor nuclear factor- κ B (NF κ B). Among all the mediators and cellular
12 effectors of inflammation, NF κ B is perhaps the central transcription factor, which regulates
13 expression of more than 400 genes [107-109]. At the same time, it is responsible for many
14 aspects of inflammatory disease and malignancy by inducing transcription of soluble
15 mediators that amplify inflammation, angiogenesis and neoplastic cell proliferation, and affect
16 progression to more aggressive disease states [97]. Members of the NF κ B family of dimeric
17 transcription factors (TF) regulate expression of a large number of genes involved in immune
18 responses, inflammation, metabolic stress, cell survival, and cancer. NF κ B family TF are
19 rapidly activated in response to various stimuli, including cytokines, infectious agents,
20 overnutrition (metabolic stress, endoplasmic reticulum stress) or danger signals (bacteria,
21 viruses, chemicals, pathogen associated molecular patterns (PAMPs), danger associated
22 molecular patterns (DAMPs), and radiation-induced DNA double-strand breaks. In
23 nonstimulated cells, some NF κ B TFs are bound to inhibitory (I) κ B proteins and are thereby
24 sequestered in the cytoplasm. Activation leads to phosphorylation of I κ B proteins and their
25 subsequent recognition by ubiquitinating enzymes. The resulting proteasomal degradation of
26 I κ B proteins liberates NF κ B TF, which translocate to the nucleus to drive expression of target
27 genes. Two protein kinases with a high degree of sequence similarity, I κ B kinase (IKK) α and
28 IKK β , mediate phosphorylation of I κ B proteins and represent a convergence point for most
29 signal transduction pathways leading to NF κ B activation. Most of the IKK α and IKK β
30 molecules in the cell are part of IKK complexes that also contain a regulatory subunit called
31 IKK γ or NF κ B-essential modulator (NEMO). Several years ago, two IKK-related kinases,
32 called IKK ϵ and TBK1 (TANK-binding kinase), were identified that exhibit structural
33 similarity to IKK α and IKK β [110]. Together, the IKKs and IKK-related kinases are
34 instrumental for inducible activation of the host defense system and controlling metabolic
35 stress [103, 105, 107, 111, 112]. Alternative to IKK, various additional kinases have been
36 identified which modulate transcriptional nuclear activity of NF κ B, including mitogen- and
37 stress-activated protein kinase (MSK), protein kinase (PK)Ac, phosphoinositide 3-kinases
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

PI3K and AKT [109, 113-116]. Furthermore, constitutive activity of NF κ B/IKK has been observed in many cancer cells, inflammatory disorders, obesity and insulin resistance [102, 103, 105-108, 112, 117-119]. During obesity, the organism needs to adapt to and function under chronic exposure to high energy and nutrient intake. To cope with these challenges, the cell has developed the endoplasmic reticulum (ER) as a key nutrition sensor of cellular metabolic parameters (hyperglycemia, fatty acid overload, hypoglycemia, oxidative stress), which participates in virtually all anabolic and catabolic branches. Failure of the ER's adaptive capacity results in activation of the unfolded protein response (UPR), which intersects with many different inflammatory and stress signaling pathways at the crossroad of inflammation, cancer and metabolic disease [17, 105, 106].

Gene-specific mechanisms operate at the level of individual genes and gene subsets. Induction of inflammatory transcriptional responses is orchestrated by many transcription factors and extragenic noncoding RNAs acting on inflammatory enhancers consistent with the complexity of the response [120-124]. Controlled expression of cytokine genes is an essential component of an immune response and is crucial for homeostasis. In order to generate an appropriate response to metabolic stress or an infectious condition, the type of cytokine, as well as the cell type, dose range and the kinetics of its expression are of critical importance [125-127]. The NF κ B family TF has a crucial role in rapid responses to metabolic stress and pathogens (innate immunity), as well as in development and differentiation of immune cells (acquired immunity). Although quite a number of genes contain NF κ B-responsive elements in their regulatory regions, their expression pattern can significantly vary from both a kinetic and quantitative point of view [10, 84, 107, 112, 125, 128-131]. At the transcription level, selectivity is conferred by the expression of specific NF κ B subunits and their respective posttranslational modifications, and by combinatorial interactions between NF κ B and other transcription enhancer factors (for example interferon regulatory factor (IRF)3, activating transcription factor (ATF)3, cAMP response element-binding protein CREB, CCAAT-enhancer binding protein C/EBP, activator protein (AP)1 (i.e. jun, fos, Fra1), Sp1, Stat3, PU.1) [109, 121, 132]. In addition to NF κ B, AP1 TF are closely involved in inflammatory disease and cancer invasive gene expression programs [133, 134]. Inflammatory transcription factors can be divided into different categories on the basis of their mode of activation and function [84]. Primary response transcription factors are constitutively expressed by many celltypes in the cytoplasm and are activated by signal-dependent posttranslational modifications, which involves their nuclear translocation, such as NF κ B, IRF, CREB. This TF are mainly responsible for the primary phase of gene induction and integrate signals from

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

diverse signaling pathways which can amplify or terminate signal-dependent transcription factor activation. Another class TF requires de novo synthesis following inflammatory stimulation, for example C/EBP δ . Most are constitutively nuclear and regulate secondary waves of gene expression. A third class of constitutively nuclear TF are expressed in a cell type-specific and differentiation-dependent manner, such as Runx, PU.1, IRF8, AP1, C/EBP [121, 128]. They establish cell type-specific patterns of gene expression and are involved in chromatin remodelling during cell differentiation and organization of higher-order chromatin structure and chromosomal domains. The transcription factors of these TF categories do not act independently, but function coordinately to control the inflammatory transcriptional response. Upon combining datasets of expression profiling of inflammatory genes and *in silico* motif scanning of promoters of these genes they can define gene clusters that are coordinately regulated and the transcription factors that are likely to control their expression [10, 84, 107, 109, 112, 125, 128-131]. Since TF bind very poorly or not at all to nucleosomal DNA, their activation is coordinated to recruitment of ATP-dependent chromatin-remodeling factors (swith/sucrose non fermentable (SWI/SNF), Brahma (Brm), brahma-related gene, (Brg1)), histone-enzyme complexes such as kinases (IKK, MSK, ataxia telangiectasia mutated (ATM), AKT, PI3K), poly(ADP-ribose) polymerase (PARP), methylases (EZH2, coactivator-associated arginine methyltransferase (CARM)1, protein arginine methyltransferases (PRMT)), demethylases (lysine specific demethylase (LSD)1, Jumonji C family histone demethylase (JMJD)3), prolyl isomerase (PIN1), acetylases (p300, CREB binding protein (CBP), p300/CBP associated factor (p/CAF)), deacetylases (HDAC, SIRT) and DNMTs [10, 43, 107, 114, 135, 136]. Parallel posttranslational modifications (phosphorylation, acetylation, methylation, ribosylation, sumoylation, ubiquitination) of histone and non-histone TF and cofactor complexes allow formation of dynamic enhanceosome complexes which establish a distinct chromatin structure. These epigenetic settings are the ultimate integration sites of both environmental and differentiative inputs, determining proper expression of each inflammatory gene [10, 107, 131, 137, 138].

49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Further investigation of epigenetic regulation of inflammatory genes, revealed different different subclasses according to chromatin activation mode and gene expression profile [128, 139]. A major class of primary response genes is characterized by CpG-island promoters, which facilitate promiscuous induction from constitutively active chromatin without a requirement for SWI/SNF nucleosome remodeling complexes. The low nucleosome occupancy at promoters in this class can be attributed to the assembly of CpG-islands into unstable nucleosomes, which may lead to SWI/SNF independence. Another major class

1 consists of non-CpG-island promoters that assemble into stable nucleosomes, resulting in
2 SWI/SNF dependence and the requirement for transcription factors that promote selective
3 nucleosome remodeling. Some inflammatory stimuli such as TNF, exhibit a strong bias
4 toward activation of SWI/SNF-independent CpG-island genes. In contrast interferon (IFN) β
5 preferentially activates SWI/SNF-dependent non-CpG-island promoters. Interestingly, by
6 activating a diverse set of transcription factors, Toll like receptors (TLR) induce both classes
7 and others for an optimal response to PAMPs and DAMPs [128]. Remarkably, DNA
8 methylation of IKK, I κ B and RelB promoters [140-142] adds another regulatory control level
9 which can act as transcription memory for repetitive pulsatile inflammatory exposures
10 (endotoxin sensitization versus tolerance) [140, 143-146].
11
12
13
14
15
16
17
18
19

20 **Chronic inflammatory disorders and epimutations: cause or consequence?**

21 Since inflammatory gene expression dynamics is highly dependent on epigenetic control
22 mechanisms [10, 84, 92, 120, 125, 137], we have compared chromatin organization in weak
23 or strong inflammatory cancer cell types with inducible or constitutive interleukin (IL)6 gene
24 expression patterns. Upon investigation of autocrine IL6 gene expression production in
25 aggressive myeloma cells or metastatic breast cancer cells, we observed euchromatin-like
26 properties and highly accessible chromatin at the IL6 gene promoter [133, 147]. Furthermore,
27 recruitment of CBP/p300 acetylases and MSK kinase seems to prevent heterochromatinisation
28 and recruitment of heterochromatin protein (HP)1 upon phosphacetylation of transcription
29 factor and histone components [114, 115, 132, 133, 148] (See Figure 2). Interestingly,
30 promoter binding activity of Sp1 and AP1 Fra1 are responsible for priming IL6 promoter
31 chromatin relaxation, which further promotes binding of NF κ B transcription factors and
32 chromatin opening for maximal expression levels [133, 147]. Interestingly, complementation
33 of low invasive cancer cells with Fra1 seems to convert the promoter chromatin architecture
34 in a highly accessible chromatin configuration. Reciprocally, highly accessible chromatin in
35 invasive cancer cells can be silenced with anti-inflammatory phytochemicals, or following
36 silencing of AP1/NF κ B transcription factors, demonstrating reversible epigenetic changes
37 towards a less aggressive phenotype [133, 149, 150]. Along the same line, we and others
38 observed DNA hypermethylation at the IL6 gene promoter in cancer cells with low
39 NF κ B/AP1 activity and inducible IL6 gene expression, as compared to DNA hypomethylation
40 in cancer cells with hyperactivated NF κ B/AP1 transcription factors and elevated constitutive
41 IL6 gene expression ([151, 152] and Figure 2). Similarly, p53 knockout cells reveal defects in
42 genomic imprinting and DNA methylation regulation [153] As such, this demonstrates that
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
inflammatory signaling and transcription factors are able to rewire epigenetic settings and amplify gene expression in an autocrine fashion [46, 154]. As another example, DNA methylation of the NFκB responsive element in the Fas (CD95, Apo-1, TNFRSF6) gene promoter, was found to silence its expression in metastatic prostate carcinoma [46, 155]. Surprisingly, depletion of NFκB can also trigger DNA demethylation and gene reactivation of the Fructose-1,6-biphosphatase-1 (FBP1) illustrating gene-specific epigenetic effects which may further depend on posttranslational NFκB modifications [136, 156].

31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Of special note, whereas studies with epigenetic drugs (azacytidin, suberoylanilide hydroxamic acid (SAHA)) frequently focus on reactivation of silenced tumor suppressor genes, these compounds also boost gene expression of inflammatory oncogenes such as IL6 which promote aggressive carcinogenesis, cancer stem cell proliferation, metastasis and hormone resistance [100, 157-164]. Furthermore, elevated IL6 gene expression is able to trigger epigenetic changes of tumor suppressor genes via regulation of DNMTs [165-168], microRNAs [158, 169, 170] and histone methyltransferases (Ezh2) [171]. This suggests that epigenetic regulators themselves and methylation of tumor suppressor genes are also susceptible to dynamic inflammatory control [158, 165, 169, 170, 172-175], which adds an extra level of complexity to the cancer-inflammation link.

66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
215

1 which in turn augments the state of inflammation. In addition, reactive nitrogen intermediates
2 such as NO and ROS, indirectly also modulate activity of epigenetic machinery which finally
3 will affect chromatin dynamics in tumor-associated immune cells [186-190].
4

5 One of the key functional characteristics of macrophages is that they can be programmed to
6 deal most effectively with a given kind of inciting stimulus. In the lung for example, resident
7 alveolar macrophages are continuously encountering inhaled substances due to their exposed
8 position in the alveolar lumen [93]. To avoid collateral damage to mucosal epithelium in
9 response to harmless antigens, they are kept in a quiescent state, producing little inflammatory
10 cytokines and displaying poor phagocytic activity. In the vessel wall, macrophages can
11 scavenge large amounts of lipid, and differentiate into lipid-laden foamy cells [179]. As
12 putative precursors for microglia and osteoclasts, monocytes may also be involved in the
13 physiology of the central nervous system and in bone remodelling. Dendritic cells (DC) are
14 also part of the monocytic phagocytic system and are the professional antigen presenting cells
15 of the immune system that bridge innate and adaptive immunity [91]. DC cells come in many
16 flavours and it is clear now that there are multiple pathways to develop DC from monocytes.
17 The complexity of the monocytic phagocytic system requires that its many functional
18 programmes are controlled coordinately in some situations, but separately in others. It is now
19 clear that epigenetic mechanisms can lead to time-dependent and stimulus-specific alterations
20 in the expression of functional modules [191]. Epigenetic modifications like histone
21 modifications and CpG-island methylation in promotor regions of critical transcription factors
22 (so called masterswitch transcription factors) of all of these functional programmes have been
23 described and are controlled by microRNAs, DNMTs, HDAC, histone acetylases (HAT
24 CBP/p300, p/CAF) histone lysine methylases (EZH2, CARM, PRMT) and demethylases
25 (LSD1, JMJD3) [192, 193]. Emerging research indicates that epigenetic mechanisms that
26 control the lineage development and polarization of monocytic phagocytic system and Th1,
27 Th2 and Treg cells can be transmitted across multiple generations [84, 89, 194, 195].
28 Smoking, diesel exhaust, diet conditions, hygiene, dust mite, obesity seem have major
29 influence on epigenetic programming of immune cell populations and inflammatory disease
30 severity in asthma, allergy, rheumatoid arthritis, bowel disease [196-202]. Similarly,
31 hypertension, obesity and lack of physical activity can have longlasting effects on epigenetic
32 reprogramming of monocytes/macrophages and endothelial cells promoting differentiation
33 into more proatherogenic phenotypes and increased risk of cardiovascular disease [68, 180,
34 182-185, 203].
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Nutri-epigenomics: lifelong remodelling of our epigenomes by nutritional, phytochemical and metabolic factors

Human epidemiological studies and appropriately designed dietary interventions in animal models have provided considerable evidence to suggest that maternal nutritional imbalance and metabolic disturbances, during critical time windows of development, may have a persistent effect on the health of offspring and may even be transmitted to the next generation [29, 32, 204-208]. This has led to the hypothesis of “fetal programming” and new term “developmental origin of health and disease” (DOHaD): common disorders, such as obesity, cardiovascular disease (CVD), diabetes, hypertension, asthma, cancer and even schizophrenia, take root in early nutrition during gestation and continues during lactation [28, 32, 209-214]. The various non-Mendelian features of metabolic disease, cancer or chronic inflammatory disorders, clinical differences between men and women or monozygotic twins and fluctuations in the course of the disease are consistent with epigenetic mechanisms in the influence of fetal and/or lifelong nutrition or stochastic events on adult phenotype [29, 32, 204-208, 215, 216]. Thus, lifetime shapes the multitude of epigenomes not only within, but also across generations [29, 207]. Interest in transgenerational epigenetic effects of food components has been fueled by observations in Agouti mice fed with a soy polyphenol diet, which revealed epigenetic changes in DNA methylation patterns in their offspring and protected against diabetes, obesity and cancer across multiple generations [217-219]. However, only weak transgenerational effects could be observed with soy polyphenols in *Daphnia Magna*, despite the presence of functional DNMTs [220]. Feeding pregnant mice with a diet rich in methyl donors altered DNA methylation at well defined CpG regions which led to decreased transcription of Runx3 and more severe asthma in offspring. Bronchial biopsies and alveolar macrophages of asthmatics revealed a disturbed epigenetic regulation of gene expression (increased HAT and decreased HDAC2 activity) [221]. Furthermore, the honeybee (*Apis mellifera*) is probably the clearest example of induction of alternative phenotypes and epigenotypes by nutrition in early life [222]. Female bees are genetic clones. However, queens are distinct from workers in their morphology, capacity to reproduce, behavior, and longevity. The difference between the queen and worker castes lies in the exposure of the genetically identical larvae to royal jelly, an as yet incompletely defined mixture of proteins, amino acids, vitamins, lipids, and other nutrients [223-225].

Studies of human populations following famine have suggested that pathologies in later life are dependent on the timing of nutritional insult during pregnancy. Follow up of the Dutch Hunger Winter cohort showed that cardiovascular disease was more prevalent in offspring of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

mothers who were severely undernourished during the first trimester of their pregnancies in 1944-1945, as compared to those born to mothers whose pregnancies were more advanced at the time of nutritional insult [226-228]. Also paternal patterns of nutrition during the prepubertal growth period in children in Överkalix, in Sweden during the nineteenth century are associated with differential risk of early cardiovascular death in their grandchildren [229, 230]. Today, various different epigenetic changes have already been characterized which are involved in atherogenesis [28, 29, 65, 67, 231]. Hypercholesterolemia, obesity, hyperhomocysteinemia, high glucose are important CVD risk factors which are implicated in enhanced inflammatory signaling and long-lasting effects are driven by epigenetic reprogramming, promoting differentiation of monocytes/macrophages into more proatherogenic phenotypes [68, 181, 182, 232]. Recent evidence suggests that the pathogenetic role of hyperhomocysteinemia in vascular diseases might be mediated via Adenosyl-homocystein (Hcy) accumulation and DNA methylation. Hcy competes with S-adenosylmethionine (SAM; the methyl-group donor) for binding on DNMT, which may lead to passive loss of methylation in replicating DNA. High blood Hcy levels correlate with DNA hypomethylation and atherosclerosis and can lead to a 35% reduction in DNA methylation status of peripheral blood lymphocytes [233-236]. Similarly, insulin, glucose folate or flavanol-rich diets interfere with the methyl-donor metabolism and the available pool of S-adenosylmethionine, resulting in DNA methylation changes [236-239]. In contrast, very few studies have focused on impact of dietary methyl-donors on histone methylation, which is also affected by alterations in SAM/SAH ratios [233, 240]. As such, specific dietary classes of functional food maybe designed as therapeutic epigenetic modulators in lifestyle disease, such as metabolic disorders (diabetes), cardiovascular disease, asthma/COPD, rheumatoid arthritis. Epidemiologic and medical anthropological studies have indicated that flavanol-rich diets are inversely associated with cardiovascular risk [241-246]. Locus-specific DNA methylation changes, both hyper- and hypomethylation, also occur at the promoter level of several genes involved in the pathogenesis of atherosclerosis, such as extracellular superoxide dismutase (SOD), hormone receptors (glucocorticoid receptor (GR), estrogen receptor (ER), peroxisome proliferator-activated receptor (PPAR), arylhydrocarbon receptor, AhR, liver X receptor (LXR)), endothelial and inducible nitric oxide synthase (iNOS/eNOS), 15-lipoxygenase (LOX), fibroblast growth factor (FGF)2, hypoxia-inducible factor (HIF)1 α , myc, insulator CCCTC binding factor (CTCF) and metalloproteases (MMPs) [68, 183, 247-249]. In a proatherogenic murine model, DNA-methylation polymorphisms preceded the appearance of histological signs of atherosclerosis [181, 182]. Interestingly, involvement of the inducible

1 JMJD3 demethylase was demonstrated to regulate monocyte/macrophage transdifferentiation
2 programs, illustrating that developmental programs are plastic and monocyte lineage
3 differentiation is susceptible to inflammatory pathways and oxidative stress [120]. A role for
4 the JMJD1A demethylase was demonstrated in metabolic gene expression and obesity
5 resistance [250]. Furthermore, it was found that knockdown of the LSD1 demethylase
6 affected monocyte adherence in a proatherogenic diabetic mouse model [251]. This suggests
7 that LSD1 contribute to metabolic memory through long-term changes in gene expression via
8 alterations in chromatin structure [146, 252].

14 Poor maternal nutrition has also been associated with increased risk of type 2 diabetes
15 mellitus over several generations in North American Indians [253, 254]. Individuals with
16 metabolic syndrome, obesity and type II diabetes (T2D) CVD may show a lifelong imbalance
17 between energy intake and energy expenditure due to incorrect epigenetic programming
18 during their early development as a result of placental insufficiency, inadequate maternal
19 nutrition, metabolic disturbances or neonatal medication [205, 253-258].

25 Recently, evidence emerged that also timing (preconception, pregnancy, lactation, neonatal
26 life, early life, pre-/post-menopause, puberty) of various dietary exposures may be vitally
27 important in determining health beneficial effects, as epigenetic plasticity changes continually
28 from conception to death [259]. In principle; epigenetic changes occurring during embryonic
29 development will have a much greater impact on the overall epigenetic status of the organism
30 because, as they can be transmitted over consecutive mitotic divisions, alterations occurring in
31 single embryonic stem cells will affect many more cells than those occurring in adult stem
32 and /or somatic cells during postnatal development [206]. Epigenetic plasticity further also
33 depends on other processes such as chromosomal instability, telomere shortening, metabolic
34 cycles, mitochondrial deteriorations, and oscillatory, circadian or seasonal rhythms of
35 systemic hormone levels (hypothalamic-pituitary-adrenal (HPA) axis) [28-30, 258-262]. In
36 addition to epigenetic imprinting during crucial periods of development, stochastic or
37 genetically and environmentally triggered epigenomic changes (epimutations) occur day after
38 day and accumulate over time, as maximal differences in DNA methylation profiles are
39 observed in aged monozygotic twins with a history of non-shared environments [6, 8].
40 Concerning nutritional transgenerational inheritance there is increasing evidence in both
41 plants and animals that, following nutritional intervention (caloric, iron and protein
42 restriction, polyphenol-, folate-, micronutrient-, fat- or carbohydrate-rich diet), maternal
43 diabetes, during pregnancy and lactation can affect the following generation(s) [38, 207, 211,
44 218, 219, 263, 264]. Although it has long been thought that the epigenomic profile is wiped
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 clean in the embryo shortly after fertilization, with the exception of imprinted genes,
2 methylation clearing is not complete after fertilization and on a global DNA level is reduced
3 to 10% [265, 266]. Alternatively, it can not be excluded that transgenerationally inherited
4 nutritional effects may also depend on polycomb proteins [207, 267-269], miRNA profiles
5 [12, 270] or epigenetic capacitor properties of hsp proteins [271-273].
6
7

8
9 A next challenge will be to determine which adverse epigenomic marks are reversible by
10 specific diets, drugs or lifestyle changes [29, 32, 38, 259]. Numerous botanical species and
11 plant parts contain a diverse array of polyphenolic phytochemicals which exert health-
12 beneficial effects in man by its anti-inflammatory, anti-oxidant, phytohormone, cardio-
13 protective, cancer preventive and anti-bacterial properties, by maintaining immune
14 homeostasis (hormesis) [274, 275]. Phytochemicals have also successfully been applied for
15 regenerative medicine and cancer stem cell therapy [173, 276-280]. Oxidative stress and
16 inflammatory damage plays an important role in epigenetic reprogramming of expression of
17 cytokines, oncogenes and tumor suppressor genes, thereby setting up a ground for chronic
18 inflammatory diseases and carcinogenesis [97, 100, 281]. As such chemoprevention, the
19 strategy to inhibit, retard, or even reverse the epigenetic stage of chronic inflammation is one
20 of the most rational approaches to reduce the global burden of non communicable lifestyle
21 diseases [3, 100, 211]. Today, various nutritional natural compounds (including
22 epigallocatechingallate, resveratrol, genistein, curcumin, isothiocyanates ...) have been
23 characterized which interfere with enzymatic activity of DNMT, Class I, II, IV HDAC, HAT
24 and Class III HDAC sirtuins (SIRT) which modulate inflammatory responses and
25 immunological senescence ([32-41, 213, 282] and references included, see Table 1). HDACs
26 are zinc metalloproteins which rely on Zn^{2+} for their activity and are divided into 4 classes
27 based on their homology with yeast HDACs. Class III HDACs, called sirtuins are zinc-
28 independent but nicotinamide adenine dinucleotide (NAD^+)-dependent. Class I, II, IV
29 HDAC inhibitors characteristically contain a Zn^{2+} chelating group consisting of a thiolate,
30 thiol, hydroxamate, carboxylate, mercaptoamide, epoxide or ketone group. Natural HDAC
31 inhibitors can be divided in following groups based on their chemical characteristics:
32 carboxylates, organosulfides, isothiocyanates, hydroamates, cyclic tetrapeptides and
33 macrocyclic depsipeptides [35]. In contrast to natural HDAC inhibitors, only few natural
34 products (i.e. niacine, vitamin B3, dihydrocoumarin) have been identified as inhibitors of
35 class III HDACs. Reciprocally, various natural flavonoids have been identified as activators
36 of class III HDACs (SIRTs). Finally, turmeric and green tea have been identified as sources of
37 natural inhibitors of p300/CBP HAT. Finally DNMT inhibitors work mainly through one of
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 the following mechanisms, either covalent trapping of DNMT through incorporation into
2 DNA (i.e. nucleoside analogues decitabine, 5-azacytidine), non-covalent blocking of DNMT
3 catalytic active site (i.e. EGCG, parthenolide), interruption of binding site of DNMT to DNA
4 (i.e. procaine), degradation of DNMT (i.e; decitabine), or suppression of DNMT expression
5 (i.e. miRNAs). Furthermore, a number of natural compounds act as multifunctional ligands by
6 simultaneously acting on nuclear hormone receptors and changing activity of histone
7 modifying enzymes and DNMTs [283-287]. Of special note, although health protective anti-
8 oxidant or anti-inflammatory effects of dietary factors and extracts have frequently been
9 demonstrated in *in vitro* experiments at concentrations which can never be achieved *in vivo*,
10 “epigenetics” sheds a more realistic light on dietary studies, as longlife exposure at
11 physiological concentrations can remodel the epigenome in a cumulative fashion by repetitive
12 effects on the epigenetic machinery [241, 288-291]. Particular attention needs to be given to
13 natural compounds which can trigger opposite effects on HDAC/HAT/DNMT or histone
14 (de)methylase (H(D)MT) depending on the concentration or cell type-specific metabolism
15 [34, 35]. It should also be stressed that it is not known whether all of them can be considered
16 authentic epigenetic modifiers because it has not yet been demonstrated whether the
17 epigenetic modifications that they induce are stable over time. Interestingly, even transient
18 exposure to a specific dietary component can induce long-lasting epigenetic changes in the
19 promoter of the NFκB subunit p65, which acts as a master switch in inflammatory gene
20 expression [146]. Alternatively, compounds may chemically interfere with histone mark
21 interacting effector domains (such as chromo-, bromo- or tudor domains) [47, 292, 293].
22 However, one should be careful with interpretation of *in vitro* compound screenings or
23 cofactor activity assays based on peptide-protein interactions, as this may not always
24 represent true targets *in vivo* [294, 295].

25 Besides specific interference of the diet with chromatin modifying enzymes and DNMTs at
26 particular target genes, global epigenetic changes can also occur following biochemical
27 metabolism of dietary factors, which can deplete cellular pools of acetyl-CoA, NAD⁺ and
28 methyl donors, resulting in unbalanced DNA methylation and/or protein acetylation or
29 methylation [282, 296, 297]. For example diets lacking in substrate or cofactors in
30 methyl donor metabolism can contribute to DNA hypomethylation by impairing synthesis of
31 SAM [234]. This methylation cycle is frequently cited to explain relations between diet and
32 epigenetic changes [233, 298]. However, even without nutritional deficiency of methyl
33 groups, impaired synthesis of SAM and perturbed DNA methylation can happen when the
34 need for the synthesis of the detoxification enzyme glutathione transferase (GSH) synthesis

1 increases [299]. Diets or nutritional compounds which affect energy metabolism or
2 mitochondrial respiration can have global epigenetic effects upon changes in NAD⁺
3 availability and SIRT activity [300]. Since SIRT activation has been linked to longevity
4 (increased lifespan and healthy aging) and mimics a caloric restricted diet, SIRT activators
5 such as resveratrol represent a major class of caloric mimetic epigenetic modulator
6 phytochemicals which could reverse metabolic disease [297].
7
8
9

10 11 12 **Conclusion**

13 The phenotype of an individual is the result of complex gene-environment interactions in the
14 current, past and ancestral environment, leading to lifelong remodelling of our epigenomes. In
15 recent years, several studies have demonstrated that disruption of epigenetic mechanisms can
16 alter immune function and and that epimutations not only contribute to various cancers but
17 also to the development of diabetes, allergy, CVD and rheumatoid arthritis. Various
18 replication-dependent and -independent epigenetic mechanisms are involved in
19 developmental programming, lifelong recording of environmental changes and transmitting
20 transgenerational effects. It is likely that understanding and manipulating the epigenome, a
21 potentially reversible source of biological variation, has great potential in chemoprevention or
22 stabilization of chronic inflammatory disorders. Much attention is currently focusing on
23 modulating hyper/hypomethylation of key inflammatory genes by dietary factors as an
24 effective approach to cure or protect against inflammatory disease [32-41, 213, 282]. In this
25 respect, “Let food be your epigenetic medicine” could represent a novel interpretation of what
26 Hippocrates said already 25 centuries ago. As such, it will be a challenge for future anti-
27 inflammatory therapeutics and preventive cancer research to identify novel epigenetic targets
28 which allow selective modulation of the inflammatory signaling network in the diseased
29 tissue and/or microenvironment [275, 301-306]. Given several encouraging trials, prevention
30 and therapy of age- and lifestyle-related diseases by individualised tailoring of optimal
31 epigenetic diets or supplements are conceivable. However, these interventions will require
32 intense efforts to unravel the complexity of these epigenetic, genetic and environment
33 interactions. Another goal is to evaluate their potential reversibility with minimal side effects
34 as diet components may reprogram malignant cells as well as the host immune system and
35 HPA-axis depending on the bioavailability of the dietary compounds [32, 133, 149, 150, 289,
36 291] (See Figure 3). There is some concern that epigenetic therapy with dietary inhibitors of
37 DNMT, HDAC, histon(de)methylases in longterm treatment setups may suffer from lack of
38 specificity [283, 292, 294]. As such, the possible alternative is to combine nonselective
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

epigenetic therapies with more targeted approaches [46]. For example, combined treatment of specific transcription factor inhibitors and/or hormone receptor ligands with epigenetic drugs may trigger synergistic activities at subsets of inflammatory genes [43, 46, 307-309]. An excellent example of cooperation between a dietary vitamin A-derivative targeting a nuclear receptor and the HDAC inhibitor butyrate has been described in the treatment of acute promyelocytic leukemias [33]. Finally, microRNA and long ncRNA pathways also hold promise to join soon the arsenal of epigenetic combination therapies, as their target sequence specificity may bridge the gap between genetic and epigenetic changes [12, 79-81]. In conclusion, studies are revealing a dazzling complexity in the mechanisms leading to dynamic alterations of the epigenome and the need of combination therapies targeting different chromatin modifiers, to reverse disease prone epigenetic alterations to preserve healthy aging. Medical benefits of dietary compounds as epigenetic modulators, especially with respect to their chronic use as nutraceutical agents, will rely on our further understanding of their epigenetic effects during embryogenesis, early life, aging as well as through different generations.

Acknowledgements

This research is partially financial supported by FP7-KBBE-2008-2B grant FLAVIOLA (“Targeted delivery of dietary flavanol for optimal human cell function: Effects on cardiovascular health (www.flaviola.org)), Research Grant from the Multiple Myeloma Research Foundation (MMRF) and Interuniversity Attraction Poles (IAP) P6/18. We regret that not all literature could be cited appropriately because of space constraints and we apologize to those authors whose original work is not mentioned.

Figure 1

1 Interplay of inflammatory and metabolic stress pathways in nutritional epigenetic effects in
2 lifestyle diseases. mut, mutation; SNP, single nucleotide polymorphism; miR, microRNA,
3 ncRNA, noncoding RNA.
4
5
6
7

Figure 2

8
9 Working model illustrating crosstalk of inflammatory and metabolic stress pathways with
10 epigenetic regulation in the nucleus. For example, transcription of the highly conserved IL6
11 gene promoter requires binding of activator protein AP1, CREB and nuclear factor NFκB
12 transcription factors in response to various stimuli [310, 311], with strict stereospecific
13 requirements for optimal cofactor recruitment, promoting a promoter enhanceosome model
14 with multiple transcription factor/cofactor interactions, in which NFκB is the primary trigger
15 for IL6 gene induction in response to TNF. Inflammatory stimuli (i.e. TNF, virus, bacteria) or
16 metabolic stress (including oxidative stress, endoplasmic reticulum (ER) stress or unfolded
17 protein response (UPR) stress, for example during obesity or malnutrition) can activate the
18 IKK-complex, responsible for IκB-phosphorylation, which allows NFκB to translocate to the
19 nucleus, following degradation of IκB in the proteasome. IL6 gene expression also requires
20 activation and recruitment of the mitogen- and stress-activated protein kinase-1 (MSK1) to
21 elicit selective chromatin relaxation at the IL6 gene promoter, upon phosphorylation of NFκB
22 p65 S276 and histone H3 S10, followed by further CBP/p300 recruitment and HAT-
23 dependent acetylation of the IL6 promoter enhanceosome/chromatin environment [115].
24 Highly elevated expression levels of AP1 Fra1 and NFκB in ER-deficient tumor cells increase
25 chromatin accessibility (DNA hypersensitive sites) in the proximal IL6 gene promoter. Of
26 particular interest, dietary anti-inflammatory phytochemicals such as soy polyphenol genistein
27 [149, 150, 274] and withanolide withaferin A [312] are able to silence the highly promiscuous
28 IL6 promoter nucleosome configuration by lowering the cellular amounts of Fra1 and NFκB
29 activity, concomitantly with decreased histone H3 phosphoacetylation levels and increased
30 DNA methylation [133, 149, 150].
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Figure 3.

52 Flowchart which summarizes the complex and multilayered regulation of nutritional
53 epigenetic effects.
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Overview of nutritional epigenetic effects by dietary phytochemicals (summarized from [32-41, 213, 282] and references included). Suffix I, inhibitor; suffix a, activator, for example HDACi, HDAC inhibitor, or SIRTa, SIRT activator

Bioactive Phytochemical	Natural Source	Disease Target	Epigenetic mechanism	<i>In vitro / In vivo</i>	Target genes / microRNAs
3,3-diindolylmethane	Broccoli	Cancer	Decreased HDAC levels	<i>In vitro / in vivo</i>	COX2
6-methoxy-2E,9E-humuladien-8-one	Ginger	Cancer	HDACi	<i>In vitro</i>	
Allylmercaptan	Garlic	Cancer	HDACi	<i>In vitro/in vivo</i>	P21/WAF
Anacardic Acid	Cashew nuts	Cancer, Leukemia	HATi	<i>In vitro</i>	
Apigenin	Parsley, celery	Cancer	DNMTi	<i>In vitro</i>	
Betainin	Beetroot red	Cancer	DNMTi	<i>In vitro</i>	
Biochanin A	Soy	Cancer	DNMTi HDACi	<i>In vitro / in vivo</i> <i>Daphnia</i>	
Butyric acid	Fermentation dietary fibers	Cancer	HDACi	<i>In vitro / In vivo</i>	
Caffeic acid	Coffea	Cancer Inflammation Energy metabolism	DNMTi HDACi	<i>In vitro</i>	RAR β , CDKN2A
Catechin	Green tea	Cancer Lymphocytes	DNMTi HATi	<i>In vitro</i>	RAR β
Chlorogenic acid	Coffea	Cancer	DNMTi HDACi	<i>In vitro</i>	RAR β , CDKN2A
Coumaric acid Cinnamic acid	Cinnamon	Cancer	DNMTi HDACi	<i>In vitro / in vivo</i>	
Coumarin analogues Dihydrocoumarin	Melilotus officinalis (Sweet clover)	Cancer, Leukemia,	SIRTi p53 acetylation	<i>In vitro</i>	
Curcumin	Turmeric (<i>Curcuma longa</i> L.)	Crohn's Disease, Ulcerative colitis Inflammatory disease Cancer, Leukemia Neurodegeneration Alzheimer Diabetes Heart Failure Epilepsy	DNMTi HATi HDACi	<i>In vitro / In vivo</i> <i>Plasmodium falciparum</i> , <i>Herpes Virus</i> <i>Mouse</i> <i>Rat</i>	GATA4, EOMES, GZMB, PRF1 Up: miR-22, miR-34a, miR-24, miR-181a, miR-21, miR-181b, miR-27a Down: miR-199a, miR-510, miR-196a, miR-7, miR-15b, miR-195, miR-374, miR-98
Cyanidin	Berries, Grapes	Cancer	DNMTi	<i>In vitro</i>	
Daidzein	Soy	Cancer	DNMTi SIRTa	<i>In vitro</i>	
Diallyl disulfide	Garlic	Inflammatory disease Cancer	HDACi	<i>In vitro / in vivo</i> <i>rat</i>	p21/WAF
Ellagic Acids	Berries	Cancer	DNMTi	<i>In vitro</i>	
Epicatechin	Apples, Cocoa, Green and black tea	Inflammatory disease, lymphocytes Cancer CVD	DNMTi HATi	<i>In vitro</i>	
Epicatechin gallate	Green tea	Cancer Lymphocytes	DNMTi HATi	<i>In vitro</i>	
Epigallocatechin	Green tea	Cancer Lymphocytes	DNMTi HATi	<i>In vitro</i>	
Epigallocatechin 3-gallate (EGCG)	Green tea	Inflammatory disease Parkinson Cancer, Leukemia CVD	DNMTi HDACi HMTi	<i>In vitro / In vivo</i> <i>Agouti</i> <i>Mouse/Human</i>	RAR β , MGMT, MLH1, CDKN2A, RECK, TERT, RXR α , CDX2, GSTP1, WIF1, NF κ B, IL6, Bmi1, Ezh2, Suz12 Up : let-7, miR-16,

		Diabetes Energy metabolism			miR-18b, miR-20a, miR-25, miR-92, miR-93, miR-221, miR-320 Down: miR-10a, miR- 18a, miR-19a, miR- 26b, miR-29b, miR-34b, miR-98, miR-129, miR-181d
Equol	Soy	Cancer	HDACi	<i>In vitro / in vivo</i> <i>Drosophila</i>	
Fisetin	Strawberries	Cancer Inflammatory disease	SIRTa	<i>In vitro</i>	
Flavone	Feijoa	Cancer	HDACi	<i>In vitro / in vivo</i> <i>Drosophila</i>	p16,p21,TRAIL
Folic Acid Folate	Leafy vegetables, Nuts, Sunflower seeds	Inflammatory disease Asthma Cancer Obesity CVD	Induction of DNA methylation	<i>In vitro / In vivo</i>	Up : miR-10a, miR- 10b, miR-9, miR-145, miR-30a-3p, miR-152, miR-122a, miR-125b Down: miR-200a, miR- 496, miR-296, miR- 30e-5p, miR-362, miR-339, miR-29c, miR-154, miR-10a
Galangin	Propolis	Cancer	DNMTi	<i>In vitro</i>	
Garcinol	Garcinia	Cancer HIV Leukemia Lymphocytes	DNMTi HATi	<i>In vitro</i>	Global downregulation gene expression
Genistein	Soy	Inflammatory disease Cancer	DNMTi HATi HDACi Increase DNA methylation Changed protein levels DNMTs, MBD1, MBD4, MeCP2	<i>In vitro / In vivo</i> <i>Agouti</i> <i>Daphnids</i>	RAR β , MGMT, CDKN2A, GSTP1, HMGN5, BTG3, TERT P21, p16,PTEN, CCLD, p53, FOXA3, SIRT1, BTG3, RAR β Up: miR-200b, miR- 200c, let-7b, let-7c, let- 7d, let-7e, miR-663, miR-146a, miR-374b Down: miR-34c, miR- 376a, miR-196a, miR- 320, miR-654, miR-34c, miR-196
Hesperidin	Citrus	Cancer	DNMTi	<i>In vitro</i>	
Isoliquiritigenin	Liquorice	Cancer	SIRTactivator	<i>In vitro / in vivo</i> <i>Drosophila</i>	
Isothiocyanate (6-Methylsulfinyl- hexylisothio- cyanate)	Broccoli, Japanese horseradish (Wasabi)	Anti-platelet effects Cancer	HDACi	<i>In vitro</i>	P21, GSTP1
Kaempferol	Apples, nuts, tea, onions	Cancer	SIRTa		
Luteolin	Parsley, celery	Cancer	DNMTi SIRTactivator	<i>In vitro / In vivo</i> <i>Drosophila</i>	
Lycopene	Tomatoes, Watermelon Apricots Pink guava Grapefruit, Rosehip	Cancer: Breast, Colon	DNMTi	<i>In vitro</i>	GSTP1, RAR β , HIN1
MCP30	Bitter melon seeds (Momordica charantia)	Cancer	HDACi	<i>In vitro</i>	
Myricetin	Berries	Cancer	DNMTi	<i>In vitro</i>	

Naringenin	Citrus	Cancer	DNMTi	<i>In vitro</i>	
Parthenolide	Feverfew (<i>Tanacetum parthenium</i>)	Arthritis Fever Lymphocytic leukemia Migraine	DNMTi HDAC depletion	<i>In vitro</i>	
Phloretin	Apples	Cancer	DNMTi	<i>In vitro</i>	
Piceatannol (Resveratrol metabolite)	Grapes, blueberries	Cancer	DNMTi SIRTa	<i>In vitro</i>	
Polyphenon B	Black and green tea	Cancer	Increased HDAC levels	<i>In vitro/in vivo</i> Rat	
Pomiferin	<i>Maclura pomifera</i>	Cancer	HDACi	<i>In vitro</i>	
Protocatechuric acid	Olives	Cancer	DNMTi	<i>In vitro</i>	
Quercetin	Citrus: Capers, Apples, Berries, Tea, Wine	Inflammatory disease Bowel inflammation Asthma Cancer CVD Pulmonary dysfunction	DNMTi HDACi SIRTactivator	<i>In vitro</i>	IL10, MIP2
Resveratrol	Grapes, Blueberries, Peanuts, Red wine	Inflammatory disease Cancer Leukemia CVD Neuroprotection	DNMTi HDACi SIRTa	<i>In vitro / In vivo</i> Yeast Mouse Rat <i>Drosophila</i>	TNF, IL8, RBP
Rosmarinic acid	Rosemary	Cancer	DNMTi	<i>In vitro</i>	
Retinoic Acid	Carrots, Spinach, Eggs	Cancer Leukemia	HDACi	<i>In vitro</i>	
S-allylmercapto-cysteine	Garlic	Cancer Leukemia	HDACi		
Sanguinarine	Opium poppy	Cancer	HMTi	<i>In vitro</i>	
Selenium	Cereals, Nuts, Legumes, Fish, Shellfish,	Cancer	DNMTi Induction of DNA methylation	<i>In vitro</i>	
Silibinin	Milk thistle (<i>Silybum marianum</i>)	Cancer Liver protection Inflammatory disease	Increased histone acetylation SIRTa	<i>In vitro</i>	p21, p27, CASP3, CASP9
Sinapic acid	Sinapis (mustard)	Cancer	DNMTi	<i>In vitro</i>	
Sulforaphane	Broccoli	Inl Cancer	DNMTi HDACi Decreased DNMT levels	<i>In vitro / In vivo</i> Mouse	RAR β , HBD2, p21, BAX
Syringic acid	Red Grape	Cancer	DNMTi	<i>In vitro</i>	
Theophylline	Black and green tea	Immune cells Pulmonary inflammation Asthma, COPD Cancer	HDACa	<i>In vitro</i>	
Ursolic acid	Basil	Leukemia	HDAC inhibitor	<i>In vitro</i>	
Withaferin A	Ashwagandha (<i>Withania somnifera</i>)	Anti-angiogenic effects Inflammatory disease Rheumatoid arthritis Asthma Cancer Leukemia Neuroprotection	Inhibition histone acetylation Chromatin silencing	<i>In vitro</i>	

References

- 1
2
3
4 [1] Boffetta P, Couto E, Wichmann J, Ferrari P, Trichopoulos D, Bueno-de-Mesquita HB,
5 et al. Fruit and vegetable intake and overall cancer risk in the European Prospective
6 Investigation into Cancer and Nutrition (EPIC). *Journal of the National Cancer*
7 *Institute* 2010;102:529-37.
- 8 [2] Bingham S, Riboli E. Diet and cancer--the European Prospective Investigation into
9 Cancer and Nutrition. *Nature reviews* 2004;4:206-15.
- 10 [3] Anand P, Kunnumakkara AB, Sundaram C, Harikumar KB, Tharakan ST, Lai OS, et
11 al. Cancer is a preventable disease that requires major lifestyle changes. *PharmRes*
12 2008;25:2097-116.
- 13 [4] Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, et al.
14 Finding the missing heritability of complex diseases. *Nature* 2009;461:747-53.
- 15 [5] Maher B. Personal genomes: The case of the missing heritability. *Nature* 2008;456:18-
16 21.
- 17 [6] Fraga MF, Ballestar E, Paz MF, Ropero S, Setien F, Ballestar ML, et al. Epigenetic
18 differences arise during the lifetime of monozygotic twins. *Proc Natl Acad Sci U S A*
19 2005;102:10604-9.
- 20 [7] Fraga MF, Esteller M. Epigenetics and aging: the targets and the marks. *Trends Genet*
21 2007;23:413-8.
- 22 [8] Christensen BC, Houseman EA, Marsit CJ, Zheng S, Wrensch MR, Wiemels JL, et al.
23 Aging and environmental exposures alter tissue-specific DNA methylation dependent
24 upon CpG island context. *PLoS genetics* 2009;5:e1000602.
- 25 [9] Lee BM, Mahadevan LC. Stability of histone modifications across mammalian
26 genomes: implications for 'epigenetic' marking. *J Cell Biochem* 2009;108:22-34.
- 27 [10] Vanden Berghe W, Ndlovu MN, Hoya-Arias R, Dijsselbloem N, Gerlo S, Haegeman
28 G. Keeping up NF-kappaB appearances: epigenetic control of immunity or
29 inflammation-triggered epigenetics. *Biochem Pharmacol* 2006;72:1114-31.
- 30 [11] Chi P, Allis CD, Wang GG. Covalent histone modifications--miswritten,
31 misinterpreted and mis-erased in human cancers. *Nature reviews* 2010;10:457-69.
- 32 [12] Guil S, Esteller M. DNA methylomes, histone codes and miRNAs: tying it all
33 together. *Int J Biochem Cell Biol* 2009;41:87-95.
- 34 [13] Davalos V, Esteller M. MicroRNAs and cancer epigenetics: a macrorevolution. *Curr*
35 *Opin Oncol* 2010;22:35-45.
- 36 [14] Plomin R, Owen MJ, McGuffin P. The genetic basis of complex human behaviors.
37 *Science* 1994;264:1733-9.
- 38 [15] Messina M, Hilakivi-Clarke L. Early intake appears to be the key to the proposed
39 protective effects of soy intake against breast cancer. *Nutr Cancer* 2009;61:792-8.
- 40 [16] Shu XO, Zheng Y, Cai H, Gu K, Chen Z, Zheng W, et al. Soy food intake and breast
41 cancer survival. *JAMA* 2009;302:2437-43.
- 42 [17] Tennant DA, Duran RV, Gottlieb E. Targeting metabolic transformation for cancer
43 therapy. *Nature reviews* 2010;10:267-77.
- 44 [18] Lujambio A, Calin GA, Villanueva A, Ropero S, Sanchez-Cespedes M, Blanco D, et
45 al. A microRNA DNA methylation signature for human cancer metastasis. *Proc Natl*
46 *Acad Sci U S A* 2008;105:13556-61.
- 47 [19] Lujambio A, Esteller M. CpG island hypermethylation of tumor suppressor
48 microRNAs in human cancer. *Cell Cycle* 2007;6:1455-9.
- 49 [20] Esteller M. Cancer epigenomics: DNA methylomes and histone-modification maps.
50 *Nat Rev Genet* 2007;8:286-98.
- 51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- [21] Esteller M. Epigenetics in cancer. *N Engl J Med* 2008;358:1148-59.
- [22] Backdahl L, Bushell A, Beck S. Inflammatory signaling as mediator of epigenetic modulation in tissue-specific chronic inflammation. *Int J Biochem Cell Biol* 2009;41:176-84.
- [23] Mulero-Navarro S, Esteller M. Chromatin remodeling factor CHD5 is silenced by promoter CpG island hypermethylation in human cancer. *Epigenetics* 2008;3:210-5.
- [24] Herceg Z. Epigenetics and cancer: towards an evaluation of the impact of environmental and dietary factors. *Mutagenesis* 2007;22:91-103.
- [25] Miller RL, Ho SM. Environmental epigenetics and asthma: current concepts and call for studies. *Am J Respir Crit Care Med* 2008;177:567-73.
- [26] Villeneuve LM, Natarajan R. The Role of Epigenetics in the Pathology of Diabetic Complications. *Am J Physiol Renal Physiol* 2010.
- [27] Schwartz DA. Epigenetics and environmental lung disease. *Proc Am Thorac Soc* 2010;7:123-5.
- [28] Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med* 2008;359:61-73.
- [29] Godfrey KM, Gluckman PD, Hanson MA. Developmental origins of metabolic disease: life course and intergenerational perspectives. *Trends Endocrinol Metab* 2010;21:199-205.
- [30] Moving AHEAD with an international human epigenome project. *Nature* 2008;454:711-5.
- [31] Altucci L, Stunnenberg HG. Time for epigenetics. *Int J Biochem Cell Biol* 2009;41:2-3.
- [32] Burdge GC, Lillycrop KA. Nutrition, Epigenetics, and Developmental Plasticity: Implications for Understanding Human Diseases. *Annual Review of Nutrition* 2010.
- [33] Delage B, Dashwood RH. Dietary manipulation of histone structure and function. *Annu Rev Nutr* 2008;28:347-66.
- [34] Link A, Balaguer F, Goel A. Cancer chemoprevention by dietary polyphenols: Promising role for epigenetics. *Biochem Pharmacol* 2010.
- [35] Folmer F, Orlikova B, Schnekenburger M, Dicato M, Diederich M. Naturally occurring regulators of histone acetylation/deacetylation. *Current Nutrition & Food Science* 2010;6:78-99.
- [36] Hauser AT, Jung M. Targeting epigenetic mechanisms: potential of natural products in cancer chemoprevention. *Planta Medica* 2008;74:1593-601.
- [37] Kontogiorgis C, Bompou E, Ntella M, Vanden Berghe W. Natural Products from Mediterranean Diet: From Anti-Inflammatory Agents to Dietary Epigenetic Modulators. *Anti-Inflammatory & Anti-Allergy Agents in Medicinal Chemistry* 2010;6:101-24.
- [38] Kirk H, Cefalu WT, Ribnicky D, Liu Z, Eilertsen KJ. Botanicals as epigenetic modulators for mechanisms contributing to development of metabolic syndrome. *Metabolism: Clinical and Experimental* 2008;57:S16-23.
- [39] Fang M, Chen D, Yang CS. Dietary polyphenols may affect DNA methylation. *The Journal of nutrition* 2007;137:223S-8S.
- [40] Suzuki T, Miyata N. Epigenetic control using natural products and synthetic molecules. *Curr Med Chem* 2006;13:935-58.
- [41] Arasaradnam RP, Commane DM, Bradburn D, Mathers JC. A review of dietary factors and its influence on DNA methylation in colorectal carcinogenesis. *Epigenetics* 2008;3:193-8.
- [42] Berger SL, Kouzarides T, Shiekhatar R, Shilatifard A. An operational definition of epigenetics. *Genes Dev* 2009;23:781-3.

- 1 [43] Perissi V, Jepsen K, Glass CK, Rosenfeld MG. Deconstructing repression: evolving
2 models of co-repressor action. *Nat Rev Genet* 2010;11:109-23.
- 3 [44] Vire E, Brenner C, Deplus R, Blanchon L, Fraga M, Didelot C, et al. The Polycomb
4 group protein EZH2 directly controls DNA methylation. *Nature* 2006;439:871-4.
- 5 [45] Brenner C, Deplus R, Didelot C, Lorient A, Vire E, De Smet C, et al. Myc represses
6 transcription through recruitment of DNA methyltransferase corepressor. *EMBO J*
7 2005;24:336-46.
- 8 [46] Hervouet E, Vallette FM, Cartron PF. Dnmt3/transcription factor interactions as
9 crucial players in targeted DNA methylation. *Epigenetics* 2009;4:487-99.
- 10 [47] Seet BT, Dikic I, Zhou MM, Pawson T. Reading protein modifications with
11 interaction domains. *Nat Rev Mol Cell Biol* 2006;7:473-83.
- 12 [48] Fischle W. Talk is cheap--cross-talk in establishment, maintenance, and readout of
13 chromatin modifications. *Genes and Development* 2008;22:3375-82.
- 14 [49] Schreiber SL, Bernstein BE. Signaling network model of chromatin. *Cell*
15 2002;111:771-8.
- 16 [50] Bird A. DNA methylation patterns and epigenetic memory. *Genes Dev* 2002;16:6-21.
- 17 [51] Jones PA, Liang G. Rethinking how DNA methylation patterns are maintained. *Nat*
18 *Rev Genet* 2009;10:805-11.
- 19 [52] Law JA, Jacobsen SE. Establishing, maintaining and modifying DNA methylation
20 patterns in plants and animals. *Nat Rev Genet* 2010;11:204-20.
- 21 [53] Lees-Murdock DJ, Walsh CP. DNA methylation reprogramming in the germ line.
22 *Advances in Experimental Medicine and Biology* 2008;626:1-15.
- 23 [54] Robertson KD. DNA methylation and human disease. *Nature Reviews Genetics*
24 2005;6:597-610.
- 25 [55] Miranda TB, Jones PA. DNA methylation: the nuts and bolts of repression. *Journal of*
26 *Cellular Physiology* 2007;213:384-90.
- 27 [56] Scarano MI, Strazzullo M, Matarazzo MR, D'Esposito M. DNA methylation 40 years
28 later: Its role in human health and disease. *Journal of Cellular Physiology*
29 2005;204:21-35.
- 30 [57] Lande-Diner L, Cedar H. Silence of the genes--mechanisms of long-term repression.
31 *Nature Reviews Genetics* 2005;6:648-54.
- 32 [58] Vaissiere T, Sawan C, Herceg Z. Epigenetic interplay between histone modifications
33 and DNA methylation in gene silencing. *Mutation Research* 2008;659:40-8.
- 34 [59] Chodavarapu RK, Feng S, Bernatavichute YV, Chen PY, Stroud H, Yu Y, et al.
35 Relationship between nucleosome positioning and DNA methylation. *Nature*
36 2010;466:388-92.
- 37 [60] Cosgrove MS, Wolberger C. How does the histone code work? *Biochemistry and Cell*
38 *Biology* 2005;83:468-76.
- 39 [61] Jenuwein T, Allis CD. Translating the histone code. *Science* 2001;293:1074-80.
- 40 [62] Margueron R, Trojer P, Reinberg D. The key to development: interpreting the histone
41 code? *Curr Opin Genet Dev* 2005;15:163-76.
- 42 [63] Robertson AG, Bilenky M, Tam A, Zhao Y, Zeng T, Thiessen N, et al. Genome-wide
43 relationship between histone H3 lysine 4 mono- and tri-methylation and transcription
44 factor binding. *Genome Research* 2008;18:1906-17.
- 45 [64] Jones RS. Epigenetics: reversing the 'irreversible'. *Nature* 2007;450:357-9.
- 46 [65] Pons D, de Vries FR, van den Elsen PJ, Heijmans BT, Quax PH, Jukema JW.
47 Epigenetic histone acetylation modifiers in vascular remodelling: new targets for
48 therapy in cardiovascular disease. *European Heart Journal* 2009;30:266-77.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [66] Bertram C, Khan O, Ohri S, Phillips DI, Matthews SG, Hanson MA. Transgenerational effects of prenatal nutrient restriction on cardiovascular and
2 hypothalamic-pituitary-adrenal function. *Journal of Physiology* 2008;586:2217-29.
- 3 [67] Kim M, Long TI, Arakawa K, Wang R, Yu MC, Laird PW. DNA methylation as a
4 biomarker for cardiovascular disease risk. *PLoS One* 5:e9692.
- 5 [68] Wierda RJ, Geutskens SB, Jukema JW, Quax PH, van den Elsen PJ. Epigenetics in
6 atherosclerosis and inflammation. *J Cell Mol Med* 2010.
- 7 [69] Perissi V, Rosenfeld MG. Controlling nuclear receptors: the circular logic of cofactor
8 cycles. *Nat Rev Mol Cell Biol* 2005;6:542-54.
- 9 [70] Nunez E, Fu XD, Rosenfeld MG. Nuclear organization in the 3D space of the nucleus
10 - cause or consequence? *Curr Opin Genet Dev* 2009;19:424-36.
- 11 [71] Nolis IK, McKay DJ, Mantouvalou E, Lomvardas S, Merika M, Thanos D.
12 Transcription factors mediate long-range enhancer-promoter interactions. *Proc Natl
13 Acad Sci U S A* 2009;106:20222-7.
- 14 [72] Fujioka M, Wu X, Jaynes JB. A chromatin insulator mediates transgene homing and
15 very long-range enhancer-promoter communication. *Development* 2009;136:3077-87.
- 16 [73] Metivier R, Reid G, Gannon F. Transcription in four dimensions: nuclear receptor-
17 directed initiation of gene expression. *EMBO Rep* 2006;7:161-7.
- 18 [74] Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragozy T, Telling A,
19 et al. Comprehensive mapping of long-range interactions reveals folding principles of
20 the human genome. *Science* 2009;326:289-93.
- 21 [75] Taft RJ, Pang KC, Mercer TR, Dinger M, Mattick JS. Noncoding RNAs: regulators of
22 disease. *J Pathol* 2009;220:126-39.
- 23 [76] Taft RJ, Glazov EA, Cloonan N, Simons C, Stephen S, Faulkner GJ, et al. Tiny RNAs
24 associated with transcription start sites in animals. *Nat Genet* 2009;41:572-8.
- 25 [77] Mattick JS, Taft RJ, Faulkner GJ. A global view of genomic information - moving
26 beyond the gene and the master regulator. *Trends Genet* 2009.
- 27 [78] Mattick JS, Amaral PP, Dinger ME, Mercer TR, Mehler MF. RNA regulation of
28 epigenetic processes. *Bioessays* 2009;31:51-9.
- 29 [79] Tsai MC, Manor O, Wan Y, Mosammaparast N, Wang JK, Lan F, et al. Long
30 Noncoding RNA as Modular Scaffold of Histone Modification Complexes. *Science*
31 2010.
- 32 [80] Gupta RA, Shah N, Wang KC, Kim J, Horlings HM, Wong DJ, et al. Long noncoding
33 RNA HOTAIR reprograms chromatin state to promote cancer metastasis. *Nature*
34 2010;464:1071-6.
- 35 [81] De Santa F, Barozzi I, Mietton F, Ghisletti S, Polletti S, Tusi BK, et al. A large
36 fraction of extragenic RNA pol II transcription sites overlap enhancers. *PLoS Biology*
37 2010;8:e1000384.
- 38 [82] Lujambio A, Esteller M. How epigenetics can explain human metastasis: a new role
39 for microRNAs. *Cell Cycle* 2009;8:377-82.
- 40 [83] Mahfouz MM. RNA-directed DNA methylation: Mechanisms and functions. *Plant*
41 *Signal Behav* 2010;5.
- 42 [84] Medzhitov R, Horng T. Transcriptional control of the inflammatory response. *Nat Rev*
43 *Immunol* 2009;9:692-703.
- 44 [85] Pasparakis M. Regulation of tissue homeostasis by NF-kappaB signaling: implications
45 for inflammatory diseases. *Nat Rev Immunol* 2009;9:778-88.
- 46 [86] Medzhitov R. Origin and physiological roles of inflammation. *Nature* 2008;454:428-
47 35.
- 48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [87] Cobb BS, Nesterova TB, Thompson E, Hertweck A, O'Connor E, Godwin J, et al. T
2 cell lineage choice and differentiation in the absence of the RNase III enzyme Dicer. *J*
3 *Exp Med* 2005;201:1367-73.
- 4 [88] Palstra RJ, Tolhuis B, Splinter E, Nijmeijer R, Grosveld F, de Laat W. The beta-globin
5 nuclear compartment in development and erythroid differentiation. *Nat Genet*
6 2003;35:190-4.
- 7 [89] Wilson CB, Rowell E, Sekimata M. Epigenetic control of T-helper-cell differentiation.
8 *Nature Reviews Immunology* 2009;9:91-105.
- 9 [90] Lambrecht BN, Hammad H. Taking our breath away: dendritic cells in the
10 pathogenesis of asthma. *Nature Reviews Immunology* 2003;3:994-1003.
- 11 [91] Hammad H, Lambrecht BN. Dendritic cells and epithelial cells: linking innate and
12 adaptive immunity in asthma. *Nature Reviews Immunology* 2008;8:193-204.
- 13 [92] Messi M, Giacchetto I, Nagata K, Lanzavecchia A, Natoli G, Sallusto F. Memory and
14 flexibility of cytokine gene expression as separable properties of human T(H)1 and
15 T(H)2 lymphocytes. *Nat Immunol* 2003;4:78-86.
- 16 [93] Shaheen SO, Adcock IM. The developmental origins of asthma: does epigenetics hold
17 the key? *American Journal of Respiratory and Critical Care Medicine* 2009;180:690-1.
- 18 [94] Adcock IM, Ford P, Ito K, Barnes PJ. Epigenetics and airways disease. *Respiratory*
19 *Research* 2006;7:21.
- 20 [95] Lambrecht BN, Hammad H. Biology of lung dendritic cells at the origin of asthma.
21 *Immunity* 2009;31:412-24.
- 22 [96] Hammad H, Chieppa M, Perros F, Willart MA, Germain RN, Lambrecht BN. House
23 dust mite allergen induces asthma via Toll-like receptor 4 triggering of airway
24 structural cells. *Nature Medicine* 2009;15:410-6.
- 25 [97] Grivennikov SI, Karin M. Inflammation and oncogenesis: a vicious connection. *Curr*
26 *Opin Genet Dev* 2010;20:65-71.
- 27 [98] Balkwill F, Charles KA, Mantovani A. Smoldering and polarized inflammation in the
28 initiation and promotion of malignant disease. *Cancer Cell* 2005;7:211-7.
- 29 [99] Balkwill F, Mantovani A. Cancer and inflammation: implications for pharmacology
30 and therapeutics. *Clin Pharmacol Ther* 2010;87:401-6.
- 31 [100] Aggarwal BB, Gehlot P. Inflammation and cancer: how friendly is the relationship for
32 cancer patients? *Curr Opin Pharmacol* 2009;9:351-69.
- 33 [101] Solinas G, Karin M. JNK1 and IKK{beta}: molecular links between obesity and
34 metabolic dysfunction. *FASEB J* 2010;in press.
- 35 [102] Park EJ, Lee JH, Yu GY, He G, Ali SR, Holzer RG, et al. Dietary and genetic obesity
36 promote liver inflammation and tumorigenesis by enhancing IL-6 and TNF
37 expression. *Cell* 2010;140:197-208.
- 38 [103] Olefsky JM. IKKepsilon: a bridge between obesity and inflammation. *Cell*
39 2009;138:834-6.
- 40 [104] Zhang X, Zhang G, Zhang H, Karin M, Bai H, Cai D. Hypothalamic IKKbeta/NF-
41 kappaB and ER stress link overnutrition to energy imbalance and obesity. *Cell*
42 2008;135:61-73.
- 43 [105] Hotamisligil GS. Endoplasmic reticulum stress and the inflammatory basis of
44 metabolic disease. *Cell* 2010;140:900-17.
- 45 [106] Mandl J, Meszaros T, Banhegyi G, Hunyady L, Csala M. Endoplasmic reticulum:
46 nutrient sensor in physiology and pathology. *Trends Endocrinol Metab* 2009;20:194-
47 201.
- 48 [107] Ghosh S, Hayden MS. New regulators of NF-kappaB in inflammation. *Nat Rev*
49 *Immunol* 2008;8:837-48.
- 50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [108] Karin M, Greten FR. NF-kappaB: linking inflammation and immunity to cancer
2 development and progression. *Nat Rev Immunol* 2005;5:749-59.
- 3 [109] Chen LF, Greene WC. Shaping the nuclear action of NF-kappaB. *Nat Rev Mol Cell*
4 *Biol* 2004;5:392-401.
- 5 [110] Chau TL, Gioia R, Gatot JS, Patrascu F, Carpentier I, Chapelle JP, et al. Are the IKKs
6 and IKK-related kinases TBK1 and IKK-epsilon similarly activated? *Trends Biochem*
7 *Sci* 2008;33:171-80.
- 8 [111] Chiang SH, Bazuine M, Lumeng CN, Geletka LM, Mowers J, White NM, et al. The
9 protein kinase IKKepsilon regulates energy balance in obese mice. *Cell* 2009;138:961-
10 75.
- 11 [112] Perkins ND. Integrating cell-signaling pathways with NF-kappaB and IKK function.
12 *Nat Rev Mol Cell Biol* 2007;8:49-62.
- 13 [113] Viatour P, Merville MP, Bours V, Chariot A. Phosphorylation of NF-kappaB and
14 IkappaB proteins: implications in cancer and inflammation. *Trends Biochem Sci*
15 2005;30:43-52.
- 16 [114] Vermeulen L, Berghe WV, Beck IM, De Bosscher K, Haegeman G. The versatile role
17 of MSKs in transcriptional regulation. *Trends Biochem Sci* 2009;34:311-8.
- 18 [115] Vermeulen L, De Wilde G, Van Damme P, Vanden Berghe W, Haegeman G.
19 Transcriptional activation of the NF-kappaB p65 subunit by mitogen- and stress-
20 activated protein kinase-1 (MSK1). *EMBO J* 2003;22:1313-24.
- 21 [116] Edmunds JW, Mahadevan LC. Cell signaling. Protein kinases seek close encounters
22 with active genes. *Science* 2006;313:449-51.
- 23 [117] Nakanishi C, Toi M. Nuclear factor-kappaB inhibitors as sensitizers to anticancer
24 drugs. *Nature reviews* 2005;5:297-309.
- 25 [118] Karin M. Nuclear factor-kappaB in cancer development and progression. *Nature*
26 2006;441:431-6.
- 27 [119] Arkan MC, Hevener AL, Greten FR, Maeda S, Li ZW, Long JM, et al. IKK-beta links
28 inflammation to obesity-induced insulin resistance. *Nat Med* 2005;11:191-8.
- 29 [120] De Santa F, Totaro MG, Prosperini E, Notarbartolo S, Testa G, Natoli G. The histone
30 H3 lysine-27 demethylase JMJD3 links inflammation to inhibition of polycomb-
31 mediated gene silencing. *Cell* 2007;130:1083-94.
- 32 [121] Ghisletti S, Barozzi I, Mietton F, Polletti S, De Santa F, Venturini E, et al.
33 Identification and characterization of enhancers controlling the inflammatory gene
34 expression program in macrophages. *Immunity* 2010;32:317-28.
- 35 [122] Ghisletti S, Huang W, Ogawa S, Pascual G, Lin ME, Willson TM, et al. Parallel
36 SUMOylation-dependent pathways mediate gene- and signal-specific transrepression
37 by LXRs and PPARgamma. *Mol Cell* 2007;25:57-70.
- 38 [123] Smale ST. Seq-ing LPS-induced enhancers. *Immunity* 2010;32:296-8.
- 39 [124] Xu J, Watts JA, Pope SD, Gadue P, Kamps M, Plath K, et al. Transcriptional
40 competence and the active marking of tissue-specific enhancers by defined
41 transcription factors in embryonic and induced pluripotent stem cells. *Genes Dev*
42 2009;23:2824-38.
- 43 [125] Natoli G, Sacconi S, Bosisio D, Marazzi I. Interactions of NF-kappaB with chromatin:
44 the art of being at the right place at the right time. *Nat Immunol* 2005;6:439-45.
- 45 [126] Hager GL, McNally JG, Misteli T. Transcription dynamics. *Mol Cell* 2009;35:741-53.
- 46 [127] Sung MH, Salvatore L, De Lorenzi R, Indrawan A, Pasparakis M, Hager GL, et al.
47 Sustained oscillations of NF-kappaB produce distinct genome scanning and gene
48 expression profiles. *PLoS One* 2009;4:e7163.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [128] Ramirez-Carrozzi VR, Braas D, Bhatt DM, Cheng CS, Hong C, Doty KR, et al. A
2 unifying model for the selective regulation of inducible transcription by CpG islands
3 and nucleosome remodeling. *Cell* 2009;138:114-28.
- 4 [129] O'Dea E, Hoffmann A. The regulatory logic of the NF-kappaB signaling system. *Cold
5 Spring Harb Perspect Biol* 2010;2:a000216.
- 6 [130] Werner SL, Barken D, Hoffmann A. Stimulus specificity of gene expression programs
7 determined by temporal control of IKK activity. *Science* 2005;309:1857-61.
- 8 [131] Hayden MS, Ghosh S. Shared principles in NF-kappaB signaling. *Cell* 2008;132:344-
9 62.
- 10 [132] Vanden Berghe W, De Bosscher K, Boone E, Plaisance S, Haegeman G. The nuclear
11 factor-kappaB engages CBP/p300 and histone acetyltransferase activity for
12 transcriptional activation of the interleukin-6 gene promoter. *J Biol Chem*
13 1999;274:32091-8.
- 14 [133] Ndlovu N, Van Lint C, Van Wesemael K, Callebert P, Chalbos D, Haegeman G, et al.
15 Hyperactivated NF- κ B and AP-1 transcription factors promote highly
16 accessible chromatin and constitutive transcription across the interleukin-6 gene
17 promoter in metastatic breast cancer cells. *Mol Cell Biol* 2009;29:5488-504.
- 18 [134] Eferl R, Wagner EF. AP-1: a double-edged sword in tumorigenesis. *Nature reviews*
19 2003;3:859-68.
- 20 [135] Rosenfeld MG, Lunyak VV, Glass CK. Sensors and signals: a
21 coactivator/corepressor/epigenetic code for integrating signal-dependent programs of
22 transcriptional response. *Genes Dev* 2006;20:1405-28.
- 23 [136] Dong J, Jimi E, Zhong H, Hayden MS, Ghosh S. Repression of gene expression by
24 unphosphorylated NF-kappaB p65 through epigenetic mechanisms. *Genes and
25 Development* 2008;22:1159-73.
- 26 [137] Natoli G, De Santa F. Shaping alternative NF-kappaB-dependent gene expression
27 programs: new clues to specificity. *Cell Death Differ* 2006;13:693-6.
- 28 [138] Ford E, Thanos D. The transcriptional code of human IFN-beta gene expression.
29 *Biochimica et biophysica acta* 2010.
- 30 [139] Ramirez-Carrozzi VR, Nazarian AA, Li CC, Gore SL, Sridharan R, Imbalzano AN, et
31 al. Selective and antagonistic functions of SWI/SNF and Mi-2beta nucleosome
32 remodeling complexes during an inflammatory response. *Genes Dev* 2006;20:282-96.
- 33 [140] Puto LA, Reed JC. Daxx represses RelB target promoters via DNA methyltransferase
34 recruitment and DNA hypermethylation. *Genes Dev* 2008;22:998-1010.
- 35 [141] Maeda G, Chiba T, Kawashiri S, Satoh T, Imai K. Epigenetic inactivation of IkappaB
36 Kinase-alpha in oral carcinomas and tumor progression. *Clinical Cancer Research*
37 2007;13:5041-7.
- 38 [142] O'Gorman A, Colleran A, Ryan A, Mann J, Egan LJ. Regulation of NF- κ B
39 responses by epigenetic suppression of I κ B α expression in HCT116
40 intestinal epithelial cells. *American Journal of Physiology Gastrointestinal and Liver
41 Physiology* 2010.
- 42 [143] Ashall L, Horton CA, Nelson DE, Paszek P, Harper CV, Sillitoe K, et al. Pulsatile
43 stimulation determines timing and specificity of NF-kappaB-dependent transcription.
44 *Science* 2009;324:242-6.
- 45 [144] El Gazzar M, Yoza BK, Chen X, Garcia BA, Young NL, McCall CE. Chromatin-
46 specific remodeling by HMGB1 and linker histone H1 silences proinflammatory genes
47 during endotoxin tolerance. *Mol Cell Biol* 2009;29:1959-71.
- 48 [145] El Gazzar M, Yoza BK, Hu JY, Cousart SL, McCall CE. Epigenetic silencing of
49 tumor necrosis factor alpha during endotoxin tolerance. *J Biol Chem* 2007;282:26857-
50 64.
- 51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [146] El-Osta A, Brasacchio D, Yao D, Poci A, Jones PL, Roeder RG, et al. Transient high
2 glucose causes persistent epigenetic changes and altered gene expression during
3 subsequent normoglycemia. *J Exp Med* 2008;205:2409-17.
- 4 [147] Gerlo S, Haegeman G, Vanden Berghe W. Transcriptional regulation of autocrine IL-6
5 expression in multiple myeloma cells. *Cell Signal* 2008;20:1489-96.
- 6 [148] Boeke J, Regnard C, Cai W, Johansen J, Johansen KM, Becker PB, et al.
7 Phosphorylation of SU(VAR)3-9 by the chromosomal kinase JIL-1. *PLoS ONE*
8 2010;5:e10042.
- 9 [149] Dijsselbloem N, Goriely S, Albarani V, Gerlo S, Francoz S, Marine JC, et al. A
10 critical role for p53 in the control of NF-kappaB-dependent gene expression in TLR4-
11 stimulated dendritic cells exposed to Genistein. *J Immunol* 2007;178:5048-57.
- 12 [150] Vanden Berghe W, Dijsselbloem N, Vermeulen L, Ndlovu N, Boone E, Haegeman G.
13 Attenuation of Mitogen- and Stress-Activated Protein Kinase-1-Driven Nuclear
14 Factor- κ B Gene Expression by Soy Isoflavones Does Not Require Estrogenic
15 Activity. *Cancer Res* 2006;66:4852-62.
- 16 [151] Armenante F, Merola M, Furia A, Palmieri M. Repression of the IL-6 gene is
17 associated with hypermethylation. *Biochem Biophys Res Commun* 1999;258:644-7.
- 18 [152] Dandrea M, Donadelli M, Costanzo C, Scarpa A, Palmieri M. MeCP2/H3meK9 are
19 involved in IL-6 gene silencing in pancreatic adenocarcinoma cell lines. *Nucleic Acids*
20 *Res* 2009;37:6681-90.
- 21 [153] Aaltonen T, Adelman J, Akimoto T, Albrow MG, Alvarez Gonzalez B, Amerio S, et
22 al. Cross-section-constrained top-quark mass measurement from dilepton events at the
23 Tevatron. *Physical Review Letters* 2008;100:062005.
- 24 [154] Liu S, Liu Z, Xie Z, Pang J, Yu J, Lehmann E, et al. Bortezomib induces DNA
25 hypomethylation and silenced gene transcription by interfering with Sp1/NF-kappaB-
26 dependent DNA methyltransferase activity in acute myeloid leukemia. *Blood*
27 2008;111:2364-73.
- 28 [155] Santourlidis S, Warskulat U, Florl AR, Maas S, Pulte T, Fischer J, et al.
29 Hypermethylation of the tumor necrosis factor receptor superfamily 6 (APT1, Fas,
30 CD95/Apo-1) gene promoter at rel/nuclear factor kappaB sites in prostatic carcinoma.
31 *Mol Carcinog* 2001;32:36-43.
- 32 [156] Liu X, Wang X, Zhang J, Lam EK, Shin VY, Cheng AS, et al. Warburg effect
33 revisited: an epigenetic link between glycolysis and gastric carcinogenesis. *Oncogene*
34 2010;29:442-50.
- 35 [157] Yu H, Pardoll D, Jove R. STATs in cancer inflammation and immunity: a leading role
36 for STAT3. *Nature Reviews Cancer* 2009;9:798-809.
- 37 [158] Iliopoulos D, Hirsch HA, Struhl K. An epigenetic switch involving NF-kappaB,
38 Lin28, Let-7 MicroRNA, and IL6 links inflammation to cell transformation. *Cell*
39 2009;139:693-706.
- 40 [159] Wang H, Lathia JD, Wu Q, Wang J, Li Z, Heddleston JM, et al. Targeting interleukin
41 6 signaling suppresses glioma stem cell survival and tumor growth. *Stem cells*
42 (Dayton, Ohio) 2009;27:2393-404.
- 43 [160] Min J, Zaslavsky A, Fedele G, McLaughlin SK, Reczek EE, De Raedt T, et al. An
44 oncogene-tumor suppressor cascade drives metastatic prostate cancer by coordinately
45 activating Ras and nuclear factor-kappaB. *Nature Medicine* 2010;16:286-94.
- 46 [161] Maeda S, Hikiba Y, Sakamoto K, Nakagawa H, Hirata Y, Hayakawa Y, et al. Ikappa
47 B kinasebeta/nuclear factor-kappaB activation controls the development of liver
48 metastasis by way of interleukin-6 expression. *Hepatology* 2009;50:1851-60.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [162] Grivennikov S, Karin E, Terzic J, Mucida D, Yu GY, Vallabhapurapu S, et al. IL-6
2 and Stat3 are required for survival of intestinal epithelial cells and development of
3 colitis-associated cancer. *Cancer Cell* 2009;15:103-13.
- 4 [163] Naugler WE, Karin M. The wolf in sheep's clothing: the role of interleukin-6 in
5 immunity, inflammation and cancer. *Trends in Molecular Medicine* 2008;14:109-19.
- 6 [164] Suttana W, Mankhetkorn S, Poompimon W, Palagani A, Zhokhov S, Gerlo S, et al.
7 Differential chemosensitization of P-glycoprotein overexpressing K562/Adr cells by
8 withaferin A and Siamois polyphenols. *MolCancer* 2010;9:99.
- 9 [165] Peng B, Hodge DR, Thomas SB, Cherry JM, Munroe DJ, Pompeia C, et al. Epigenetic
10 silencing of the human nucleotide excision repair gene, hHR23B, in interleukin-6-
11 responsive multiple myeloma KAS-6/1 cells. *J Biol Chem* 2005;280:4182-7.
- 12 [166] Pompeia C, Hodge DR, Plass C, Wu YZ, Marquez VE, Kelley JA, et al. Microarray
13 analysis of epigenetic silencing of gene expression in the KAS-6/1 multiple myeloma
14 cell line. *Cancer Res* 2004;64:3465-73.
- 15 [167] Hodge DR, Peng B, Pompeia C, Thomas S, Cho E, Clausen PA, et al. Epigenetic
16 Silencing of Manganese Superoxide Dismutase (SOD-2) in KAS 6/1 Human Multiple
17 Myeloma Cells Increases Cell Proliferation. *Cancer Biol Ther* 2005;4.
- 18 [168] Hodge DR, Cho E, Copeland TD, Guszczynski T, Yang E, Seth AK, et al. IL-6
19 enhances the nuclear translocation of DNA cytosine-5-methyltransferase 1 (DNMT1)
20 via phosphorylation of the nuclear localization sequence by the AKT kinase. *Cancer*
21 *Genomics Proteomics* 2007;4:387-98.
- 22 [169] Braconi C, Huang N, Patel T. MicroRNA-dependent regulation of DNA
23 methyltransferase-1 and tumor suppressor gene expression by interleukin-6 in human
24 malignant cholangiocytes. *Hepatology* 2010;51:881-90.
- 25 [170] Meng F, Wehbe-Janek H, Henson R, Smith H, Patel T. Epigenetic regulation of
26 microRNA-370 by interleukin-6 in malignant human cholangiocytes. *Oncogene*
27 2008;27:378-86.
- 28 [171] Croonquist PA, Van Ness B. The polycomb group protein enhancer of zeste homolog
29 2 (EZH 2) is an oncogene that influences myeloma cell growth and the mutant ras
30 phenotype. *Oncogene* 2005;24:6269-80.
- 31 [172] Mathews LA, Crea F, Farrar WL. Epigenetic gene regulation in stem cells and
32 correlation to cancer. *Differentiation* 2009;78:1-17.
- 33 [173] Kawasaki BT, Hurt EM, Mistree T, Farrar WL. Targeting cancer stem cells with
34 phytochemicals. *Mol Interv* 2008;8:174-84.
- 35 [174] Hodge DR, Peng B, Cherry JC, Hurt EM, Fox SD, Kelley JA, et al. Interleukin 6
36 supports the maintenance of p53 tumor suppressor gene promoter methylation. *Cancer*
37 *Res* 2005;65:4673-82.
- 38 [175] Hodge DR, Xiao W, Clausen PA, Heidecker G, Szyf M, Farrar WL. Interleukin-6
39 regulation of the human DNA methyltransferase (HDNMT) gene in human
40 erythroleukemia cells. *J Biol Chem* 2001;276:39508-11.
- 41 [176] Kim KC, Friso S, Choi SW. DNA methylation, an epigenetic mechanism connecting
42 folate to healthy embryonic development and aging. *J Nutr Biochem* 2009;20:917-26.
- 43 [177] Sharma SV, Lee DY, Li B, Quinlan MP, Takahashi F, Maheswaran S, et al. A
44 chromatin-mediated reversible drug-tolerant state in cancer cell subpopulations. *Cell*
45 2010;141:69-80.
- 46 [178] Galkina E, Ley K. Vascular adhesion molecules in atherosclerosis. *Arteriosclerosis,*
47 *Thrombosis, and Vascular Biology* 2007;27:2292-301.
- 48 [179] Galkina E, Ley K. Immune and inflammatory mechanisms of atherosclerosis (*).
49 *Annual Review of Immunology* 2009;27:165-97.
- 50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [180] Lund G, Zaina S. Atherosclerosis, lipids, inflammation and epigenetics. *Current*
2 *Opinion in Lipidology* 2007;18:699-701.
- 3 [181] Lund G, Zaina S. Atherosclerosis risk factors can impose aberrant DNA methylation
4 patterns: a tale of traffic and homocysteine. *Current Opinion in Lipidology*
5 2009;20:448-9.
- 6 [182] Lund G, Zaina S. Epigenetics, transgenerational effects and risk factors for
7 atherosclerosis. *Current Opinion in Lipidology* 2009;20:150-1.
- 8 [183] Turunen MP, Aavik E, Yla-Herttuala S. Epigenetics and atherosclerosis. *Biochimica*
9 *et biophysica acta* 2009;1790:886-91.
- 10 [184] Yu W, Fu YC, Chen CJ, Wang X, Wang W. SIRT1: a novel target to prevent
11 atherosclerosis. *J Cell Biochem* 2009;108:10-3.
- 12 [185] Zaina S, Lindholm MW, Lund G. Nutrition and aberrant DNA methylation patterns in
13 atherosclerosis: more than just hyperhomocysteinemia? *Journal of Nutrition*
14 2005;135:5-8.
- 15 [186] Franco R, Schoneveld O, Georgakilas AG, Panayiotidis MI. Oxidative stress, DNA
16 methylation and carcinogenesis. *Cancer Lett* 2008;266:6-11.
- 17 [187] Forneris F, Binda C, Battaglioli E, Mattevi A. LSD1: oxidative chemistry for
18 multifaceted functions in chromatin regulation. *Trends in Biochemical Sciences*
19 2008;33:181-9.
- 20 [188] Illi B, Colussi C, Grasselli A, Farsetti A, Capogrossi MC, Gaetano C. NO sparks off
21 chromatin: tales of a multifaceted epigenetic regulator. *Pharmacol Ther* 2009;123:344-
22 52.
- 23 [189] Brewer GJ. Epigenetic oxidative redox shift (EORS) theory of aging unifies the free
24 radical and insulin signaling theories. *Experimental Gerontology* 2010;45:173-9.
- 25 [190] Carta S, Castellani P, Delfino L, Tassi S, Vene R, Rubartelli A. DAMPs and
26 inflammatory processes: the role of redox in the different outcomes. *J Leukoc Biol*
27 2009;86:549-55.
- 28 [191] Wen H, Schaller MA, Dou Y, Hogaboam CM, Kunkel SL. Dendritic cells at the
29 interface of innate and acquired immunity: the role for epigenetic changes. *Journal of*
30 *Leukocyte Biology* 2008;83:439-46.
- 31 [192] Ng SS, Yue WW, Oppermann U, Klose RJ. Dynamic protein methylation in
32 chromatin biology. *Cellular and Molecular Life Sciences* 2009;66:407-22.
- 33 [193] Klose RJ, Zhang Y. Regulation of histone methylation by demethylination and
34 demethylation. *Nature Reviews Molecular Cell Biology* 2007;8:307-18.
- 35 [194] Jhunjhunwala S, van Zelm MC, Peak MM, Murre C. Chromatin architecture and the
36 generation of antigen receptor diversity. *Cell* 2009;138:435-48.
- 37 [195] Parra M. Epigenetic events during B lymphocyte development. *Epigenetics*
38 2009;4:462-8.
- 39 [196] Martino DJ, Prescott SL. Silent mysteries: epigenetic paradigms could hold the key to
40 conquering the epidemic of allergy and immune disease. *Allergy* 2010;65:7-15.
- 41 [197] Prescott SL, Clifton V. Asthma and pregnancy: emerging evidence of epigenetic
42 interactions in utero. *Current Opinion in Allergy and Clinical Immunology*
43 2009;9:417-26.
- 44 [198] Karouzakis E, Gay RE, Gay S, Neidhart M. Epigenetic control in rheumatoid arthritis
45 synovial fibroblasts. *Nature Reviews Rheumatology* 2009;5:266-72.
- 46 [199] Patel DR, Richardson BC. Epigenetic mechanisms in lupus. *Current Opinion in*
47 *Rheumatology* 2010.
- 48 [200] Rodriguez-Canales J, Hanson JC, Tangrea MA, Erickson HS, Albert PS, Wallis BS, et
49 al. Identification of a unique epigenetic sub-microenvironment in prostate cancer.
50 *Journal of Pathology* 2007;211:410-9.
- 51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [201] Hewagama A, Richardson B. The genetics and epigenetics of autoimmune diseases. *Journal of Autoimmunity* 2009;33:3-11.
- 2 [202] Barnett M, Bermingham E, McNabb W, Bassett S, Armstrong K, Rounce J, et al. Investigating micronutrients and epigenetic mechanisms in relation to inflammatory
3 bowel disease. *Mutation Research* 2010.
- 4 [203] Zhang QJ, Wang Z, Chen HZ, Zhou S, Zheng W, Liu G, et al. Endothelium-specific
5 overexpression of class III deacetylase SIRT1 decreases atherosclerosis in
6 apolipoprotein E-deficient mice. *Cardiovasc Res* 2008;80:191-9.
- 7 [204] Cooney CA. Germ cells carry the epigenetic benefits of grandmother's diet. *Proc Natl
8 Acad Sci U S A* 2006;103:17071-2.
- 9 [205] Weaver IC. Shaping adult phenotypes through early life environments. *Birth Defects
10 Research Part C, Embryo Today* 2009;87:314-26.
- 11 [206] Aguilera O, Fernandez AF, Munoz A, Fraga MF. Epigenetics and environment: a
12 complex relationship. *J Appl Physiol* 2010;109:243-51.
- 13 [207] Youngson NA, Whitelaw E. Transgenerational epigenetic effects. *Annual Review of
14 Genomics and Human Genetics* 2008;9:233-57.
- 15 [208] Gallou-Kabani C, Vige A, Gross MS, Junien C. Nutri-epigenomics: lifelong
16 remodelling of our epigenomes by nutritional and metabolic factors and beyond. *Clin
17 Chem Lab Med* 2007;45:321-7.
- 18 [209] Anway MD, Cupp AS, Uzumcu M, Skinner MK. Epigenetic transgenerational actions
19 of endocrine disruptors and male fertility. *Science* 2005;308:1466-9.
- 20 [210] Anway MD, Skinner MK. Epigenetic transgenerational actions of endocrine
21 disruptors. *Endocrinology* 2006;147:S43-9.
- 22 [211] Jirtle RL, Skinner MK. Environmental epigenomics and disease susceptibility. *Nat
23 Rev Genet* 2007;8:253-62.
- 24 [212] Barker DJ, Martyn CN. The maternal and fetal origins of cardiovascular disease.
25 *Journal of Epidemiology and Community Health* 1992;46:8-11.
- 26 [213] Jackson AA, Burdge GC, Lillycrop KA. Diet, nutrition and modulation of genomic
27 expression in fetal origins of adult disease. *World Review of Nutrition and Dietetics*
28 2010;101:56-72.
- 29 [214] Chmurzynska A. Fetal programming: link between early nutrition, DNA methylation,
30 and complex diseases. *Nutrition Reviews* 2010;68:87-98.
- 31 [215] Kaminsky ZA, Tang T, Wang SC, Ptak C, Oh GH, Wong AH, et al. DNA methylation
32 profiles in monozygotic and dizygotic twins. *Nature Genetics* 2009;41:240-5.
- 33 [216] Petronis A. Epigenetics and twins: three variations on the theme. *Trends in Genetics*
34 2006;22:347-50.
- 35 [217] Dolinoy DC, Weidman JR, Waterland RA, Jirtle RL. Maternal genistein alters coat
36 color and protects Avy mouse offspring from obesity by modifying the fetal
37 epigenome. *Environ Health Perspect* 2006;114:567-72.
- 38 [218] Dolinoy DC, Jirtle RL. Environmental epigenomics in human health and disease.
39 *Environ Mol Mutagen* 2008;49:4-8.
- 40 [219] Waterland RA. Is epigenetics an important link between early life events and adult
41 disease? *Horm Res* 2009;71 Suppl 1:13-6.
- 42 [220] Vandegheuchte MB, Lemiere F, Vanhaecke L, Vanden Berghe W, Janssen CR. Direct
43 and transgenerational impact on *Daphnia magna* of chemicals with a known effect on
44 DNA methylation. *Comparative Biochemistry and Physiology Toxicology and
45 Pharmacology* 2010;151:278-85.
- 46 [221] Miller RL. Prenatal maternal diet affects asthma risk in offspring. *J Clin Invest*
47 2008;118:3265-8.
- 48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [222] Gabor Miklos GL, Maleszka R. Epigenomic communication systems in humans and
2 honey bees: From molecules to behavior. *Horm Behav* 2010.
- 3 [223] Kucharski R, Maleszka J, Foret S, Maleszka R. Nutritional control of reproductive
4 status in honeybees via DNA methylation. *Science* 2008;319:1827-30.
- 5 [224] Maleszka R. Epigenetic integration of environmental and genomic signals in honey
6 bees: the critical interplay of nutritional, brain and reproductive networks. *Epigenetics*
7 2008;3:188-92.
- 8 [225] Foret S, Kucharski R, Pittelkow Y, Lockett GA, Maleszka R. Epigenetic regulation of
9 the honey bee transcriptome: unravelling the nature of methylated genes. *BMC*
10 *Genomics* 2009;10:472.
- 11 [226] Painter RC, Osmond C, Gluckman P, Hanson M, Phillips DI, Roseboom TJ.
12 Transgenerational effects of prenatal exposure to the Dutch famine on neonatal
13 adiposity and health in later life. *BJOG* 2008;115:1243-9.
- 14 [227] Lumey LH, Stein AD. Transgenerational effects of prenatal exposure to the Dutch
15 famine. *BJOG* 2009;116:868; author reply
- 16 [228] Roseboom T, de Rooij S, Painter R. The Dutch famine and its long-term consequences
17 for adult health. *Early Human Development* 2006;82:485-91.
- 18 [229] Whitelaw E. Epigenetics: sins of the fathers, and their fathers. *European Journal of*
19 *Human Genetics* 2006;14:131-2.
- 20 [230] Kaati G, Bygren LO, Edvinsson S. Cardiovascular and diabetes mortality determined
21 by nutrition during parents' and grandparents' slow growth period. *European Journal of*
22 *Human Genetics* 2002;10:682-8.
- 23 [231] Hanson MA, Gluckman PD. Developmental processes and the induction of
24 cardiovascular function: conceptual aspects. *J Physiol* 2005;565:27-34.
- 25 [232] Waldo SW, Li Y, Buono C, Zhao B, Billings EM, Chang J, et al. Heterogeneity of
26 human macrophages in culture and in atherosclerotic plaques. *American Journal of*
27 *Pathology* 2008;172:1112-26.
- 28 [233] Sharma P, Senthilkumar RD, Brahmachari V, Sundaramoorthy E, Mahajan A, Sharma
29 A, et al. Mining literature for a comprehensive pathway analysis: a case study for
30 retrieval of homocysteine related genes for genetic and epigenetic studies. *Lipids*
31 *Health Dis* 2006;5:1.
- 32 [234] Chen NC, Yang F, Capecci LM, Gu Z, Schafer AI, Durante W, et al. Regulation of
33 homocysteine metabolism and methylation in human and mouse tissues. *FASEB*
34 *Journal* 2010.
- 35 [235] Jiang Y, Sun T, Xiong J, Cao J, Li G, Wang S. Hyperhomocysteinemia-mediated
36 DNA hypomethylation and its potential epigenetic role in rats. *Acta Biochim Biophys*
37 *Sin (Shanghai)* 2007;39:657-67.
- 38 [236] Kim JM, Hong K, Lee JH, Lee S, Chang N. Effect of folate deficiency on placental
39 DNA methylation in hyperhomocysteinemic rats. *J Nutr Biochem* 2009;20:172-6.
- 40 [237] Bistulfi G, Vandette E, Matsui S, Smiraglia DJ. Mild folate deficiency induces genetic
41 and epigenetic instability and phenotype changes in prostate cancer cells. *BMC Biol*
42 2010;8:6.
- 43 [238] Ghoshal K, Li X, Datta J, Bai S, Pogribny I, Pogribny M, et al. A folate- and methyl-
44 deficient diet alters the expression of DNA methyltransferases and methyl CpG
45 binding proteins involved in epigenetic gene silencing in livers of F344 rats. *The*
46 *Journal of nutrition* 2006;136:1522-7.
- 47 [239] Ulrich CM, Reed MC, Nijhout HF. Modeling folate, one-carbon metabolism, and
48 DNA methylation. *Nutr Rev* 2008;66 Suppl 1:S27-30.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [240] Pogribny IP, Tryndyak VP, Muskhelishvili L, Rusyn I, Ross SA. Methyl deficiency,
2 alterations in global histone modifications, and carcinogenesis. *The Journal of*
3 *nutrition* 2007;137:216S-22S.
- 4 [241] Manach C, Mazur A, Scalbert A. Polyphenols and prevention of cardiovascular
5 diseases. *Curr Opin Lipidol* 2005;16:77-84.
- 6 [242] Fisher ND, Hollenberg NK. Aging and vascular responses to flavanol-rich cocoa.
7 *Journal of Hypertension* 2006;24:1575-80.
- 8 [243] Sies H, Schewe T, Heiss C, Kelm M. Cocoa polyphenols and inflammatory mediators.
9 *American Journal of Clinical Nutrition* 2005;81:304S-12S.
- 10 [244] Heiss C, Kleinbongard P, Dejam A, Perre S, Schroeter H, Sies H, et al. Acute
11 consumption of flavanol-rich cocoa and the reversal of endothelial dysfunction in
12 smokers. *Journal of the American College of Cardiology* 2005;46:1276-83.
- 13 [245] Baur JA, Sinclair DA. Therapeutic potential of resveratrol: the in vivo evidence. *Nat*
14 *Rev Drug Discov* 2006;5:493-506.
- 15 [246] Bertelli AA, Das DK. Grapes, wines, resveratrol, and heart health. *J Cardiovasc*
16 *Pharmacol* 2009;54:468-76.
- 17 [247] van Straten EM, Bloks VW, Huijkman NC, Baller JF, Meer H, Lutjohann D, et al. The
18 liver X-receptor gene promoter is hypermethylated in a mouse model of prenatal
19 protein restriction. *American Journal of Physiology Regulatory, Integrative and*
20 *Comparative Physiology* 2010;298:R275-82.
- 21 [248] Burdge GC, Slater-Jefferies J, Torrens C, Phillips ES, Hanson MA, Lillycrop KA.
22 Dietary protein restriction of pregnant rats in the F0 generation induces altered
23 methylation of hepatic gene promoters in the adult male offspring in the F1 and F2
24 generations. *Br J Nutr* 2007;97:435-9.
- 25 [249] Lillycrop KA, Phillips ES, Jackson AA, Hanson MA, Burdge GC. Dietary protein
26 restriction of pregnant rats induces and folic acid supplementation prevents epigenetic
27 modification of hepatic gene expression in the offspring. *The Journal of nutrition*
28 2005;135:1382-6.
- 29 [250] Tateishi K, Okada Y, Kallin EM, Zhang Y. Role of Jhdm2a in regulating metabolic
30 gene expression and obesity resistance. *Nature* 2009;458:757-61.
- 31 [251] Reddy MA, Villeneuve LM, Wang M, Lanting L, Natarajan R. Role of the lysine-
32 specific demethylase 1 in the proinflammatory phenotype of vascular smooth muscle
33 cells of diabetic mice. *Circulation Research* 2008;103:615-23.
- 34 [252] Brasacchio D, Okabe J, Tikellis C, Balcerzyk A, George P, Baker EK, et al.
35 Hyperglycemia induces a dynamic cooperativity of histone methylase and
36 demethylase enzymes associated with gene-activating epigenetic marks that coexist on
37 the lysine tail. *Diabetes* 2009;58:1229-36.
- 38 [253] Attig L, Gabory A, Junien C. Nutritional developmental epigenomics: immediate and
39 long-lasting effects. *Proceedings of the Nutrition Society* 2010;69:221-31.
- 40 [254] Gallou-Kabani C, Junien C. Nutritional epigenomics of metabolic syndrome: new
41 perspective against the epidemic. *Diabetes* 2005;54:1899-906.
- 42 [255] Karemaker R, Karemaker JM, Kavelaars A, Tersteeg-Kamperman M, Baerts W, Veen
43 S, et al. Effects of neonatal dexamethasone treatment on the cardiovascular stress
44 response of children at school age. *Pediatrics* 2008;122:978-87.
- 45 [256] Barres R, Osler ME, Yan J, Rune A, Fritz T, Caidahl K, et al. Non-CpG methylation
46 of the PGC-1alpha promoter through DNMT3B controls mitochondrial density. *Cell*
47 *Metabolism* 2009;10:189-98.
- 48 [257] Plagemann A, Roepke K, Harder T, Brunn M, Harder A, Wittrock-Staar M, et al.
49 Epigenetic malprogramming of the insulin receptor promoter due to developmental
50 overfeeding. *Journal of Perinatal Medicine* 2010.
- 51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [258] Plagemann A, Harder T, Brunn M, Harder A, Roepke K, Wittrock-Staar M, et al.
2 Hypothalamic proopiomelanocortin promoter methylation becomes altered by early
3 overfeeding: an epigenetic model of obesity and the metabolic syndrome. *Journal of*
4 *Physiology* 2009;587:4963-76.
- 5 [259] Burdge GC, Lillycrop KA, Jackson AA. Nutrition in early life, and risk of cancer and
6 metabolic disease: alternative endings in an epigenetic tale? *British Journal of*
7 *Nutrition* 2009;101:619-30.
- 8 [260] De Assis S, Hilakivi-Clarke L. Timing of dietary estrogenic exposures and breast
9 cancer risk. *Ann N Y Acad Sci* 2006;1089:14-35.
- 10 [261] Guerrero-Bosagna CM, Sabat P, Valdovinos FS, Valladares LE, Clark SJ. Epigenetic
11 and phenotypic changes result from a continuous pre and post natal dietary exposure
12 to phytoestrogens in an experimental population of mice. *BMC Physiol* 2008;8:17.
- 13 [262] Hunter P. We are what we eat. The link between diet, evolution and non-genetic
14 inheritance. *EMBO Rep* 2008;9:413-5.
- 15 [263] Waterland RA, Travisano M, Tahiliani KG, Rached MT, Mirza S. Methyl donor
16 supplementation prevents transgenerational amplification of obesity. *Int J Obes (Lond)*
17 2008;32:1373-9.
- 18 [264] Waterland RA, Jirtle RL. Early nutrition, epigenetic changes at transposons and
19 imprinted genes, and enhanced susceptibility to adult chronic diseases. *Nutrition*
20 2004;20:63-8.
- 21 [265] Surani MA, Ancelin K, Hajkova P, Lange UC, Payer B, Western P, et al. Mechanism
22 of mouse germ cell specification: a genetic program regulating epigenetic
23 reprogramming. *Cold Spring Harbor Symposia on Quantitative Biology* 2004;69:1-9.
- 24 [266] Hajkova P, Ancelin K, Waldmann T, Lacoste N, Lange UC, Cesari F, et al. Chromatin
25 dynamics during epigenetic reprogramming in the mouse germ line. *Nature*
26 2008;452:877-81.
- 27 [267] Chong S, Youngson NA, Whitelaw E. Heritable germline epimutation is not the same
28 as transgenerational epigenetic inheritance. *Nature Genetics* 2007;39:574-5; author
29 reply 5-6.
- 30 [268] Blewitt ME, Vickaryous NK, Paldi A, Koseki H, Whitelaw E. Dynamic
31 Reprogramming of DNA Methylation at an Epigenetically Sensitive Allele in Mice.
32 *PLoS genetics* 2006;2:e49.
- 33 [269] Bracken AP, Helin K. Polycomb group proteins: navigators of lineage pathways led
34 astray in cancer. *Nature reviews* 2009;9:773-84.
- 35 [270] Li Y, Kong D, Wang Z, Sarkar FH. Regulation of microRNAs by Natural Agents: An
36 Emerging Field in Chemoprevention and Chemotherapy Research. *Pharmaceutical*
37 *Research* 2010;27:1027-41.
- 38 [271] Ruden DM, Xiao L, Garfinkel MD, Lu X. Hsp90 and environmental impacts on
39 epigenetic states: a model for the trans-generational effects of diethylstilbestrol on
40 uterine development and cancer. *Hum Mol Genet* 2005;14 Spec No 1:R149-55.
- 41 [272] Sollars V, Lu X, Xiao L, Wang X, Garfinkel MD, Ruden DM. Evidence for an
42 epigenetic mechanism by which Hsp90 acts as a capacitor for morphological
43 evolution. *Nature Genetics* 2003;33:70-4.
- 44 [273] Ruden DM, De Luca M, Garfinkel MD, Bynum KL, Lu X. *Drosophila* nutrigenomics
45 can provide clues to human gene-nutrient interactions. *Annu Rev Nutr* 2005;25:499-
46 522.
- 47 [274] Dijsselbloem N, Vanden Berghe W, De Naeyer A, Haegeman G. Soy isoflavone
48 phyto-pharmaceuticals in interleukin-6 affections: multi-purpose nutraceuticals at the
49 crossroad of hormone replacement, anti-cancer and anti-inflammatory therapy.
50 *Biochem Pharmacol* 2004;68:1171-85.
- 51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [275] Surh YJ. Cancer chemoprevention with dietary phytochemicals. *Nature reviews*
2 2003;3:768-80.
- 3 [276] Blanpain C, Fuchs E. Epidermal homeostasis: a balancing act of stem cells in the skin.
4 *Nat Rev Mol Cell Biol* 2009;10:207-17.
- 5 [277] Crea F, Mathews LA, Farrar WL, Hurt EM. Targeting prostate cancer stem cells.
6 *Anticancer Agents Med Chem* 2009;9:1105-13.
- 7 [278] Shytle RD, Ehrhart J, Tan J, Vila J, Cole M, Sanberg CD, et al. Oxidative stress of
8 neural, hematopoietic, and stem cells: protection by natural compounds. *Rejuvenation*
9 *Res* 2007;10:173-8.
- 10 [279] Bickford PC, Tan J, Shytle RD, Sanberg CD, El-Badri N, Sanberg PR. Nutraceuticals
11 synergistically promote proliferation of human stem cells. *Stem Cells Dev*
12 2006;15:118-23.
- 13 [280] Zhou J, Zhang H, Gu P, Bai J, Margolick JB, Zhang Y. NF-kappaB pathway inhibitors
14 preferentially inhibit breast cancer stem-like cells. *Breast Cancer Res Treat*
15 2008;111:419-27.
- 16 [281] Aggarwal BB. Inflammation, a silent killer in cancer is not so silent! *Curr Opin*
17 *Pharmacol* 2009;9:347-50.
- 18 [282] Vaquero A, Reinberg D. Calorie restriction and the exercise of chromatin. *Genes Dev*
19 2009;23:1849-69.
- 20 [283] Mai A, Cheng D, Bedford MT, Valente S, Nebbioso A, Perrone A, et al. Epigenetic
21 multiple ligands: mixed histone/protein methyltransferase, acetyltransferase, and class
22 III deacetylase (sirtuin) inhibitors. *J Med Chem* 2008;51:2279-90.
- 23 [284] Kuniyasu H. The Roles of Dietary PPARgamma Ligands for Metastasis in Colorectal
24 Cancer. *PPAR research* 2008;2008:529720.
- 25 [285] Denison MS, Nagy SR. Activation of the aryl hydrocarbon receptor by structurally
26 diverse exogenous and endogenous chemicals. *Annual review of pharmacology and*
27 *toxicology* 2003;43:309-34.
- 28 [286] Darbre PD, Charles AK. Environmental oestrogens and breast cancer: evidence for
29 combined involvement of dietary, household and cosmetic xenoestrogens. *Anticancer*
30 *Res* 2010;30:815-27.
- 31 [287] Newbold RR, Padilla-Banks E, Jefferson WN. Environmental estrogens and obesity.
32 *Mol Cell Endocrinol* 2009;304:84-9.
- 33 [288] Manach C, Donovan JL. Pharmacokinetics and metabolism of dietary flavonoids in
34 humans. *Free Radic Res* 2004;38:771-85.
- 35 [289] Manach C, Williamson G, Morand C, Scalbert A, Remesy C. Bioavailability and
36 bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am J*
37 *Clin Nutr* 2005;81:230S-42S.
- 38 [290] Manach C, Scalbert A, Morand C, Remesy C, Jimenez L. Polyphenols: food sources
39 and bioavailability. *Am J Clin Nutr* 2004;79:727-47.
- 40 [291] Williamson G, Manach C. Bioavailability and bioefficacy of polyphenols in humans.
41 II. Review of 93 intervention studies. *Am J Clin Nutr* 2005;81:243S-55S.
- 42 [292] Zheng YG, Wu J, Chen Z, Goodman M. Chemical regulation of epigenetic
43 modifications: opportunities for new cancer therapy. *Med Res Rev* 2008;28:645-87.
- 44 [293] Wigle TJ, Herold JM, Senisterra GA, Vedadi M, Kireev DB, Arrowsmith CH, et al.
45 Screening for inhibitors of low-affinity epigenetic peptide-protein interactions: an
46 AlphaScreen-based assay for antagonists of methyl-lysine binding proteins. *J Biomol*
47 *Screen* 15:62-71.
- 48 [294] Altucci L, Minucci S. Epigenetic therapies in haematological malignancies: searching
49 for true targets. *Eur J Cancer* 2009;45:1137-45.
- 50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [295] Pacholec M, Bleasdale JE, Chrnyk B, Cunningham D, Flynn D, Garofalo RS, et al.
2 SRT1720, SRT2183, SRT1460, and resveratrol are not direct activators of SIRT1. *J*
3 *Biol Chem* 2010;285:8340-51.
- 4 [296] Ladurner AG. Chromatin places metabolism center stage. *Cell* 2009;138:18-20.
- 5 [297] Imai S, Guarente L. Ten years of NAD-dependent SIR2 family deacetylases:
6 implications for metabolic diseases. *Trends Pharmacol Sci* 2010;31:212-20.
- 7 [298] Chiang EP, Wang YC, Chen WW, Tang FY. Effects of insulin and glucose on cellular
8 metabolic fluxes in homocysteine transsulfuration, remethylation, S-
9 adenosylmethionine synthesis, and global deoxyribonucleic acid methylation. *Journal*
10 *of Clinical Endocrinology and Metabolism* 2009;94:1017-25.
- 11 [299] Lee DH, Jacobs DR, Jr., Porta M. Hypothesis: a unifying mechanism for nutrition and
12 chemicals as lifelong modulators of DNA hypomethylation. *Environ Health Perspect*
13 2009;117:1799-802.
- 14 [300] Whittle JR, Powell MJ, Popov VM, Shirley LA, Wang C, Pestell RG. Sirtuins, nuclear
15 hormone receptor acetylation and transcriptional regulation. *Trends Endocrinol Metab*
16 2007;18:356-64.
- 17 [301] Paul AT, Gohil VM, Bhutani KK. Modulating TNF-alpha signaling with natural
18 products. *Drug Discov Today* 2006;11:725-32.
- 19 [302] Rios JL, Recio MC, Escandell JM, Andujar I. Inhibition of transcription factors by
20 plant-derived compounds and their implications in inflammation and cancer. *Curr*
21 *Pharm Des* 2009;15:1212-37.
- 22 [303] Deorukhkar A, Krishnan S, Sethi G, Aggarwal BB. Back to basics: how natural
23 products can provide the basis for new therapeutics. *Expert Opin Investig Drugs*
24 2007;16:1753-73.
- 25 [304] Khanna D, Sethi G, Ahn KS, Pandey MK, Kunnumakkara AB, Sung B, et al. Natural
26 products as a gold mine for arthritis treatment. *Curr Opin Pharmacol* 2007;7:344-51.
- 27 [305] Bremner P, Heinrich M. Natural products as targeted modulators of the nuclear factor-
28 kappaB pathway. *J Pharm Pharmacol* 2002;54:453-72.
- 29 [306] Karin M, Yamamoto Y, Wang QM. The IKK NF-kappa B system: a treasure trove for
30 drug development. *Nat Rev Drug Discov* 2004;3:17-26.
- 31 [307] Biddie SC, John S, Hager GL. Genome-wide mechanisms of nuclear receptor action.
32 *Trends Endocrinol Metab* 2010;21:3-9.
- 33 [308] Fiskus W, Wang Y, Sreekumar A, Buckley KM, Shi H, Jillella A, et al. Combined
34 epigenetic therapy with the histone methyltransferase EZH2 inhibitor 3-
35 deazaneplanocin A and the histone deacetylase inhibitor panobinostat against human
36 AML cells. *Blood* 2009;114:2733-43.
- 37 [309] Di Croce L, Raker VA, Corsaro M, Fazi F, Fanelli M, Faretta M, et al.
38 Methyltransferase recruitment and DNA hypermethylation of target promoters by an
39 oncogenic transcription factor. *Science* 2002;295:1079-82.
- 40 [310] Vanden Berghe W, Francesconi E, De Bosscher K, Resche-Rigon M, Haegeman G.
41 Dissociated glucocorticoids with anti-inflammatory potential repress interleukin-6
42 gene expression by a nuclear factor-kappaB-dependent mechanism. *Mol Pharmacol*
43 1999;56:797-806.
- 44 [311] Vanden Berghe W, Plaisance S, Boone E, De Bosscher K, Schmitz ML, Fiers W, et al.
45 p38 and extracellular signal-regulated kinase mitogen-activated protein kinase
46 pathways are required for nuclear factor-kappaB p65 transactivation mediated by
47 tumor necrosis factor. *J Biol Chem* 1998;273:3285-90.
- 48 [312] Kaileh M, Vanden Berghe W, Heyerick A, Horion J, Piette J, Libert C, et al.
49 Withaferin a strongly elicits IkappaB kinase beta hyperphosphorylation concomitant
50 with potent inhibition of its kinase activity. *J Biol Chem* 2007;282:4253-64.
- 51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

