



**HAL**  
open science

## MELATONIN: A PLEIOTROPIC MOLECULE REGULATING INFLAMMATION

Flavia Radogna, Marc Diederich, Lina Ghibelli

► **To cite this version:**

Flavia Radogna, Marc Diederich, Lina Ghibelli. MELATONIN: A PLEIOTROPIC MOLECULE REGULATING INFLAMMATION. *Biochemical Pharmacology*, 2010, 80 (12), pp.1844. 10.1016/j.bcp.2010.07.041 . hal-00637156

**HAL Id: hal-00637156**

**<https://hal.science/hal-00637156>**

Submitted on 31 Oct 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Accepted Manuscript

Title: MELATONIN: A PLEIOTROPIC MOLECULE  
REGULATING INFLAMMATION

Authors: Flavia Radogna, Marc Diederich, Lina Ghibelli

PII: S0006-2952(10)00579-4  
DOI: doi:10.1016/j.bcp.2010.07.041  
Reference: BCP 10674

To appear in: *BCP*

Received date: 8-6-2010  
Revised date: 22-7-2010  
Accepted date: 27-7-2010

Please cite this article as: Radogna F, Diederich M, Ghibelli L, MELATONIN:  
A PLEIOTROPIC MOLECULE REGULATING INFLAMMATION, *Biochemical  
Pharmacology* (2010), doi:10.1016/j.bcp.2010.07.041

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**MELATONIN: A PLEIOTROPIC MOLECULE REGULATING INFLAMMATION**

1  
2  
3  
4  
5  
6  
7 Flavia Radogna, Marc Diederich<sup>°</sup> and Lina Ghibelli\*

8  
9  
10 Dipartimento di Biologia, Università di Roma Tor Vergata

11  
12  
13  
14 <sup>°</sup> Laboratoire de Biologie Moléculaire et Cellulaire du Cancer, Hôpital Kirchberg, 9 rue Edward  
15 Steichen, L-2540 Luxembourg, Luxembourg  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 **Address correspondence to:** Prof Lina Ghibelli, Dipartimento di Biologia, Università di Roma  
32 “Tor Vergata”, Via della Ricerca Scientifica, I-00133 Rome, Italy. E-mail: ghibelli@uniroma2.it  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**ABSTRACT**

1  
2  
3  
4 Melatonin is a neurohormone produced by the pineal gland that regulates sleep and circadian  
5 functions. Melatonin also regulates inflammatory and immune processes acting as both an activator  
6 and inhibitor of these responses. Melatonin demonstrates endocrine, but also paracrine and  
7 autocrine effects in the leukocyte compartment: on one side, leukocytes respond to melatonin in a  
8 circadian fashion; on the other side, leukocytes are able to synthesize melatonin by themselves.  
9 With its endocrine and paracrine effects, melatonin differentially modulates pro-inflammatory  
10 enzymes, controls production of inflammatory mediators such as cytokines and leukotrienes and  
11 regulates the lifespan of leukocytes by interfering with apoptotic processes. Moreover, its potent  
12 anti-oxidant ability allows scavenging of oxidative stress in the inflamed tissues. The interesting  
13 timing of pro- and anti-inflammatory effects, such as those affecting lipoxygenase activity, suggests  
14 that melatonin might promote early phases of inflammation on one hand and contribute to its  
15 attenuation on the other hand, in order to avoid complications of chronic inflammation. This review  
16 aims at giving a comprehensive overview of the various inflammatory pathways regulated by this  
17 pleiotropic hormone.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

**KEY WORDS**

32  
33  
34  
35  
36 Melatonin; inflammation; oxidative stress; reactive oxygen species; lipoxygenase; NF- $\kappa$ B  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1. Introduction

1 Inflammation is an essential response to tissue injuries induced by physical, chemical or  
2 biological insults. Acute and chronic inflammation are essential to restore homeostasis [1,2].  
3 However, the mechanisms by which activated leukocytes fight germs and tumor cells or eliminate  
4 tissue debris in the inflamed area, lead to production of oxidants and/or of cytotoxic cytokines [2,3].  
5 These processes are finely regulated, but may occasionally escape control and prolong the  
6 inflammatory response creating a self-perpetuating condition that produces further damage to the  
7 injured tissue. Since chronic inflammation is a major cause of many serious pathological conditions,  
8 and is considered by many scientists as a strong tumor promoter [4]. It is crucial to investigate  
9 novel pharmacological approaches that allow to reduce pathologic chronic inflammation without  
10 impairing the physiological inflammatory response, since many synthetic anti-inflammatory agents  
11 generate considerable side effects.

12 Pharmaceutical compounds modulating their action in subtle ways (which may be defined as  
13 smart compounds), more specifically of natural origins, able to control, regulate and defeat chronic  
14 inflammation, are within the focus of nowadays medicinal chemistry research. Endogenous  
15 molecules may be exogenously supplemented, thus increasing overall body levels, and changing  
16 timing or localization within the body with respect to their endogenous production.

17 Melatonin is very attractive from this point of view as it may act as a regulator of the  
18 inflammatory cell compartment with a potent anti-oxidant potential able to reduce the oxidative  
19 environment of chronic inflammation and to regulate leukocyte function and number [6,7].

20 We analyze here data describing the possible modulation of inflammatory functions with a  
21 focus on the anti-oxidant effects of melatonin on its ability to alter production of pro-inflammatory  
22 molecules *in vivo* and *in vitro*. Moreover, this review will cover the effects that melatonin exerts on  
23 the abundance of leukocytes *via* alteration of their proliferation and apoptosis.

## 2. Melatonin synthesis and its targets

24 Melatonin (N-acetyl-5-methoxytryptamine) is derived from tryptophan *via* enzymatic  
25 conversion reactions (see **Figure 1** for details) and is mainly produced by the pineal gland.  
26 Melatonin plasma concentration varies according to a circadian cycle and this cyclic production  
27 determines the periodic effects at a systemic level [8]. Melatonin is evolutionary conserved and  
28 exerts many regulatory functions by modulating cellular behavior *via* binding to specific receptors  
29 and intracellular targets (**Figure 2**)[9, 10].

30 Target cells express specific high affinity receptors, namely MT1/MT2, on plasma  
31 membrane [11], able to trigger intracellular signaling by adenylate cyclase or G-proteins (**Figure**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**2A**), depending on the intra- and extra-cellular environment. Even though the mechanisms through which melatonin stimulates these signal transduction pathways have been extensively studied [12-21], little is known about the resulting cellular effects: MT1/MT2 receptors have been shown to mediate melatonin-dependent decrease of tumor cell proliferation [22] and anti-apoptotic effects [23-25]; MT1 seems to be responsible for the general response to photoperiodic signals [26].

High affinity nuclear receptors to melatonin have also been identified (**Figure 2B**). They belong to the RZR/ROR nuclear hormone receptor family; melatonin binds with high affinity to RZR-beta [27, 28]. Upon activation, this complex down-regulates the 5-lipoxygenase (LOX) whose promoter possesses an RZR-beta-repressor binding site [29]; subsequently, 5-LOX expression is reduced by melatonin in cells bearing this RZR-beta receptor [30].

Cells expressing either one of those receptors are targets of the hormonal effect of melatonin and respond to its nocturnal low nanomolar plasma levels. Additional binding to intracellular targets in the micromolar range, such as the enzymes hydroquinone [31] and calmodulin [32] (**Figure 2C**) have been reported. These interactions trigger signaling functions including microtubule regulation [33] and possibly phospholipase A2 activation [34]; since hydroquinone and calmodulin are ubiquitous, most cells are targets of melatonin; however, low affinity allows productive interaction only at concentrations much higher than those produced by the pineal gland, suggesting that other mechanisms of melatonin production or accumulation may take place.

In addition to the pineal gland, which is the major source of melatonin, other tissues like retina [35], gastrointestinal tract [36, 37], skin [38] and leukocytes [6] (both in peripheral blood [39] and in bone marrow [40, 41]) synthesize melatonin. Melatonin produced by these non-endocrine organs is not regulated by circadian cycles but rather respond to other signals, exerting in fact a paracrine or autocrine effect [6]. Local melatonin levels may differ from those regulated by its neuro-endocrine functions; indeed, melatonin reaches the highest body concentration in the bone marrow (micromolar concentrations), exceeding plasma levels of at least two orders of magnitude [42]. This accumulation may result from the intrinsic melatonin synthesis machinery of the leukocyte.

As a matter of fact, melatonin finely regulates leukocyte function and number and contributes to the control of inflammatory tissue [43]. The effects of melatonin on leukocytes imply paracrine or autocrine responses that superimpose on the neuro-endocrine hormone response. In *vivo* studies with pinealectomized animals [44] allow to discriminate between hormonal and paracrine responses, whereas the study of melatonin synthesis and functions in cultured non-pineal cells [6] provides information of the autocrine effects of melatonin.

### 3. Neuro-endocrine melatonin: systemic effects of pineal melatonin on immune system

1 It is an established notion that the nervous and endocrine systems can interact with the  
2 immune system in order to modulate its function [46]; indeed, many neurotransmitters, neuro-  
3 endocrine factors and hormones can alter immune function [47]. Also melatonin is able to modulate  
4 immune functions through its neuro-endocrine action. Melatonin acts as a regulator of circadian  
5 rhythms [48] in a hormone-like fashion by affecting target cells and by modulating other functions  
6 depending on the photoperiod including regulation of photoperiodic oscillations of the  
7 immune/inflammatory response [45, 48]. Leukocytes possess melatonin specific receptors including  
8 both MT1/MT2 and nuclear RZR, and this provides the molecular basis for the sensitivity of  
9 leukocytes to melatonin.  
10

11 A strong indication of the effect of melatonin produced by the pineal gland on inflammatory  
12 cells *via* its endocrine activity was obtained in the Gerbil and in the Syrian Hamster. Here the  
13 artificial shortening of day lengths or alternatively melatonin injections induce an increase in  
14 thymus weight [49] and spleen hypertrophy [50], implying a melatonin-mediated photoperiodic  
15 effect on the immune system. Furthermore, it was demonstrated that melatonin controls diurnal and  
16 seasonal rhythms of leukocyte proliferation [51], of cytokine production [44, 45] and NK cell  
17 activity [52] in mammalian bone marrow cells. In other studies, the nocturnal peak of melatonin  
18 was associated with the proliferation peak of granulocytes and macrophage progenitor cells in mice  
19 [53].  
20

21 Data from the studies of the effects of experimental pinealectomy further contribute to a  
22 better understanding of the underlying mechanisms. It was described that the removal of the pineal  
23 gland causes partial (and transient) impairment of immune response in rats [44, 53, 54]; moreover,  
24 NK activity and IL-2 production were described to be reduced in rats after pinealectomy [44].  
25 Interestingly, melatonin administration in pinealectomized animals reverses the adverse effects on  
26 immune response. For example, Martins *et al.*, in an experimental model of allergic airway  
27 inflammation, showed that melatonin administration to pinealectomized rats restores the ability of  
28 cells to migrate from the bone marrow to the broncho-alveolar fluid, which was prevented after  
29 pineal gland removal, implying that neuro-endocrine melatonin is important in the control of cell  
30 recruitment from the bone marrow and the migration of those cells to the lung [54]  
31

32 As the melatonin-dependent effects of pinealectomy on the immune response appear to be  
33 partial, this further indicates that other compartments can overcome or contribute to  
34 immunomodulatory effects triggered by melatonin.  
35

### 4. Paracrine effects of melatonin on the white blood cell compartment

1 Leucocytes possess the enzymatic machinery necessary to synthesize melatonin.  
2 Accordingly melatonin may act as a paracrine or autocrine regulator in leukocyte communication  
3 independent of the pineal gland. Indeed Jurkat T cells produce interleukine-2 (IL-2) as a result of  
4 endogenous melatonin signaling and Lardone *et al.* demonstrated that IL-2 production is hampered  
5 by specifically inhibiting melatonin signaling by the MT1/MT2 antagonist luzindole [55]. Recently  
6 it was discovered that in bone marrow, melatonin reaches micromolar concentrations, that is much  
7 higher than those found in the blood stream, suggesting that the paracrine/autocrine immuno-  
8 regulatory role of melatonin might be also mediated by binding to the low affinity targets; a  
9 putative signal transduction pathway triggered by melatonin binding to calmodulin (affinity = 63  
10  $\mu\text{M}$  [56]) leads to transient activation of phospholipase A2 and lipoxygenase activation [34]  
11 (Figure 3).  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

## 22 5. Effect of exogenous melatonin on inflammatory parameters

23 The potential use of melatonin as a pharmaceutical agent derives from numerous *in vitro*  
24 studies where melatonin doses exceeding the nocturnal plasma levels are required to exert clear  
25 effects. At supra-physiological concentrations, melatonin induces T-cell proliferation and up-  
26 regulation of pro-inflammatory cytokines [43, 57]. Exogenous melatonin administration increases  
27 the proliferative response of rat lymphocytes [58], increases the number of NK cells [59, 60],  
28 stimulates the pro-inflammatory cytokines IL-1 and TNF- $\alpha$  [61,62] and enhances phagocytosis  
29 [63]. Melatonin exerts a concentration-dependent effect on the immune system. Indeed, increasing  
30 concentrations of melatonin induce T-cell proliferation in a dose-dependent way. In addition, it was  
31 demonstrated that pharmacological doses of melatonin inhibit INF- $\gamma$  production in a range of 0.1 to  
32 1mM [65]. In some systems, the modulation of apoptosis requires high melatonin doses [23, 24].  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 Whether or not these effects have a physiological significance, occurring as a paracrine or  
43 autocrine response, *i.e.*, in microenvironments with an elevated melatonin concentration like in the  
44 bone marrow, is presently investigated. In any case, these findings designate melatonin as a  
45 potential exogenous pharmacological modulator of the inflammatory response.  
46  
47  
48  
49  
50

## 51 6. Melatonin, ROS and inflammation

52 Oxidative stress can be defined as the imbalance between cellular oxidant species like  
53 reactive oxygen species (ROS) production and antioxidant potential. ROS are involved in a variety  
54 of different cellular processes ranging from physiological to pathological responses. It is well  
55 known that ROS can promote cell survival, proliferation and differentiation at physiological levels,  
56 but also cell death by apoptosis or necrosis at higher levels [66]. Oxidative stress has a direct toxic  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 effect on cells, which leads to lipid peroxidation, protein oxidation or DNA damage; it plays a  
2 causative or adjuvant role in almost all human pathologies, including cancer and neurodegeneration,  
3 and is involved in aging and chronic inflammatory pathologies. To cope with oxidative stress, a  
4 battery of enzymes able to scavenge ROS are up-regulated by intracellular signaling [67].  
5  
6

7 Oxidative phenomena play a key role in signaling and management of the initial phase of  
8 inflammation. Activated neutrophils generate an oxidative burst that directs toxicity towards  
9 invading microbes [68]; ROS generation in the damaged tissue produce a concentration gradient  
10 that directs leukocyte recruitment to the site of tissue injury [69]. Moreover, ROS induce monocyte  
11 maturation [70] and promote adhesion to endothelial cells allowing diapedesis. Thus, beyond its  
12 anti-septic action, ROS generation can regulate leukocyte recruitment and maturation.  
13  
14  
15  
16  
17

18 ROS generation also activates the intracellular inflammatory signaling pathways leading to  
19 the release of the inflammatory mediators, including cytokines produced by the activation of  
20 Redox-regulated transcription factors such as NF- $\kappa$ B [72,73]. Thus, ROS can promote and drive the  
21 inflammatory response in various tissues and can stimulate the signaling pathways triggered by  
22 inflammatory conditions [74].  
23  
24  
25  
26

27 If tissue healing is delayed or fails, inflammation may become chronic, increase tissue  
28 damage and induce the recruitment of additional leukocytes, producing a vicious cycle and leading  
29 to organ failure or eventually to cancer. This is characterized by chronic ROS production. An  
30 efficient anti-inflammatory therapeutic strategy requires anti-oxidants to stop this vicious cycle and  
31 to reduce the burden of inflammatory mediators at the inflammatory site.  
32  
33  
34  
35

36 Melatonin is a powerful anti-oxidant [75] as it has been shown to scavenge different types of  
37 free radicals *in vitro* [76], in body fluids [77] and in cells [78]. Furthermore, melatonin plays an  
38 important role in activating anti-oxidant defenses such as superoxide dismutase (SOD), catalase  
39 (CAT), glutathione peroxidase (GSH-Px), glutathione reductase (GSH-Rd) and glucose-6-  
40 phosphate dehydrogenase (G6PD) [79, 80].  
41  
42  
43  
44

45 Both effects allow melatonin to reduce the extent of ROS, improving oxidative related  
46 pathologies such as hypertension [81], atherosclerosis [82], cancer [83], ischemia [84] or  
47 neurodegenerative diseases [85]. Moreover it was suggested that melatonin prevents aging [86].  
48  
49  
50  
51  
52

## 53 **7. Effects of melatonin on the arachidonic acid metabolism**

54 The inflammatory response starts with an insult that activates and recruits leukocytes to the  
55 inflamed tissue through recruitment of different molecular mediators; an important role in the early  
56 phase is played by phospholipase A2 (PLA2), which cleaves membrane phospholipids liberating  
57 membrane bound arachidonic acid (AA); arachidonic acid is then processed by cyclooxygenases  
58  
59  
60  
61  
62  
63  
64  
65

(COX) and LOX to produce key inflammatory mediators such as prostaglandins and leukotrienes [87], respectively. After experimental inflammation induction *in vivo* and *ex vivo*, melatonin was shown to prevent or reduce the inflammatory-derived activation of PLA2 [88], LOX [89] and COX [90]. The reduction of PLA2 [88] and LOX [89] up-regulation by melatonin occurs *via* non-oxidative mechanisms, implying the engagement of MT1/MT2 [88] or ROR/RZR [89] receptors; thus, many different actions of melatonin including receptor stimulation or radical scavenging seem to cooperate to achieve the same final goal, that is to interfere with the inflammatory cascade.

Moreover, recent evidence shows that melatonin activates PLA2 and 5-LOX [34] as a consequence to its binding to calmodulin [34, 91], leading to mobilization of arachidonic acid (AA) from phospholipids. Arachidonic acid then activates 5-LOX triggering production of endogenous monohydroxy-eicosatetraenoic acids (HETEs) [34]. Interestingly, the down-regulation of 5-LOX (and PLA2) requires the engagement of RZR (or MT1/MT2), receptors [88, 89] but its activation instead requires melatonin to bind to calmodulin (**Figure 3**). We can conclude that the stimulation by melatonin of different targets produces different effects, altogether suggesting a sophisticated regulation.

The activation of 5-LOX and PLA2 activity by melatonin is a transient phenomenon and extinguishes within 2-3 hrs, possibly because of the down-regulation of either 5-LOX or PLA2. These observations are consistent with the view that melatonin favors the early phases of inflammation on one side, in fact, LOX induces the production of leukotriene B, which promotes diapedesis by vessel permeabilization, and acts as a chemo-attractant [92, 93]. On the other side, melatonin reduces the inflammatory process when it proceeds towards the chronic phase.

Melatonin also inhibits the activity of the enzyme nitric oxide synthase (NOS) [94-96], which catalyzes the formation of NO•. In inflamed tissues, NO released after cytokine stimulation contributes to the development of micro-vessel damage and vascular hyporeactivity [97]. Thus, melatonin-related down-regulation of NOS could contribute to melatonin's prevention of inflammation.

It was reported by that melatonin inhibits iNOS expression *via* down-regulation of NF-κB [98] on one hand but is also able to activate NF-κB, thus regulating the expression of adhesion molecules on circulating leukocytes on the other hand [99, 100]. Thus, melatonin plays a multifaceted role depending largely on the targets involved.

## 8. Role of melatonin in leukocyte survival or apoptosis

Leukocytes have to perform briefly but efficiently in an oxidative environment that may cause mutations or cell death. It is nevertheless important that cells warrant a prolonged anti-

1 bacterial effect in the injured area in spite of cellular injuries. To this purpose, the organism  
2 generated cellular resistance strategies contributing to extend leukocyte lifetime by avoiding or  
3 delaying cell death mechanisms.  
4

5 Apoptosis plays an important role in inflammatory responses but also in the regulation of the  
6 maturation rate of B and T cells [101, 102]. In particular, melatonin was shown to regulate the  
7 apoptosis of B and T cells [103, 104]. Indeed, melatonin inhibits apoptosis of precursor B cells in  
8 mouse bone marrow [105]. Moreover, an anti-apoptotic effect on different leukocytes was also  
9 demonstrated [23, 102].  
10

11 The mechanisms of the anti-apoptotic action of melatonin in leukocytes have been  
12 investigated. Studies in peripheral blood have suggested that melatonin inhibits apoptotic processes  
13 *via* its antioxidant properties [106], which is in line with many evidences showing that in some  
14 instances the apoptotic process occurs *via* oxidative signaling [107]. Alternative mechanisms have  
15 been proposed; in U937 monocytes, melatonin reduces apoptosis *via* intracellular signal  
16 transduction stimulated by interaction with MT1/MT2 plasma membrane receptors [23]. However,  
17 the majority of reports indicate that the anti-apoptotic of melatonin occurs by acting on the  
18 abundance of the two main members of the Bcl-2 family, namely the pro-apoptotic Bax and the  
19 anti-apoptotic Bcl-2 proteins [108].  
20

21 Bcl-2 exerts its well-known anti-apoptotic effect by inhibiting the action of Bax [109],  
22 which induces apoptosis by forming pores in the outer mitochondrial membrane, permeabilizing it  
23 to the passage of pro-apoptotic factors such as cytochrome c [109]. The balance between Bax and  
24 Bcl-2 determines the propensity of cells to respond to a given insult by apoptosis or survival [109].  
25 Melatonin is able to shift the balance towards a cell-protective state [108]. Indeed, many studies  
26 have demonstrated that its anti-apoptotic effect involves the down-regulation of Bax [110-113] and  
27 /or the up-regulation of Bcl-2 [110, 114, 115] that occurs *via* activation of NF- $\kappa$ B [115]. Besides  
28 altering their abundance, melatonin could alter Bax and Bcl-2 activity in other ways. Oxidative  
29 alterations cause Bax disulfide dimerization, and the resulting conformational changes are sufficient  
30 for its translocation to mitochondria independently of apoptosis [116]. However, melatonin  
31 maintains Bax in a monomeric state within mitochondria [108]. The inhibition of Bax activation  
32 may be due to melatonin's anti-oxidant ability, and/or to the simultaneous re-localization of Bcl-2  
33 [108] to mitochondria, thus co-localizing with (and sequestering) Bax. Indeed, the persistence of  
34 Bax monomers in cells induced to apoptosis in the presence of melatonin results from the  
35 prevalence of a Bax-Bcl-2 heterodimer over the active Bax dimers within mitochondria, thus,  
36 blocking both cytochrome C release and apoptosis.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 It is interesting to note that melatonin has developed various independent strategies to  
2 inhibit the intrinsic pathway of apoptosis at the level of Bax activation, inducing a pro-survival  
3 pathway and by this way maintaining viability of cells at the inflammatory site.  
4  
5  
6

### 7 **9. Role of melatonin in maintaining and restoring human health**

8 Melatonin controls the onset and progression of many human pathological states by multiple  
9 mechanisms as described by studies reporting effects of clinical trials.  
10

11 Alterations of circadian rhythms implying variations of the 24-h light/dark cycle, play a critical role  
12 in the development and progression of many diseases. Several mediators of the inflammatory  
13 response, including cytokines and hormones, have been observed to undergo significant diurnal  
14 plasma concentration variations [45-48]. Indeed, a mathematical model validated by experimental  
15 data and clinical observations, demonstrates the interplay between inflammation and circadian  
16 rhythms [117]. Melatonin is synthesized in the suprachiasmatic nucleus according to a circadian  
17 rhythm, thus causing diurnal oscillation of its blood levels. This regulation allows melatonin to  
18 coordinate circadian rhythms, including the sleep-wake cycle, thus possessing a beneficial and  
19 therapeutic effect as demonstrated by the reports of clinical trials with administration of melatonin  
20 [118].  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 Melatonin is widely used to treat various sleep disorders [119-130], including disorders associated  
32 with delayed phase sleep syndrome [119-121], periodic sleep disorders in blind patients [122-125]  
33 and sleep and behavioral disorders in children with severe brain damage [126-129]. In a randomized  
34 double-blind, placebo-controlled study [130], melatonin was shown to improve sleep efficiency and  
35 sleep latency [130] in cystic fibrosis (CF) patients, a chronic progressive disorder characterized by  
36 repeated episodes of respiratory distress and sleep disturbances [130].  
37  
38  
39  
40  
41

42 However, most of the clinical trials involving melatonin exploit its anti-oxidant properties.  
43 Melatonin was shown to reduce nitrite levels in exhaled breath condensate in CF patients [130].  
44

45 Sepsis is a severe condition associated with a significant imbalance of the intracellular redox  
46 state resulting from an increased production of oxidant species and a decrease in endogenous  
47 antioxidant defenses. In critically ill patients, sources of oxidative stress include the mitochondrial  
48 respiratory electron transport chain [131], the respiratory burst associated with neutrophil activation  
49 [68], and arachidonic acid metabolism [132]. Melatonin exerts well-documented protective effects  
50 against septic shock in both animals and humans. Melatonin was investigated in a prospective  
51 clinical trial regarding antioxidant therapy to inhibit the action of nitro-oxidative stress [133]. It was  
52 demonstrated that nitric oxide (NO) plays an important role in septic shock, an acute inflammatory  
53 response [134]. Indeed, melatonin improves the survival of mice with LPS-induced shock by  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

reducing NO synthesis *via* its antioxidant and anti-apoptotic properties [135]. It was demonstrated that antioxidant treatment improves respiratory syncytial virus (RSV)-induced pulmonary inflammation. Interestingly, the administration of melatonin reverses the oxidative stress induced by RSV by reducing NO and hydroxyl radical (OH) levels and by increasing glutathione (GSH) level and SOD activity [136]. At the same time, melatonin inhibits the production of pro-inflammatory cytokines such as TNF- $\alpha$  improving RSV-induced lung inflammatory injury in mice.

Melatonin prevents multiple organ failure, circulatory failure, and mitochondrial damage in model of experimental sepsis [137], and reduces mortality in septic children together with lipid peroxidation as an index of inflammation [138]. Melatonin was shown to improve clinical outcome in terms of infection and organ failure in randomized clinical trials, thus being a promising antioxidant molecule in antioxidant therapy to combat sepsis [132-134,137].

Administration of melatonin to humans and animals at both physiological and pharmacological concentrations is essentially non-toxic; it can be easily synthesized in a pharmacologically pure form or extracted by natural sources; it possesses outstanding versatility in reducing oxidative stress and inflammation. Altogether, these features make of melatonin a potential therapeutic tool to improve human health.

## 10. Conclusions and perspectives

Commonly used anti-inflammatory agents hardly distinguish between physiological and pathological inflammation; the indiscriminate use of anti-inflammatory agents can lead to a de-regulation of the inflammatory response, with the risk to fail to fight pathogens or injuries. The differential pro- and anti-inflammatory roles of melatonin might promote, regulate, and counteract inflammation simultaneously.

Indeed, melatonin, as an anti-inflammatory molecule, could reduce oxidative tissue injuries by its anti-oxidant properties and selectively inhibit the late/chronic phases of the inflammatory response. Furthermore, melatonin could act on early phases of inflammation by stimulating pro-inflammatory mediators such as arachidonic acid and 5-HETE *via* activation of PLA2 and 5-LOX, respectively [34]. Interestingly, effects of 5-LOX extinguish within 2-3 h, since a down-regulation of the same enzymes is also triggered by melatonin as a later event induced by different mechanisms [88, 89].

Thus, melatonin might facilitate the promotion of the inflammatory response, at the same time limiting it by providing to its resolution *via* down-regulation of 5-LOX and/or PLA2. Altogether melatonin could then avoid complications of chronic inflammation.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Freshly explanted blood monocytes require LOX-derived signals and also ROS to maximize *in vitro* activation and viability, two events that require the over-expression of Bcl-2 [70]. The role of LOX in maintaining cell viability is not unusual; 5-LOX expression is required to maintain viability of some tumor cells; in addition, in EBV-converted B lymphocytes, 5-LOX inhibition is a rapid and efficient mean of killing EBV+ cells [139].

It has been long debated that melatonin in some instances may exert a pro-apoptotic role [106]; the mechanisms and rationale of this effect are unclear. However, upon Bcl-2 silencing, melatonin acquires pro-apoptotic abilities [108], indicating that the very finely regulated changes in Bcl-2 levels during lymphocyte maturation may confer different susceptibility to melatonin, modulating a pro- or anti-apoptotic effect according to the intracellular amount of Bcl-2.

Thus, melatonin could promote acute inflammation recruiting leukocytes *via* an indirect pro-oxidant effect consisting of 5-LOX activation and promotion of inflammatory products, and prolonging the lifespan of the recruited leukocytes; it may also prevent chronic inflammation by its anti-oxidant ability. This includes a reduction of oxidative damage in the injured tissue as well as the inhibition of pro-inflammatory mediators; break the vicious cycle of oxidation/leukocyte recruitment, and promoting leukocyte apoptosis (**Figure 4**).

Altogether these investigations open new important avenues for the pharmacological use of melatonin in inflammation as the stimulation of different targets, combined with differential cellular sensitivities generates a variety of effects witnessing sophisticated regulatory mechanisms.

## Acknowledgements

F.R. is a recipient of a postdoctoral “Recherches Scientifiques Luxembourg (RSL)” grant. M.D.’s lab is financed by “Recherches Scientifiques Luxembourg (RSL)”, “Recherche Cancer et Sang” foundation, by “Een Haerz fir kriibskrank Kanner” association, by the Action Lions “Vaincre le Cancer” association and by Télévie Luxembourg.

**References**

- 1  
2 [1] Iwasaki A, Medzhitov R. Regulation of adaptive immunity by the innate immune system.  
3 Science 2010; 327:291-5.  
4
- 5 [2] Cain D, Kondo M, Chen H, Kelsoe G. Effects of acute and chronic inflammation on B-cell  
6 development and differentiation. *J Invest Dermatol.* 2009;129:266-77.  
7
- 8 [3] Davì G, Falco A. Oxidant stress, inflammation and atherogenesis. *Lupus.* 2005;14:760-4.  
9
- 10 [4] Federico A, Morgillo F, Tuccillo C, Ciardiello F, Loguercio C. Chronic inflammation and  
11 oxidative stress in human carcinogenesis. *Int J Cancer.* 2007;121:2381-6.  
12
- 13 [5] Rinaldi S, Landucci F, De Gaudio AR. Antioxidant therapy in critically septic patients. *Curr*  
14 *Drug Targets.* 2009;10:872-80.  
15
- 16 [6] Carrillo-Vico A, Calvo JR, Abreu P, Lardone PJ, Garcia-Morino S, Reiter RJ et al. Evidence of  
17 melatonin synthesis in human lymphocytes and its physiological significance: possible role as  
18 intracrine, autocrine and/or paracrine substance. *FASEB J* 2004;18:537–539.  
19
- 20 [7] Tan DX, Manchester LC, Hardeland R, Lopez-Burillo S, Mayo JC, Sainz RM, et al. Melatonin:  
21 a hormone, a tissue factor, an autocoid, a paracoid, and an antioxidant vitamin. *J Pineal Res* 2003;  
22 34:75–78.  
23
- 24 [8] Reiter RJ. Pineal melatonin: cell biology of its synthesis and of its physiological interactions.  
25 *Endocr. Rev.* 1999;12:151–180.  
26
- 27 [9] Pandi-Perumal SR, Trakht I, Srinivasan V, Spence DW, Maestroni GJ, Zisapel N, et al.  
28 Physiological effects of melatonin: role of melatonin receptors and signal transduction pathways.  
29 *Prog Neurobiol.* 2008; 85:335-53.  
30
- 31 [10] Carlson, L. L., D. R. Weaver, And S. M. Reppert. Melatonin signal transduction in hamster  
32 brain: inhibition of adenylyl cyclase by a pertussis toxin-sensitive G protein. *Endocrinology* 1989;  
33 125: 2670-2676.  
34
- 35 [11] Reppert Sm, Weaver Dr, Ebisawa T. Cloning and characterization of a mammalian melatonin  
36 receptor that mediates reproductive and circadian responses. *Neuron* 1994; 13: 1177–1185.  
37
- 38 [12] Hardeland R. Melatonin: signaling mechanisms of a pleiotropic agent. *Biofactors.*  
39 2009;35:183-92.  
40
- 41 [13] Rodriguez C, Mayo JC, Sainz RM, Antolín I, Herrera F, Martín V, Reiter RJ. Regulation of  
42 antioxidant enzymes: a significant role for melatonin. *J. Pineal Res.* 2004;36:1-9.  
43
- 44 [14] Vanecek, J, Vollrath L. Melatonin inhibits cyclic AMP and cyclic GMP accumulation in the rat  
45 pituitary. *Brain Res.* 1989;505: 157-159.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2 [15] McArthur, A J, Hunt A E, Gillette MU. Melatonin action and signal transduction in the rat  
3 suprachiasmatic circadian clock: activation of protein kinase C at dusk and dawn. *Endocrinology*  
4 1997; 138: 627-634.
- 5 [16] Vanecek, J, Klein DC. Melatonin inhibits gonadotropin-releasing hormone-induced elevation  
6 of intracellular Ca<sup>2+</sup> in neonatal rat pituitary cells. *Endocrinology* 1992; 130: 701-707.
- 7 [17] Vanecek, J, Vollrath L. Melatonin inhibits cyclic AMP and cyclic GMP accumulation in the rat  
8 pituitary. *Brain Res.* 1989; 505: 157-159.
- 9 [18] Vanecek, J, Vollrath L. Melatonin modulates diacylglycerol and arachidonic acid metabolism  
10 in the anterior pituitary of immature rats. *Neurosci. Lett.* 1990; 110: 199-203,
- 11 [19] Rodriguez C, Mayo JC, Sainz RM, Antolín I, Herrera F, Martín V, Regulation of antioxidant  
12 enzymes: a significant role for melatonin. *J Pineal Res.* 2004; 36:1-9.
- 13 [20] Chan A, Lai F, Lo R et al. Melatonin mt1 and MT2 receptors stimulate c-Jun N-terminal  
14 kinase via pertussis toxin-sensitive and -insensitive G proteins. *Cell Signal* 2002;14: 249–257.
- 15 [21] Pablos MI, Guerrero JM, Ortiz GG, Agapito MT, Reiter RJ. Both melatonin and putative  
16 nuclear melatonin receptor agonist CGP 52608 stimulate glutathione peroxidase and glutathione  
17 reductase activities in mouse brain in vivo. *Neuroendocrinol Lett* 1997;18: 49–58.
- 18 [22] Hill SM, Frasch T, Xiang S, Yuan L, Duplessis T, Mao L. Molecular mechanisms of melatonin  
19 anticancer effects. *Integr Cancer Ther.* 2009; 8: 337-46.
- 20 [23] Radogna F, Paternoster L, Albertini MC, Cerella C, Accorsi A, Bucchini A, et al. Melatonin  
21 antagonizes apoptosis via receptor interaction in U937 monocytic cells. *J Pineal Res.* 2007; 43:154-  
22 62.
- 23 [24] Radogna F, Nuccitelli S, Mengoni F, Ghibelli L. Neuroprotection by melatonin on astrocytoma  
24 cell death. *Ann N Y Acad Sci.* 2009;1171:509-13.
- 25 [25] Das A, McDowell M, Pava MJ, Smith JA, Reiter RJ, Woodward JJ, et al. The inhibition of  
26 apoptosis by melatonin in VSC4.1 motoneurons exposed to oxidative stress, glutamate  
27 excitotoxicity, or TNF-alpha toxicity involves membrane melatonin receptors. *J Pineal Res.* 2010;  
28 48:157-69.
- 29 [26] Yasuo S, Yoshimura T, Ebihara S, Korf HW. Melatonin transmits photoperiodic signals  
30 through the MT1 melatonin receptor. *J Neurosci.* 2009; 29:2885-9.
- 31 [27] Becker-André M, Wiesenberg I, Schaeren-Wiemers N, André E, Missbach M, Saurat JH et al.  
32 Pineal gland hormone melatonin binds and activates an orphan of the nuclear receptor superfamily.  
33 *J Biol Chem.* 1994; 269:28531-4. Erratum in: *J Biol Chem* 1997; 272:16707.
- 34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



- 1 [28] Wiesenberg I, Missbach M, Kahlen JP, Schröder M, Carlberg C. Transcriptional activation of  
2 the nuclear receptor RZR alpha by the pineal gland hormone melatonin and identification of CGP  
3 52608 as a synthetic ligand. *Nucleic Acids Res.* 1995; 23:327-33.  
4
- 5 [29] Carlberg C, Wiesenberg I. The orphan receptor family RZR/ROR, melatonin and 5-  
6 lipoygenase: an unexpected relationship. *J Pineal Res.* 1995;18:171-8.  
7
- 8 [30] Steinhilber D, Brungs M, Werz O, Wiesenberg I, Danielsson C, Kahlen JP et al. The nuclear  
9 receptor for melatonin repress 5-lipoxygenase gene expression in human B lymphocytes. *J Biol*  
10 *Chem* 1995; 270:7037–7040.  
11
- 12 [31] Boutin JA. Melatonin binding site MT3 is QR2: state of the art *J Soc Biol.* 2007; 201:97-103.  
13
- 14 [32] Benitez-King G, Huerto-Delgadillo L, Anton-Tay F. Binding of 3H-melatonin to calmodulin.  
15 *Life Sci* 1993;53:201–207.  
16
- 17 [33] Benítez-King G, Antón-Tay F. Calmodulin mediates melatonin cytoskeletal effects.  
18 *Experientia.* 1993;49:635-41.  
19
- 20 [34] Radogna F, Sestili P, Martinelli C, Paolillo M, Paternoster L, Albertini MC, et al.  
21 Lipoygenase-mediated pro-radical effect of melatonin via stimulation of arachidonic acid  
22 metabolism. *Toxicol Appl Pharmacol.* 2009; 238: 170-7.  
23
- 24 [35] Faillace MP, Cutrera R, Keller-Sarmento MI, Rosenstein RE. Evidence for local synthesis of  
25 melatonin in golden hamster retina. *Neuroreport* 1995; 6: 2093–2095.  
26
- 27 [36] Huether G, Poeggeler B, Reimer A, George A. Effect of tryptophan administration on  
28 circulating melatonin levels in checks and rats: evidence for stimulation of melatonin synthesis and  
29 release in the gastrointestinal tract. *Life Sci.* 1992; 52:945–953.  
30
- 31 [37] Bubenik GA. Gastrointestinal melatonin: localization, function and clinical relevance. *Dig Dis*  
32 *Sci* 2002; 47:2336–2348.  
33
- 34 [38] Iominski A, Pisarchik A, Semak I, Sweatman T, Wortsman J, Szczesniewski A, et al.  
35 Serotonergic and melatonergic systems are fully expressed in human skin. *FASEB J* 2002;  
36 16:896–898.  
37
- 38 [39] Lopez-Gonzalez MA, Calvo JR, Segura JJ, Guerrero JM. Characterization of melatonin binding  
39 sites in human peripheral blood neutrophils. *Biotechnol Ther.* 1993;4: 253-62.  
40
- 41 [40] Conti A, Conconi S, Hertens E, Skwarlo-Sonta K, Markowska M, Maestroni GJM. Evidence  
42 for melatonin synthesis in mouse and human bone marrow cells. *J Pineal Res* 2000; 28:193–202.  
43
- 44 [41] Tan DX, Manchester LC, Reiter RJ, Qi W, Zhang M, Weintraub ST, et al. Identification of  
45 highly elevated levels of melatonin in bone marrow: its origin and significance. *Biochim Biophys*  
46 *Acta* 1999; 1472:206–214.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2 [42] Tan DX, Manchester LC, Reiter RJ, Qi W, Zhang M, Weintraub ST, et al. Identification of  
3 highly elevated levels of melatonin in bone marrow: its origin and significance. *Biochim Biophys*  
4 *Acta* 1999; 1472:206–214,
- 5 [43] Carrillo-Vico A, Guerrero JM, Lardone PJ, Reiter RJ. A review of the multiple actions of  
6 melatonin on the immune system. *Endocrine*. 2005; 27:189-200.
- 7 [44] Del Gobbo V, Libri V, Villani N, Calì R, Nisticò G. Pinealectomy inhibits interleukin-2  
8 production and natural killer activity in mice. *Int J Immunopharmacol*. 1989; 11:567-73.
- 9 [45] Scheff JD, Calvano SE, Lowry SF, Androulakis IP. Modeling the influence of circadian  
10 rhythms on the acute inflammatory response. *J Theor Biol*. 2010; 264:1068-76.
- 11 [46] Blalock JE. The syntax of immune-neuroendocrine communication. *Immunol Today* 1994; 15:  
12 504-511.
- 13 [47] Ader R., Choen N., Felten D. Psychoneuroimmunology: interactions between the nervous  
14 system and the immune system. *Lancet* 1995; 345: 99-103.
- 15 [48] Malpoux B, Migaud M, Tricoire H, Chemineau P. Biology of mammalian photoperiodism and  
16 the critical role of the pineal gland and melatonin. *J Biol Rhythms* 2001;16:336-347.
- 17 [49] Vaughan MK, Vaughan GM, Reiter RJ, Blask DE. Influence of melatonin constant light, or  
18 blinding on reproductive system of gerbils (*Meriones unguiculatus*). *Experientia* 1976; 32:1341-  
19 1342.
- 20 [50] Vaughan MK, Hubbard GB, Champney TH, Vaughan GM, Little JC, Reiter RJ. Splenic  
21 hypertrophy and extramedullary hematopoiesis induced in male Syrian hamster by short  
22 photoperiod or melatonin injection and reversed by melatonin pellets or pinealectomy. *Am.J. Anat.*  
23 1987;179:131-136.
- 24 [51] Drazen DL, Bilu D, Bilbo SD, Nelson RJ. Melatonin enhancement of splenocyte proliferation  
25 is attenuated by luzindole, a melatonin receptor antagonist. *Am J Physiol Regul Integr Comp*  
26 *Physiol*. 2001;280:1476-82.
- 27 [52] Matsumoto Y, Mishima K, Satoh K, Tozawa T, Mishima Y, Shimizu T et al. Total sleep  
28 deprivation induces an acute and transient increase in NK cell activity in healthy young volunteers.  
29 *Sleep*. 2001;24:804-9.
- 30 [53] Haldar C, Häussler D, Gupta D. Effect of the pineal gland on circadian rhythmicity of colony  
31 forming units for granulocytes and macrophages (CFU-GM) from rat bone marrow cell cultures. *J*  
32 *Pineal Res*. 1992;12:79-83.
- 33 [54] Martins E Jr, Ligeiro de Oliveira AP, Fialho de Araujo AM, Tavares de Lima W, Cipolla-Neto  
34 J, Costa Rosa LF. Melatonin modulates allergic lung inflammation. *J Pineal Res*. 2001; 31:363-9.
- 35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2 [55] Lardone PJ, Carrillo-Vico A, Molinero P, Rubio A, Guerrero JM A novel interplay between  
3 membrane and nuclear melatonin receptors in human lymphocytes: significance in IL-2  
4 production. *Cell Mol Life Sci.* 2009; 66:516-25.
- 5 [56] Romero, M.P., García-Pergañeda, A., Guerrero, J.M., Osuna, C. Membrane-bound in *Xenopus*  
6 *laevis* oocytes as a novel binding site for melatonin. *FASEB J.* 1998; 12; 1401-1408.
- 7 [57] Currier NL, Sun LZ, Miller SC. Exogenous melatonin: quantitative enhancement in vivo of  
8 cells mediating non-specific immunity. *J Neuroimmunol.* 2000;104:101-8.
- 9 [58] Pahlavani MA, Harris MD. In vitro effects of melatonin on mitogen-induced lymphocyte  
10 proliferation and cytokine expression in young and old rats. *Immunopharmacol Immunotoxicol.*  
11 1997; 19:327-37.
- 12 [59] Angeli A, Gatti G, Sartori ML. Effect of exogenous melatonin on human natural killer (NK)  
13 cell activity. An approach to the immunomodulatory role of the pineal gland. *Neuroendocrinol Lett*  
14 1987; 9:286.
- 15 [60] Huang YS, Jiang JW, Wu GC, Cao XD. Effect of melatonin and electroacupuncture (EA) on  
16 NK cell activity, interleukin-2 production and POMC-derived peptides in traumatic rats. *Acupunct*  
17 *Electrother Res.* 2002; 27:95-105.
- 18 [61] Shafer LL, McNulty JA, Young MR. Assessment of melatonin's ability to regulate cytokine  
19 production by macrophage and microglia cell types. *J Neuroimmunol.* 2001;120:84-93.
- 20 [62] Pioli C, Caroleo MC, Nistico G, Doria G. Melatonin increases antigen presentation and  
21 amplifies specific and non specific signals for T-cell proliferation. *Int J Immunopharmacol.*  
22 1993;15:463-8.
- 23 [63] Sanchez S, Paredes SD, Sanchez CL, Barriga C, Reiter RJ, Rodriguez AB. Tryptophan  
24 administration in rats enhances phagocytic function and reduces oxidative metabolism. *Neuro*  
25 *Endocrinol Lett.* 2008; 29:1026-32.
- 26 [64] Shaji AV, Kulkarni SK, Agrewala JN. Regulation of secretion of IL-4 and IgG1 isotype by  
27 melatonin-stimulated ovalbumin-specific T cells. *Clin Exp Immunol.* 1998;111:181-5.
- 28 [65] Arzt ES, Fernández-Castelo S, Finocchiaro LM, Criscuolo ME, Díaz A, Finkielman S, et al.  
29 Immunomodulation by indoleamines: serotonin and melatonin action on DNA and interferon-  
30 gamma synthesis by human peripheral blood mononuclear cells. *J Clin Immunol.* 1988; 8:513-20.
- 31 [66] Yao H, Guo L, Jiang BH, Luo J, Shi X. Oxidative stress and chromium(VI) carcinogenesis. *J*  
32 *Environ Pathol Toxicol Oncol.* 2008;27:77-88.
- 33 [67] Dröge W. Free radicals in the physiological control of cell function. *Physiol Rev.* 2002; 82:47-  
34 95.
- 35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2 [68] Bogdan C, Röllinghoff M, Diefenbach A. Reactive oxygen and reactive nitrogen intermediates  
3 in innate and specific immunity. *Curr Opin Immunol.* 2000;12:64-76.
- 4 [69] Niethammer P, Grabher C, Look AT, Mitchison TJ. A tissue-scale gradient of hydrogen  
5 peroxide mediates rapid wound detection in zebrafish. *Nature.* 2009;459:996-9.
- 6  
7 [70] Cristofanon S, Nuccitelli S, D'Alessio M, Dicato M, Diederich M, Ghibelli L. Oxidation-  
8 dependent maturation and survival of explanted blood monocytes via Bcl-2 up-regulation. *Biochem*  
9 *Pharmacol.* 2008;76:1533-43.
- 10  
11 [71] Rosseau S, Selhorst J, Wiechmann K, Leissner K, Maus U, Mayer K, et al. Monocyte  
12 migration through the alveolar epithelial barrier: adhesion molecule mechanisms and impact of  
13 chemokines. *Immunol.* 2000;164:427-35.
- 14  
15 [72] Kim H, Hwang JS, Woo CH, Kim EY, Kim TH, Cho KJ, et al. TNF-alpha-induced up-  
16 regulation of intercellular adhesion molecule-1 is regulated by a Rac-ROS-dependent cascade in  
17 human airway epithelial cells. *Exp Mol Med.* 2008; 40:167-75.
- 18  
19 [73] Cristofanon S, Morceau F, Scovassi AI, Dicato M, Ghibelli L, Diederich M. Oxidative,  
20 multistep activation of the noncanonical NF-kappaB pathway via disulfide Bcl-3/p50  
21 complex. *FASEB J.* 2009;23:45-57.
- 22  
23 [74] Martinon F. Signaling by ROS drives inflammasome activation. *Eur J Immunol.* 2010; 40:  
24 616-9.
- 25  
26 [75] Tan D-X, Chen LD, Poeggeler B, Manchester LC, Reiter RJ. Melatonin: A potent, endogenous  
27 hydroxyl radical scavenger. *Endocr J* 1993; 1:57-60.
- 28  
29 [76] Poeggeler B, Reiter RJ, Hardeland R, Tan DX, Barlow-Walden LR. Melatonin and structurally  
30 related endogenous indoles act as potent electron donors and radical scavengers in vitro. *Redox Rep*  
31 1996; 2:179-184.
- 32  
33 [77] Du Plessis SS, Hagenaar K, Lampiao F. The in vitro effects of melatonin on human sperm  
34 function and its scavenging activities on NO and ROS. *Andrologia.* 2010; 42:112-6.
- 35  
36 [78] Zavodnik IB, Domanski AV, Lapshina EA, Bryszewska M, Reiter RJ. Melatonin directly  
37 scavenges free radicals generated in red blood cells and a cell-free system: chemiluminescence  
38 measurements and theoretical calculations. *Life Sci.* 2006; 79:391-400.
- 39  
40 [79] Bharti VK, Srivastava RS. Pineal proteins upregulate specific antioxidant defense systems in  
41 the brain. *Oxid Med Cell Longev.* 2009; 2:88-92.
- 42  
43 [80] Hardeland R. Neuroprotection by radical avoidance: search for suitable agents. *Molecules.*  
44 2009;14:5054-102.
- 45  
46 [81] Reiter RJ, Tan DX, Korkmaz A. The circadian melatonin rhythm and its modulation: possible  
47 impact on hypertension. *J Hypertens Suppl.* 2009;27:S17-20.
- 48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2 [82] Broncel M, Koziróg-Kołacińska M, Chojnowska-Jeziarska J. Melatonin in the treatment of  
3 atherosclerosis. *Pol Merkur Lekarski*. 2007;23:124-7.
- 4 [83] Ravindra T, Lakshmi NK, Ahuja YR. Melatonin in pathogenesis and therapy of cancer. *Indian*  
5 *J Med Sci*. 2006;60:523-35.
- 6 [84] Reiter RJ, Tan DX, Leon J, Kilic U, Kilic E. When melatonin gets on your nerves: its  
7 beneficial actions in experimental models of stroke. *Exp Biol Med*. 2005; 230:104-17.
- 8 [85] Skaper SD, Floreani M, Ceccon M, Facci L, Giusti P. Excitotoxicity, oxidative stress, and the  
9 neuroprotective potential of melatonin. *Ann N Y Acad Sci*. 1999;890:107-18.
- 10 [86] Reiter RJ, Guerrero JM, Garcia JJ, Acuña-Castroviejo D. Reactive oxygen intermediates,  
11 molecular damage, and aging. Relation to melatonin. *Ann N Y Acad Sci*. 1998;854:410-24.
- 12 [87] González-Pérez A, Clària J. New approaches to the modulation of the cyclooxygenase-2 and 5-  
13 lipoygenase pathways. *Curr Top Med Chem*. 2007;7:297-309.
- 14 [88] Li, B., Zhang, H., Akbar, M., Kim, H.Y. 2000. Negative regulation of cytosolic phospholipase  
15 A(2) by melatonin in the rat pineal gland. *Biochem J*. 2000;351:709-716.
- 16 [89] Steinhilber D, Brungs M, Werz O et al. The nuclear receptor for melatonin repress 5-  
17 lipoygenase gene expression in human B lymphocytes. *J Biol Chem* 1995; 270: 7037–7040.
- 18 [90] Deng WG, Tang ST, Tseng HP, Wu KK. Melatonin suppresses macrophage cyclooxygenase-2  
19 and inducible nitric oxide synthase expression by inhibiting p52 acetylation and binding. *Blood*.  
20 2006;108:518-24.
- 21 [91] Jenkins CM, Wolf MJ, Mancuso DJ, Gross RW. Identification of the calmodulin-binding  
22 domain of recombinant calcium-independent phospholipase A2beta. Implications for structure and  
23 function. *J Biol Chem*. 2001; 276:7129-35.
- 24 [92] Pidgeon GP, Lysaght J, Krishnamoorthy S, Reynolds JV, O'Byrne K, Nie D et al.  
25 Lipoygenase metabolism: roles in tumor progression and survival. *Cancer Metastasis Rev*.  
26 2007;26:503-24.
- 27 [93] Islam SA, Thomas SY, Hess C, Medoff BD, Means TK, Brander C, et al. The leukotriene B4  
28 lipid chemoattractant receptor BLT1 defines antigen-primed T cells in humans. *Blood*.  
29 2006;107:444-53.
- 30 [94] Bettahi I, Pozo D, Osuna C, Reiter RJ, Acuna-Castroviejo D, Guerrero JM. Physiological  
31 concentrations of melatonin inhibit nitric oxide synthase activity in the rat hypothalamus. *J Pineal*  
32 *Res* 1996;20:205–210.
- 33 [95] Guerrero JM, Reiter RJ, Ortiz GG, Pablos MI, Sewerynek E, Chuang JI. Melatonin prevents  
34 increases in neural nitric oxide and cycle GMP production after transient brain ischemia and  
35 reperfusion in the Mongolian gerbil (*Meriones unguiculatus*). *J Pineal Res* 1997; 23:24–31.
- 36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2 [96] Pozo D, Reiter RJ, Calvo JM, Guerrero JM. Inhibition of cerebellar nitric oxide synthase and  
3 cyclic GMP production by melatonin via complex formation with calmodulin. *J Cell Biochem*  
4 1997;65:430–442.
- 5 [97] Tesse A, Martínez MC, Hugel B, Chalupsky K, Muller CD, Meziani F, Upregulation of  
6 proinflammatory proteins through NF-kappaB pathway by shed membrane microparticles results in  
7 vascular hyporeactivity. *Arterioscler Thromb Vasc Biol.* 2005; 25:2522-7.
- 8  
9 [98] Alonso M, Collado PS, González-Gallego J. Melatonin inhibits the expression of the inducible  
10 isoform of nitric oxide synthase and nuclear factor kappa B activation in rat skeletal muscle. *J*  
11 *Pineal Res.* 2006; ;41:8-14.
- 12  
13 [99] Cristofanon S, Uguccioni F, Cerella C, Radogna F, Dicato M, Ghibelli L, et al. Intracellular  
14 prooxidant activity of melatonin induces a survival pathway involving NF-kappaB activation. *Ann*  
15 *N Y Acad Sci.* 2009;1171:472-8.
- 16  
17 [100] Cecon E, Fernandes PA, Pinato L, Ferreira ZS, Markus RP. Daily variation of constitutively  
18 activated nuclear factor kappa B (NFkB) in rat pineal gland. *Chronobiol Int.* 2010; 27:52-67.
- 19  
20 [101] Wesche-Soldato De, Swan Rz, Chung Cs, Ayala A. The apoptotic pathway as a therapeutic  
21 target in sepsis. *Curr Drug Targets.* 2007; 8:493-500.
- 22  
23 [102] Bolitho P, Voskoboinik I, Trapani Ja, Smyth Mj. Apoptosis induced by the lymphocyte  
24 effector molecule perforin. *Curr Opin Immunol.* 2007; 19:339-347.
- 25  
26 [103] Pedrosa AM, Weinlich R, Mognol GP, Robbs BK, Viola JP, Campa A Melatonin protects  
27 CD4+ T cells from activation-induced cell death by blocking NFAT-mediated CD95 ligand  
28 upregulation. *J Immunol.* 2010;184:3487-94.
- 29  
30 [104] Paternoster L, Radogna F, Accorsi A, Cristina Albertini M, Gualandi G, et al. Melatonin as a  
31 modulator of apoptosis in B-lymphoma cells. *Ann N Y Acad Sci.* 2009;1171:345-9.
- 32  
33 [105] Yu Q, Miller SC, Osmond DG. Melatonin inhibits apoptosis during early B-cell development  
34 in mouse bone marrow. *J Pineal Res.* 2000; 29:86-93.
- 35  
36 [106] Sainz RM, Mayo JC, Rodriguez C, Tan DX, Lopez-Burillo S, Reiter RJ. Melatonin and cell  
37 death: differential actions on apoptosis in normal and cancer cells. *Cell Mol Life Sci.*  
38 2003;60:1407-26.
- 39  
40 [107] Ortona E, Margutti P, Matarrese P, Franconi F, Malorni W. Redox state, cell death and  
41 autoimmune diseases: a gender perspective. *Autoimmun Rev.* 2008;7:579-84.
- 42  
43 [108] Radogna F, Cristofanon S, Paternoster L, D'Alessio M, De Nicola M, Cerella C, et al.  
44 Melatonin antagonizes the intrinsic pathway of apoptosis via mitochondrial targeting of Bcl-2. *J*  
45 *Pineal Res.* 2008;44:316-25.
- 46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2 [109] Lucken-Ardjomande S, Martinou JC. Regulation of Bcl-2 proteins and of the permeability of  
the outer mitochondrial membrane. *C R Biol.* 2005;328:616-31.
- 3  
4 [110] Maity P, Bindu S, Dey S, Goyal M, Alam A, Pal C, et al. Melatonin reduces indoethacin-  
5 induced gastric mucosal cell apoptosis by preventing mitochondrial oxidative stress and the  
6 activation of mitochondrial pathway of apoptosis. *J Pineal Res.* 2009;46:314-23.
- 7  
8 [111] Chetsawang B, Putthaprasart C, Phansuwan-pujito P, Govitrapong P. Melatonin protects  
9 against hydrogen peroxide-induced cell death signaling in SH-SY5Y cultured cells: involvement of  
10 nuclear factor kappa B, Bax and Bcl-2. *J Pineal Res.* 2006 ; 41:116-23.
- 11  
12 [112] Juknat AA, Mendez Mdel V, Quaglino A, et al. Melatonin prevents hydrogen peroxide-  
13 induced Bax expression in cultured rat astrocytes. *J Pineal Res.* 2005; 38:84-92.
- 14  
15 [113] Espino J, Bejarano I, Redondo PC, Rosado JA, Barriga C, Reiter RJ, et al. Melatonin reduces  
16 apoptosis induced by calcium signaling in human leukocytes: Evidence for the involvement of  
17 mitochondria and Bax activation. *J Membr Biol.* 2010;233:105-18.
- 18  
19 [114] Ling X, Zhang LM, Lu SD, Li XJ, Sun FY. Protective effect of melatonin on injured cerebral  
20 neurons is associated with bcl-2 protein over-expression. *Zhongguo Yao Li Xue Bao.* 1999; 20:409-  
21 414.
- 22  
23 [115] Cristofanon S, Uguccioni F, Cerella C, Radogna F, Dicato M, Ghibelli L, et al. Intracellular  
24 prooxidant activity of melatonin induces a survival pathway involving NF-kappaB activation. *Ann*  
25 *N Y Acad Sci.* 2009 ;1171:472-8.
- 26  
27 [116] D'Alessio M, De Nicola M, Coppola S, Gualandi G, Pugliese L, Cerella C, et al. Oxidative  
28 Bax dimerization promotes its translocation to mitochondria independently of apoptosis. *FASEB J.*  
29 2005;19:1504-6.
- 30  
31 [117] Scheff JD, Calvano SE, Lowry SF, Androulakis IP. Modeling the influence of circadian  
32 rhythms on the acute inflammatory response. *J Theor Biol.* 2010; 264:1068-76.
- 33  
34 [118] Boeve BF, Silber MH, Ferman TJ. Melatonin for treatment of REM sleep behavior disorder in  
35 neurologic disorders: results in 14 patients. *Sleep Med.* 2003;4:281-4.
- 36  
37 [119] Dahlitz M, Alvarez B, Vignau J, English J, Arendt J, Parkes JD. Delayed sleep phase  
38 syndrome response to melatonin. *Lancet* 1991;337:1121-1124.
- 39  
40 [120] Munday K, Benloucif S, Harsanyi K, Dubocovich ML, Zee PC. Phase-dependent treatment of  
41 delayed sleep phase syndrome with melatonin. *Sleep.* 2005; 28:1271-8.
- 42  
43 [121] Oldani A, Ferini-Strambi L, Zucconi M, Stankov B, Fraschini F, Smirne S: Melatonin and  
44 delayed sleep phase syndrome: Ambulatory polygraphic evaluation. *Neuroreport* 1994;6:132-134.
- 45  
46 [122] Lapiere O, Dumont M. Melatonin treatment of a non-24-hour sleep-wake cycle in a blind  
47 retarded child. *Biol Psychiatry* 1995;38:119-122.
- 48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1 [123] Folkard S, Arendt J, Aldhous M, Kennett H: Melatonin stabilises sleep onset time in a blind  
2 man without entrainment of cortisol or temperature rhythms. *Neurosci Lett* 1990;113:193–198.
- 3 [124] Arendt J, Aldhous M, Wright J. Synchronisation of a disturbed sleep-wake cycle in a blind  
4 man by melatonin treatment (letter). *Lancet* 1988;1:772–773.
- 5 [125] Sack RL, Lewy AJ, Blood ML, Stevenson J, Keith LD: Melatonin administration to blind  
6 people: Phase advances and entrainment. *J Biol Rhythms* 1991;6:249–261.
- 7 [126] Palm L, Blennow G, Wetterberg L: Correction of non-24-hour sleep/wake cycle by melatonin  
8 in a blind retarded boy. *Ann Neurol* 1991;29:336–339.
- 9 [127] Jan JE, Espezel H, Appleton RE: The treatment of sleep disorders with melatonin. *Dev Med*  
10 *Child Neurol* 1994;36:97–107.
- 11 [128] Jan JE, O'Donnell ME: Use of melatonin in the treatment of paediatric sleep disorders.  
12 *J.Pineal Res* 1996;21:193–199.
- 13 [129] Jan JE, Abroms IF, Freeman RD, Brown GM, Espezel H, Connolly MB: Rapid cycling in  
14 severely multidisabled children: A form of bipolar affective disorder? *Pediatr Neurol* 1994;10:34–  
15 39.
- 16 [130] De Castro-Silva C, de Bruin VM, Cunha GM, Nunes DM, Medeiros CA, de Bruin PF.  
17 Melatonin improves sleep and reduces nitrite in the exhaled breath condensate in cystic fibrosis--a  
18 randomized, double-blind placebo-controlled study. *J Pineal Res.* 2010;48:65-71.
- 19 [131] Fredriksson K, Hammarqvist F, Strigård K, Hultenby K, Ljungqvist O, Wernerman J, et al.  
20 Derangements in mitochondrial metabolism in intercostal and leg muscle of critically ill patients  
21 with sepsis-induced multiple organ failure. *Am J Physiol Endocrinol Metab.* 2006; 291:E1044-50.
- 22 [132] Rinaldi S, Landucci F, De Gaudio AR. Antioxidant therapy in critically septic patients. *Curr*  
23 *Drug Targets.* 2009;10: 872-80.
- 24 [133] 133 Korkmaz A, Reiter RJ, Topal T, Manchester LC, Oter S, Tan DX. Melatonin: an  
25 established antioxidant worthy of use in clinical trials. *Mol Med.* 2009;15:43-50.
- 26 [134] Bubenik GA, Blask DE, Brown GM, Maestroni GJ, Pang SF, Reiter RJ, et al. Prospects of the  
27 clinical utilization of melatonin. *Biol Signals Recept.* 1998;7:195-219.
- 28 [135] Carrillo-Vico A, Lardone PJ, Naji L, Fernández-Santos JM, Martín-Lacave I, Guerrero JM, et  
29 al. Beneficial pleiotropic actions of melatonin in an experimental model of septic shock in mice:  
30 regulation of pro-/anti-inflammatory cytokine network, protection against oxidative damage and  
31 anti-apoptotic effects. *J Pineal Res.* 2005; 39:400-8.
- 32 [136] Huang SH, Cao XJ, Liu W, Shi XY, Wei W. Inhibitory effect of melatonin on lung oxidative  
33 stress induced by respiratory syncytial virus infection in mice. *J Pineal Res.* 2010; 48:109-16.
- 34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 [137] Germaine Escames<sup>2</sup>, Darío Acuña-Castroviejo<sup>1</sup>, Luis Carlos López<sup>1</sup>, Dun-xian Tan<sup>2</sup>, Maria  
2 Dolores Maldonado<sup>2</sup>, Marina Sánchez-Hidalgo<sup>2</sup>, et al. Pharmacological utility of melatonin in the  
3 treatment of septic shock: experimental and clinical evidence. *J. Pharm. Pharmacol.*  
4 2006;58:1153-65.  
5

6  
7 [138] Gitto E, Karbownik M, Reiter RJ, Tan DX, Cuzzocrea S, Chiurazzi P, et al. Effects of  
8 melatonin treatment in septic newborns. *Pediatr Res.* 2001; 50:756-60.  
9

10 [139] Belfiore, M.C., Natoni, A., Barzellotti, R., Merendino, N., Pessina, G., Ghibelli, L., et al..  
11 Involvement of 5-lipoxygenase in survival of Epstein-Barr virus (EBV)-converted B lymphoma  
12 cells. *Cancer Lett.* 2007; 254: 236-243.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65


## Legends to figures

**Figure 1:** Biosynthesis of melatonin


**Figure 2:** Intracellular melatonin targets. Melatonin interacts with plasma membrane (MT1/MT2) and nuclear (ROR/RZR) receptors at high affinity. Additional intracellular targets at lower affinity are the enzymes calmodulin and hydroquinone. The affinity for melatonin and the relative reference is indicated.

**Figure 3:** Dual effects of melatonin on lipoxygenase. Melatonin both down-regulates lipoxygenase gene expression through RZR-beta (A) and activates 5-LOX *via* PLA2 activation *via* calmodulin-dependent signaling (B).

**Figure 4:** Pleiotropic effects of melatonin on different steps of inflammation. Summary of the multiple effects that melatonin exerts at different steps of the inflammatory response, indicating a pro-inflammatory role at an early phase, and an antagonist role at later phases. This evokes a smart behavior where melatonin may favor the inflammatory healing processes while contrasting pathologically chronic or deregulated inflammation, thus potentially being an ideal compound to treat inflammatory disturbances.




A.




Cytosol


C.

CALMODULIN (63  $\mu$ M affinity [56])


B.

ROR/RZR (pM - nM affinity [27])







# Melatonin



Early phase  
of inflammation



Chronic  
inflammation




De-regulation of  
inflammation  
(Risk of cancer; autoimmunity)

Activates  
pro-inflammatory mediators  
such as PLA2, LOX,  
Cytokines (IL1-TNF $\alpha$ )

1. Down-regulates pro-inflammatory mediators such as PLA2, LOX, cytokines (IL1-TNF $\alpha$ )
2. Contrasts oxidative tissue damage by its anti-oxidant properties
3. Induces a survival pathway in leukocytes


Induces apoptosis

### Molecular structure of melatonin



### Intracellular melatonin targets.

A.



B.

ROR/RZR (pM - nM affinity [27])



### Melatonin as a smart drug modulating inflammation.

