



HAL
open science

Exponential convergence for a convexifying equation

Guillaume Carlier, Alfred Galichon

► **To cite this version:**

Guillaume Carlier, Alfred Galichon. Exponential convergence for a convexifying equation. ESAIM: Control, Optimisation and Calculus of Variations, 2012, pp.x-xx. 10.1051/cocv/2011163. hal-00637398

HAL Id: hal-00637398

<https://hal.science/hal-00637398>

Submitted on 1 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exponential convergence for a convexifying equation

G. Carlier ^{*}, A. Galichon [†]

April 7, 2011

Abstract

We consider an evolution equation similar to that introduced by Vese in [12] and whose solution converges in large time to the convex envelope of the initial datum. We give a stochastic control representation for the solution from which we deduce, under quite general assumptions that the convergence in the Lipschitz norm is in fact exponential in time.

Keywords: convex envelope, viscosity solutions, stochastic control representation, nonautonomous gradient flows.

1 Introduction

In an interesting paper [12], L.Vese considered the following PDE:

$$\partial_t u = \sqrt{1 + |\nabla u|^2} \min(0, \lambda_1(D^2u)), \quad u|_{t=0} = u_0 \quad (1.1)$$

where $\lambda_1(D^2u)$ denotes the smallest eigenvalue of the Hessian matrix D^2u . Vese proved, under quite general assumptions on the initial condition u_0 , that the viscosity solution of (1.1) converges as $t \rightarrow \infty$ to u_0^{**} the convex envelope of u_0 . Starting from this result, Vese developed an original and purely PDE approach to approximate convex envelopes (which is in general a delicate problem as soon as the space dimension is larger than 2). More recently,

^{*}CEREMADE, UMR CNRS 7534, Université Paris IX Dauphine, Pl. de Lattre de Tassigny, 75775 Paris Cedex 16, France. carlier@ceremade.dauphine.fr

[†]Département d'Economie, UMR CNRS 7176, Ecole polytechnique, 91128 Palaiseau Cedex. alfred.galichon@polytechnique.edu.

A. Oberman [7], [8], [9], noticed that the convex envelope can be directly characterized via a nonlinear elliptic PDE of obstacle type and developed this idea for numerical computation of convex envelopes as well. As noticed by Oberman, the solution of the PDE he introduced naturally has a stochastic control representation. This is of course also the case for the evolutionary equation and, as we shall see, this representation will turn out to be very useful to obtain convergence estimates.

In the present paper, we will focus on an evolution equation similar to (1.1) and will study some of its properties thanks to the stochastic control representation of the solution.

The paper is organized as follows. In section 2, we introduce the *convexifying evolution equation* and recall some basic facts about convex envelopes. In section 3, we give a stochastic control representation for the solution of the convexifying evolution equation. Section 4 gives some regularity properties of the solution. Our exponential convergence result is then proved in section 5 by simple probabilistic arguments. Finally, section 6 is devoted to some concluding remarks. We apply our exponential convergence result to study the behavior of some nonautonomous gradient flow and we finally make a link with some geometric flow by considering a family of interpolating problems.

2 A convexifying evolution equation

In the present paper, we will consider a slight variant of (1.1), namely:

$$\partial_t u(t, x) = \min(0, \lambda_1(D^2 u(t, x))), \quad (t, x) \in (0; \infty) \times \mathbb{R}^d, \quad u|_{t=0} = u_0 \quad (2.1)$$

From a PDE point of view, it may be worthwhile to observe that the nonlinear operator involved in the equation may be considered, in a degenerate case, as one of the Pucci extremal operators, which have been extensively studied in the literature on fully nonlinear PDE's (see for instance the book by Caffarelli and Cabré [2]).

In the sequel, we shall refer to (2.1) as the *convexifying evolution equation*. Following the same arguments of the proof of Vese [12] (also see remark 3.4 below), one can prove under mild assumptions on u_0 that the solution converges pointwise to the convex envelope u_0^{**} of the initial condition. Our aim will be to quantify this convergence and this goal will be achieved rather easily by using a stochastic representation formula for the solution of (2.1). Before we do so, let us recall some basic facts about the convex envelope.

Given a continuous (say) and bounded from below function u_0 defined on \mathbb{R}^d , the convex envelope of u_0^{**} is the largest convex function that is

everywhere below u_0 . The convex envelope is a very natural object in many contexts and in particular in optimization since u_0 and u_0^{**} have the same infimum but u_0^{**} is in principle much simpler to minimize since it is convex. One can also define u_0^{**} as the supremum of all affine functions that are below u_0 and thus define u_0^{**} as the “Legendre Transform of the Legendre Transform” of u_0 (and this is where the notation “**” comes from). Rather than iterating the Legendre transform, let us recall the well-known formula:

$$u_0^{**}(x) = \inf \left\{ \sum_{i=1}^{d+1} \lambda_i u_0(x_i) : \lambda_i \geq 0, \sum_{i=1}^{d+1} \lambda_i = 1, \sum_{i=1}^{d+1} \lambda_i x_i = x \right\}, \forall x \in \mathbb{R}^d \quad (2.2)$$

(the fact that one can restrict to $d + 1$ points follows from Carathéodory’s theorem) which can also be written in probabilistic terms as

$$u_0^{**}(x) = \inf \left\{ \mathbb{E}(u_0(x + X)) : \mathbb{E}(X) = 0 \right\}. \quad (2.3)$$

The latter formula strongly suggests that a good approximation for the convex envelope should be

$$u(t, x) := \inf_{\sigma : |\sigma| \leq 1} \left\{ \mathbb{E} \left(u_0 \left(x + \int_0^t \sqrt{2} \sigma_s dW_s \right) \right) \right\} \quad (2.4)$$

for large t where $(W_s)_{s \geq 0}$ is a standard Brownian motion, σ_s is a $d \times d$ -matrix valued process that is adapted to the Brownian filtration and $|\sigma|$ stands for the matrix norm $|\sigma| := \sqrt{\text{Tr}(\sigma \sigma^T)}$.

In order to keep things as elementary as possible, from now on, we shall always assume that u_0 satisfies:

$$u_0 \in C^{1,1}(\mathbb{R}^d), \quad \lim_{|x| \rightarrow \infty} \frac{u_0(x)}{|x|} = +\infty, \quad \exists R_0 > 0 : u_0 = u_0^{**} \text{ outside } \overline{B}_{R_0}. \quad (2.5)$$

The coercivity assumption guarantees that the infimum in formula (2.2) is actually achieved. The assumption that u_0 and u_0^{**} agree outside of some ball, will be convenient and allow us to work mainly on a ball instead of on the whole space. Indeed, thanks to (2.5), we claim that there exists $R \geq R_0$ such that, for any $x \in \overline{B}_{R_0}$, in the formula (2.2), it is enough to restrict the minimization to points x_i in \overline{B}_R :

$$u_0^{**}(x) = \inf \left\{ \sum_{i=1}^{d+1} \lambda_i u_0(x_i) : \lambda_i \geq 0, \sum_{i=1}^{d+1} \lambda_i = 1, \sum_{i=1}^{d+1} \lambda_i x_i = x, x_i \in \overline{B}_R \right\}, \quad (2.6)$$

for every $x \in \overline{B}_{R_0}$. To prove this claim, it is enough to remark that whenever the $\lambda_i > 0$'s and x_i 's solve (2.2) then u_0^{**} is actually affine on the convex hull of the points x_i , and then to note that the sets on which u_0^{**} is affine are necessarily bounded thanks to the coercivity of u_0 and the fact that $u_0 = u_0^{**}$ outside \overline{B}_{R_0} . Finally, the assumption that u_0 is $C^{1,1}$ implies that so is u_0^{**} (see [6]) and we will see that it also implies that $u(t, \cdot)$ remains $C^{1,1}$.

3 Stochastic control representation

As we shall see (but this should already be clear to stochastic control-oriented readers), the value function of (2.4) is in fact characterized by the PDE:

$$\partial_t v = \min(0, \lambda_1(D^2 v)) \quad (3.1)$$

in the viscosity sense that we now recall (for the sake of simplicity, we will restrict ourselves to the framework of continuous solutions which is sufficient in our context):

Definition 3.1. *Let Ω be some open subset of \mathbb{R}^d and let v be continuous on $(0, +\infty) \times \Omega$, then v is:*

- *a viscosity subsolution of (3.1) on $(0, +\infty) \times \Omega$ if for every smooth function $\varphi \in C^2((0, +\infty) \times \Omega)$ and every $(t_0, x_0) \in (0, +\infty) \times \Omega$ such that $(v - \varphi)(t_0, x_0) = \max_{(0, +\infty) \times \Omega}(v - \varphi)$ one has*

$$\partial_t \varphi(t_0, x_0) \leq \min(0, \lambda_1(D^2 \varphi(t_0, x_0))),$$

- *a viscosity supersolution of (3.1) on $(0, +\infty) \times \Omega$ if for every smooth function $\varphi \in C^2((0, +\infty) \times \Omega)$ and every $(t_0, x_0) \in (0, +\infty) \times \Omega$ such that $(v - \varphi)(t_0, x_0) = \min_{(0, +\infty) \times \Omega}(v - \varphi)$ one has*

$$\partial_t \varphi(t_0, x_0) \geq \min(0, \lambda_1(D^2 \varphi(t_0, x_0))),$$

- *a viscosity solution of (3.1) on $(0, +\infty) \times \Omega$ if it is both a viscosity subsolution and a viscosity supersolution.*

We then have the following stochastic representation formula for (2.1):

Theorem 3.2. *Under assumption (2.5), there is a unique continuous function u on $[0, +\infty) \times \mathbb{R}^d$ that agrees with u_0 at $t = 0$, that is a viscosity solution*

of (2.1) and that agrees with u_0^{**} outside B_{R_0} . It admits the following representation

$$u(t, x) = \inf_{\sigma: |\sigma| \leq 1} \left\{ \mathbb{E} \left(u_0 \left(x + \int_0^t \sqrt{2} \sigma_s dW_s \right) \right) \right\}, \quad t \geq 0, \quad x \in \mathbb{R}^d \quad (3.2)$$

where $(W_s)_{s \geq 0}$ is a standard Brownian motion and $|\sigma|$ stands for the matrix norm $|\sigma| := \sqrt{\text{Tr}(\sigma \sigma^T)}$.

Proof. Recalling that for every symmetric matrix S one has

$$\min(0, \lambda_1(S)) = \min_{|\sigma| \leq 1} \text{Tr}(\sigma \sigma^T S),$$

the fact that formula (3.2) actually defines a viscosity solution of (2.1) is a classical fact from stochastic control theory (see for instance [5] or [11]), assumption (2.5) guarantees that the function defined by (3.2) agrees with u_0^{**} outside B_{R_0} . Continuity (local Lipschitz continuity in fact) of the value function defined by (3.2) will be established in section 4. It follows from well-known parabolic comparison principles (e.g Theorem 4.1 in [3]) that if $T > 0$, v_1 (respectively v_2) is a viscosity subsolution (respectively supersolution) of (2.1) on $(0, T) \times B_{R_0}$ such that $v_1(t, x) \leq v_2(t, x)$ for $(t, x) \in \{0\} \times B_{R_0} \cup [0, T] \times \partial B_{R_0}$ then $v_1 \leq v_2$ on $[0, T] \times B_{R_0}$. This proves the uniqueness of a viscosity solution of (2.1) which equals u_0 at $t = 0$ and that agrees with u_0^{**} outside B_{R_0} . □

Remark 3.3. Optimal feedback control. Very formally, if the solution u of the PDE were very well-behaved then, as usual in control theory, one could find an optimal feedback (Markov) control depending on D^2u (since there is no drift). Introduce a time-dependent vector field $Z = Z(t, y)$, as follows. If $\lambda_1(D^2u(t, y)) < 0$, then let $Z(t, y)$ be a unit eigenvector associated to $\lambda_1(D^2u(t, y))$ and let $Z(t, y) = 0$ otherwise. So that in any case:

$$\text{Tr}(\sigma(t, y) \sigma(t, y)^T D^2u(t, y)) = \min(0, \lambda_1(D^2u(t, y))), \quad \text{and} \quad |\sigma(t, y)| \leq 1,$$

where σ is the projector

$$\sigma(t, y) := Z(t, y) \otimes Z(t, y).$$

Of course the problem is that σ is not well-defined: not only u does not need to be C^2 but also it may be the case that $\lambda_1 < 0$ has multiplicity larger than 2. Ignoring those serious issues, let us consider the SDE:

$$dY_t = \sqrt{2} \sigma(t, Y_t) dW_t$$

then σ is (again very formally) an optimal feedback control. We then have for $t > s \geq 0$

$$u(t, y) = \mathbb{E}[u(s, Y_t) | Y_s = y],$$

and, formally, the envelope theorem gives

$$\nabla u(t, y) = \mathbb{E}[\nabla u(s, Y_t) | Y_s = y].$$

Finally, notice that the drift of $u(t, Y_t)$ is the nonpositive quantity given by

$$\partial_t u(t, Y_t) + \text{Tr}(\sigma \sigma^T D^2 u(t, Y_t)) = 2 \min(0, \lambda_1(D^2 u(t, Y_t))).$$

Remark 3.4. One has $u_0^{**} \leq u(t, \cdot) \leq u_0$ and $u(\cdot, x)$ is nonincreasing and thus monotonically converges to $v(x) := \lim_{t \rightarrow \infty} u(t, x) = \inf_{t > 0} u(t, x)$. Now, as shown by Vese in [12], v is necessarily convex (it is a viscosity solution of the stationary equation) and since $u_0^{**} \leq u(t, \cdot) \leq u_0$ this gives $v = u_0^{**}$. In other words, u pointwise monotonically converges to the convex envelope of the initial condition. Of course, in view of the representation formula (3.2) and (2.2) this convergence is not surprising. We shall see in the next sections how (3.2) can easily give much more precise informations and provide in a simple way very strong convergence estimates.

4 Regularity properties of u

Lemma 4.1. *If $M > 0$ is such that $u_0 - \frac{M}{2}|\cdot|^2$ is concave then $u(t, \cdot) - \frac{M}{2}|\cdot|^2$ is concave for every $t > 0$.*

Proof. Set $v_0 := u_0 - \frac{M}{2}|\cdot|^2$ and let $(X_\alpha)_{\alpha \in A}$ be a family of centered, \mathbb{R}^d -valued, square integrable random variables, then define

$$\varphi(x) = \inf_{\alpha \in A} \mathbb{E}(u_0(x + X_\alpha)), \quad x \in \mathbb{R}^d$$

we then have

$$\varphi(x) - \frac{M}{2}|x|^2 = \inf_{\alpha \in A} \{\mathbb{E}(v_0(x + X_\alpha) + \frac{M}{2}|X_\alpha|^2)\}$$

so that $\varphi - \frac{M}{2}|\cdot|^2$ is concave as an infimum of concave functions. This proves the desired claim. □

Proposition 4.2. *Let u_0 be $C^{1,1}$ and let $M := \|D^2u_0\|_\infty$, then for every $(t, s) \in (0, +\infty)$ and every $x \in \mathbb{R}^d$ one has*

$$|u(t, x) - u(s, x)| \leq M|s - t| \quad (4.1)$$

and $u(t, \cdot)$ is $C^{1,1}$ for every t and more precisely, one has $\|D^2u(t, \cdot)\|_\infty \leq M$.

Proof. Let $0 < t < s$, we already know that $u(t, \cdot) \geq u(s, \cdot)$. Let us assume for a moment that u_0 is smooth and let $x \in \mathbb{R}^d$ and let σ be an adapted process with values in the set of matrices with norm less than 1 such that

$$\mathbb{E}\left(u_0\left(x + \int_0^s \sqrt{2}\sigma dW\right)\right) \leq u(s, x) + \varepsilon$$

then defining

$$Y_h := x + \int_0^h \sqrt{2}\sigma dW, \quad Z_h := u_0(Y_h), \quad s \geq h \geq 0$$

thanks to Itô's formula, we thus get:

$$\begin{aligned} u(t, x) &\leq \mathbb{E}(Z_t) \leq u(s, x) + \varepsilon - \mathbb{E}(Z_s - Z_t) \\ &= u(s, x) + \varepsilon - \mathbb{E}\left(\int_t^s \text{Tr}(\sigma\sigma^T D^2u_0(Y_h))dh\right) \\ &\leq u(s, x) + \varepsilon + M(s - t) \end{aligned}$$

and we conclude that (4.1) holds by letting $\varepsilon \rightarrow 0^+$. In the general case, one applies the same argument to the regularization $\rho_n \star u_0$ (where ρ_n is, as usual, a sequence of mollifiers) and then passes to the limit to obtain (4.1).

Using (2.1), we then quite easily obtain that

$$\lambda_1(D^2v(t, x)) \geq 0 \text{ on } (0, +\infty) \times \mathbb{R}^d, \text{ with } v(t, x) := u(t, x) + \frac{M}{2}|x|^2$$

in the viscosity sense which means that as soon as φ is smooth and $v - \varphi$ has a (local or global) maximum at $(t_0, x_0) \in (0, +\infty) \times \mathbb{R}^d$ then $\lambda_1(D^2\varphi(t_0, x_0)) \geq 0$. To see that this implies that $v(t, \cdot)$ is convex, we invoke the same arguments as in Lemma 1 in [1]. Assume on the contrary that there are $t_0 > 0$, x_0, y_0 in \mathbb{R}^d and $\lambda \in (0, 1)$ such that $v(t_0, \lambda x_0 + (1 - \lambda)y_0) > \lambda v(t_0, x_0) + (1 - \lambda)v(t_0, y_0)$. Without loss of generality, denoting elements of \mathbb{R}^d as (x_1, x') in $\mathbb{R} \times \mathbb{R}^{d-1}$, we may assume that $y_0 = (0, 0)$, $x_0 = (1, 0)$, we may also assume (adding if necessary some suitable affine function to v) that $v(t_0, 0) = v(t_0, (1, 0)) < 0$ and $v(t_0, (\lambda, 0)) > 0$. We then choose $h \in (0, t_0)$ and $r > 0$ such that

$$v(t, (0, x')) < 0, \quad v(t, (1, x')) < 0, \quad \forall (t, x') \in [t_0 - h, t_0 + h] \times B_r. \quad (4.2)$$

We then define

$$\Omega := \{(x_1, x') \in (0, 1) \times B_r\}, \quad Q := (t_0 - h, t_0 + h) \times \Omega$$

and choose $\alpha > 0$ such that $v(t_0, (\lambda, 0)) > \frac{\alpha\lambda(1-\lambda)}{2}$. We then define

$$\varphi(t, (x_1, x')) := \frac{\alpha}{2}x_1(1-x_1) + \frac{\beta}{2}|x'|^2 + \frac{\gamma}{2}(t-t_0)^2$$

with β and γ chosen so that

$$\beta r^2 \geq 2 \max_{\overline{Q}} v, \quad \gamma h^2 \geq 2 \max_{\overline{Q}} v. \quad (4.3)$$

We then have $v(t_0, (\lambda, 0)) - \varphi(t_0, (\lambda, 0)) > 0$ and by (4.2)-(4.3), $v - \varphi \leq 0$ on ∂Q , hence $v - \varphi$ achieves its maximum on \overline{Q} at an interior point of Q , but at this point one should have $0 \leq \lambda_1(D^2\varphi) = -\alpha$ which gives the desired contradiction. This proves that $u(t, \cdot) + \frac{M}{2}|\cdot|^2$ is convex for every t . Together with lemma 4.1 this enables us to conclude that u remains semiconvex and semiconcave is hence $C^{1,1}$ with the estimate $\|D^2u(t, \cdot)\|_\infty \leq M$. \square

Proceeding as in the proof of the two previous results and using the fact that the PDE is autonomous, one gets:

Corollary 4.3. *Suppose u_0 satisfies (2.5), then $\text{Essinf} \lambda_1(D^2u(t, \cdot))$ is non-decreasing with respect to t and $\text{Esssup} \lambda_d(D^2u(t, \cdot))$ is nonincreasing with respect to t (where λ_d stands for the largest eigenvalue).*

5 Exponential convergence to the convex envelope

Before we state our result concerning the convergence of $u(t, \cdot)$ to u_0^{**} , we need two elementary lemmas.

Lemma 5.1. *Let $v : \mathbb{R}^d \rightarrow \mathbb{R}$ and $M \geq 0$ be such that $v + \frac{M}{2}|\cdot|^2$ is convex, then for every $r > 0$ one has:*

$$\|\nabla v\|_{L^\infty(B_r)} \leq 2 \left(M \|v\|_{L^\infty(B_{r+r'})} \right)^{1/2} \text{ with } r' = \frac{2}{Mr} \|v\|_{L^\infty(B_{2r})} + \frac{r}{2}. \quad (5.1)$$

Proof. Let $r > 0$, $R > 0$, for $x \in B_r$ a point of differentiability of v (which is a.e. the case) and $h \in B_R$ in \mathbb{R}^d , one first has

$$2\|v\|_{L^\infty(B_{r+R})} \geq v(x+h) - v(x) \geq \nabla v(x) \cdot h - \frac{M}{2}|h|^2. \quad (5.2)$$

Taking $r = R$, $h = r\nabla v(x)/|\nabla v(x)|$ and maximizing with respect to $x \in B_r$ thus gives

$$\|\nabla v\|_{L^\infty(B_r)} \leq \frac{2}{r}\|v\|_{L^\infty(B_{2r})} + \frac{Mr}{2}. \quad (5.3)$$

We then take $R = r'$ with r' defined by (5.1) and set $h = \nabla v(x)/M$, thanks to (5.3), $h \in B_R$, using (5.2) again, we then get

$$\frac{|\nabla v(x)|^2}{2M} \leq 2\|v\|_{L^\infty(B_{r+r'})}, \quad \forall x \in B_r$$

which finally gives (5.1). □

Lemma 5.2. *Let (B_t) be a standard one-dimensional Brownian motion, let $r > 0$, $x \in (-r, r)$ and*

$$\tau := \inf\{t > 0 : x + B_t \notin [-r, r]\}$$

then for every $t > 0$, one has

$$\mathbb{P}(\tau \geq t) \leq q(r)^{t-1}, \quad \text{with } q(r) := \frac{1}{\sqrt{2\pi}} \int_{-2r}^{2r} e^{-\frac{s^2}{2}} ds.$$

Proof. Let n be the integer part of t , we then have

$$\mathbb{P}(\tau \geq t) \leq \mathbb{P}(|B_k - B_{k-1}| \leq 2r, \quad \text{for } k = 1, \dots, n)$$

and since $(B_k - B_{k-1})_{k=1, \dots, n}$ are independent and normally distributed random variables, we immediately get the desired estimate. □

Our main result then reads as

Theorem 5.3. *Under assumption (2.5), there exists $C \geq 0$ and $\lambda > 0$ such that*

$$\|u(t, \cdot) - u_0^{**}\|_{L^\infty} \leq Ce^{-\lambda t}, \quad \forall t \geq 0, \quad (5.4)$$

and

$$\|\nabla u(t, \cdot) - \nabla u_0^{**}\|_{L^\infty} \leq Ce^{-\lambda t}, \quad \forall t \geq 0. \quad (5.5)$$

Proof. First let us remark that if $x \notin \overline{B}_{R_0}$, there is nothing to prove. Let us then recall (2.6), thanks to (2.5), there is some ball \overline{B}_R containing \overline{B}_{R_0} , such that for any $x \in \overline{B}_{R_0}$, in the formula (2.2), it is enough to restrict the minimization to points x_i in \overline{B}_R . Let then $x \in B_{R_0}$, let $(x_1, \dots, x_{d+1}) \in \overline{B}_R^{d+1}$ and $(\lambda_1, \dots, \lambda_{d+1})$ be nonnegative such that

$$\sum_{i=1}^{d+1} \lambda_i = 1, \quad \sum_{i=1}^{d+1} \lambda_i x_i = x, \quad \sum_{i=1}^{d+1} \lambda_i u_0(x_i) = u_0^{**}(x). \quad (5.6)$$

We shall also assume that the points (x_1, \dots, x_{d+1}) are affinely independent (and this is actually without loss of generality for what follows), the coefficients λ_i are then uniquely defined and are the unique barycentric coordinates of x in the simplex K which is the convex hull of the points (x_1, \dots, x_{d+1}) . We shall also assume that all the coefficients λ_i are strictly positive (again this is not a restriction).

Let $\varepsilon > 0$ and let $\sigma_s = 0$, for $s \in [0, \varepsilon]$, then set

$$v_1 := \frac{W_\varepsilon}{|W_\varepsilon|}, \tau_1 := \inf\{t \geq \varepsilon : x + \sqrt{2}v_1 \otimes v_1(W_t - W_\varepsilon) \notin K\}$$

and $\sigma_s = v_1 \otimes v_1$ for $s \in (\varepsilon, \tau_1]$. By construction, $x + \sqrt{2}v_1 \otimes v_1(W_{\tau_1} - W_\varepsilon)$ a.s. belongs to a facet of K of dimension $d-1$. Let us denote by K_1 this facet and by E_1 the hyperplane parallel to this facet. Let then v_2 be \mathcal{F}_ε -measurable and uniformly distributed on $S^d \cap E_1$ and define

$$\tau_2 := \inf\{t \geq \tau_1 : x + \sqrt{2}v_1 \otimes v_1(W_{\tau_1} - W_\varepsilon) + \sqrt{2}v_2 \otimes v_2(W_t - W_{\tau_1}) \notin K_1\}$$

and $\sigma_s = v_2 \otimes v_2$ for $s \in (\tau_1, \tau_2]$.

We repeat inductively this construction d times and define successive (random and adapted) times τ_k , $k = 1, \dots, d$, directions v_1, \dots, v_k , and a piecewise constant control $\sigma_s = v_k \otimes v_k$ for $s \in (\tau_{k-1}, \tau_k]$, in such a way that $x + \int_0^t \sqrt{2}\sigma_s dW_s$ belongs to a facet K_k of dimension $d-k$ for $t \in [\tau_k, \tau_{k+1}]$. Let us extend the control σ by 0 after time τ_d and set

$$Y_t := x + \sqrt{2} \int_0^t \sigma_s dW_s = Y_{t \wedge \tau_d}.$$

and remark that at time τ_d the previous process has hit one of the vertices of K . By construction $(Y_t)_t$ is a continuous martingale and it is bounded since it takes values in the compact K , it therefore converges to Y_{τ_d} which is a discrete random variables with values in the vertices of K , $\{x_1, \dots, x_{d+1}\}$, we then have

$$\mathbb{E}(Y_{\tau_d}) = x = \sum_{i=1}^{d+1} \mathbb{P}(Y_{\tau_d} = x_i) x_i$$

which implies that $\mathbb{P}(Y_{\tau_d} = x_i) = \lambda_i$ by uniqueness of the barycentric coordinates. We thus have:

$$u_0^{**}(x) = \mathbb{E}(u_0(Y_{\tau_d})) \quad (5.7)$$

and then using the fact that Y_t takes values in K and that u_0 is locally Lipschitz:

$$\begin{aligned} u(t, x) &\leq \mathbb{E}(u_0(Y_t)) \leq \mathbb{E}(u_0(Y_{\tau_d})) + \|\nabla u_0\|_{L^\infty(K)} \mathbb{E}(|Y_t - Y_{\tau_d}|) \\ &\leq u_0^{**}(x) + \text{diam}(K) \|\nabla u_0\|_{L^\infty(K)} \mathbb{P}(\tau_d \geq t) \end{aligned}$$

We then remark that

$$\{\tau_d \geq t\} \subset \bigcup_{k=1}^d \left\{ T_k \geq \frac{t - \varepsilon}{d} \right\}$$

where the T_k 's are the times the process $(Y_s)_s$ spends on the (random) facet K_{k-1} (setting $K_0 = K$), the previous probabilities can therefore be estimated by the probability that a one-dimensional Brownian motion spends more than $\frac{t-\varepsilon}{d}$ time in the interval $[-\text{diam}(K), \text{diam}(K)]$. Using lemma 5.2, we thus get

$$\mathbb{P}(\tau_d \geq t) \leq M e^{-\lambda(t-\varepsilon)}$$

for constants M and $\lambda > 0$ that depend only on d and $\text{diam}(K)$. Letting $\varepsilon \rightarrow 0$, we then obtain

$$u(t, x) \leq u_0^{**}(x) + \text{diam}(K) \|\nabla u_0\|_{L^\infty(K)} M e^{-\lambda t}$$

since we already know that $u(t, \cdot) \geq u_0^{**}$, this terminates the proof of (5.4).

Finally, the estimate (5.5) easily follows from (5.4), lemma 5.1 and the fact that $u(t, \cdot) - u_0^{**}$ remains uniformly semiconcave thanks to lemma 4.1. \square

Remark 5.4. Let us remark that in the inequality (5.4) in theorem 5.3, the constant λ only depends on the dimension and the diameter of the *faces* of the convex envelope on the set where $\{u_0 > u_0^{**}\}$ whereas the constant C also depends on the Lipschitz constant of u_0 on the set of such faces. In (5.5), C also depends on $\|D^2 u_0\|_{L^\infty}$.

6 Concluding remarks

6.1 A nonautonomous gradient flow

As a possible application of theorem 5.3, one can prove convergence results for the Cauchy problem for the non-autonomous gradient flow:

$$\dot{x}(t) = -\nabla u(t, x(t)), \quad t > 0, \quad x(0) = x_0 \quad (6.1)$$

where $x_0 \in \mathbb{R}^d$ is an arbitrary initial position. Thanks to proposition 4.2, the previous Cauchy problem possesses a unique solution that is defined for all positive times. An easy consequence of theorem 5.3 then reads

Theorem 6.1. *Let $x_0 \in \mathbb{R}^d$, and let $x(\cdot)$ be the solution of the Cauchy problem (6.1), then $x(t)$ converges as $t \rightarrow \infty$ to some point y_∞ that is a (global) minimum of u_0^{**} .*

Proof. Let us denote by F the (convex and compact) set where u_0^{**} attains its minimum. Let $y \in F$, since $\nabla u_0^{**}(y) = 0$, using the convexity of u_0^{**} and (5.5), we get

$$\begin{aligned} \frac{d}{dt} \left(\frac{1}{2} |x(t) - y|^2 \right) &= \langle \nabla u_0^{**}(y) - \nabla u_0^{**}(x(t)), x(t) - y \rangle + \\ &\quad \langle \nabla u_0^{**}(x(t)) - \nabla u(t, x(t)), x(t) - y \rangle \\ &\leq C e^{-\lambda t} |x(t) - y|. \end{aligned}$$

From which we easily deduce that $|x(t) - y| + \frac{C}{\lambda} e^{-\lambda t}$ is nonincreasing for every $y \in F$ so that $\min_{y \in F} |x(t) - y| + \frac{C}{\lambda} e^{-\lambda t}$ is also nonincreasing. There exists therefore some $d_\infty \geq 0$ such that

$$d(x(t), F) := \min_{y \in F} |x(t) - y| \rightarrow d_\infty \text{ as } t \rightarrow \infty.$$

Now we claim that $d_\infty = 0$; assume on the contrary that $d_\infty > 0$ and let $y \in F$, we then have

$$\delta := \min \{ \langle \nabla u_0^{**}(x) - \nabla u_0^{**}(y), x - y \rangle : d(x, F) = d_\infty \} > 0$$

so that by the same computations as above, we obtain that for large enough t , one has

$$\frac{d}{dt} \left(\frac{1}{2} |x(t) - y|^2 \right) \leq -\frac{\delta}{2}$$

which contradicts the convergence of $|x(t) - y|$ as $t \rightarrow +\infty$. We thus have proved that $d(x(t), F) \rightarrow 0$ as $t \rightarrow +\infty$ so that all limit points of the trajectory $x(\cdot)$ belong to F . Let $y_1 = \lim_n x(t_n)$ and $y_2 = \lim_n x(s_n)$ with $t_n, s_n \rightarrow \infty$ be two such limit points, since $|x(t) - y_i|$ converges as $t \rightarrow \infty$ for $i = 1, 2$, we deduce that $|y_1 - y_2| = \lim_n |x(t_n) - y_2| = \lim_n |x(s_n) - y_2| = 0$. Together with the compactness of F , this proves that $x(t)$ converges to some $y_\infty \in F$ as $t \rightarrow \infty$. \square

6.2 Interpolation with a geometric flow

This final section is devoted to an informal discussion on a natural interpolation between the HJB equation (2.1) and some geometric flow. Let us indeed remark that the HJB equation

$$\partial_t u = \min(0, \lambda_1(D^2 u)), \quad u|_{t=0} = u_0 \quad (6.2)$$

can be imbedded into the family of PDE depending on a parameter $\theta \geq 0$:

$$\partial_t v = \min(0, \lambda_1(D^2 v + \theta \nabla v \otimes \nabla v)), \quad v|_{t=0} = v_0. \quad (6.3)$$

Now, let us formally remark that for $\theta > 0$, v solves (6.3) if and only if $u := e^{\theta v}$ solves (6.2) with initial condition $u_0 := e^{\theta v_0}$ and therefore admits the representation formula:

$$v(t, x) = \frac{1}{\theta} \log \left(\inf_{\sigma: |\sigma| \leq 1} \mathbb{E} \left(\exp(\theta v_0(x + \int_0^t \sqrt{2} \sigma_s dW_s)) \right) \right), \quad (6.4)$$

and, as $t \rightarrow \infty$, $e^{\theta v(t, \cdot)}$ converges to the convex envelope of $e^{\theta v_0}$. Letting θ tend to $+\infty$ in (6.3), we formally obtain:

$$\partial_t v = \min(0, \lambda_1(D^2 v|_{\nabla v^\perp})), \quad v|_{t=0} = v_0 \quad (6.5)$$

which is a geometric flow of motion by the principal negative curvature. One of course expects that the evolution along this flow will tend to convexify the level-sets of v_0 . It is therefore natural to view (6.5) as a *quasi-convexification* equation. Equation (6.3) therefore interpolates between a convexifying and a quasi-convexifying evolution. There is also a stochastic representation formula for (6.5) due to Soner and Touzi [10], which relies on a *stochastic target* problem different in nature from (6.4).

Acknowledgments

G.C. wishes to thank the support of the ANR through the projects ANR-07-BLAN-0235 OTARIE and ANR-09-JCJC-0096-01 EVAMEF. A.G. gratefully acknowledges support from Chaire EDF-Calyon *Finance and Développement Durable* and FiME, Laboratoire de Finance des Marchés de l'Énergie (www.fime-lab.org), and Chaire Axa *Assurance et Risques Majeurs*. Both authors are grateful to an anonymous referee for several suggestions that led to an improved presentation.

References

- [1] O. Alvarez, J.-M. Lasry, P.-L. Lions, Convex viscosity solutions and state constraints. *J. Math. Pures Appl.* (9) 76 (1997), no. 3, 265-288.

- [2] L. Caffarelli, X. Cabré , Fully nonlinear elliptic equations, American Mathematical Society Colloquium Publications, 43, American Mathematical Society, Providence, RI, 1995.
- [3] Y.G. Chen, Y. Giga, S. Goto, Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. *J. Differential Geom.* 33 (1991), no. 3, 749-786.
- [4] M.G. Crandall, H. Ishii, P.-L. Lions, User's guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc.* 27 (1992), 1-67.
- [5] W. H. Fleming, H.M. Soner, *Controlled Markov Processes and Viscosity Solutions*, Graduate Studies in Mathematics, 58, Springer-Verlag, Applications of Mathematics, (1993).
- [6] B. Kirchheim, J. Kristensen, Differentiability of convex envelopes. *C. R. Acad. Sci. Paris Sér. I Math.* 333 (2001), no. 8, 725-728.
- [7] A. Oberman, The convex envelope is the solution of a nonlinear obstacle problem, *Proc. Amer. Math. Soc.* 135 (2007), no. 6, 1689-1694.
- [8] A. Oberman, Computing the convex envelope using a nonlinear partial differential equation, *Mathematical Models and Methods in Applied Sciences (M3AS)*, Vol. 18. No 5 (2008) 759-780.
- [9] A. Oberman, L. Silvestre, The Dirichlet Problem for the Convex Envelope, to appear in *Trans. Amer. Math. Soc.*
- [10] H.M. Soner, N. Touzi, Stochastic representation of mean curvature type geometric flows, *Annals of Probability*, 31 (2003), 1145-1165.
- [11] N. Touzi, Stochastic control and application to Finance, Lecture Notes available at <http://www.cmap.polytechnique.fr/~touzi/>.
- [12] L. Vese, A method to convexify functions via curve evolution. *Comm. Partial Differential Equations* 24 (1999), no. 9-10, 1573-1591.