

BRAIN ABSCESS ASSOCIATED WITH AGGREGATIBACTER ACTINOMYCETEMCOMITANS: CASE REPORT AND REVIEW OF LITERATURE

Janette Rahamat-Langendoen, Marit van Vonderen, Lex Engström, Willem Manson, Arie Jan van Winkelhoff, Esther Mooi-Kokenberg

▶ To cite this version:

Janette Rahamat-Langendoen, Marit van Vonderen, Lex Engström, Willem Manson, Arie Jan van Winkelhoff, et al.. BRAIN ABSCESS ASSOCIATED WITH AGGREGATIBACTER ACTINO-MYCETEMCOMITANS: CASE REPORT AND REVIEW OF LITERATURE. Journal of Clinical Periodontology, 2011, 38 (8), pp.702. 10.1111/j.1600-051X.2011.01737.x. hal-00637819

HAL Id: hal-00637819

https://hal.science/hal-00637819

Submitted on 3 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



BRAIN ABSCESS ASSOCIATED WITH AGGREGATIBACTER ACTINOMYCETEMCOMITANS: CASE REPORT AND REVIEW OF LITERATURE

Journal:	Journal of Clinical Periodontology
Manuscript ID:	CPE-01-11-3091.R1
Manuscript Type:	Case Report
Date Submitted by the Author:	04-Apr-2011
Complete List of Authors:	Rahamat-Langendoen, Janette; University Medical Centre Groningen, Groningen University, Department of Microbiology van Vonderen, Marit; Medical Centre Leeuwarden, Department of Internal Medicine Engström, Lex; Medical Centre Leeuwarden, Department of Neurology Manson, Willem; University Medical Centre Groningen, Groningen University, Department of Microbiology van Winkelhoff, Arie; Groningen University, Department of Microbiology Mooi-Kokenberg, Esther; Izore, Centre for Infectious Diseases Friesland, Department of Microbiology
Topic:	Diagnosis
Keywords:	Periodontal disease
Main Methodology:	Microbiology

SCHOLARONE™ Manuscripts

BRAIN ABSCESS ASSOCIATED WITH AGGREGATIBACTER

ACTINOMYCETEMCOMITANS: CASE REPORT AND REVIEW OF LITERATURE

Janette C. Rahamat-Langendoen*, †, Marit G.A. van Vonderen§, Lex J. Engström[∥], Willem L.

Manson[†], Arie Jan van Winkelhoff[†], [‡], Esther A.N.M. Mooi-Kokenberg^{*}

*Department of Microbiology, Izore, Centre for Infectious Diseases Friesland, Leeuwarden, the

Netherlands

[†]University Medical Centre Groningen, Groningen University, Department of Medical

Microbiology, Groningen, the Netherlands

[‡]University Medical Centre Groningen, Groningen University, Centre for Dentistry and Oral

Hygiene, Groningen, the Netherlands

§Department of Internal Medicine, Medical Centre Leeuwarden, Leeuwarden, the Netherlands

Department of Neurology, Medical Centre Leeuwarden, Leeuwarden, the Netherlands

Corresponding author:

J.C. Rahamat-Langendoen

University Medical Centre Groningen

Department of Medical Microbiology

P.O. Box 30 001

3700 RB Groningen

The Netherlands

e-mail: j.c.rahamat@mmb.umcg.nl

phone: + 31 50 36 13480

fax: +31 50 36 33 528

Deleted: DUE TO

Deleted: Arie Jan van Winkelhoff^{†,‡},

Word count: <u>1625</u>

Figures: 2

Tables: 1

Short running title:

Poor dentition risk for severe infection

mycetemcon. Keywords: Aggregatibacter actinomycetemcomitans, periodontal disease, brain abscess, extra oral infection, sequence analysis

Financial support: no financial support was received.

Conflict of interest: none



Abstract

Introduction

Aggregatibacter actinomycetemcomitans is considered a major pathogen in localized and generalized aggressive periodontitis. A. actinomycetemcomitans has been found in various extra oral infections and most frequently in endocarditis. We report a patient with multiple brain abscesses due to infection with A. actinomycetemcomitans and review the English language literature related to this subject.

Case report

A 42-year old patient with no underlying medical conditions presented with multiple brain lesions initially thought to be metastatic lesions of a tumor of unknown origin. Findings during drainage and subsequent histopathological conclusions made infection more likely. Culture of drained material remained negative; however 16S rDNA PCR and sequence analysis on direct material revealed *A. actinomycetemcomitans* as the causative agent of the infection. The most likely source of infection was the poor dentition of the patient. After repeated drainage of the lesions and antibiotic treatment the patient gradually improved, although cognitive impairment remained.

Conclusions

Our report illustrates that a poor dental condition, notably destructive periodontal disease, can be a risk for <u>life threatening extra oral disease</u>, and thus contributes to the total inflammatory burden of the body.

Deleted: intracranial abscess formation. We have also shown that in cases of culture-negative results, molecular detection and identification can be an essential adjunctive tool.

Clinical relevance

Scientific rationale for study

Aggregatibacter actinomycetemcomitans is a major pathogen in periodontitis. In this case report we show the relevancy of oral health in relation to life threatening extra oral infection.

Principal findings

A 42-year old patient with poor dentition was found to have multiple brain abscesses caused by

Aggregatibacter actinomycetemcomitans, detected by 16S rDNA PCR and sequence analysis on direct material.

Practical implications

Poor oral health is an important contributing factor to severe extra oral infection. We illustrate the

diagnostic value of gene amplification and sequencing from direct specimens for patients with culture negative infections.

Deleted: Poor o

Deleted: ral health is a risk factor for intracranial infection by a periodontal pathogen.

Introduction

Dental condition has been linked to several non-oral diseases such as infectious bowel diseases, atherosclerosis, coronary heart diseases, stroke, diabetes and rheumatoid arthritis. The link between destructive periodontal disease and these chronic disorders can be explained by common risk factors such as smoking, stress and possibly genetic traits.

Periodontitis is a bacterial infection of the tooth-supporting tissues characterized by a large inflamed and necrotic area associated with a high bacterial load and destruction of the alveolar bone. (Nesse et al 2008) The infected periodontium can act as a focus for extra oral infections by dissemination of periodontal bacteria to non-oral body sites. Periodontitis may thus contribute to the total inflammatory burden of the body.

Aggregatibacter actinomycetemcomitans is considered a major pathogen in localized and generalized aggressive periodontitis.(Zambon 1985) A. actinomycetemcomitans has been found in various extra oral infections, most frequently in endocarditis.(van Winkelhoff & Slots 1999) We report a patient with multiple brain abscesses due to infection with A. actinomycetemcomitans and review the English language literature related to this subject.

Case report

A 42-year old man was admitted to the Medical Centre Leeuwarden with a 3 days history of confusion and reduced consciousness. Besides heavy smoking and alcohol abuse there was no relevant medical history. Physical examination revealed a slowly reacting man, who did not respond adequately. Oral examination showed severe candidiasis and the orthopantogram revealed caries profunda in multiple teeth and severe periodontal breakdown. (Figure 1) There was no heart murmur and petechiae were not seen. No focal signs were detected at neurological examination. Laboratory data revealed an elevated white blood count (17.1 x 10⁹/L) and a moderately elevated C-reactive protein (34 mg/L). Liver enzymes and renal function tests showed normal values. Computed Tomography (CT) of the brain showed multiple intracerebral lesions with, after injection of intravenous contrast, ring enhancement surrounded by oedema. (Figure 2) The lesions were primarily thought to be brain metastases of an unidentified tumour. Magnetic Resonance Imaging (MRI) confirmed the findings with no signs of lesions elsewhere in the body. A chest X-ray provided no indications for malignancy in the lungs. Serological tests for infection with HIV and Toxoplasma were negative. Blood cultures taken at admission were negative. The patient was treated with dexamethasone. A follow-up CT after 4 days showed no improvement regarding the cerebral lesions and the surrounding oedema. The patient was transferred to the University Medical Centre Groningen for stereotactic biopsy and further treatment. Upon biopsying the lesions appeared to be abscesses and pus was collected from a lesion in the right frontal region. Histopathological examination revealed reactive inflammatory changes with necrosis and no signs of malignancy. At gram staining no microorganisms and few leucocytes were seen. Blood, chocolate, and MacConkey's 3 agar and Fastidious Broth were inoculated and incubated in air with 5% CO₂ at 35°C. Anaerobic cultures were performed on selective media and incubated in 80% N₂, 10% H₂, and 10% CO₂. The plates were examined for the presence of colonies over the course of 10 days. Post-operative empiric intravenous antibiotic

treatment was initiated with ceftriaxone 2 gram bid, metronidazole 500 milligram bid and penicillin 6 times a day 2 million units. Aerobic and anaerobic culture remained negative and 16S rDNA PCR and sequence analysis on direct material was performed four days after drainage. Sequence data were compared with known sequences of 16S rDNA in the National Centre for Biotechnology Information Genbank. The highest similarity (99.8%) was found with A. actinomycetemcomitans. No other micro-organisms were detected. The patient continued to receive ceftriaxone and metronidazole intravenously, because a mixed anaerobic infection could not be excluded definitively. During his stay at the hospital the patient did not develop signs of endocarditis. A trans-thoracic echocardiogram made one week after operation showed no valvular vegetations. Besides his poor oral health no other focus of infection could be found. Nine days after the operation the patient deteriorated with reduced consciousness. CT of the brain showed no signs of expansion of the lesions, but because of the deterioration it was decided to drain four lesions. During this operation the oral surgeon extracted five teeth because of advanced periodontal breakdown. Although the patient gradually improved, supported by diminishment of the lesions on CT scan, cognitive impairment remained. In the reconvalescent period the patient developed focal epilepsy with secondary generalisation. Treatment with sodium valproate was started. Parenteral antibiotic treatment was continued for 6 weeks, after which the patient was transferred to a rehabilitation centre for intensive rehabilitation. His condition improved rapidly and after four weeks he was able to return to his home. Besides epileptic seizures which made four short term admissions to the hospital necessary during a one year follow up, he showed no

other sequelae of his disease.

Deleted:

Discussion

A. actinomycetemcomitans is a non motile, facultative anaerobic small gram negative coccobacillus.(von Graevenitz & Zbinden 2007) Its fastidious slow growing nature makes it difficult to isolate through culture, especially because it is often part of mixed infection with other bacteria. Incubation on blood and chocolate agar for at least 48 hours is needed in enhanced atmosphere with CO2. A. actinomycetemcomitans was first described as a human pathogen in 1912 by Klinger (Klinger 1912) It was initially isolated in conjunction with Actinomyces israelii, leading to the speculation that A. actinomycetemcomitans itself was not capable of causing disease. After the introduction of penicillin, which eradicated *Actinomyces*, pure cultures of A. actinomycetemcomitans were increasingly reported from various body sites and organs. (Page & King 1966) It is recognized as one of the major pathogens in periodontitis. Besides, it is reported as cause of soft tissue infection (usually in association with A. isrealii or A. meyeri) and endocarditis. Rarely, A. actinomycetemcomitans has been isolated from patients with endophthalmitis, septic arthritis, osteomyelitis, pneumonia, empyema, infected pacemaker wire and spinal epidural abscess. (Kaplan et al 1989; Yuan et al 1992; van Winkelhoff et al 1993; Binder et al 2003) Extra oral infections are thought to occur due to hematogenous dissemination from the infected periodontium. (van Winkelhoff & Slots 1999) Indeed, the poor dentition of our patient is thought to be the most likely source of infection, although we have no information regarding the presence or absence of A. actinomycetemcomitans in material of the extracted teeth.. Also, the oral candidiasis may have been a factor that has facilitated invasion of A. actinomycetemcomitans. Endocarditis was considered unlikely because of the absence of clinical signs of endocarditis and the absence of vegetations at echocardiography.

A review of the English language literature reveals 12 reports of brain abscesses due to infection

with A. actinomycetemcomitans. (Martin et al 1967; Burgher et al 1973; Brewer et al 1975; Ingham

et al 1978; Garner 1979; Churton & Greer 1980; Kaplan, Weber, Oddone, & Perfect 1989; Zijlstra

Formatted: Font: Not Italic

et al 1992; Kuijper et al 1992; Renton et al 1996; Stepanovic et al 2005) (Table 1) All but two reports are of male patients in the age of 19 to 62 years. In 6 patients the most probable source of infection was poor dentition or recent dental therapy, emphasizing the importance of considering the dental region as the origin of brain abscesses due to *A. actinomycetemcomitans*. Also, in 6 patients the insidious onset of symptoms was more characteristic of a neoplastic rather than an infectious process, as was the initial suspicion in our patient. Culture of the brain lesions revealed mixed infection with associated aerobic and anaerobic micro-organisms in six patients; not surprisingly, considering the source of infection, the identified micro-organisms were mainly common pathogens of the oro-pharyngeal region.

The optimal treatment of brain abscesses consists of surgical drainage combined with antibiotic therapy. *A. actinomycetemcomitans* is usually susceptible to cephalosporins (especially third generation cephalosporins), ampicillin, rifampin, fluoroquinolones, trimethoprim-sulfamethoxazole, aminoglycosides and chloramphenicol. In vitro, there is variable susceptibility to penicillin. Clindamycin and metronidazole have poor activity against *A. actinomycetemcomitans*, but metronidazole in combination with amoxicilline has shown strong synergistic effect. (Steinberg & Burd 2009)

The multiplicity of antibiotic regimes that has been used and the small number of patients do not permit identification of the ideal therapeutic regimen. Because of the potential of beta-lactamase production and the difficulties with susceptibility testing because of fastidious growth, third generation cephalosporins are now considered the drugs of choice. The empirical treatment of our patient consisted of ceftriaxone, metronidazole and penicillin, accounting for the microorganisms to be suspected in a brain abscess (streptococci, gram negative aerobes, and strict anaerobes). Based on the outcome of 16S rDNA PCR and sequence analysis, in combination with the most likely periodontal source of infection, it was decided to continue therapy for 6 weeks with ceftriaxone and metronidazole intravenously. To achieve definite improvement however a

second surgical drainage was necessary, which emphasizes the importance of surgical treatment of brain abscesses.

Our report also illustrates the diagnostic value of gene amplification and sequencing from direct specimens for patients with culture negative infections who have a high likelihood of a bacterial infection. Microbiological detection of organisms in brain abscesses primarily relies on microscopic examination and culturing of abscess pus specimen collected during neurosurgical drainage. However, the incidence of negative cultures can be high, especially in patients in whom antibiotic therapy started before collection of the specimen. (Tseng & Tseng 2006) Indeed, despite meticulous handling of the specimen and prolonged incubation in aerobic as well as anaerobic conditions, the cultures from our patient remained negative. Molecular detection of microorganisms in direct material can be used to overcome the limitations of culture-based bacteria detection. (Tsai et al 2004;Al Masalma et al 2009) Although susceptibility testing is not feasible for culture negative, PCR positive specimens, molecular identification can guide the choice of antimicrobial treatment, thereby shortening the period of empirical treatment with multiple broad spectrum antibiotics.

Conclusion

We describe a patient with multiple culture negative brain abscesses initially thought to be brain metastases of an unidentified tumour. Detection and identification of the oral pathogen A. actinomycetemcomitans was established by 16S rDNA sequencing on direct material, giving guidance to diagnosis and therapy. The patient's well documented destructive periodontal disease is considered the most likely focus of infection. Also, the oral cavity has been identified as the primary ecological niche of A. actinomycetemcomitans. This report illustrates that severe periodontal infection can be a risk for serious and life threatening extra oral disease.

Deleted: We have shown that poor oral health can be a risk factor for life threatening intracranial infection by a periodontal pathogen.



Reference List

- Al Masalma, M., Armougom, F., Scheld, W.M., Dufour, H., Roche, P.H., Drancourt, M., & Raoult, D. (2009) The expansion of the microbiological spectrum of brain abscesses with use of multiple 16S ribosomal DNA sequencing. *Clinical Infectious Diseases* 48, 1169-1178.
- 2. Binder, M.I., Chua, J., Kaiser, P.K., Mehta, N., & Isada, C.M. (2003) *Actinobacillus actinomycetemcomitans* endogenous endophthalmitis: report of two cases and review of the literature. *Scandinavian Journal of Infectious Diseases* **35**, 133-136.

Formatted: Font: Italic

- 3. Brewer, N.S., MacCarty, C.S., & Wellman, W.E. (1975) Brain abscess: a review of recent experience. *Annals of Internal Medicine* **82**, 571-576.
- 4. Burgher, L.W., Loomis, G.W., & Ware, F. (1973) Systemic infection due to <u>Actinobacillus</u> actinomycetemcomitans. American Journal of Clinical Pathology **60**, 412-415.

Formatted: Font: Italic

- Churton,M.C. & Greer,N.D. (1980) Intracranial abscess secondary to dental infection. *The New Zealand Dental Journal* 76, 58-60.
- 6. Garner, J.G. (1979) Isolation of *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* at Auckland Hospital. *New Zealand Dental Journal* **89**, 384-386.

Formatted: Font: Italic
Formatted: Font: Italic

- 7. Ingham, H.R., Kalbag, R.M., Tharagonnet, D., High, A.S., Sengupta, R.P., & Selkon, J.B. (1978) Abscesses of the frontal lobe of the brain secondary to covert dental sepsis. *The Lancet* **2**, 497-499.
- 8. Kaplan, A.H., Weber, D.J., Oddone, E.Z., & Perfect, J.R. (1989) Infection due to *Actinobacillus actinomycetemcomitans*: 15 cases and review. *Review of Infectious Diseases* 11, 46-63.

Formatted: Font: Italic

- 9. Klinger, R. Untersuchungen ubermenschliche Aktinomykose. (1912) *Zentralblad Bakteriologie* **62**, 191-200.
- 10. Kuijper, E.J., Wiggerts, H.O., Jonker, G.J., Schaal, K.P., & de Gans, J. (1992) Disseminated actinomycosis due to *Actinomyces meyeri* and *Actinobacillus actinomycetemcomitans*. *Scandinavian Journal of Infectious Diseases* 24, 667-672.

Formatted: Font: Italic

Formatted: Font: Italic

11. Martin,B.F., Derby,B.M., Budzilovich,G.N., & Ransohoff,J. (1967) Brain abscess due to *Actinobacillus actinomycetemcomitans*. *Neurology* **17**, 833-837.

Formatted: Font: Italic

- 12. Nesse, W., Abbas, F., van der Ploeg, I., Spijkervet, F.K., Dijkstra, P.U., & Vissink, A. (2008) Periodontal inflamed surface area: quantifying inflammatory burden. *Journal of Clinical Periodontology*. **35**, 668-673.
- 13. Page, M.I. & King, E.O. (1966) Infection due to Actinobacillus actinomycetemcomitans and Haemophilus aphrophilus. New England Journal of Medicine 275, 181-188.

Formatted: Font: Italic

Formatted: Font: Italic

- 14. Renton, T.F., Danks, J., & Rosenfeld, J.V. (1996) Cerebral abscess complicating dental treatment. Case report and review of the literature. *Australian Dental Journal* **41**, 12-15.
- 15. Steinberg, J. P. and Burd, E. M. (2009) Other Gram-Negative and Gram-Variable Bacilli. In Principles and Practice of Infectious Diseases, eds. Mandell, G. L, Bennet, J. E., and Dolin R. pp 3015-3017. Philadelphia, Churchill Livingstone Elsevier.
- 16. Stepanovic, S., Tosic, T., Savic, B., Jovanovic, M., K'ouas, G., & Carlier, J.P. (2005) Brain abscess due to *Actinobacillus actinomycetemcomitans*. *APMIS* 113, 225-228.

Formatted: Font: Italic

- 17. Tsai,J.C., Teng,L.J., & Hsueh,P.R. (2004) Direct detection of bacterial pathogens in brain abscesses by polymerase chain reaction amplification and sequencing of partial 16S ribosomal deoxyribonucleic acid fragments. *Neurosurgery* **55**, 1154-1162.
- 18. Tseng, J.H. & Tseng, M.Y. (2006) Brain abscess in 142 patients: factors influencing outcome and mortality. *Surgical Neurology* **65**, 557-562.
- 19. van Winkelhoff, A.J., Overbeek, B.P., Pavicic, M.J.A.M.P., van den Bergh, J.P., Ernst, J.P., & de Graaff, J. (1993) Long-standing bacteremia caused by oral *Actinobacillus actinomycetemcomitans* in a patient with a pacemaker. *Clinical Infectious Diseases* 16, 216-218.

Formatted: Font: Italic

20. van Winkelhoff, A.J. & Slots, J. (1999) *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* in nonoral infections. *Periodontology* **20**, 122-135.

Formatted: Font: Italic
Formatted: Font: Italic

 von Graevenitz, A. and Zbinden, R. (2007) Actinobacillus, Capnocytophaga, Eikenella, Kingella, Pasteurella, and other fastidious or rarely encountered gram-negative rods. In Manual of Clinical Microbiology, eds Murray, P. R. and Baron, E. J., pp 621-636. Washington, ASM Press. Formatted: Font: Italic

22. Yuan, A., Yang, P.C., Lee, L.N., Chang, D.B., Kuo, S.H., & Luh, K.T. (1992) *Actinobacillus actinomycetemcomitans* pneumonia with chest wall involvement and rib destruction. *Chest* 101, 1450-1452.

Formatted: Font: Italic

23. Zambon, J.J. (1985) *Actinobacillus actinomycetemcomitans* in human periodontal disease. *Journal of Clinical Periodontology* **12**, 1-20.

Formatted: Font: Italic

24. Zijlstra, E.E., Swart, G.R., Godfroy, F.J., & Degener, J.E. (1992) Pericarditis, pneumonia and brain abscess due to a combined *Actinomyces--Actinobacillus actinomycetemcomitans* infection. *Journal of Infection* **25**, 83-87.

Formatted: Font: Italic

Formatted: Font: Italic

Tables and figure legends

Figure 1

Orthopantogram revealing caries profunda in multiple teeth and severe periodontal breakdown

Figure 2

Computed Tomogram brain. Abscesses visible in right and left cerebrum.

es due to Table 1. Reported cases of cerebral abscesses due to infection with A. actinomycetemcomitans.

Pen = penicillin

Chlo = chloramphenicol

Co = colistin

Amp = ampicillin

Strep = streptomycin

Ge = gentamycin

Me = metronidazole

Amox = amoxicillin

Cefo = cefotaxime

Wk = week

Mo = month

M = male


F = female

sex/age (yr)	underlying illness	source	presenting symptoms	localisation	infection	mixed with	antibiotic treatment	duration	outcome	author/year
M/62	Carcinoma left ear	Otogenous	Drainage left ear, confusion	Posterior inferior temporal region	Single		Pen + Chlo +Co	3 wks (until death)	Died	Martin B.F., 1967
M/31	None	?	Right hemianopsia	Left parietal- occipital	Single		Amp + strep	2 wks	Recovered	Burgher L.W.,1973
?	?	?			Single		?	?	?	Brewer, 1975
?	?	?				?	?	?	?	Brewer, 1975
M/34	None	Dental	Headache, vomiting	Left frontal region		Anaerobic streptococci, Fusobacterium polymorphum, Bacteroides melaninogenicus, S. milleri	Amp + Ge + Me	?	Recovered	Ingham H.R., 1978
F/7	Cyanotic congenital heart disease	?	Headache, irritability	Left frontal region	Single		Clo	At least 4 wks	Recovered	Garner J.G., 1979
M/36	None	Dental	Headache, convulsion	Right frontal lobe		Fusobacterium nucleatum	Me + Chlo	?	Recovered	Churton M.C., 1980
F/15	Neurosurgery for meningioma	Nasopharynx	Headache, fever	Right frontal region		Eikenella corrodens, H. aphrophilus	Chlo + Me	1 mo	Recovered	Kaplan A.H., 1989
M/44	Alcohol abuse, emphysema	Dental	Right sided weakness, speeking disorder	Left fronto- parietal region, right occipital		Actinomyces meyeri	Amox	12 mo	Recovered	Kuijper E.J., 1992
M/41	None	Dental	Convulsion	Left frontoparietal region		Actinomyces	Chlo	6 wks	Recovered	Zijlstra E.E., 1992
M/19	None	Dental	Headache, vomiting, right sided weakness	Left parietal region	Single		Chlo + Cefo	10 wks	Recovered	Renton T.F., 1996
M/47	Alcohol abuse	Dental	Headache, vomiting, left sided weakness	Right occipital lobe	Single		Ceftriaxon	4 wks	Recovered	Stepanovic S., 2005



424x200mm (72 x 72 DPI)





262x262mm (72 x 72 DPI)



sex/age (yr)	underlying illness	source	presenting symptoms	localisation	infection	mixed with	antibiotic treatment	duration	outcome	author/year
M/62	Carcinoma left ear	Otogenous	Drainage left ear, confusion	Posterior inferior temporal region	Single		Pen + Chlo +Co	3 wks (until death)	Died	Martin B.F., 1967
M/31	None	?	Right hemianopsia	Left parietal- occipital	Single		Amp + strep	2 wks	Recovered	Burgher L.W.,1973
?	?	?			Single		?	?	?	Brewer, 1975
?	?	?				?	?	?	?	Brewer, 1975
M/34	None	Dental	Headache, vomiting	Left frontal region		Anaerobic streptococci, Fusobacterium polymorphum, Bacteroides melaninogenicus, S. milleri	Amp + Ge + Me	?	Recovered	Ingham H.R., 1978
F/7	Cyanotic congenital heart disease	?	Headache, irritability	Left frontal region	Single		Clo	At least 4 wks	Recovered	Garner J.G., 1979
M/36	None	Dental	Headache, convulsion	Right frontal lobe		Fusobacterium nucleatum	Me + Chlo	?	Recovered	Churton M.C., 1980
F/15	Neurosurgery for meningioma	Nasopharynx	Headache, fever	Right frontal region		Eikenella corrodens, H. aphrophilus	Chlo + Me	1 mo	Recovered	Kaplan A.H., 1989
M/44	Alcohol abuse, emphysema	Dental	Right sided weakness, speeking disorder	Left fronto- parietal region, right occipital		Actinomyces meyeri	Amox	12 mo	Recovered	Kuijper E.J., 1992
M/41	None	Dental	Convulsion	Left frontoparietal region		Actinomyces	Chlo	6 wks	Recovered	Zijlstra E.E., 1992
M/19	None	Dental	Headache, vomiting, right sided weakness	Left parietal region	Single		Chlo + Cefo	10 wks	Recovered	Renton T.F., 1996
M/47	Alcohol abuse	Dental	Headache, vomiting, left sided weakness	Right occipital lobe	Single		Ceftriaxon	4 wks	Recovered	Stepanovic S., 2005

Table I. Reported cases of cerebral abscesses due to infection with A. actinomycetemcomitans.

