



HAL
open science

Two-Scale Macro-Micro decomposition of the Vlasov equation with a strong magnetic field

Nicolas Crouseilles, Emmanuel Frenod, Sever Adrian Hirstoaga, Alexandre Mouton

► **To cite this version:**

Nicolas Crouseilles, Emmanuel Frenod, Sever Adrian Hirstoaga, Alexandre Mouton. Two-Scale Macro-Micro decomposition of the Vlasov equation with a strong magnetic field. 2011. hal-00638617v1

HAL Id: hal-00638617

<https://hal.science/hal-00638617v1>

Preprint submitted on 6 Nov 2011 (v1), last revised 5 Nov 2012 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Two-Scale Macro-Micro decomposition of the Vlasov equation with a strong magnetic field

Nicolas Crouseilles* Emmanuel FRÉNOD† Sever HIRSTOAGA‡
Alexandre MOUTON§

Abstract

In this paper, we build a Two-Scale Macro-Micro decomposition of the Vlasov equation with a strong magnetic field. This consists in writing the solution of this equation as a sum of two oscillating functions with circumscribed oscillations. The first of these functions has a shape which is close to the shape of the Two-Scale limit of the solution and the second one is a correction built to offset this imposed shape. The aim of such a decomposition is to be the starting point for the construction of Two-Scale Asymptotic-Preserving Schemes.

1 Introduction

The goal of this paper is to make the first step towards the setting out of a new class of numerical methods: the *Two-Scale Asymptotic-Preserving Schemes* or TSAPS. We intend to use these methods in order to treat problems involving the following two numerical difficulties: first, dealing efficiently on long time scales with solutions having high frequency oscillations (the *Two-Scale* approach) and second, an accurate and stable managing of the transition between different regimes, using a unified model (the *Asymptotic-Preserving* approach). These problems naturally occur in solving a Vlasov equation implying some small parameter, for the modelling of the dynamics of charged particles in the presence of a strong magnetic field. More precisely, we are interested to develop a model whose discretization will be able to simulate both, the regime when the parameter is not small (*i.e.* the magnetic field is not large), and the limit regime obtained when the parameter is small. The discrete scheme will automatically shift from one regime to the other. In addition, in the limit of the small parameter, the particles mean behaviour will be efficiently described by the two-scale limit while the micro behaviour will be expressed by some correctors.

To make the purpose more precise, we first refer to [4, 8, 13, 14, 15, 16] and the references therein as previous works about different regimes, by taking limits of Vlasov or Vlasov-Poisson equations corresponding to small parameters, having in common the framework of a large magnetic field. Then, we refer to [1, 2, 9, 11, 21, 22, 23] for a detailed description of Two-Scale Numerical Methods. As for the Asymptotic-Preserving schemes, although there is an abundant literature about, we first cite the classical paper [17] and then [3, 6, 7, 19, 20] which treat problems close to our, without considering the two-scale aspect.

Next we start to detail the method, by setting a generic and formal problem

$$E^\epsilon u^\epsilon = 0, \tag{1.1}$$

*INRIA-Rennes Bretagne Atlantique, IPSO Project & IRMAR (UMR CNRS 6625) Université de Rennes 1.

†Université Européenne de Bretagne, Lab-STICC (UMR CNRS 3192), Université de Bretagne-Sud & INRIA-Nancy Grand Est, CALVI Project & IRMA (UMR CNRS 7501), Université de Strasbourg.

‡INRIA-Nancy Grand Est, CALVI Project & IRMA (UMR CNRS 7501), Université de Strasbourg.

§CNRS & Laboratoire Painlevé (UMR CNRS 8524) Université de Lille 1.

in which E^ϵ is a partial differential operator depending on a parameter ϵ which induces oscillations of typical size ϵ in the solution $u^\epsilon = u^\epsilon(z)$. In the case where ϵ is uniformly small over the domain on which (1.1) is posed, a Two-Scale approach may give good results. To summarize, this approach consists in noticing that, in some sense,

$$u^\epsilon(z) \approx U(z, \frac{z}{\epsilon}), \quad (1.2)$$

where $U = U(z, \zeta)$ is the Two-Scale limit of u^ϵ (Two-Scale convergence goes back to [23]), and that U is the solution of a well-posed problem of the form

$$\mathcal{E}U = 0, \quad (1.3)$$

where \mathcal{E} does not induce high frequency oscillations. Building a Two-Scale Numerical Method consists then in solving, via an usual numerical method, model (1.3) and rebuilding u^ϵ using an algorithm based on (1.2).

In the case where ϵ is not a uniform parameter over the domain, in the sense that it may be small in some sub-region and of order 1 in other ones, Asymptotic-Preserving schemes are intended to be used, since they allow to simulate both sub-regions with the same method. The methodology for these numerical schemes is based on the fact that u^ϵ may converge, in a strong or a weak topology, towards a limit u which is solution of

$$Eu = 0. \quad (1.4)$$

Then, an Asymptotic-Preserving scheme consists in solving

$$E_{\Delta z}^\epsilon u_{\Delta z}^\epsilon = 0, \quad (1.5)$$

such that

$$u_{\Delta z}^\epsilon \longrightarrow u^\epsilon, \quad \text{as } \Delta z \rightarrow 0, \quad (1.6)$$

and

$$u_{\Delta z}^\epsilon \longrightarrow u_{\Delta z}, \quad \text{as } \epsilon \rightarrow 0, \quad (1.7)$$

where $u_{\Delta z}$ is a numerical approximation of u , meaning that

$$u_{\Delta z} \longrightarrow u, \quad \text{as } \Delta z \rightarrow 0, \quad (1.8)$$

and is the solution of

$$E_{\Delta z} u_{\Delta z} = 0, \quad (1.9)$$

where $E_{\Delta z}$ is a numerical operator. Notice that the property (1.6)-(1.7)-(1.8) is called the *Asymptotic-Preserving Property*.

All what was just said may be summarized in the following diagram:

$$\begin{array}{ccc}
 \begin{array}{c} u^\epsilon \text{ solution of} \\ E^\epsilon u^\epsilon = 0 \end{array} & \xrightarrow{\epsilon \rightarrow 0} & \begin{array}{c} u \text{ solution of} \\ Eu = 0 \end{array} \\
 \uparrow \Delta z \rightarrow 0 & & \uparrow \Delta z \rightarrow 0 \\
 \begin{array}{c} u_{\Delta z}^\epsilon \text{ solution of} \\ E_{\Delta z}^\epsilon u_{\Delta z}^\epsilon = 0 \end{array} & \xrightarrow{\epsilon \rightarrow 0} & \begin{array}{c} u_{\Delta z} \text{ solution of} \\ E_{\Delta z} u_{\Delta z} = 0 \end{array}
 \end{array} \quad (1.10)$$

In the case when u^ϵ contains high frequency oscillations with high amplitude when ϵ is small, solving (1.5) can be time consuming to capture the oscillations. In this case, the convergence

$$u^\epsilon \rightarrow u, \quad (1.11)$$

occurs in a weak sense only. Besides this, the Two-Scale limit $U = U(z, \zeta)$ may describe very well the oscillations of u^ϵ in the sense that the convergence

$$U(z, \frac{z}{\epsilon}) - u^\epsilon(z) \rightarrow 0, \quad (1.12)$$

occurs in a stronger sense than (1.11). This is particularly the case when E^ϵ generates pseudo-periodic oscillations, essentially at a sole period, of size ϵ in the solution u^ϵ when ϵ becomes small. Moreover, u and U are linked by a relation of the kind

$$u(z) = \frac{1}{|\mathcal{Z}|} \int_{\mathcal{Z}} U(z, \zeta) d\zeta, \quad (1.13)$$

where \mathcal{Z} is the cell such that $\zeta \mapsto U(z, \zeta)$ is \mathcal{Z} -periodic for any z , and $|\mathcal{Z}|$ is its measure.

In this case, we propose to add a layer in the diagram (1.10) which summarizes the ideas on which Two-Scale Asymptotic-Preserving Scheme is based:




Diagram (1.14) may be looked at as a prism. At the bottom of the front edge stands the TSAPS and the back rectangle is made of the Asymptotic-Preserving scheme diagram of (1.10). The intermediate layer gathers the Two-Scale limit problem, at the top, and the Two-Scale Numerical Method, at the bottom.

The solution $U_{\Delta z}^\epsilon$ computed with the TSAPS must be close to u^ϵ when it is computed in $\zeta = \frac{z}{\epsilon}$ and when Δz converges to 0. This is the first part of the Two-Scale Asymptotic-Preserving Property and it is symbolized by the arrow going from $U_{\Delta z}^\epsilon$ to $u_{\Delta z}^\epsilon$ and by the one linking $u_{\Delta z}^\epsilon$ and u^ϵ .

The second part of the Two-Scale Asymptotic-Preserving Property is symbolized by the right part of the diagram (1.14) : $U_{\Delta z}^\epsilon$ must be close to the numerical approximation $U_{\Delta z}$ of the Two-Scale limit of u_ϵ when $\epsilon \rightarrow 0$ and a numerical integration (symbolized by $\frac{1}{|\mathcal{Z}|} \int_{\mathcal{Z}}^{\text{NUM}} d\zeta$) of $U_{\Delta z}$ needs to recover an

approximation $u_{\Delta z}$ of the weak-* limit u of u^ϵ .

We will deduce the Two-Scale Asymptotic-Preserving model on the base of a Two-Scale Macro-Micro decomposition that yields an equation which stands at the top of the front edge. Arrows linking U^ϵ and u^ϵ , U^ϵ and U and u may be seen as the continuous Two-Scale Asymptotic-Preserving Property. They are constraints for the Macro-Micro decomposition. We refer to [3, 6, 18, 20] for the use of micro-macro decomposition in order to design an asymptotic-preserving scheme.

The present paper implements the Two-Scale Macro-Micro decomposition in the case of the linear Vlasov equation with a strong magnetic field. This equation is the following:

$$\begin{cases} \frac{\partial f^\epsilon}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}} f^\epsilon + \left(\mathbf{E}^\epsilon + \mathbf{v} \times \left(\mathbf{B}^\epsilon + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot \nabla_{\mathbf{v}} f^\epsilon = 0, \\ f^\epsilon(t=0, \mathbf{x}, \mathbf{v}) = f_0(\mathbf{x}, \mathbf{v}), \end{cases} \quad (1.15)$$

where $f^\epsilon = f^\epsilon(t, \mathbf{x}, \mathbf{v})$ is the distribution function of an ion gaz submitted to external electric and magnetic field $\mathbf{E}^\epsilon = \mathbf{E}^\epsilon(t, \mathbf{x})$ and $\mathbf{B}^\epsilon = \mathbf{B}^\epsilon(t, \mathbf{x})$, where $t \in \mathbb{R}^+$, $\mathbf{x} \in \mathbb{R}^3$, $\mathbf{v} \in \mathbb{R}^3$, and where $\mathcal{M} = 2\pi \mathbf{e}_1$, *i.e.* the first vector of the canonical base $(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ of \mathbb{R}^3 .

The paper is organized as follows: in section 2, we recall the results of Frénod & Sonnendrücker [12] giving the Two-Scale convergence of f^ϵ . Roughly speaking they claim that, when ϵ is small,

$$f^\epsilon(t, \mathbf{x}, \mathbf{v}) \text{ is close to } G\left(t, \mathbf{x}, \mathbf{u}\left(\frac{t}{\epsilon}, \mathbf{v}\right)\right) \text{ in a strong sense,} \quad (1.16)$$

$$f^\epsilon(t, \mathbf{x}, \mathbf{v}) \text{ is close to } f(t, \mathbf{x}, \mathbf{v}) \text{ in a weak sense,} \quad (1.17)$$

and the equations for G and f are known ($\mathbf{u}(\tau, \mathbf{v})$ is defined by (2.9)). In particular the \mathbf{v} -dependence of f is only through $v_{\parallel} = \mathbf{v} \cdot \mathbf{e}_1$, $v_{\perp} = \sqrt{(\mathbf{v} \cdot \mathbf{e}_2)^2 + (\mathbf{v} \cdot \mathbf{e}_3)^2}$ and not through the angle α , which is such that $\mathbf{v} \cdot \mathbf{e}_2 = v_{\perp} \cos \alpha$. Based on this fact, in section 3, we build a classical Macro-Micro decomposition of (1.15) by following the same ideas of Lemou & Mieussens [20]. It consists in decomposing f^ϵ in the following way

$$f^\epsilon(t, \mathbf{x}, \mathbf{v}) = m^\epsilon(t, \mathbf{x}, v_{\parallel}, v_{\perp}) + n^\epsilon(t, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) \quad (1.18)$$

with $m^\epsilon \in \text{Ker}((\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}})$ and $n^\epsilon \in \text{Im}((\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}})$. Indeed, we will see in details that this decomposition is unique so that it is valid for every ϵ , small or not. Note that m^ϵ does not depend on α , as f , while n^ϵ does. Equations for m^ϵ and n^ϵ are set out, making up the Macro-Micro model. The drawback of this model is that, because of (1.17), the exact equality (1.18) implies that function n^ϵ is still highly oscillating, as f^ϵ . To avoid this drawback, we develop the Two-Scale Macro-Micro framework. This is done in sections 4, 5, and 6, where we build the Two-Scale Macro-Micro approximation of f^ϵ . First, based on results of Frénod, Raviart & Sonnendrücker [10] we improve (1.16) by using a first order approximation, claiming that when ϵ is small

$$f^\epsilon(t, \mathbf{x}, \mathbf{v}) \text{ is close to } G\left(t, \mathbf{x}, \mathbf{u}\left(\frac{t}{\epsilon}, \mathbf{v}\right)\right) + \epsilon G_1\left(t, \mathbf{x}, \mathbf{u}\left(\frac{t}{\epsilon}, \mathbf{v}\right)\right) + \epsilon l\left(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}\right) \text{ in a strong sense,} \quad (1.19)$$

where G_1 is the solution of a partial differential equation and l is given by a formula (see [10]). Inspired by (1.19), we construct our Two-Scale Macro-Micro decomposition in the following way

$$f^\epsilon(t, \mathbf{x}, \mathbf{v}) = G\left(t, \mathbf{x}, \mathbf{u}\left(\frac{t}{\epsilon}, \mathbf{v}\right)\right) + \epsilon G_1^\epsilon\left(t, \mathbf{x}, \mathbf{u}\left(\frac{t}{\epsilon}, \mathbf{v}\right)\right) + \epsilon l\left(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}\right) + \epsilon h^\epsilon\left(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}\right), \quad (1.20)$$

where G_1^ϵ is intended to be close to G_1 when ϵ is small and h^ϵ is the corrector to be taken into account when the order of magnitude of ϵ is 1. In other words, we use the Macro-Micro decomposition in order to make precise for every ϵ the approximation in (1.19) which is valid only for small ϵ . Then, we need to

show that the decomposition in (1.20) is unique. This is done by paying attention to the properties of the operator

$$\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}, \quad (1.21)$$

since the expression (1.20) may also be set in the following form:

$$f^\epsilon(t, \mathbf{x}, \mathbf{v}) = \mathcal{A}^\epsilon(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}) + \mathcal{B}^\epsilon(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}), \quad (1.22)$$

with

$$\mathcal{A}^\epsilon = \mathcal{A}^\epsilon(t, \tau, \mathbf{x}, \mathbf{v}) \in \text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right) \quad \text{and} \quad \mathcal{B}^\epsilon = \mathcal{B}^\epsilon(t, \tau, \mathbf{x}, \mathbf{v}) \in \left(\text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)\right)^\perp. \quad (1.23)$$

Then, we obtain the equations for the Macro part G_1^ϵ and the Micro one h^ϵ . In section 7, we underline the continuous asymptotic-preserving property of the Two-Scale Macro-Micro decomposition. We close the paper with two appendices about some properties of the operator in (1.21) leading mainly to the uniqueness of the decomposition in (1.22).

2 Two-Scale convergence of f^ϵ

In this section, we briefly recall the results of Frénod & Sonnendrücker [12] where the asymptotic behaviour of (1.15) is studied. What is known and taken for granted concerning (1.15) is that, considering the following hypothesis

$$f_0 \geq 0, \quad f_0 \in L^2(\mathbb{R}^6), \quad (2.1)$$

and

$$\begin{aligned} \mathbf{E}^\epsilon &\rightarrow \mathbf{E} \quad \text{strong in } L^\infty(0, T; L^2_{\text{loc}}(\mathbb{R}^3)), \\ \mathbf{B}^\epsilon &\rightarrow \mathbf{B} \quad \text{strong in } L^\infty(0, T; L^2_{\text{loc}}(\mathbb{R}^3)), \end{aligned} \quad (2.2)$$

we claim that, for any $T > 0$, $(f^\epsilon)_{\epsilon > 0}$ is bounded in $L^\infty(0, T; L^2(\mathbb{R}^6))$. Moreover, as $\epsilon \rightarrow 0$, up to a subsequence, $(f^\epsilon)_{\epsilon > 0}$ Two-Scale converges towards the Two-Scale limit (see [2, 23])

$$F = F(t, \tau, \mathbf{x}, \mathbf{v}) \in L^\infty(0, T; L^2_{\#_1}(\mathbb{R}^+; L^2(\mathbb{R}^6))), \quad (2.3)$$

where the subscript $\#_1$ indicates that $\tau \mapsto F(t, \tau, \mathbf{x}, \mathbf{v})$ is 1-periodic for any $(t, \mathbf{x}, \mathbf{v})$. Introduced in [23] and further analyzed and used in several homogenization problems in [2], the Two-Scale convergence means that for any regular function $\psi = \psi(t, \tau, \mathbf{x}, \mathbf{v})$ such that $(t, \mathbf{x}, \mathbf{v}) \mapsto \psi(t, \tau, \mathbf{x}, \mathbf{v})$ is compactly supported in $\Omega = [0, T] \times \mathbb{R}^6$ for any τ , and $\tau \mapsto \psi(t, \tau, \mathbf{x}, \mathbf{v})$ is 1-periodic for any $(t, \mathbf{x}, \mathbf{v})$, we have

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega} f^\epsilon (\psi)^\epsilon dt d\mathbf{x} d\mathbf{v} = \int_{\Omega} \int_0^1 F \psi d\tau dt d\mathbf{x} d\mathbf{v}, \quad (2.4)$$

where

$$(\psi)^\epsilon = (\psi)^\epsilon(t, \mathbf{x}, \mathbf{v}) = \psi(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}). \quad (2.5)$$

It was shown in [12] that the Two-Scale limit F has the following property:

$$F \in \text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right), \quad (2.6)$$

where $\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}$ is seen as anti-symmetric, non-bounded, with open range from $L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6)))$ on itself. The direct consequence of (2.6) is that, introducing the rotation matrix r as

$$r(\tau) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(2\pi\tau) & -\sin(2\pi\tau) \\ 0 & \sin(2\pi\tau) & \cos(2\pi\tau) \end{pmatrix}, \quad (2.7)$$

the Two-Scale limit F expresses itself in terms of a function $G = G(t, \mathbf{x}, \mathbf{u}) \in L^\infty(0, T; L^2(\mathbb{R}^3))$ by

$$F(t, \tau, \mathbf{x}, \mathbf{v}) = G(t, \mathbf{x}, \mathbf{u}(\tau, \mathbf{v})), \quad (2.8)$$

where

$$\mathbf{u}(\tau, \mathbf{v}) = r(\tau) \mathbf{v}. \quad (2.9)$$

Using oscillating test functions defined by (2.5) for regular functions ψ which are moreover in $\text{Ker}(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}})$, it may be brought out that G is the solution of

$$\begin{cases} \frac{\partial G}{\partial t} + \mathbf{u}_{\parallel} \cdot \nabla_{\mathbf{x}} G + (\mathbf{E}_{\parallel} + \mathbf{u} \times \mathbf{B}_{\parallel}) \cdot \nabla_{\mathbf{u}} G = 0, \\ G(t = 0, \mathbf{x}, \mathbf{u}) = f_0(\mathbf{x}, \mathbf{u}), \end{cases} \quad (2.10)$$

where the notation $\mathbf{u}_{\parallel} = (\mathbf{u} \cdot \mathbf{e}_1) \mathbf{e}_1$ is considered.

Having in mind the diagram (1.14), we can say at this point that equation (1.15) plays the role of $E^\epsilon u^\epsilon = 0$, and (2.8)-(2.10) plays the role of $\mathcal{E}U = 0$. So, if we build a numerical method to approximate (2.10), this numerical method, coupled with (2.8) would play the role of $\mathcal{E}_{\Delta z} U_{\Delta z} = 0$.

In [12], it was also proved that the weak-* limit f of f^ϵ is the solution to

$$\begin{cases} \frac{\partial f}{\partial t} + \mathbf{v}_{\parallel} \cdot \nabla_{\mathbf{x}} f + (\mathbf{E}_{\parallel} + \mathbf{v} \times \mathbf{B}_{\parallel}) \cdot \nabla_{\mathbf{v}} f = 0, \\ f(t = 0, \mathbf{x}, \mathbf{v}) = \int_0^1 f_0(\mathbf{x}, \mathbf{u}(\tau, \mathbf{v})) d\tau. \end{cases} \quad (2.11)$$

This equation is gotten from (2.8)-(2.10) by performing an integration in τ and by using the link between the weak-* limit f and the Two-Scale limit F given by

$$f(t, \mathbf{x}, \mathbf{v}) = \int_0^1 F(t, \tau, \mathbf{x}, \mathbf{v}) d\tau. \quad (2.12)$$

Then, equation (2.11) plays the role of $E u = 0$ in the diagram (1.14) and a numerical approximation of it would play the role of $E_{\Delta z} u_{\Delta z} = 0$.

3 Classical Macro-Micro decomposition

3.1 Rewriting the weak-* limit model

Before going further, we will rewrite model (2.11) in a form involving cylindrical variables in velocity. In this coordinate system, the weak-* limit is independent of the angle variable, and consequently it is easily gotten that the $\mathbf{v} \times \mathbf{B}_{\parallel}$ piece in the force term in (2.11) is removed.

Because of the average over τ and of the definition of $\mathbf{u}(\tau, \mathbf{v})$, it is obvious to see that the initial data in (2.11) only depends on \mathbf{x} , \mathbf{v}_{\parallel} and $v_{\perp} = \sqrt{v_2^2 + v_3^2}$. In other words, writing the velocity in the following

cylindrical variables

$$\begin{cases} v_{\parallel} \in \mathbb{R} \text{ such that } v_{\parallel} = \mathbf{v} \cdot \mathbf{e}_1, \\ v_{\perp} \in \mathbb{R}^+ \text{ such that } v_{\perp} = \sqrt{v_2^2 + v_3^2}, \\ \alpha \in [0, 2\pi] \text{ such that } v_2 = v_{\perp} \cos \alpha \text{ and } v_3 = v_{\perp} \sin \alpha, \end{cases} \quad (3.1)$$

there exists a function $m_0 = m_0(\mathbf{x}, v_{\parallel}, v_{\perp})$ such that

$$m_0(\mathbf{x}, v_{\parallel}, v_{\perp}) = \frac{1}{2\pi} \int_0^{2\pi} f_0(\mathbf{x}, \mathbf{u}(\tau, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3)) d\alpha. \quad (3.2)$$

The quickest way to get this is to notice that

$$\mathbf{u}(\tau, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) = \begin{pmatrix} v_{\parallel} \\ v_{\perp} \cos(\alpha + 2\pi\tau) \\ v_{\perp} \sin(\alpha + 2\pi\tau) \end{pmatrix}, \quad (3.3)$$

and that, consequently, integrating in τ suppresses the α -dependency. Hence, (3.2) writes also

$$m_0(\mathbf{x}, v_{\parallel}, v_{\perp}) = \frac{1}{2\pi} \int_0^{2\pi} f_0(\mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) d\alpha. \quad (3.4)$$

Concerning the function f itself, we introduce the function $m = m(t, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha)$ linked with f by

$$m(t, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) = f(t, \mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3). \quad (3.5)$$

Because of (2.12) and (2.8), and using the same argument as just above, it is easily seen that m does not depend on α and that

$$\begin{aligned} m(t, \mathbf{x}, v_{\parallel}, v_{\perp}) &= \int_0^1 G(t, \mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos(\alpha + 2\pi\tau) \mathbf{e}_2 + v_{\perp} \sin(\alpha + 2\pi\tau) \mathbf{e}_3) d\tau \\ &= \frac{1}{2\pi} \int_0^{2\pi} G(t, \mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) d\alpha. \end{aligned} \quad (3.6)$$

Noticing that

$$\begin{aligned} \frac{\partial m}{\partial v_{\perp}}(t, \mathbf{x}, v_{\parallel}, v_{\perp}) &= \cos \alpha \frac{\partial f}{\partial v_2}(t, \mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) \\ &\quad + \sin \alpha \frac{\partial f}{\partial v_3}(t, \mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3), \\ \frac{\partial m}{\partial \alpha}(t, \mathbf{x}, v_{\parallel}, v_{\perp}) &= -v_{\perp} \sin \alpha \frac{\partial f}{\partial v_2}(t, \mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) \\ &\quad + v_{\perp} \cos \alpha \frac{\partial f}{\partial v_3}(t, \mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3), \end{aligned} \quad (3.7)$$

the equation for m can be deduced from (2.11). For this purpose, we develop $\mathbf{v} \times \mathbf{B}_{\parallel}$ in the new variables:

$$v_{\perp} (\cos \alpha \mathbf{e}_2 + \sin \alpha \mathbf{e}_3) \times \mathbf{B}_{\parallel} = |B_{\parallel}| v_{\perp} (\sin \alpha \mathbf{e}_2 - \cos \alpha \mathbf{e}_3). \quad (3.8)$$

As a consequence, we have

$$\begin{aligned} (v_{\perp} (\cos \alpha \mathbf{e}_2 + \sin \alpha \mathbf{e}_3) \times \mathbf{B}_{\parallel}) \cdot \nabla_{\mathbf{v}} f(t, \mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) \\ = -|B_{\parallel}| \frac{\partial m}{\partial \alpha}(t, \mathbf{x}, v_{\parallel}, v_{\perp}) = 0, \end{aligned} \quad (3.9)$$

meaning that the force term in (2.11) may be reduced to \mathbf{E}_{\parallel} . Then, m is the solution of

$$\begin{cases} \frac{\partial m}{\partial t} + v_{\parallel} \frac{\partial m}{\partial x_{\parallel}} + E_{\parallel} \frac{\partial m}{\partial v_{\parallel}} = 0, \\ m(t = 0, \mathbf{x}, v_{\parallel}, v_{\perp}) = m_0(\mathbf{x}, v_{\parallel}, v_{\perp}). \end{cases} \quad (3.10)$$

3.2 Macro-Micro decomposition

Inspired by (2.11) or more precisely by (3.10), we now write a classical Macro-Micro decomposition of the solution f^ϵ of (1.15). The aim of this decomposition is (see [3, 20]) to lead to a new model which will be equivalent to (1.15) when $\epsilon \sim 1$ and will formally give the weak-* limit model (3.10) when $\epsilon \rightarrow 0$. It is evident that this last property is not satisfied by the model (1.15). The new model will be the base for the build of an Asymptotic-Preserving scheme which will have to satisfy diagram (1.10).

The main idea of the Macro-Micro decomposition is to write f^ϵ as a sum of a function being in $\text{Ker}((\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}})$ and a function being in $(\text{Ker}((\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}))^\perp$. Adapting this idea to our context and having in mind (3.10), we seek $m_1^\epsilon = m_1^\epsilon(t, \mathbf{x}, v_{||}, v_\perp) \in L^\infty(0, T; L^2(\mathbb{R}^3 \times \mathbb{R} \times \mathbb{R}^+; v_\perp d\mathbf{x} dv_{||} dv_\perp))$ and $p^\epsilon = p^\epsilon(t, \mathbf{x}, v_{||}, v_\perp, \alpha) \in L^\infty(0, T; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^3 \times \mathbb{R} \times \mathbb{R}^+; v_\perp d\mathbf{x} dv_{||} dv_\perp)))$ such that

$$n^\epsilon(t, \mathbf{x}, v_{||}, v_\perp, \alpha) = \frac{\partial p^\epsilon}{\partial \alpha}(t, \mathbf{x}, v_{||}, v_\perp, \alpha), \quad (3.11)$$

and

$$f^\epsilon(t, \mathbf{x}, v_{||} \mathbf{e}_1 + v_\perp \cos \alpha \mathbf{e}_2 + v_\perp \sin \alpha \mathbf{e}_3) = m(t, \mathbf{x}, v_{||}, v_\perp) + m_1^\epsilon(t, \mathbf{x}, v_{||}, v_\perp) + n^\epsilon(t, \mathbf{x}, v_{||}, v_\perp, \alpha). \quad (3.12)$$

The subscript $\#2\pi$ indicates that $\alpha \mapsto p^\epsilon(t, \mathbf{x}, v_{||}, v_\perp, \alpha)$ is 2π -periodic for any $(t, \mathbf{x}, v_{||}, v_\perp)$.

Considering the anti-symmetric, non-bounded operator with closed range¹

$$\begin{aligned} \frac{\partial}{\partial \alpha} : L^\infty(0, T; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^3 \times \mathbb{R} \times \mathbb{R}^+; v_\perp d\mathbf{x} dv_{||} dv_\perp))) \\ \longrightarrow L^\infty(0, T; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^3 \times \mathbb{R} \times \mathbb{R}^+; v_\perp d\mathbf{x} dv_{||} dv_\perp))), \end{aligned} \quad (3.13)$$

we clearly have that

$$\begin{aligned} m &\in \text{Ker}\left(\frac{\partial}{\partial \alpha}\right), \\ m_1^\epsilon &\in \text{Ker}\left(\frac{\partial}{\partial \alpha}\right), \\ n^\epsilon &\in \left(\text{Ker}\left(\frac{\partial}{\partial \alpha}\right)\right)^\perp = \text{Im}\left(\frac{\partial}{\partial \alpha}\right). \end{aligned} \quad (3.14)$$

Then, if we inject the decomposition (3.12) in Vlasov equation (1.15), we obtain

$$\begin{aligned} \frac{\partial m}{\partial t} + \frac{\partial m_1^\epsilon}{\partial t} + \frac{\partial n^\epsilon}{\partial t} + \begin{pmatrix} v_{||} \\ v_\perp \cos \alpha \\ v_\perp \sin \alpha \end{pmatrix} \cdot \nabla_{\mathbf{x}}(m + m_1^\epsilon + n^\epsilon) \\ + \begin{pmatrix} E_{||}^\epsilon + v_\perp (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \\ \cos \alpha E_2^\epsilon + \sin \alpha E_3^\epsilon - v_{||} (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \\ \frac{1}{v_\perp} (-\sin \alpha E_2^\epsilon + \cos \alpha E_3^\epsilon + v_{||} (\sin \alpha B_3^\epsilon + \cos \alpha B_2^\epsilon)) - B_{||}^\epsilon - \frac{1}{\epsilon} \end{pmatrix} \\ \cdot \begin{pmatrix} \frac{\partial m}{\partial v_{||}} + \frac{\partial m_1^\epsilon}{\partial v_{||}} + \frac{\partial n^\epsilon}{\partial v_{||}} \\ \frac{\partial m}{\partial v_\perp} + \frac{\partial m_1^\epsilon}{\partial v_\perp} + \frac{\partial n^\epsilon}{\partial v_\perp} \\ \frac{\partial n^\epsilon}{\partial \alpha} \end{pmatrix} = 0. \end{aligned} \quad (3.15)$$

¹See Appendix A

Combining this equation with (3.10), we finally obtain

$$\begin{aligned}
& \frac{\partial m_1^\epsilon}{\partial t} + \frac{\partial n^\epsilon}{\partial t} + \begin{pmatrix} 0 \\ v_\perp \cos \alpha \\ v_\perp \sin \alpha \end{pmatrix} \cdot \nabla_{\mathbf{x}} m + \begin{pmatrix} v_\parallel \\ v_\perp \cos \alpha \\ v_\perp \sin \alpha \end{pmatrix} \cdot \nabla_{\mathbf{x}} (m_1^\epsilon + n^\epsilon) \\
& + \begin{pmatrix} E_\parallel^\epsilon - E_\parallel + v_\perp (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \\ \cos \alpha E_2^\epsilon + \sin \alpha E_3^\epsilon - v_\parallel (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \\ \frac{1}{v_\perp} (-\sin \alpha E_2^\epsilon + \cos \alpha E_3^\epsilon + v_\parallel (\sin \alpha B_3^\epsilon + \cos \alpha B_2^\epsilon)) - B_\parallel^\epsilon - \frac{1}{\epsilon} \end{pmatrix} \cdot \begin{pmatrix} \frac{\partial m}{\partial v_\parallel} \\ \frac{\partial m}{\partial v_\perp} \\ 0 \end{pmatrix} \\
& + \begin{pmatrix} E_\parallel^\epsilon + v_\perp (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \\ \cos \alpha E_2^\epsilon + \sin \alpha E_3^\epsilon - v_\parallel (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \\ \frac{1}{v_\perp} (-\sin \alpha E_2^\epsilon + \cos \alpha E_3^\epsilon + v_\parallel (\sin \alpha B_3^\epsilon + \cos \alpha B_2^\epsilon)) - B_\parallel^\epsilon - \frac{1}{\epsilon} \end{pmatrix} \cdot \begin{pmatrix} \frac{\partial m_1^\epsilon}{\partial v_\parallel} + \frac{\partial n^\epsilon}{\partial v_\parallel} \\ \frac{\partial m_1^\epsilon}{\partial v_\perp} + \frac{\partial n^\epsilon}{\partial v_\perp} \\ \frac{\partial n^\epsilon}{\partial \alpha} \end{pmatrix} = 0.
\end{aligned} \tag{3.16}$$

By projecting (3.15) onto $\text{Ker}(\frac{\partial}{\partial \alpha})$ and $(\text{Ker}(\frac{\partial}{\partial \alpha}))^\perp = \text{Im}(\frac{\partial}{\partial \alpha})$, we will obtain the Macro-Micro decomposition of f^ϵ . We refer to Appendix A and [5] to get that projecting a function onto $\text{Ker}(\frac{\partial}{\partial \alpha})$ consists in computing its average in α and projecting it onto $\text{Im}(\frac{\partial}{\partial \alpha})$ consists in subtracting from it its average value. This is what we do in the next lines.

Since m , m_1^ϵ , \mathbf{E} , \mathbf{E}^ϵ , \mathbf{B} and \mathbf{B}^ϵ do not depend on α , and recalling the definition of n^ϵ given in (3.11), the projection of (3.15) onto $\text{Ker}(\frac{\partial}{\partial \alpha})$ gives

$$\begin{aligned}
& \frac{\partial m_1^\epsilon}{\partial t} + v_\parallel \frac{\partial m_1^\epsilon}{\partial x_\parallel} + E_\parallel^\epsilon \frac{\partial m_1^\epsilon}{\partial v_\parallel} \\
& = -(E_\parallel^\epsilon - E_\parallel) \frac{\partial m}{\partial v_\parallel} - \frac{v_\perp}{2\pi} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 0 \\ \int_0^{2\pi} \sin \alpha \frac{\partial p^\epsilon}{\partial x_2} d\alpha \\ -\int_0^{2\pi} \cos \alpha \frac{\partial p^\epsilon}{\partial x_3} d\alpha \end{pmatrix} \\
& - \frac{1}{2\pi} \mathbf{E}^\epsilon \cdot \begin{pmatrix} 0 \\ \int_0^{2\pi} \sin \alpha \left(\frac{\partial p^\epsilon}{\partial v_\perp} + \frac{p^\epsilon}{v_\perp} \right) d\alpha \\ -\int_0^{2\pi} \cos \alpha \left(\frac{\partial p^\epsilon}{\partial v_\perp} + \frac{p^\epsilon}{v_\perp} \right) d\alpha \end{pmatrix} \\
& + \frac{1}{2\pi} \begin{pmatrix} -v_\perp \\ v_1 B_3^\epsilon \\ v_1 B_2^\epsilon \end{pmatrix} \cdot \begin{pmatrix} \int_0^{2\pi} (\cos \alpha B_2^\epsilon + \sin \alpha B_3^\epsilon) \frac{\partial p^\epsilon}{\partial v_\parallel} d\alpha \\ \int_0^{2\pi} \sin \alpha \left(\frac{\partial p^\epsilon}{\partial v_\perp} + \frac{p^\epsilon}{v_\perp} \right) d\alpha \\ \int_0^{2\pi} \cos \alpha \left(\frac{\partial p^\epsilon}{\partial v_\perp} + \frac{p^\epsilon}{v_\perp} \right) d\alpha \end{pmatrix}.
\end{aligned} \tag{3.17}$$

In the second, third and last terms of the right hand side of (3.17), we use integrations by parts to tackle α -integrations. Finally, we perform the projection onto $\text{Im}(\frac{\partial}{\partial \alpha})$ by subtracting (3.17) from (3.15), we

obtain a second order equation for p^ϵ in which m and m_1^ϵ appear as right hand side members:

$$\begin{aligned}
& \frac{\partial^2 p^\epsilon}{\partial t \partial \alpha} + \begin{pmatrix} v_{\parallel} \\ v_{\perp} \\ v_{\perp} \end{pmatrix} \cdot \begin{pmatrix} \frac{\partial^2 p^\epsilon}{\partial x_{\parallel} \partial \alpha} \\ \cos \alpha \frac{\partial^2 p^\epsilon}{\partial x_2 \partial \alpha} - \frac{1}{2\pi} \int_0^{2\pi} \cos \alpha \frac{\partial^2 p^\epsilon}{\partial x_2 \partial \alpha} d\alpha \\ \sin \alpha \frac{\partial^2 p^\epsilon}{\partial x_3 \partial \alpha} - \frac{1}{2\pi} \int_0^{2\pi} \sin \alpha \frac{\partial^2 p^\epsilon}{\partial x_3 \partial \alpha} d\alpha \end{pmatrix} \\
& + \mathbf{E}^\epsilon \cdot \begin{pmatrix} \frac{\partial^2 p^\epsilon}{\partial v_{\parallel} \partial \alpha} \\ \cos \alpha \frac{\partial^2 p^\epsilon}{\partial v_{\perp} \partial \alpha} - \frac{1}{2\pi} \int_0^{2\pi} \cos \alpha \frac{\partial^2 p^\epsilon}{\partial v_{\perp} \partial \alpha} d\alpha - \frac{1}{v_{\perp}} \sin \alpha \frac{\partial^2 p^\epsilon}{\partial \alpha^2} + \frac{1}{2\pi v_{\perp}} \int_0^{2\pi} \sin \alpha \frac{\partial^2 p^\epsilon}{\partial \alpha^2} d\alpha \\ \sin \alpha \frac{\partial^2 p^\epsilon}{\partial v_{\perp} \partial \alpha} - \frac{1}{2\pi} \int_0^{2\pi} \sin \alpha \frac{\partial^2 p^\epsilon}{\partial v_{\perp} \partial \alpha} d\alpha + \frac{1}{v_{\perp}} \cos \alpha \frac{\partial^2 p^\epsilon}{\partial \alpha^2} - \frac{1}{2\pi v_{\perp}} \int_0^{2\pi} \cos \alpha \frac{\partial^2 p^\epsilon}{\partial \alpha^2} d\alpha \end{pmatrix} \\
& + \begin{pmatrix} v_{\perp} \\ -v_{\parallel} B_3^\epsilon \\ v_{\parallel} B_2^\epsilon \end{pmatrix} \cdot \begin{pmatrix} (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \frac{\partial^2 p^\epsilon}{\partial v_{\parallel} \partial \alpha} - \frac{1}{2\pi} \int_0^{2\pi} (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \frac{\partial^2 p^\epsilon}{\partial v_{\parallel} \partial \alpha} d\alpha \\ \cos \alpha \frac{\partial^2 p^\epsilon}{\partial v_{\perp} \partial \alpha} - \frac{1}{v_{\perp}} \sin \alpha \frac{\partial^2 p^\epsilon}{\partial \alpha^2} - \frac{1}{2\pi} \int_0^{2\pi} \left(\cos \alpha \frac{\partial^2 p^\epsilon}{\partial v_{\perp} \partial \alpha} - \frac{1}{v_{\perp}} \sin \alpha \frac{\partial^2 p^\epsilon}{\partial \alpha^2} \right) d\alpha \\ \sin \alpha \frac{\partial^2 p^\epsilon}{\partial v_{\perp} \partial \alpha} + \frac{1}{v_{\perp}} \cos \alpha \frac{\partial^2 p^\epsilon}{\partial \alpha^2} - \frac{1}{2\pi} \int_0^{2\pi} \left(\sin \alpha \frac{\partial^2 p^\epsilon}{\partial v_{\perp} \partial \alpha} + \frac{1}{v_{\perp}} \cos \alpha \frac{\partial^2 p^\epsilon}{\partial \alpha^2} \right) d\alpha \end{pmatrix} \\
& - (B_{\parallel}^\epsilon + \frac{1}{\epsilon}) \frac{\partial^2 p^\epsilon}{\partial \alpha^2} \\
& = - \begin{pmatrix} 0 \\ v_{\perp} \cos \alpha \\ v_{\perp} \sin \alpha \end{pmatrix} \cdot \nabla_{\mathbf{x}} m - \mathbf{E}^\epsilon \cdot \begin{pmatrix} 0 \\ \cos \alpha \frac{\partial m}{\partial v_{\perp}} \\ \sin \alpha \frac{\partial m}{\partial v_{\perp}} \end{pmatrix} \\
& - \begin{pmatrix} v_{\perp} (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \\ -v_{\parallel} B_3^\epsilon \\ v_{\parallel} B_2^\epsilon \end{pmatrix} \cdot \begin{pmatrix} \frac{\partial m}{\partial v_{\parallel}} \\ \cos \alpha \frac{\partial m}{\partial v_{\perp}} \\ \sin \alpha \frac{\partial m}{\partial v_{\perp}} \end{pmatrix} \\
& - \begin{pmatrix} 0 \\ v_{\perp} \cos \alpha \\ v_{\perp} \sin \alpha \end{pmatrix} \cdot \nabla_{\mathbf{x}} m_1^\epsilon - \mathbf{E}^\epsilon \cdot \begin{pmatrix} 0 \\ \cos \alpha \frac{\partial m_1^\epsilon}{\partial v_{\perp}} \\ \sin \alpha \frac{\partial m_1^\epsilon}{\partial v_{\perp}} \end{pmatrix} \\
& - \begin{pmatrix} v_{\perp} (\cos \alpha B_3^\epsilon - \sin \alpha B_2^\epsilon) \\ -v_{\parallel} B_3^\epsilon \\ v_{\parallel} B_2^\epsilon \end{pmatrix} \cdot \begin{pmatrix} \frac{\partial m_1^\epsilon}{\partial v_{\parallel}} \\ \cos \alpha \frac{\partial m_1^\epsilon}{\partial v_{\perp}} \\ \sin \alpha \frac{\partial m_1^\epsilon}{\partial v_{\perp}} \end{pmatrix}.
\end{aligned} \tag{3.18}$$

Concerning the initial conditions for m_1^ϵ and p^ϵ , we invoke the initial distribution f_0 as follows:

$$\begin{aligned}
& f_0(\mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) \\
&= \frac{1}{2\pi} \int_0^{2\pi} f_0(\mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \theta \mathbf{e}_2 + v_{\perp} \sin \theta \mathbf{e}_3) d\theta \\
&\quad + \left[f_0(\mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) \right. \\
&\quad \quad \left. - \frac{1}{2\pi} \int_0^{2\pi} f_0(\mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \theta \mathbf{e}_2 + v_{\perp} \sin \theta \mathbf{e}_3) d\theta \right] \\
&= m_0(\mathbf{x}, v_{\parallel}, v_{\perp}) \\
&\quad + [f_0(\mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) - m_0(\mathbf{x}, v_{\parallel}, v_{\perp})].
\end{aligned} \tag{3.19}$$

Since $m_{|t=0} = m_0$, we have

$$\begin{aligned}
& m_1^\epsilon(t=0, \mathbf{x}, v_{\parallel}, v_{\perp}) + \frac{\partial p^\epsilon}{\partial \alpha}(t=0, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) \\
&= f_0(\mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \alpha \mathbf{e}_2 + v_{\perp} \sin \alpha \mathbf{e}_3) - m_0(\mathbf{x}, v_{\parallel}, v_{\perp}).
\end{aligned} \tag{3.20}$$

By integrating this equation in α , we find

$$m_1^\epsilon(t=0, \mathbf{x}, v_{\parallel}, v_{\perp}) = 0, \tag{3.21}$$

so we also have

$$p^\epsilon(t=0, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) = \int_0^\alpha f_0(\mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp} \cos \theta \mathbf{e}_2 + v_{\perp} \sin \theta \mathbf{e}_3) d\theta - \alpha m_0(\mathbf{x}, v_{\parallel}, v_{\perp}). \tag{3.22}$$

Remark 3.1 (Conceiving Asymptotic-Preserving schemes). *Building a somehow discretization of the coupled problem (3.10), (3.17), (3.21), (3.18), and (3.22) would give a scheme which will be consistent with (1.15) when $\epsilon > 0$ and consistent with its weak-* limit (2.11) when ϵ is small. This scheme would play the role of the problem $E_{\Delta z}^\epsilon u_{\Delta z}^\epsilon = 0$ in the diagrams shown in (1.10) and (1.14).*

4 Two-Scale Macro-Micro decomposition: preliminaries

The aim exposed in the first paragraph of subsection 3.2 will now be changed in order to obtain the Two-Scale limit model (2.10) when $\epsilon \rightarrow 0$. Eventually, we will see in section 7.2, that the Two-Scale Macro-Micro decomposition described in the following sections, will lead by integration in τ to the classical Macro-Micro decomposition.

We assume from now that the input electric field and the input magnetic field do not depend on ϵ , *i.e.* we take

$$\mathbf{E}^\epsilon = \mathbf{E}, \quad \mathbf{B}^\epsilon = \mathbf{B}. \tag{4.1}$$

4.1 Complements on convergence results

Before entering in the core of the Two-Scale Macro-Micro decomposition, we integrate some knowledge which can be deduced from Frénod, Raviart & Sonnendrücker [10] where the asymptotic expansion of f^ϵ is presented. From the hypotheses (2.1) and (2.2) and adding some regularity assumptions for \mathbf{E} and \mathbf{B} , we can claim that

$$\frac{1}{\epsilon} \left(f^\epsilon(t, \mathbf{x}, \mathbf{v}) - F(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}) \right) \text{ Two-Scale converges to } \tilde{F}(t, \tau, \mathbf{x}, \mathbf{v}) \text{ as } \epsilon \rightarrow 0, \tag{4.2}$$

and that \tilde{F} expresses in terms of other functions \tilde{G} and l in the following way:

$$\tilde{F}(t, \tau, \mathbf{x}, \mathbf{v}) = \tilde{G}(t, \mathbf{x}, \mathbf{u}(\tau, \mathbf{v})) + l(t, \tau, \mathbf{x}, \mathbf{v}), \quad (4.3)$$

where

$$\begin{aligned} l(t, \tau, \mathbf{x}, \mathbf{v}) = & \left(r(\tau + \frac{1}{4}) - r(\frac{1}{4}) \right) \mathbf{v} \cdot \nabla_{\mathbf{x}_\perp} G(t, \mathbf{x}, \mathbf{u}(\tau, \mathbf{v})) \\ & + \left[\left(r(\tau + \frac{1}{4}) - r(\frac{1}{4}) \right) \mathbf{E}(t, \mathbf{x}) + \mathbf{u}(\tau, \mathbf{v}) \times \left(\left(r(\tau + \frac{1}{4}) - r(\frac{1}{4}) \right) \mathbf{B}(t, \mathbf{x}) \right) \right] \\ & \cdot \nabla_{\mathbf{u}_\perp} G(t, \mathbf{x}, \mathbf{u}(\tau, \mathbf{v})), \end{aligned} \quad (4.4)$$

and where \tilde{G} satisfies an equation of the form

$$\frac{\partial \tilde{G}}{\partial t} + u_{\parallel} \frac{\partial \tilde{G}}{\partial x_{\parallel}} + (\mathbf{E}_{\parallel} + \mathbf{u} \cdot \mathbf{B}_{\parallel}) \cdot \nabla_{\mathbf{v}} \tilde{G} = RHS(t, \mathbf{x}, \mathbf{v}, \mathbf{E}, \mathbf{B}, G), \quad (4.5)$$

which right hand side can be explicitly computed like that in Theorem 4.2 of [10].

In these equations, G is the solution of (2.10), r is defined in (2.7), and the gradients $\nabla_{\mathbf{x}_\perp}$ and $\nabla_{\mathbf{u}_\perp}$ are defined as

$$\nabla_{\mathbf{x}_\perp} = \begin{pmatrix} 0 \\ \frac{\partial}{\partial x_2} \\ \frac{\partial}{\partial x_3} \end{pmatrix}, \quad \nabla_{\mathbf{u}_\perp} = \begin{pmatrix} 0 \\ \frac{\partial}{\partial v_2} \\ \frac{\partial}{\partial v_3} \end{pmatrix}. \quad (4.6)$$

4.2 Sought shape of f^ϵ

As announced in the introduction, we want to use a shape given by (1.22)-(1.23) to f^ϵ .

We clearly have that operator

$$\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} : L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6))) \rightarrow L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6))), \quad (4.7)$$

is anti-symmetric, non bounded and with closed range (see Appendix B) and satisfies

$$\text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right) \oplus \text{Im}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right) = L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6))). \quad (4.8)$$

On another hand, we have

$$F \in \text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right). \quad (4.9)$$

Furthermore, we can easily prove that the function l defined in (4.4) is such that

$$l \in \text{Im}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right). \quad (4.10)$$

Indeed, any regular function in $\text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$ reads $\gamma(t, \mathbf{x}, \mathbf{u}(\tau, \mathbf{v}))$ for some $\gamma \in L^\infty(0, T; L^2(\mathbb{R}^6))$. Multiplying $l(t, \tau, \mathbf{x}, \mathbf{v})$ by $\gamma(t, \mathbf{x}, \mathbf{u}(\tau, \mathbf{v}))$, integrating over $\mathbb{R}^6 \times [0, 1]$ and performing the change of variables

$$\mathbf{v} \mapsto \mathbf{u} = \mathbf{u}(\tau, \mathbf{v}), \quad (4.11)$$

we obtain

$$\begin{aligned} \int_{\mathbb{R}^6} \gamma(t, \mathbf{x}, \mathbf{u}) \left\{ \left[\left(\int_0^1 r(\tau - \frac{1}{4}) - r(\frac{1}{4}) d\tau \right) \mathbf{u} \right] \cdot \nabla_{\mathbf{x}_\perp} G(t, \mathbf{x}, \mathbf{u}) \right. \\ \left. - \left[\left(\int_0^1 r(\tau - \frac{1}{4}) - r(\frac{1}{4}) d\tau \right) \mathbf{E}(t, \mathbf{x}) \right. \right. \\ \left. \left. + \mathbf{u} \times \left[\left(\int_0^1 r(\tau - \frac{1}{4}) - r(\frac{1}{4}) d\tau \right) \mathbf{B}(t, \mathbf{x}) \right] \right] \cdot \nabla_{\mathbf{u}_\perp} G(t, \mathbf{x}, \mathbf{u}) \right\} d\mathbf{x} d\mathbf{u} = 0, \end{aligned} \quad (4.12)$$

because the integrals over $[0, 1]$ all vanish.

Now, we integrate every remark we just did. We also notice that a decomposition of the type (1.22) cannot be unique because of choices concerning the place where oscillations are put: in the superscript ϵ or in variable $\frac{t}{\epsilon}$. This will also prescribe what we set now: we look for

$$\begin{aligned} G_1^\epsilon &= G_1^\epsilon(t, \mathbf{x}, \mathbf{u}) \in L^\infty(0, T; L^2(\mathbb{R}^6)), \\ k^\epsilon &= k^\epsilon(t, \tau, \mathbf{x}, \mathbf{v}) \in L^\infty(0, T; L_{\#1}^\infty(\mathbb{R}^+; L^2(\mathbb{R}^6))), \end{aligned} \quad (4.13)$$

such that, for any τ ,

$$\begin{aligned} f^\epsilon(t, \mathbf{x}, \mathbf{v}) &= G(t, \mathbf{x}, \mathbf{u}(\tau, \mathbf{v})) + \epsilon G_1^\epsilon(t, \mathbf{x}, \mathbf{u}(\tau, \mathbf{v})) \\ &+ \epsilon l(t, \tau, \mathbf{x}, \mathbf{v}) + \epsilon \left(\frac{\partial k^\epsilon}{\partial \tau}(t, \tau, \mathbf{x}, \mathbf{v}) + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} k^\epsilon(t, \tau, \mathbf{x}, \mathbf{v}) \right). \end{aligned} \quad (4.14)$$

From now, we shall denote

$$(G \circ \mathbf{u})(t, \tau, \mathbf{x}, \mathbf{v}) = G(t, \mathbf{x}, \mathbf{u}(\tau, \mathbf{v})), \quad (4.15)$$

leading, for instance, to the writing of $F = G \circ \mathbf{u}$. With this notation, we can claim that

$$\begin{aligned} (G \circ \mathbf{u}) + \epsilon (G_1^\epsilon \circ \mathbf{u}) &\in \text{Ker} \left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \right), \\ \epsilon l + \epsilon \left(\frac{\partial k^\epsilon}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} k^\epsilon \right) &\in \text{Im} \left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \right), \end{aligned} \quad (4.16)$$

and, as a consequence of (4.8), decomposition (4.14) exists and is unique since G and l are already identified as the unique couple satisfying (2.10) and (4.4).

In the previous section, when building the usual Macro-Micro decomposition, we projected the equation satisfied by f^ϵ onto the kernel and the range of the involved operator. Here, this cannot be done. Indeed, here, the involved operator is $\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}$ and the function which has to be used in equation (1.15) in order to extract some informations on G_1^ϵ and k^ϵ is

$$\begin{aligned} f^\epsilon(t, \mathbf{x}, \mathbf{v}) &= (G \circ \mathbf{u})(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}) + \epsilon (G_1^\epsilon \circ \mathbf{u})(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}) \\ &+ \epsilon l(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}) + \epsilon \left(\frac{\partial k^\epsilon}{\partial \tau}(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}) + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} k^\epsilon(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}) \right), \end{aligned} \quad (4.17)$$

where τ was replaced by $\frac{t}{\epsilon}$ in (4.14). Once this replacement done, we face with functional spaces where operator (4.7) makes no sense. In order to bypass this difficulty, we use a weak formulation of (1.15) with

oscillating test functions

$$(\psi)^\epsilon = (\psi)^\epsilon(t, \mathbf{x}, \mathbf{v}) = \psi(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}), \quad (4.18)$$

where $\psi = \psi(t, \tau, \mathbf{x}, \mathbf{v})$ is regular, 1-periodic in τ and with compact support on $[0, T) \times \mathbb{R}^3 \times \mathbb{R}^3$ for any fixed $\tau \in [0, 1]$. Writing

$$(F_1^\epsilon)^\epsilon = (F_1^\epsilon)^\epsilon(t, \mathbf{x}, \mathbf{v}) = F_1^\epsilon(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}) = (G_1^\epsilon \circ \mathbf{u})(t, \frac{t}{\epsilon}, \mathbf{x}, \mathbf{v}), \quad (4.19)$$

and using the same convention for other functions, the weak formulation with oscillating test functions reads

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^6} \left[(F)^\epsilon + \epsilon (F_1^\epsilon)^\epsilon + \epsilon (l)^\epsilon + \epsilon (h^\epsilon)^\epsilon \right] \left[\left(\frac{\partial \psi}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial \psi}{\partial \tau} \right)^\epsilon + \mathbf{v} \cdot (\nabla_{\mathbf{x}} \psi)^\epsilon \right. \\ \left. + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot (\nabla_{\mathbf{v}} \psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt = - \int_{\mathbb{R}^6} f_0(\mathbf{x}, \mathbf{v}) \psi(0, 0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v}, \end{aligned} \quad (4.20)$$

where F is linked with G by (2.8), F_1^ϵ with G_1^ϵ by (4.19), l is defined in (4.4) and h^ϵ is linked with k^ϵ by

$$h^\epsilon = \frac{\partial k^\epsilon}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} k^\epsilon. \quad (4.21)$$

The way to deduce the equation for G_1^ϵ and for k^ϵ consists in using, in a first computation, test functions $\psi \in \text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$, *i.e.* of the form

$$\psi = (\gamma \circ \mathbf{u}), \quad (4.22)$$

for regular functions $\gamma = \gamma(t, \mathbf{x}, \mathbf{u})$ compactly supported in $[0, T) \times \mathbb{R}^3 \times \mathbb{R}^3$. Then, in a second computation, test functions belonging to $\text{Im}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$, or writing

$$\psi = \frac{\partial \kappa}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \kappa, \quad (4.23)$$

for some regular functions $\kappa = \kappa(t, \tau, \mathbf{x}, \mathbf{v})$ with compact support in $[0, T) \times \mathbb{R}^3 \times \mathbb{R}^3$ for every $\tau \in [0, 1]$, will be chosen.

5 The Two-Scale Macro equation

We begin by going further in the writing of weak formulation (4.20) by using the link between G and l . We recall this weak formula:

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^6} (F)^\epsilon \left[\frac{\partial(\psi)^\epsilon}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}}(\psi)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot \nabla_{\mathbf{v}}(\psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ + \epsilon \int_0^T \int_{\mathbb{R}^6} (F_1^\epsilon)^\epsilon \left[\left(\frac{\partial \psi}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial \psi}{\partial \tau} \right)^\epsilon + \mathbf{v} \cdot (\nabla_{\mathbf{x}} \psi)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot (\nabla_{\mathbf{v}} \psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ + \epsilon \int_0^T \int_{\mathbb{R}^6} (l)^\epsilon \left[\frac{\partial(\psi)^\epsilon}{\partial t} + (\mathbf{v} \times \frac{\mathcal{M}}{\epsilon}) \cdot \nabla_{\mathbf{v}}(\psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ + \epsilon \int_0^T \int_{\mathbb{R}^6} (l)^\epsilon \left[\mathbf{v} \cdot (\nabla_{\mathbf{x}} \psi)^\epsilon + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot (\nabla_{\mathbf{v}} \psi)^\epsilon \right] dt d\mathbf{x} d\mathbf{v} \\ + \epsilon \int_0^T \int_{\mathbb{R}^6} (h^\epsilon)^\epsilon \left[\left(\frac{\partial \psi}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial \psi}{\partial \tau} \right)^\epsilon + \mathbf{v} \cdot (\nabla_{\mathbf{x}} \psi)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot (\nabla_{\mathbf{v}} \psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ = - \int_{\mathbb{R}^6} f_0(\mathbf{x}, \mathbf{v}) \psi(0, 0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v}. \end{aligned} \quad (5.1)$$

Performing an integration by parts on the first term of (5.1) gives

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^6} (F)^\epsilon \left[\frac{\partial(\psi)^\epsilon}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}}(\psi)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot \nabla_{\mathbf{v}}(\psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
&= - \int_0^T \int_{\mathbb{R}^6} \left[\left(\frac{\partial F}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial F}{\partial \tau} \right)^\epsilon + \mathbf{v} \cdot (\nabla_{\mathbf{x}} F)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot (\nabla_{\mathbf{v}} F)^\epsilon \right] (\psi)^\epsilon d\mathbf{x} d\mathbf{v} dt \\
&\quad - \int_{\mathbb{R}^6} F(0, 0, \mathbf{x}, \mathbf{v}) \psi(0, 0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v} \\
&= - \int_0^T \int_{\mathbb{R}^6} \left[\left(\frac{\partial F}{\partial t} \right)^\epsilon + \mathbf{v} \cdot (\nabla_{\mathbf{x}} F)^\epsilon + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot (\nabla_{\mathbf{v}} F)^\epsilon \right] (\psi)^\epsilon d\mathbf{x} d\mathbf{v} dt \\
&\quad - \int_{\mathbb{R}^6} f_0(\mathbf{x}, \mathbf{v}) \psi(0, 0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v},
\end{aligned} \tag{5.2}$$

since $F \in \text{Ker} \left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \right)$. We invoke the link between F and G , and we consider the change of variables

$$\mathbf{v} \mapsto \mathbf{u} = \mathbf{u} \left(\frac{t}{\epsilon}, \mathbf{v} \right) \text{ which is equivalent to } \mathbf{v} = \mathbf{u} \left(-\frac{t}{\epsilon}, \mathbf{u} \right) = \mathbf{u}^{-1} \left(\frac{t}{\epsilon}, \mathbf{u} \right). \tag{5.3}$$

We also use the following notation

$$(\psi \circ \mathbf{u}^{-1})(t, \tau, \mathbf{x}, \mathbf{u}) = \psi(t, \tau, \mathbf{x}, \mathbf{u}^{-1}(\tau, \mathbf{u})) = \psi(t, \tau, \mathbf{x}, \mathbf{u}(-\tau, \mathbf{u})). \tag{5.4}$$

Since a direct computation yields

$$(\nabla_{\mathbf{v}} F)^\epsilon = \nabla_{\mathbf{v}}(F)^\epsilon = \nabla_{\mathbf{v}}(G \circ \mathbf{u})^\epsilon = r \left(-\frac{t}{\epsilon} \right) ((\nabla_{\mathbf{u}} G) \circ \mathbf{u}), \tag{5.5}$$

(5.2) reads

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^6} (F)^\epsilon \left[\frac{\partial(\psi)^\epsilon}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}}(\psi)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot \nabla_{\mathbf{v}}(\psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
&= - \int_0^T \int_{\mathbb{R}^6} \left[\frac{\partial G}{\partial t} + \left(r \left(-\frac{t}{\epsilon} \right) \mathbf{u} \right) \cdot \nabla_{\mathbf{x}} G + \left(r \left(\frac{t}{\epsilon} \right) \left[\mathbf{E} + \left(r \left(-\frac{t}{\epsilon} \right) \mathbf{u} \right) \times \mathbf{B} \right] \right) \cdot \nabla_{\mathbf{u}} G \right] \\
&\quad (\psi \circ \mathbf{u}^{-1})^\epsilon dt d\mathbf{x} d\mathbf{u} \\
&\quad - \int_{\mathbb{R}^6} f_0(\mathbf{x}, \mathbf{v}) \psi(0, 0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v} \\
&= - \int_0^T \int_{\mathbb{R}^6} \left[\frac{\partial G}{\partial t} + \left(r \left(-\frac{t}{\epsilon} \right) \mathbf{u} \right) \cdot \nabla_{\mathbf{x}} G + \left[r \left(\frac{t}{\epsilon} \right) \mathbf{E} + \mathbf{u} \times \left(r \left(\frac{t}{\epsilon} \right) \mathbf{B} \right) \right] \cdot \nabla_{\mathbf{u}} G \right] (\psi \circ \mathbf{u}^{-1})^\epsilon dt d\mathbf{x} d\mathbf{u} \\
&\quad - \int_{\mathbb{R}^6} f_0(\mathbf{x}, \mathbf{v}) \psi(0, 0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v}.
\end{aligned} \tag{5.6}$$

Using now equation (2.10) satisfied by G , we finally obtain

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^6} (F)^\epsilon \left[\frac{\partial(\psi)^\epsilon}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}}(\psi)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot \nabla_{\mathbf{v}}(\psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
&= - \int_0^T \int_{\mathbb{R}^6} \left(\left(r \left(-\frac{t}{\epsilon} \right) \mathbf{u} \right) \cdot \nabla_{\mathbf{x}_\perp} G + \left[\left(r \left(\frac{t}{\epsilon} \right) \mathbf{E} \right)_\perp + \left[\mathbf{u} \times \left(r \left(\frac{t}{\epsilon} \right) \mathbf{B} \right) \right]_\perp \right] \cdot \nabla_{\mathbf{u}_\perp} G \right) \\
&\quad (\psi \circ \mathbf{u}^{-1})^\epsilon dt d\mathbf{x} d\mathbf{u} \\
&\quad - \int_{\mathbb{R}^6} f_0(\mathbf{x}, \mathbf{v}) \psi(0, 0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v}.
\end{aligned} \tag{5.7}$$

Notice that the last term containing f_0 will vanish with the right hand side member of (5.1).

Now, we study the third term of (5.1). For this purpose, we first notice that

$$\begin{aligned} \frac{\partial(l \circ \mathbf{u}^{-1})}{\partial \tau} &= \left(\frac{\partial l}{\partial \tau} \circ \mathbf{u}^{-1} \right) + \left((r(-\tau) \mathbf{u}) \times \mathcal{M} \right) \cdot \left((\nabla_{\mathbf{v}} l) \circ \mathbf{u}^{-1} \right) \\ &= - (r(-\tau) \mathbf{u}) \cdot \nabla_{\mathbf{x}_{\perp}} G - \left[r(\tau) \mathbf{E} + \mathbf{u} \times (r(\tau) \mathbf{B}) \right] \cdot \nabla_{\mathbf{u}_{\perp}} G, \end{aligned} \quad (5.8)$$

hence, integrating by parts the third term of (5.1) and making the change of variables defined by (5.3), we get

$$\begin{aligned} &\epsilon \int_0^T \int_{\mathbb{R}^6} (l)^\epsilon \left[\frac{\partial(\psi)^\epsilon}{\partial t} + \left(\mathbf{v} \times \frac{\mathcal{M}}{\epsilon} \right) \cdot \nabla_{\mathbf{v}}(\psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ &= - \int_0^T \int_{\mathbb{R}^6} \left[\epsilon \left(\frac{\partial l}{\partial t} \right)^\epsilon + \left(\frac{\partial l}{\partial \tau} \right)^\epsilon + \left(\mathbf{v} \times \mathcal{M} \right) \cdot \left(\nabla_{\mathbf{v}} l \right)^\epsilon \right] (\psi)^\epsilon d\mathbf{x} d\mathbf{v} dt - \int_{\mathbb{R}^6} (l)^\epsilon(0, \mathbf{x}, \mathbf{v}) (\psi)^\epsilon(0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v}, \\ &= - \int_0^T \int_{\mathbb{R}^6} \left[\epsilon \left(\frac{\partial(l \circ \mathbf{u}^{-1})}{\partial t} \right)^\epsilon + \left(\frac{\partial(l \circ \mathbf{u}^{-1})}{\partial \tau} \right)^\epsilon \right] (\psi \circ \mathbf{u}^{-1})^\epsilon d\mathbf{x} d\mathbf{u} dt \\ &= - \int_0^T \int_{\mathbb{R}^6} \epsilon \left(\frac{\partial(l \circ \mathbf{u}^{-1})}{\partial t} \right)^\epsilon (\psi \circ \mathbf{u}^{-1})^\epsilon d\mathbf{x} d\mathbf{u} dt \\ &\quad + \int_0^T \int_{\mathbb{R}^6} \left[\left(r(-\frac{t}{\epsilon}) \mathbf{u} \right) \cdot \nabla_{\mathbf{x}_{\perp}} G + \left[r(\frac{t}{\epsilon}) \mathbf{E} + \mathbf{u} \times \left(r(\frac{t}{\epsilon}) \mathbf{B} \right) \right] \cdot \nabla_{\mathbf{u}_{\perp}} G \right] (\psi \circ \mathbf{u}^{-1})^\epsilon d\mathbf{x} d\mathbf{u} dt, \end{aligned} \quad (5.9)$$

since $l(0, 0, \mathbf{x}, \mathbf{v}) = 0$ (see (4.4)). The last term of (5.9) will vanish with the first term of the right hand side member of (5.7).

Hence (5.1) has to be replaced by

$$\begin{aligned} &- \int_0^T \int_{\mathbb{R}^6} \epsilon \left(\frac{\partial(l \circ \mathbf{u}^{-1})}{\partial t} \right)^\epsilon (\psi \circ \mathbf{u}^{-1})^\epsilon d\mathbf{x} d\mathbf{u} dt \\ &+ \epsilon \int_0^T \int_{\mathbb{R}^6} (F_1^\epsilon)^\epsilon \left[\left(\frac{\partial \psi}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial \psi}{\partial \tau} \right)^\epsilon + \mathbf{v} \cdot \left(\nabla_{\mathbf{x}} \psi \right)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot \left(\nabla_{\mathbf{v}} \psi \right)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ &+ \epsilon \int_0^T \int_{\mathbb{R}^6} (l)^\epsilon \left[\mathbf{v} \cdot \left(\nabla_{\mathbf{x}} \psi \right)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \mathbf{B} \right) \cdot \left(\nabla_{\mathbf{v}} \psi \right)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ &+ \epsilon \int_0^T \int_{\mathbb{R}^6} (h^\epsilon)^\epsilon \left[\left(\frac{\partial \psi}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial \psi}{\partial \tau} \right)^\epsilon + \mathbf{v} \cdot \left(\nabla_{\mathbf{x}} \psi \right)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot \left(\nabla_{\mathbf{v}} \psi \right)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt = 0. \end{aligned} \quad (5.10)$$

Now, choosing ψ as in (4.22) yields

$$\begin{aligned} &\epsilon \int_0^T \int_{\mathbb{R}^6} (G_1^\epsilon \circ \mathbf{u})^\epsilon \left[\left(\frac{\partial \gamma}{\partial t} \right) \circ \mathbf{u} \right)^\epsilon + \mathbf{v} \cdot \left((\nabla_{\mathbf{x}} \gamma) \circ \mathbf{u} \right)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \mathbf{B} \right) \cdot \nabla_{\mathbf{v}}(\gamma \circ \mathbf{u})^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ &- \int_0^T \int_{\mathbb{R}^6} \epsilon \left(\frac{\partial(l \circ \mathbf{u}^{-1})}{\partial t} \right)^\epsilon \gamma d\mathbf{x} d\mathbf{u} dt + \int_0^T \int_{\mathbb{R}^6} \epsilon (h^\epsilon)^\epsilon \left(\frac{\partial(\gamma \circ \mathbf{u})}{\partial t} \right)^\epsilon d\mathbf{x} d\mathbf{v} dt \\ &+ \int_0^T \int_{\mathbb{R}^6} \left(\epsilon (l)^\epsilon + \epsilon (h^\epsilon)^\epsilon \right) \left[\mathbf{v} \cdot \left(\nabla_{\mathbf{x}}(\gamma \circ \mathbf{u}) \right)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \mathbf{B} \right) \cdot \left(\nabla_{\mathbf{v}}(\gamma \circ \mathbf{u}) \right)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt = 0. \end{aligned} \quad (5.11)$$

Making in the first, third and fourth terms the change of variables $\mathbf{v} \mapsto \mathbf{u}(\frac{t}{\epsilon}, v)$, replacing h^ϵ by its

expression in terms of k^ϵ and dividing (5.11) by ϵ finally gives

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^6} G_1^\epsilon \left[\frac{\partial \gamma}{\partial t} + (r(-\frac{t}{\epsilon}) \mathbf{u}) \cdot \nabla_{\mathbf{x}} \gamma + [r(\frac{t}{\epsilon}) \mathbf{E} + \mathbf{u} \times (r(\frac{t}{\epsilon}) \mathbf{B})] \cdot \nabla_{\mathbf{u}} \gamma \right] d\mathbf{x} d\mathbf{u} dt \\
& - \int_0^T \int_{\mathbb{R}^6} \left(\frac{\partial(l \circ \mathbf{u}^{-1})}{\partial t} \right)^\epsilon \gamma d\mathbf{x} d\mathbf{u} dt \\
& + \int_0^T \int_{\mathbb{R}^6} \left[\left(\left(\frac{\partial k^\epsilon}{\partial \tau} \right) \circ \mathbf{u}^{-1} \right)^\epsilon + (r(-\frac{t}{\epsilon}) \mathcal{M}) \cdot \left((\nabla_{\mathbf{v}} k^\epsilon) \circ \mathbf{u}^{-1} \right)^\epsilon \right] \frac{\partial \gamma}{\partial t} d\mathbf{x} d\mathbf{u} dt \\
& + \int_0^T \int_{\mathbb{R}^6} \left[(l \circ \mathbf{u}^{-1})^\epsilon + \left(\left(\frac{\partial k^\epsilon}{\partial \tau} \right) \circ \mathbf{u}^{-1} \right)^\epsilon + (r(-\frac{t}{\epsilon}) \mathcal{M}) \cdot \left((\nabla_{\mathbf{v}} k^\epsilon) \circ \mathbf{u}^{-1} \right)^\epsilon \right] \\
& \quad \times \left[(r(-\frac{t}{\epsilon}) \mathbf{u}) \cdot \nabla_{\mathbf{x}} \gamma + [r(\frac{t}{\epsilon}) \mathbf{E} + \mathbf{u} \times (r(\frac{t}{\epsilon}) \mathbf{B})] \cdot \nabla_{\mathbf{u}} \gamma \right] d\mathbf{x} d\mathbf{u} dt = 0,
\end{aligned} \tag{5.12}$$

which is the Two-Scale Macro equation of the model.

6 The Two-Scale Micro equation

In this section, we use in the weak formulation (4.20) oscillating test functions defined by (4.23). The computation leading the formula (5.10) is valid for any oscillating function ψ so we use it as a starting point for finding the Micro equation.

Recalling that $F_1^\epsilon \in \text{Ker}(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}})$, the second term in (5.10) yields

$$\begin{aligned}
& \epsilon \int_0^T \int_{\mathbb{R}^6} (F_1^\epsilon)^\epsilon \left[\left(\frac{\partial \psi}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial \psi}{\partial \tau} \right)^\epsilon + \mathbf{v} \cdot (\nabla_{\mathbf{x}} \psi)^\epsilon + \left(\mathbf{E} + \mathbf{v} \times \left(\mathbf{B} + \frac{\mathcal{M}}{\epsilon} \right) \right) \cdot (\nabla_{\mathbf{v}} \psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
& = \epsilon \int_0^T \int_{\mathbb{R}^6} (F_1^\epsilon)^\epsilon \left[\frac{\partial(\psi)^\epsilon}{\partial t} + (\mathbf{v} \times \frac{\mathcal{M}}{\epsilon}) \cdot \nabla_{\mathbf{v}}(\psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
& \quad + \epsilon \int_0^T \int_{\mathbb{R}^6} (F_1^\epsilon)^\epsilon \left[\mathbf{v} \cdot (\nabla_{\mathbf{x}} \psi)^\epsilon + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot (\nabla_{\mathbf{v}} \psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
& = -\epsilon \int_0^T \int_{\mathbb{R}^6} \left(\frac{\partial F_1^\epsilon}{\partial t} \right)^\epsilon (\psi)^\epsilon d\mathbf{x} d\mathbf{v} dt + \epsilon \int_{\mathbb{R}^6} F_1^\epsilon(0, 0, \mathbf{x}, \mathbf{v}) \psi(0, 0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v} \\
& \quad + \epsilon \int_0^T \int_{\mathbb{R}^6} (F_1^\epsilon)^\epsilon \left[\mathbf{v} \cdot (\nabla_{\mathbf{x}} \psi)^\epsilon + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot (\nabla_{\mathbf{v}} \psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
& = -\epsilon \int_0^T \int_{\mathbb{R}^6} \left(\frac{\partial F_1^\epsilon}{\partial t} \right)^\epsilon \left[\left(\frac{\partial \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
& \quad + \epsilon \int_{\mathbb{R}^6} F_1^\epsilon(0, 0, \mathbf{x}, \mathbf{v}) \left[\frac{\partial \kappa}{\partial \tau}(0, 0, \mathbf{x}, \mathbf{v}) + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \kappa(0, 0, \mathbf{x}, \mathbf{v}) \right] d\mathbf{x} d\mathbf{v} \\
& \quad + \epsilon \int_0^T \int_{\mathbb{R}^6} (F_1^\epsilon)^\epsilon \left[\mathbf{v} \cdot \left(\frac{\partial \nabla_{\mathbf{x}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot \left((\nabla_{\mathbf{x}} \nabla_{\mathbf{v}} \kappa)^\epsilon \mathbf{v} \right) \right. \\
& \quad \quad \quad \left. + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{E} \times \mathcal{M} + \mathbf{v} \times \mathbf{B} \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \right. \\
& \quad \quad \quad \left. + (\mathbf{E} + \mathbf{v} \times \mathcal{M}) \cdot \left((\nabla_{\mathbf{v}}^2 \kappa)^\epsilon (\mathbf{v} \times \mathcal{M}) \right) \right] d\mathbf{x} d\mathbf{v} dt.
\end{aligned} \tag{6.1}$$

The first term of (5.10) reads

$$\begin{aligned}
& - \int_0^T \int_{\mathbb{R}^6} \epsilon \left(\frac{\partial(l \circ \mathbf{u}^{-1})}{\partial t} \right)^\epsilon (\psi \circ \mathbf{u}^{-1})^\epsilon d\mathbf{x} d\mathbf{v} dt \\
& = -\epsilon \int_0^T \int_{\mathbb{R}^6} \left(\frac{\partial(l \circ \mathbf{u}^{-1})}{\partial t} \right)^\epsilon \left[\left(\frac{\partial \kappa}{\partial \tau} \circ \mathbf{u}^{-1} \right)^\epsilon + \left[r \left(-\frac{t}{\epsilon} \right) \mathbf{v} \times \mathcal{M} \right] \cdot ((\nabla_{\mathbf{v}} \kappa) \circ \mathbf{u}^{-1})^\epsilon \right] d\mathbf{x} d\mathbf{v} dt
\end{aligned} \tag{6.2}$$

The third term gives

$$\begin{aligned}
& \epsilon \int_0^T \int_{\mathbb{R}^6} (l)^\epsilon \left[\mathbf{v} \cdot (\nabla_{\mathbf{x}} \psi)^\epsilon + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot (\nabla_{\mathbf{v}} \psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
& = \epsilon \int_0^T \int_{\mathbb{R}^6} (l)^\epsilon \left[\mathbf{v} \cdot \left(\frac{\partial \nabla_{\mathbf{x}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot ((\nabla_{\mathbf{x}} \nabla_{\mathbf{v}} \kappa)^\epsilon \mathbf{v}) + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial \tau} \right)^\epsilon \right. \\
& \quad \left. + (\mathbf{E} \times \mathcal{M} + \mathbf{v} \times \mathbf{B} \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot ((\nabla_{\mathbf{v}}^2 \kappa)^\epsilon (\mathbf{v} \times \mathcal{M})) \right] d\mathbf{x} d\mathbf{v} dt.
\end{aligned} \tag{6.3}$$

Concerning the last term of (5.10), we have

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^6} (h^\epsilon)^\epsilon \left(\left(\frac{\partial \psi}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial \psi}{\partial \tau} \right)^\epsilon \right) d\mathbf{x} d\mathbf{v} dt \\
& = \int_0^T \int_{\mathbb{R}^6} (h^\epsilon)^\epsilon \frac{\partial(\psi)^\epsilon}{\partial t} d\mathbf{x} d\mathbf{v} dt \\
& = - \int_0^T \int_{\mathbb{R}^6} \frac{\partial(h^\epsilon)^\epsilon}{\partial t} (\psi)^\epsilon d\mathbf{x} d\mathbf{v} dt + \int_{\mathbb{R}^6} h^\epsilon(0, 0, \mathbf{x}, \mathbf{v}) \psi(0, 0, \mathbf{x}, \mathbf{v}) d\mathbf{x} d\mathbf{v} \\
& = - \int_0^T \int_{\mathbb{R}^6} \left(\left(\frac{\partial h^\epsilon}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial h^\epsilon}{\partial \tau} \right)^\epsilon \right) \left[\left(\frac{\partial \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
& \quad + \int_{\mathbb{R}^6} h^\epsilon(0, 0, \mathbf{x}, \mathbf{v}) \left[\frac{\partial \kappa}{\partial \tau}(0, 0, \mathbf{x}, \mathbf{v}) + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \kappa(0, 0, \mathbf{x}, \mathbf{v}) \right] d\mathbf{x} d\mathbf{v} \\
& = - \int_0^T \int_{\mathbb{R}^6} \left(\left(\frac{\partial^2 k^\epsilon}{\partial t \partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial^2 k^\epsilon}{\partial \tau^2} \right)^\epsilon + \frac{1}{\epsilon} (\mathbf{v} \times \mathcal{M}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial \tau} \right)^\epsilon \right) \\
& \quad \times \left[\left(\frac{\partial \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
& \quad + \int_{\mathbb{R}^6} \left[\frac{\partial k^\epsilon}{\partial \tau}(0, 0, \mathbf{x}, \mathbf{v}) + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} k^\epsilon(0, 0, \mathbf{x}, \mathbf{v}) \right] \\
& \quad \times \left[\frac{\partial \kappa}{\partial \tau}(0, 0, \mathbf{x}, \mathbf{v}) + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \kappa(0, 0, \mathbf{x}, \mathbf{v}) \right] d\mathbf{x} d\mathbf{v},
\end{aligned} \tag{6.4}$$

on the one hand, and

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^6} (h^\epsilon)^\epsilon \left[\mathbf{v} \cdot (\nabla_{\mathbf{x}} \psi)^\epsilon + (\mathbf{E} + \mathbf{v} \times (\mathbf{B} + \frac{\mathcal{M}}{\epsilon})) \cdot (\nabla_{\mathbf{v}} \psi)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\
& = \int_0^T \int_{\mathbb{R}^6} \left(\left(\frac{\partial k^\epsilon}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} k^\epsilon)^\epsilon \right) \left[\mathbf{v} \cdot \left(\frac{\partial \nabla_{\mathbf{x}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot ((\nabla_{\mathbf{x}} \nabla_{\mathbf{v}} \kappa)^\epsilon \mathbf{v}) \right. \\
& \quad + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{E} \times \mathcal{M} + (\mathbf{v} \times \mathbf{B}) \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \\
& \quad + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot ((\nabla_{\mathbf{v}}^2 \kappa)^\epsilon (\mathbf{v} \times \mathcal{M})) + \frac{1}{\epsilon} (\mathbf{v} \times \mathcal{M}) \cdot \left(\frac{\partial \nabla_{\mathbf{x}} \kappa}{\partial \tau} \right)^\epsilon \\
& \quad \left. + ((\mathbf{v} \times \mathcal{M}) \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot ((\nabla_{\mathbf{v}}^2 \kappa)^\epsilon (\mathbf{v} \times \mathcal{M})) \right] d\mathbf{x} d\mathbf{v} dt,
\end{aligned} \tag{6.5}$$

on the other hand. Remarking that

$$((\mathbf{v} \times \mathcal{M}) \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon = -(\mathbf{v} \times \mathcal{M}) \cdot ((\nabla_{\mathbf{v}} \kappa)^\epsilon \times \mathcal{M}) = -\mathbf{v} \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \quad (6.6)$$

and dividing by ϵ , we finally get the weak formulation of the Two-Scale Micro equation:

$$\begin{aligned} & - \int_0^T \int_{\mathbb{R}^6} \left(\left(\frac{\partial^2 k^\epsilon}{\partial t \partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial t} \right)^\epsilon + \frac{1}{\epsilon} \left(\frac{\partial^2 k^\epsilon}{\partial \tau^2} \right)^\epsilon + \frac{1}{\epsilon} (\mathbf{v} \times \mathcal{M}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial \tau} \right)^\epsilon \right) \\ & \quad \left[\left(\frac{\partial \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ & + \int_{\mathbb{R}^6} \left[\frac{\partial k^\epsilon}{\partial \tau} (0, 0, \mathbf{x}, \mathbf{v}) + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} k^\epsilon (0, 0, \mathbf{x}, \mathbf{v}) \right] \\ & \quad \left[\frac{\partial \kappa}{\partial \tau} (0, 0, \mathbf{x}, \mathbf{v}) + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \kappa (0, 0, \mathbf{x}, \mathbf{v}) \right] d\mathbf{x} d\mathbf{v} \\ & + \int_0^T \int_{\mathbb{R}^6} \left(\left(\frac{\partial k^\epsilon}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} k^\epsilon)^\epsilon \right) \left[\mathbf{v} \cdot \left(\frac{\partial \nabla_{\mathbf{x}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot ((\nabla_{\mathbf{x}} \nabla_{\mathbf{v}} \kappa)^\epsilon \mathbf{v}) \right. \\ & \quad + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{E} \times \mathcal{M} + (\mathbf{v} \times \mathbf{B}) \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \\ & \quad + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot ((\nabla_{\mathbf{v}}^2 \kappa)^\epsilon (\mathbf{v} \times \mathcal{M})) + \frac{1}{\epsilon} (\mathbf{v} \times \mathcal{M}) \cdot \left(\frac{\partial \nabla_{\mathbf{x}} \kappa}{\partial \tau} \right)^\epsilon \\ & \quad \left. - \mathbf{v} \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot ((\nabla_{\mathbf{v}}^2 \kappa)^\epsilon (\mathbf{v} \times \mathcal{M})) \right] d\mathbf{x} d\mathbf{v} dt \\ & - \int_0^T \int_{\mathbb{R}^6} \left(\frac{\partial (l \circ \mathbf{u}^{-1})}{\partial t} \right)^\epsilon \left[\left(\frac{\partial \kappa}{\partial \tau} \circ \mathbf{u}^{-1} \right)^\epsilon + \left[(r(-\frac{t}{\epsilon}) \mathbf{v}) \times \mathcal{M} \right] \cdot ((\nabla_{\mathbf{v}} \kappa) \circ \mathbf{u}^{-1})^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \quad (6.7) \\ & + \int_0^T \int_{\mathbb{R}^6} (l)^\epsilon \left[\mathbf{v} \cdot \left(\frac{\partial \nabla_{\mathbf{x}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot ((\nabla_{\mathbf{x}} \nabla_{\mathbf{v}} \kappa)^\epsilon \mathbf{v}) + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial \tau} \right)^\epsilon \right. \\ & \quad \left. + (\mathbf{E} \times \mathcal{M} + (\mathbf{v} \times \mathbf{B}) \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot ((\nabla_{\mathbf{v}}^2 \kappa)^\epsilon (\mathbf{v} \times \mathcal{M})) \right] d\mathbf{x} d\mathbf{v} dt \\ & - \int_0^T \int_{\mathbb{R}^6} \left(\frac{\partial F_1^\epsilon}{\partial t} \right)^\epsilon \left[\left(\frac{\partial \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \right] d\mathbf{x} d\mathbf{v} dt \\ & + \int_{\mathbb{R}^6} F_1^\epsilon (0, 0, \mathbf{x}, \mathbf{v}) \left[\frac{\partial \kappa}{\partial \tau} (0, 0, \mathbf{x}, \mathbf{v}) + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \kappa (0, 0, \mathbf{x}, \mathbf{v}) \right] d\mathbf{x} d\mathbf{v} \\ & + \int_0^T \int_{\mathbb{R}^6} (F_1^\epsilon)^\epsilon \left[\mathbf{v} \cdot \left(\frac{\partial \nabla_{\mathbf{x}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{v} \times \mathcal{M}) \cdot ((\nabla_{\mathbf{x}} \nabla_{\mathbf{v}} \kappa)^\epsilon \mathbf{v}) \right. \\ & \quad + (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \left(\frac{\partial \nabla_{\mathbf{v}} \kappa}{\partial \tau} \right)^\epsilon + (\mathbf{E} \times \mathcal{M} + (\mathbf{v} \times \mathbf{B}) \times \mathcal{M}) \cdot (\nabla_{\mathbf{v}} \kappa)^\epsilon \\ & \quad \left. + (\mathbf{E} + \mathbf{v} \times \mathcal{M}) \cdot ((\nabla_{\mathbf{v}}^2 \kappa)^\epsilon (\mathbf{v} \times \mathcal{M})) \right] d\mathbf{x} d\mathbf{v} dt = 0. \end{aligned}$$

Remark 6.1. We are aware that the Two-Scale Macro-Micro system, (5.12)-(6.7), has not necessarily a unique solution. In future work we will look for an additional condition for G_1^ϵ and k^ϵ leading to the uniqueness of the whole solution $(G_1^\epsilon, k^\epsilon)$. Such a condition will be useful for the conception of the numerical scheme for the Two-Scale Macro-Micro model in order to be sure that when $\epsilon \sim 1$ we will approximate the solution of the starting Vlasov equation (1.15).

7 Asymptotic-Preserving Properties

7.1 The Two-Scale Macro-Micro problem

The Two-Scale Macro-Micro problem, *i.e.* the system of equations (2.10)-(4.4)-(5.12)-(6.7), is more complicated to solve than the original problem (1.15), but has two advantages over the latter. First, when $\epsilon \rightarrow 0$ the Two-Scale Macro-Micro model reduces to the limit model in (2.10) while equation (1.15) becomes singular at the limit. Second, the Two-Scale Macro-Micro model decomposes the original solution f^ϵ into a macro part, $G \circ \mathbf{u} + \epsilon G_1^\epsilon \circ \mathbf{u}$, and a micro part, $\epsilon l + \epsilon h_\epsilon$. This question is relevant when doing the transition from the very small ϵ regime to the one of $\epsilon \sim 1$, since our model describes separately the evolution at the macroscopic time (of $G \circ \mathbf{u}$ and $G_1^\epsilon \circ \mathbf{u}$) which contains essential oscillation through \mathbf{u} , and the evolution of oscillation corrections (l and h_ϵ) at the microscopic time.

Now it is easy to see that, under the hypothesis of uniqueness of the solution of equations (5.12) and (6.7) (see Remark 6.1), the Two-Scale Macro-Micro problem (2.10)-(4.4)-(5.12)-(6.7) is equivalent to the original problem (1.15). Indeed, let f^ϵ be the solution of (1.15). We have seen, by using Lemma B.1, that the decomposition (4.14) exists and is unique. Then, by the calculations we did in the previous sections, we obtain that G is solution to (2.10), l is given by the formula (4.4), G_1^ϵ is solution to (5.12) and k^ϵ to (6.7). In particular, we have proved that the equation system (5.12)-(6.7) has solution.

Conversely, assume that the solution of the system (5.12)-(6.7) is unique. Let $(G, l, G_1^\epsilon, k^\epsilon)$ be the solution of (2.10)-(4.4)-(5.12)-(6.7). Then constructing f^ϵ by (4.17), we will obtain the solution of (1.15), since this problem has unique solution.

7.2 Macro-Micro Decomposition *vs* Two-Scale Macro-Micro Decomposition

Property 7.1. *Integrating in τ the Two-Scale Macro-Micro decomposition leads to the classical Macro-Micro model developed in Section 3.2.*

Indeed, let us recall that the unique decomposition made in (4.14) is of the kind

$$f^\epsilon(t, \mathbf{x}, \mathbf{v}) = F(t, \tau, \mathbf{x}, \mathbf{v}) + F_1^\epsilon(t, \tau, \mathbf{x}, \mathbf{v}) + H_1^\epsilon(t, \tau, \mathbf{x}, \mathbf{v}), \quad (7.1)$$

for some $F_1^\epsilon \in \text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$ and $H_1^\epsilon \in \text{Im}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$. Then, integrating (7.1) in τ , we obtain using (2.12)

$$f^\epsilon(t, \mathbf{x}, \mathbf{v}) = f(t, x, \mathbf{v}) + \int_0^1 F_1^\epsilon(t, \tau, \mathbf{x}, \mathbf{v}) d\tau + \int_0^1 H_1^\epsilon(t, \tau, \mathbf{x}, \mathbf{v}) d\tau. \quad (7.2)$$

Now, changing the variable \mathbf{v} in $(v_{||}, v_{\perp}, \alpha)$ defined in (3.1), and using Lemma B.2, we obtain that the function $(v_{\perp}, \alpha) \mapsto \int_0^1 F_1^\epsilon(t, \tau, \mathbf{x}, v_{||} \mathbf{e}_1 + v_{\perp}(\cos \alpha \mathbf{e}_2 + \sin \alpha \mathbf{e}_3)) d\tau$ is in $\text{Ker}\left(\frac{\partial}{\partial \alpha}\right)$. Similarly, $(v_{\perp}, \alpha) \mapsto \int_0^1 H_1^\epsilon(t, \tau, \mathbf{x}, v_{||} \mathbf{e}_1 + v_{\perp}(\cos \alpha \mathbf{e}_2 + \sin \alpha \mathbf{e}_3)) d\tau$ is in $\text{Im}\left(\frac{\partial}{\partial \alpha}\right)$. Next, writing (7.2) in the new variables and recalling the decomposition of $f^\epsilon(t, \mathbf{x}, \mathbf{v})$ in (3.12), we deduce from (3.5) and the uniqueness of such a decomposition (stated in Lemma A.1(iii)) that

$$\begin{aligned} m_1^\epsilon(t, \mathbf{x}, v_{||}, v_{\perp}) &= \int_0^1 F_1^\epsilon(t, \tau, \mathbf{x}, v_{||} \mathbf{e}_1 + v_{\perp}(\cos \alpha \mathbf{e}_2 + \sin \alpha \mathbf{e}_3)) d\tau, \\ n^\epsilon(t, \mathbf{x}, v_{||}, v_{\perp}, \alpha) &= \int_0^1 H_1^\epsilon(t, \tau, \mathbf{x}, v_{||} \mathbf{e}_1 + v_{\perp}(\cos \alpha \mathbf{e}_2 + \sin \alpha \mathbf{e}_3)) d\tau. \end{aligned} \quad (7.3)$$

7.3 Convergence of the Two-Scale Macro-Micro problem

Let us recall that our Two-Scale Macro-Micro decomposition (4.14) is based on the convergence result in (4.2). More precisely, keeping in mind that $G + \epsilon \tilde{G} + \epsilon l$ is the first order approximation of f^ϵ , then (4.14) can be seen as a Macro-Micro decomposition at the first order level of approximation of f^ϵ . Next, we justify this approximation using the Two-Scale convergence.

Theorem 7.2. *We assume that $f_0 \in L^2(\mathbb{R}^6)$, $\mathbf{E} \in W^{1,\infty}(\mathbb{R}^3)$, $\mathbf{B} \in W^{1,\infty}(\mathbb{R}^3)$, and $\partial l/\partial t, \nabla_{\mathbf{x},\mathbf{v}} l \in L^\infty(0, T; L^\infty(0, 1; L^2(\mathbb{R}^6)))$. Then, when $\epsilon \rightarrow 0$, the solutions $G_1^\epsilon \circ \mathbf{u}$ of (5.12) Two-Scale converges to $\tilde{G} \in L^\infty(0, T; L^2(\mathbb{R}^6))$, the solution of (4.5). When $\epsilon \rightarrow 0$, the solutions k^ϵ of (6.7) Two-Scale converges to 0.*

Proof. Recall from Theorem 1.5 of [10] that when $\epsilon \rightarrow 0$, $\left(\frac{1}{\epsilon}(f^\epsilon - G \circ \mathbf{u}) - l\right)_{\epsilon>0}$ Two-Scale converges to $\tilde{G} \circ \mathbf{u} \in \text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$. Then, since (4.17) implies that

$$\frac{1}{\epsilon}(f^\epsilon - G \circ \mathbf{u}) - l = G_1^\epsilon \circ \mathbf{u} + h^\epsilon,$$

and since $G_1^\epsilon \circ \mathbf{u} \in \text{Ker}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$ and $h^\epsilon \in \text{Im}\left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$, the theorem's conclusion is clearly true. \square

Appendices

In the first Appendix we characterize the projection onto $\text{Ker}\left(\frac{\partial}{\partial \alpha}\right)$. We establish that it consists in computing its average in α and projecting it onto $\text{Im}\left(\frac{\partial}{\partial \alpha}\right)$ consists in subtracting from it its average value. Then, in the second one, we do the same with operator $\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}$. We also establish the link that exists between these two operators.

A About the operator $\frac{\partial}{\partial \alpha}$

Lemma A.1. *Let the unbounded operator $\frac{\partial}{\partial \alpha} : L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp)) \rightarrow L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$. Then, it has a closed image in $L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$ and*

$$(i) \text{Ker}\left(\frac{\partial}{\partial \alpha}\right)^\perp = \text{Im}\left(\frac{\partial}{\partial \alpha}\right).$$

(ii) *For any function $f \in L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$, the projection of f on $\text{Ker}\left(\frac{\partial}{\partial \alpha}\right)$ is the function*

$$Pf : (v_\perp, \alpha) \mapsto \frac{1}{2\pi} \int_0^{2\pi} f(v_\perp, \theta) d\theta. \quad (\text{A.1})$$

$$(iii) L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp)) = \text{Ker}\left(\frac{\partial}{\partial \alpha}\right) \oplus \text{Im}\left(\frac{\partial}{\partial \alpha}\right).$$

Proof. (i) Let $q \in \left(\text{Ker}\left(\frac{\partial}{\partial\alpha}\right)\right)^\perp$. Then, for any $g \in \text{Ker}\left(\frac{\partial}{\partial\alpha}\right)$, i.e. $g = g(v_\perp)$, we have

$$\int_0^{2\pi} \int_0^{+\infty} q(v_\perp, \alpha) g(v_\perp) v_\perp dv_\perp d\alpha = 0. \quad (\text{A.2})$$

Remarking that q can be written as

$$q(v_\perp, \alpha) = \frac{\partial}{\partial\alpha} \left(\int_0^\alpha q(v_\perp, \theta) d\theta \right), \quad (\text{A.3})$$

we only have to prove that the function K defined by

$$K(v_\perp, \alpha) = \int_0^\alpha q(v_\perp, \theta) d\theta, \quad (\text{A.4})$$

is in $L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$, i.e. K is 2π -periodic in α and $\int_0^{2\pi} \int_0^{+\infty} |K(v_\perp, \alpha)|^2 v_\perp dv_\perp d\alpha < +\infty$.

First, we have

$$K(v_\perp, \alpha + 2\pi) - K(v_\perp, \alpha) = \int_0^{2\pi} q(v_\perp, \theta) d\theta = 0, \quad (\text{A.5})$$

by (A.2). Secondly, using the Jensen inequality, we remark that

$$|K(v_\perp, \alpha)|^2 \leq |\alpha| \int_0^\alpha |q(v_\perp, \theta)|^2 d\theta \leq 2\pi \int_0^{2\pi} |q(v_\perp, \theta)|^2 d\theta. \quad (\text{A.6})$$

Then

$$\int_0^{2\pi} \int_0^{+\infty} |K(v_\perp, \alpha)|^2 v_\perp dv_\perp d\alpha \leq 4\pi^2 \int_0^{2\pi} \int_0^{+\infty} |q(v_\perp, \theta)|^2 v_\perp dv_\perp d\theta < +\infty, \quad (\text{A.7})$$

since $q \in L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$. Thus $q \in \text{Im}\left(\frac{\partial}{\partial\alpha}\right)$ and so $\text{Ker}\left(\frac{\partial}{\partial\alpha}\right)^\perp \subset \text{Im}\left(\frac{\partial}{\partial\alpha}\right)$. Since the converse inclusion is obvious, (i) is proved and the image of the operator is indeed closed.

(ii) It is clear that $f \in L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$ implies that $Pf \in L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$. Then, using the projection characterization, we have for any function $\psi \in L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$ such that $\frac{\partial\psi}{\partial\alpha} = 0$,

$$\begin{aligned} & \int_0^{2\pi} \int_0^{+\infty} \left(f(v_\perp, \alpha) - \frac{1}{2\pi} \int_0^{2\pi} f(v_\perp, \theta) d\theta \right) \psi(v_\perp) v_\perp dv_\perp d\alpha \\ &= \int_0^{+\infty} \left(\int_0^{2\pi} f(v_\perp, \alpha) d\alpha \right) \psi(v_\perp) v_\perp dv_\perp - \left(\frac{1}{2\pi} \int_0^{2\pi} d\alpha \right) \int_0^{+\infty} \left(\int_0^{2\pi} f(v_\perp, \theta) d\theta \right) \psi(v_\perp) v_\perp dv_\perp \\ &= 0. \end{aligned}$$

Hence Pf defined in (A.1) is the projection of f on $\text{Ker}\left(\frac{\partial}{\partial\alpha}\right)$.

(iii) Using item (i) we have only to prove that any $f \in L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$ writes as a sum $f = g + h$ with $g \in \text{Ker}\left(\frac{\partial}{\partial\alpha}\right)$ and $h \in \text{Im}\left(\frac{\partial}{\partial\alpha}\right)$. Given $f \in L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_\perp dv_\perp))$, let g be the projection of f on $\text{Ker}\left(\frac{\partial}{\partial\alpha}\right)$ and let $h = f - g$. Then, by item (ii) we only need to show that $h \in \text{Im}\left(\frac{\partial}{\partial\alpha}\right)$. This can be done using a similar way as in the proof of item (i). This concludes the proof of the Lemma. \square

B The operator $\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}$

Lemma B.1. *Let the unbounded operator*

$$\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} : L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6))) \rightarrow L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6))).$$
 Then we have

$$L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6))) = \text{Ker} \left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \right) \oplus \text{Im} \left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \right). \quad (\text{B.1})$$

Proof. Let $f \in L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6)))$. Through the change of variables

$$\begin{cases} v_{\parallel} \in \mathbb{R} \text{ such that } v_{\parallel} = \mathbf{v} \cdot \mathbf{e}_1, \\ v_{\perp} \in \mathbb{R}^+ \text{ such that } v_{\perp} = \sqrt{v_2^2 + v_3^2}, \\ \alpha \in [0, 2\pi] \text{ such that } v_2 = v_{\perp} \cos \alpha \text{ and } v_3 = v_{\perp} \sin \alpha, \end{cases} \quad (\text{B.2})$$

\tilde{f} (where $\tilde{f}(t, \tau, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) = f(t, \tau, \mathbf{x}, v_{\parallel} \mathbf{e}_1 + v_{\perp}(\cos \alpha \mathbf{e}_2 + \sin \alpha \mathbf{e}_3))$) is in $L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^3 \times \mathbb{R} \times \mathbb{R}^+; v_{\perp} d\mathbf{x} dv_{\parallel} dv_{\perp})))$) while the operator $\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}$ writes $\frac{\partial}{\partial \tau} - 2\pi \frac{\partial}{\partial \alpha}$. Next, performing the second change of variables

$$\begin{cases} \sigma = \tau + \frac{1}{2\pi} \alpha, \\ \beta = \tau - \frac{1}{2\pi} \alpha, \end{cases} \quad (\text{B.3})$$

\check{f} (where $\check{f}(t, \sigma, \mathbf{x}, v_{\parallel}, v_{\perp}, \beta) = \tilde{f}(t, (\sigma + \beta)/2, \mathbf{x}, v_{\parallel}, v_{\perp}, \pi(\sigma - \beta))$) stays in $L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^3 \times \mathbb{R} \times \mathbb{R}^+; v_{\perp} d\mathbf{x} dv_{\parallel} dv_{\perp})))$) and the operator $\frac{\partial}{\partial \tau} - 2\pi \frac{\partial}{\partial \alpha}$ (and consequently $\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}$) becomes $2 \frac{\partial}{\partial \beta}$. From now on, let us fix $(t, \mathbf{x}) \in [0, T] \times \mathbb{R}^3$. Considering

$$\begin{aligned} & \frac{\partial}{\partial \beta} : L^\infty_{\#1}(\mathbb{R}^+; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^3 \times \mathbb{R} \times \mathbb{R}^+; v_{\perp} d\mathbf{x} dv_{\parallel} dv_{\perp}))) \\ & \rightarrow L^\infty_{\#1}(\mathbb{R}^+; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^3 \times \mathbb{R} \times \mathbb{R}^+; v_{\perp} d\mathbf{x} dv_{\parallel} dv_{\perp}))), \end{aligned} \quad (\text{B.4})$$

we know, by Lemma A.1, that $L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R}^+; v_{\perp} dv_{\perp})) = \text{Ker} \left(\frac{\partial}{\partial \beta} \right) \oplus \text{Im} \left(\frac{\partial}{\partial \beta} \right)$. Hence, because $L^\infty_{\#1}(\mathbb{R}^+; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R} \times \mathbb{R}^+; v_{\perp} dv_{\parallel} dv_{\perp}))) \subset L^2_{\#1}(\mathbb{R}^+; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R} \times \mathbb{R}^+; v_{\perp} dv_{\parallel} dv_{\perp})))$, we can write $\check{f} = P\check{f} + (\check{f} - P\check{f})$, where $P\check{f}$ is defined as in (A.1). Since \check{f} was chosen in $L^\infty_{\#1}(\mathbb{R}^+; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R} \times \mathbb{R}^+; v_{\perp} dv_{\parallel} dv_{\perp})))$ then it is easy to see that $P\check{f}$ belongs to this same space. Obviously $\check{f} - P\check{f} \in L^\infty_{\#1}(\mathbb{R}^+; L^2_{\#2\pi}(\mathbb{R}; L^2(\mathbb{R} \times \mathbb{R}^+; v_{\perp} dv_{\parallel} dv_{\perp})))$ and thus (B.1) is proved. \square

Lemma B.2. *Let $f \in L^\infty(0, T; L^\infty_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6)))$ and let \tilde{f} be defined as in the proof of Lemma B.1. Then*

(i) *if $f \in \text{Ker} \left(\frac{\partial}{\partial \tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}} \right)$ then for any fixed $(t, \mathbf{x}, v_{\parallel}) \in [0, T] \times \mathbb{R}^3 \times \mathbb{R}$ we have*

$$\mathcal{F} : (v_{\perp}, \alpha) \mapsto \int_0^1 \tilde{f}(t, \tau, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) d\tau \in \text{Ker} \left(\frac{\partial}{\partial \alpha} \right)$$

(ii) if $f \in \text{Im}\left(\frac{\partial}{\partial\tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$ then for any fixed $(t, \mathbf{x}, v_{\parallel}) \in [0, T] \times \mathbb{R}^3 \times \mathbb{R}$ we have

$$\mathcal{F} : (v_{\perp}, \alpha) \mapsto \int_0^1 \tilde{f}(t, \tau, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) d\tau \in \text{Im}\left(\frac{\partial}{\partial\alpha}\right).$$

Proof. (i) Let $f \in \text{Ker}\left(\frac{\partial}{\partial\tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$ and fix $(t, \mathbf{x}, v_{\parallel}) \in [0, T] \times \mathbb{R}^3 \times \mathbb{R}$. Thus, by the proof of Lemma B.1, $\tilde{f} \in \text{Ker}\left(\frac{\partial}{\partial\tau} - 2\pi\frac{\partial}{\partial\alpha}\right)$. We then derive \mathcal{F} :

$$\begin{aligned} \frac{\partial\mathcal{F}}{\partial\alpha}(v_{\perp}, \alpha) &= \int_0^1 \frac{\partial\tilde{f}}{\partial\alpha}(t, \tau, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) d\tau \\ &= \frac{1}{2\pi} \int_0^1 \frac{\partial\tilde{f}}{\partial\tau}(t, \tau, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) d\tau \\ &= 0, \end{aligned} \tag{B.5}$$

by the 1-periodicity in τ of \tilde{f} . Thus, $\mathcal{F} \in \text{Ker}\left(\frac{\partial}{\partial\alpha}\right)$. (ii) Now let $f \in \text{Im}\left(\frac{\partial}{\partial\tau} + (\mathbf{v} \times \mathcal{M}) \cdot \nabla_{\mathbf{v}}\right)$. Then there exists a function $h \in L^{\infty}(0, T; L^{\infty}_{\#1}(\mathbb{R}^+; L^2(\mathbb{R}^6)))$ such that $\tilde{f} = \frac{\partial\tilde{h}}{\partial\tau} - 2\pi\frac{\partial\tilde{h}}{\partial\alpha}$. Fix $(t, \mathbf{x}, v_{\parallel}) \in [0, T] \times \mathbb{R}^3 \times \mathbb{R}$. Integrating in τ from 0 to 1, we obtain, using that h is 1-periodic in τ ,

$$\mathcal{F}(v_{\perp}, v_{\parallel}) = -2\pi\frac{\partial}{\partial\alpha} \int_0^1 \tilde{h}(t, \tau, \mathbf{x}, v_{\parallel}, v_{\perp}, \alpha) d\tau,$$

leading to $\mathcal{F} \in \text{Im}\left(\frac{\partial}{\partial\alpha}\right)$ and thus concluding the Lemma. \square

References

- [1] ALLIOT, P., FRÉNOT, E., MONBET, V., *Long term object drift in the ocean with tide and wind*, Multiscale Model. Simul. **5**-2 (2006), 514-531.
- [2] ALLAIRE, G., *Homogenization and two-scale convergence*, SIAM J. Math. Anal. **23**-6 (1992), 1482-1518.
- [3] BENNOUNE, M., LEMOU M., MIEUSSENS L., *Uniformly stable numerical schemes for the Boltzmann equation preserving the compressible Navier-Stokes asymptotics*, J. Comput. Phys. **227**-8 (2008), 3781-3803.
- [4] BOSTAN, M., *The Vlasov-Poisson system with strong external magnetic field. Finite Larmor radius regime*, Asymptot. Anal. **61**-2 (2009), 91-123.
- [5] BOSTAN, M., *Transport equations with disparate advection fields. Application to the gyrokinetic models in plasma physics*, J. Diff. Eq., **249** (2010), 1620-1663.
- [6] CROUSEILLES, N., LEMOU, M., *An asymptotic preserving scheme based on a micro-macro decomposition for Collisional Vlasov equations: diffusion and high-field scaling limits*, Kinet. Relat. Models **4**-2 (2011), 441-477.
- [7] FILBET, F., JIN, S., *A class of asymptotic preserving schemes for kinetic equations and related problems with stiff sources*, J. Comp. Phys. **229**-20 (2010), 7625-7648.
- [8] FRÉNOT, E., MOUTON. A., *Two-dimensional Finite Larmor Radius approximation in canonical gyrokinetic coordinates*, J. Pure Appl. Math. Adv. Appl. **4**-2 (2010), 135-166.

- [9] FRÉNOT, E., MOUTON, A., SONNENDRÜCKER, E., *Two-scale numerical simulation of the weakly compressible 1D isentropic Euler equations*, Numer. Math. **108**-2 (2007), 263-293.
- [10] FRÉNOT, E., RAVIART, P.-A., SONNENDRÜCKER, E., *Two-scale expansion of a singularly perturbed convection equation*, J. Math. Pures Appl. **80**-8 (2001), 815-843.
- [11] FRÉNOT, E., SALVARANI, F., SONNENDRÜCKER, E., *Long time simulation of a beam in a periodic focusing channel via a two-scale PIC-method*, Math. Models Methods Appl. Sci. **19**-2 (2009), 175-197.
- [12] FRÉNOT, E., SONNENDRÜCKER, E., *Homogenization of the Vlasov equation and of the Vlasov-Poisson system with a strong external magnetic field*, Asymptot. Anal. **18**-3-4 (1998), 193-214.
- [13] FRÉNOT, E., SONNENDRÜCKER, E., *The Finite Larmor Radius approximation*, SIAM J. Math. Anal. **32**-6 (2001), 1227-1247.
- [14] GOLSE, F., SAINT-RAYMOND, L., *The Vlasov-Poisson system with strong magnetic field*, J. Math. Pures Appl. **78** (1999), 791-817.
- [15] GOLSE, F., SAINT-RAYMOND, L., *The Vlasov-Poisson system with strong magnetic field in quasineutral regime*, Math. Models Methods Appl. Sci. **13**-5 (2003), 661-714.
- [16] HAN-KWAN, D., *The three-dimensional finite Larmor radius approximation*, Asymptot. Anal. **66**-1 (2010), 9-33.
- [17] JIN S., *Efficient asymptotic-preserving (AP) schemes for some multiscale kinetic equations*, SIAM J. Sci. Comput. **21**-2 (1999), 441-454.
- [18] JIN, S., SHI Y., *A Micro-Macro Decomposition-Based Asymptotic-Preserving Scheme for the Multispecies Boltzmann Equation*, SIAM J. Sci. Comp. **31**-6 (2010), 4580-4606.
- [19] KLAR, A., *A Numerical Method for Kinetic Semiconductor Equations in the Drift Diffusion Limit*, SIAM J. Sci. Comp. **20**-5 (1999), 1696-1712.
- [20] LEMOU, M., MIEUSSENS, L., *A New Asymptotic Preserving Scheme Based on Micro-Macro Formulation for Linear Kinetic Equations in the Diffusion Limit*, SIAM J. Sci. Comp. **31**-1 (2008), 334-368.
- [21] MOUTON, A., *Two-scale semi-lagrangian simulation of a charged particles beam in a periodic focusing channel*, Kinet. Relat. Models **2**-2 (2009), 251-274.
- [22] MOUTON, A., *Approximation multi-échelles de l'équation de Vlasov*, Thèse de l'Université de Strasbourg (2009).
- [23] NGUETSENG, G., *A general convergence result for a functional related to the theory of homogenization*, SIAM J. Math. Anal. **20**-3 (1989), 608-623.