



HAL
open science

**Refinement of the alpha-U₄O₉ crystalline structure:
New insight into the U₄O₉-> U₃O₈
transformation**

Lionel Desgranges, Gianguido Baldinozzi, David Simeone, H. E. Fischer

► **To cite this version:**

Lionel Desgranges, Gianguido Baldinozzi, David Simeone, H. E. Fischer. Refinement of the alpha-U₄O₉ crystalline structure: New insight into the U₄O₉-> U₃O₈ transformation. *Inorganic Chemistry*, 2011, 50, pp.6146-6151. 10.1021/ic200316b . hal-00639190

HAL Id: hal-00639190

<https://hal.science/hal-00639190>

Submitted on 23 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Refinement of the $r\text{-U}_4\text{O}_9$ Crystalline Structure: New Insight into the U_4O_9 to U_3O_8 Transformation

L. Desgranges,^{*,†} G. Baldinozzi,^{‡,§} D. Siméone,^{‡,§} and H. E. Fischer[^]

[†]CEA/DEN/DEC, Bat 352, Cadarache 13108, Saint Paul lez Durance, France


[‡]Matériaux Fonctionnels pour l'Energie, SPMS CNRS-Ecole Centrale Paris, 92295 Châtenay-Malabry, France

[§]CEA/DEN/DANS/DMN/SRMA/LA2M, 91191 Gif-sur-Yvette, France

[^]Institut Laue-Langevin, 6 rue Jules Horowitz, B.P. 156, 38042 Grenoble Cedex, France

b Supporting Information

ABSTRACT:



The oxidation reaction of UO_2 into U_3O_8 is studied as a function of the crystalline distortion of interstitial oxygen clusters, named cuboctahedra, which appear in U_4O_9 and U_3O_7 intermediate phases. For that purpose, the refinement of $R\text{-U}_4\text{O}_9$ was performed because this phase undergoes a trigonal distortion from cubic $\beta\text{-U}_4\text{O}_9$ when the temperature is decreased. In $R\text{-U}_4\text{O}_9$, the cuboctahedra can be described as crumpled sheets taken from a fragment of U_3O_8 . The manner by which the accumulation of crumpled sheets can lead to the formation of U_3O_8 is discussed.

INTRODUCTION

The reaction of oxidation of UO_2 into U_3O_8 is of technological importance for the nuclear industry, both for reprocessing of fresh defective fuel and for dry storage of used nuclear fuel.¹ The kinetics of this reaction was characterized 40 years ago,² but the underlying atomic mechanisms are still under discussion. Allen and Holmes³ first proposed that this reaction occurred as a shear transformation of the (111) dense planes of the face-centered-cubic (fcc) crystalline structure of UO_2 . This interpretation was consistent with the crystallographic positions of the uranium atoms in both UO_2 and U_3O_8 structures. At that time, however,

no information was available on the positions of the oxygen atoms in U_4O_9 and U_3O_7 , the intermediate phases formed during

the oxidation of UO_2 at temperatures of less than 400 °C.² Since then, following Bevan et al.'s pioneering work,⁶ Cooper and Willis⁷ determined the crystal structure of $\beta\text{-U}_4\text{O}_9$ (stable between 353 and 873 K), and Desgranges et al.⁸ proposed a crystal structure for $\beta\text{-U}_3\text{O}_7$. These crystallographic refinements shed light on the existence of a complex cluster, named a cuboctahedron, that includes five oxygen interstitials,⁹ with one

being at the cluster's center (Figure 1). The stability of the cuboctahedron was discussed in comparison with single oxygen interstitials using first-principle methods^{10–12} and molecular

dynamics.¹³ However, no clear relationship exists between fluorite-derived crystal structures (large supercells) and the crystal structure of U_3O_8 because the organization in the dense planes proposed by Allen and Holmes is not related to the distribution of cuboctahedra.

In fact, transformation from UO_2 into U_3O_8 results from the convolution of two different mechanisms: the accumulation of oxygen in the solid and the distortion of its crystalline lattice. In this paper, we will focus on the distortion of the crystalline lattice only. For that purpose, we decided to study the crystalline structure of $R\text{-U}_4\text{O}_9$ (space group $R3c$), which is a trigonal

distortion of the cubic $\beta\text{-U}_4\text{O}_9$ (space group $F43d$). This trigonal distortion privileges the family of UO_2 dense planes perpendi-

cular to the rhombohedral axis. Thus, information could be gained on how the crystalline lattice behaves during the transformation to U_3O_8 .

Conradson et al.¹⁴ performed X-ray absorption spectroscopy (XAS) experiments on several samples, having compositions ranging from UO_2 to U_4O_9 , in order to study the uranium atom's local environment. Garrido et al.¹⁵ performed neutron diffraction

Received: February 16, 2011




Figure 1. Comparison between the UO₂ lattice (left part) and a cuboctahedron in the B-U₄O₉ lattice (right part). Uranium atoms are gray, oxygen atoms are red, and the oxygen atom at the center of the cuboctahedron is yellow.

experiments at room temperature on a R-U₄O₉ sample, and they derived its pair distribution function (PDF). The PDF is a function of the interatomic distance r and relates the probability of finding an atom at a distance r from an average atom at the origin. However, the PDF averages over all atomic species and also over all crystallographic directions and thus provides some structural information, but no detailed crystal structure of R-U₄O₉ has been proposed so far.

Although R-U₄O₉ is a crystalline compound, it was not possible to use standard Rietveld analysis of neutron diffraction patterns that had been successful in the case of B-U₄O₉ because of the number of parameters to be refined is larger. In a low-symmetry structure, even though the crystallographic data included in the diffraction pattern are essentially the same in both phases, the PDF refinement is more robust than the Rietveld refinement. In the case of U₄O₉, PDF analysis is very interesting because the central oxygen atom has a very different local environment compared with those of other oxygen atoms. The central oxygen atom has a characteristic distance with other oxygen atoms at 3.3 Å, the distance of which does not overlap with those for oxygen atoms in the regular UO₂ lattice.

In this paper, we present the measurement and refinement of the PDF for U₄O₉ at room temperature. The results are discussed with respect to the transformation of UO₂ into U₃O₈.

EXPERIMENTAL SECTION

Sample Preparation. The U₄O₉ sample was prepared by heat treatment of a mixture of UO₂ and U₃O₈ powder. The relative mass fraction of UO₂ and U₃O₈ was chosen in order to get, on average, a UO_{2.23} composition. The two powders were carefully mixed. They were loaded into a quartz tube. The quartz tube was vacuum-pumped and sealed. Then this powder underwent a heat treatment at 1050 °C. The obtained powder had a dark color. Its quality was checked by X-ray diffraction. The powder had a grain size of a few micrometers, as measured by optical microscopy.

Neutron Diffraction and PDF Determination. The neutron diffraction experiment was performed on the D4c neutron diffractometer¹⁶ at the Institute Laue-Langevin (ILL), Grenoble, France. A mass of 14.04 g of U₄O₉ powder was put into a sample container consisting of an 8-mm-diameter vertical cylinder of 0.1-mm-thick vanadium foil with niobium stoppers at the top and bottom. This sample container was sufficiently sealed to prevent the powder from escaping but not sufficiently sealed to allow a vacuum to be pumped. The sample environment in the diffractometer was therefore ambient air. Several diffraction patterns were acquired in order to measure accurately the various sources of background scattering: first with no sample at all




Figure 2. R-U₄O₉ neutron diffraction pattern corrected from air and sample holder diffusion.




Figure 3. R-U₄O₉ $D(r)$ obtained by Fourier transformation (see the text for details).

(i.e., an empty diffractometer), then with an empty sample container, and last with the sample container filled with U₄O₉ powder, each for respective counting times of 4, 4, and 24 h. The stability of the diffraction pattern was checked by comparing data acquired at the same position and at different times. The neutron wavelength was determined by refinement of the diffraction pattern of a standard nickel sample and was found to be 0.4970 Å. The diffraction pattern was recorded up to $Q_{\text{max}} = 23.5 \text{ \AA}^{-1}$ (Figure 2).

The obtained diffraction pattern was treated to subtract the background due to air scattering and that due to scattering from the vanadium container, as well as the isotropic scattering coming from incoherent scattering that is subject to distortion from inelastic scattering effects (Placzek correction). A measurement of the sample height in the container gave the packing fraction, and by measurement of the diffraction from a vanadium standard, the diffraction pattern for the sample could be normalized onto an absolute scale of barns/sr/atom. The differential correlation function, $D(r)$, contains information relating to differences between the average density at a distance, r , from an average origin atom and the corresponding radial density for a homogeneous material. It was obtained for R-U₄O₉ (Figure 3) by Fourier transformation of this corrected diffraction pattern. The quality of this $D(r)$, as well as its amplitude normalization, was confirmed by comparing the slope of the PDF at $r = 0 \text{ \AA}$ to the density of atoms per volume unit in the sample. The total distribution function $T(r)$ obtained from the R-U₄O₉ $D(r)$ is presented in Figure 4. A peak in $T(r)$ indicates an interatomic distance that occurs frequently in the sample, and its area is directly related to the coordination number for that distance.

Our $D(r)$ and $T(r)$ are very similar to those obtained by Garrido et al. measured at ISIS¹⁵ although both experiments used very different experimental procedures. On the basis of these observations, our results were considered to be of sufficient quality for refinement of the R-U₄O₉ PDF.




Figure 4. Zoom from 0 to 7 Å of the calculated R-U₄O₉ $T(r)$ (see the text for details).

Refinement Procedure. Our PDF for R-U₄O₉ was refined with

PDFGUI software.¹⁷ In such a minimization procedure, the crystalline structure used as the starting point has to be as close as possible to the actual crystalline structure.

According to Belbeoch et al.,¹⁸ the room temperature phase of U₄O₉ is the result of a distortion of the cubic β phase that is very sensitive to the grain size. On the one hand, in fine-grain-size powder samples (<1 μm), the 4-fold superstructure lines, which are characteristic of the ordering of the cuboctahedra, reversibly disappear when the sample temperature is lowered below the R \rightarrow β transition temperature with a significant trigonal distortion. On the other hand, in coarse-grained samples ($\geq 1 \mu\text{m}$), the lattice superstructure does not vanish and a smaller trigonal distortion is observed. Our sample belongs to this latter category; therefore, it seems reasonable to start with the high-temperature phase for determination of the room temperature structure. We have established the group—subgroup relationship between the cubic ($Ia3d$) and

most symmetric trigonal phase ($R3c$) space groups compatible with the diffraction pattern and used the lost symmetry elements to generate the atomic positions in the room temperature structure. In this structure, there are 75 independent atomic sites occupied by 22 uranium atoms and 48 oxygen atoms and 4 sites with $1/4$ occupancy for the oxygen atom at the center of the cuboctahedron. This crystalline structure was used to model the R-U₄O₉ neutron diffraction pattern, refining only the lattice parameters, which led to $R_{\text{Bragg}} = 6.5\%$. That provided a good starting point from the point of view of Rietveld analysis. Unfortunately, no satisfactory refinement of the crystalline structure could be performed because of the large number of parameters.

In PDF analysis, two choices are possible for the position of the oxygen atom at the center of the cuboctahedron: either it shares disordered positions or it occupies only one of those possible sites. We chose to order this atom: therefore, the starting crystalline structure in the $R3c$ space group consists of 22 independent uranium atoms and 49 independent oxygen atoms.

In order to reduce the number of parameters to be refined, all uranium atoms were given a common isotropic thermal motion and all oxygen atoms were also given another common isotropic thermal motion. Nevertheless, the number of parameters associated with the atomic positions was still 207. The parameters describing instrumenta-

tion parameters were fixed to values appropriate for the D4 instrument: Q_{broad} , which describes the experimental peak broadening, was set to 0.03, and Q_{damp} , which describes the intensity dampening with increasing r , was set to 0.047. The high Q cutoff Q_{max} was set to 23.5 Å. Moreover, the ripples of the PDF for $r < 1.6$ Å were removed and

RESULTS

A first refinement was performed in which the positions of the 22 uranium and 49 oxygen atoms were refined with uranium and oxygen thermal motion. Refinement was performed up to $\chi^2 = 1.38$ (χ^2 is a statistical indicator that should reach 1 for best refinements), but at that point, it led to too short distances to be physically relevant. One oxygen atom had a 1.4 Å distance with a uranium atom and a 1.7 Å distance with another oxygen atom. This can be explained as a result of displacement of the central oxygen atom of the cuboctahedron, which leaves the central position and repels another oxygen atom of the cuboctahedron. The latter oxygen atom has less space than that in β -U₄O₉ because the other atoms are already closely packed, and this leads to too short distances to the surrounding atoms.

Hence, this crystallographic model based on the $R3c$ space group does not, obviously, correspond well to the actual crystal structure of R-U₄O₉, but it reflects some of its main characteristics because the quality of the fit is rather good with $\chi^2 = 1.38$.

This can be interpreted in at least two different ways.

First, the $R3c$ space group might not be the actual space group of the crystal structure of our sample. Belbeoch et al. proposed the $R3c$ space group for the R-U₄O₉ sample whose unit cell parameters would lead to a splitting of the diffraction lines that we do not observe for our sample.¹⁸ One reason for that could be the existence of correlations between the cuboctahedra that could be fully achieved or frustrated as a function of the sample history. Using the $R3c$ space group implies that displacement of the central oxygen atom is correlated for all cuboctahedra in the crystal. A space group with a lower symmetry or a larger supercell would be necessary to account for an uncorrelated central oxygen atom and could induce sufficient relaxation of the crystalline structure so that short atomic distances could be avoided.

Second, the oxygen content of our sample could be slightly

different from the one expected from β -U₄O₉. U₄O₉ is known to

be a nonstoichiometric compound whose oxygen/uranium ratio can vary from 2.23 to 2.25, with the exact ratio of β -U₄O₉ being 2.234375. This difference in the oxygen content could create different local concentrations, or frustration.

In order to obtain results that can be discussed independently of both the oxygen content and local symmetry, we chose to remove the oxygen atoms with unrealistic short bonds from the description of R-U₄O₉ in the $R3c$ space group. This does not provide the exact crystal structure of R-U₄O₉, but it gives at least an approximate description of frustration-free local areas.

Because of symmetry constraints, six oxygen atoms were removed. Thus, the initial chemical formula U₁₂₈O₂₈₆ of β -U₄O₉ became U₁₂₈O₂₈₀ corresponding to a -2% decrease in the oxygen content. A second refinement was then performed starting from the atomic positions previously refined in which these six oxygen atoms were removed. In this refinement, the previously refined thermal motion parameters were fixed: 0.0016 for u_{U} and 0.005 for u_{O} . Refinement was performed up to $\chi^2 =$

replaced by a straight line, so that the PDF is essentially flat from 0 to 2 Å. With this procedure, we wanted to remove the unphysical ripples at low r that result from Fourier transformation of data having finite Q_{max} .

1.21 The calculated PDF is compared to the measured one in the upper part of Figure 5. A detail plot between 0.5 and 7.5 Å is presented in the lower part of Figure 5.

The refined atomic positions of R-U₄O₉ are slightly displaced from the positions in β-U₄O₉, but the shift of the central atom associated with the suppression of six oxygen atoms leads to an arrangement of the oxygen atoms of the cuboctahedron that has a lot of similarities with the U₃O₈ crystal structure.




Figure 5. Comparison between the fitted PDF (purple line) and experimental PDF (crosses): top, the whole refined pattern; bottom, a zoom for r between 0 and 7.5 Å. The difference between the experimental and calculated points is given by a black line, which was shifted by -4 for better clarity.

The displacement of the oxygen atom located in the center of the cuboctahedron in β - U_4O_9 induces a new organization of the atoms surrounding the cuboctahedron, which is depicted in Figure 6. In this figure, the six nearest uranium atoms are located on two dense planes, derived from UO_2 (111) planes, below and above the central oxygen atom, marked in yellow. Each uranium atom is surrounded by a distorted pentagon made of oxygen atoms more or less parallel to the dense planes. Each uranium atom is also connected to other oxygen atoms, forming $U-O$ bonds perpendicular to the dense planes or linking uranium atoms from the two dense planes. Three $U-O$ bonds, shown in orange, are formed at 1.8 Å; this bond length is usually observed for the UO_6 cation in uranium oxo compounds.¹⁹ This configuration of the six uranium atoms and their coordination polyhedra makes them resemble crumpled sheets taken from a fragment of U_3O_8 . The corresponding fragment of U_3O_8 is represented in Figure 6. In U_3O_8 , oxygen atoms form pentagons in the $a-b$ plane and UO_2 chains along the c axis.

DISCUSSION

The atomic positions obtained by refinement of the PDF give some new insight into the mechanism by which U_4O_9 is transformed into U_3O_8 . The similarity between the U_4O_9




Figure 6. Comparison between the U_3O_8 encapsulated cluster in U_4O_9 (upper part) and the U_3O_8 fragment (lower part), which could be transformed by a shear strain. Uranium atoms are gray, and oxygen atoms are red, are orange for 1.8 Å short $U-O$ bonds, and are yellow for the atom initially at the cuboctahedron center.

crumpled sheets and the U_3O_8 fragments shown in Figure 7 could indicate that transformation from U_4O_9 to U_3O_8 consists of stretching U_4O_9 crumpled sheets, which is consistent with the mechanism proposed by Allen and Holmes.³ Because the sheets derived from U_3O_8 are crumpled in U_4O_9 , they can be considered as mechanically strained by the surrounding atoms, which approximately keep the UO_2 crystallographic positions. Hence, the shear transformation from the fluorite-derived structure to the U_3O_8 -layered structure should be viewed as a competition between frustrated crumpled sheets and the remaining UO_2 lattice shown in Figure 6. Two conditions are required for such a transformation to occur. First, the crumpled sheets must lie in the same plane, so that the shear transformation creates U_3O_8 in a sufficiently large domain. Second, the energetic balance must be in favor of U_3O_8 , meaning that a minimum concentration of crumpled sheets should be reached. These conditions are discussed below for three different cases.

(1) $R-U_4O_9$ cannot undergo a shear transformation because the crumpled sheets are disoriented with respect to each other, at least in our data. Nevertheless, Belbeoch et al.¹⁸ observed $R-U_4O_9$ with a large rhombohedral distortion with a grain size of under 1 μm (the exact value is not known). This size effect could be understood by considering that, below a minimal size, the orientation of the crumpled sheets is not imposed by the bulk




Figure 7. Coordination polyhedra of the remaining uranium atoms around the crumpled sheets. Uranium atoms are gray, and the oxygen atoms that they are linked to are represented as small red dots. The oxygen atom initially at the center of the cuboctahedron is yellow. The coordination polyhedron of uranium atoms are dark blue when they share one face with the cuboctahedron and light blue when they share no face with it.

symmetry but by surface effects. Thus, they could be aligned, leading to a cooperative effect that would stretch the crumpled sheets. This size effect could also explain the different PDFs measured on R-U₄O₉. In our samples, the grain size is more than 1 μm. Garrido prepared his sample by a high-temperature route in a manner similar to what we did. Consistently, his results, cubic R-U₄O₉, and his PDF are very similar to ours and can be associated with grain sizes bigger than 1 μm. Conradson prepared his XAS samples by a high-temperature route, followed by a grinding phase. The grain size of the obtained powder is not given in the paper, but if it was less than 1 μm, in the PDF he obtained a significant number of short U–O bonds that could be related to the heavily distorted lattice, as first shown by Belbeoch et al. The importance of the route used for preparing the samples was also evidenced by Raman spectroscopy data. Manara and Renker²⁰ and He and Shoosmith²¹ both prepared samples of overstoichiometric UO₂ by heat treating UO₂ specimens in an oxidizing atmosphere at high temperature. He and Shoosmith found a 630 cm⁻¹ band in their Raman spectrum, while Manara and Renker did not.

(2) The formation of U₃O₈ involves different crystalline phases as a function of the temperature. At high temperature, U₃O₈ forms directly on UO₂. This can be understood by considering that thermal motion makes UO₂ dense layers loose and that a cuboctahedron induces shear transformation of UO₂ as soon as it is created. At a temperature lower than 400 °C, U₃O₈ appears only after a layer of U₃O₇ is formed. The U₃O₇ crystal structure is a UO₂-derived structure with 16 cuboctahedra per unit cell instead of 12 cuboctahedra per unit cell as in β-U₄O₉. U₃O₇ is a metastable compound stabilized by the mechanical stresses induced by its formation on the UO₂ surface.²² This can be understood by considering the energy balance inside the U₃O₇ layer: more energy is needed to achieve a shear transformation that is hindered by the tensile stress in the layer. This shear transformation can only occur when this tensile stress is relaxed by cracking the U₃O₇ layer.²²

(3) Doping UO₂ with trivalent cations is known to limit U₃O₈ formation. Assuming that doping leads to a random distribution

of trivalent cations, it will hinder the formation of cuboctahedra because it requires six surrounding uranium atoms with 5+ oxidation number. Thus, cuboctahedra cannot be formed at any position, and their cooperative effect for shear transformation is thereby reduced.

CONCLUSION

The neutron PDF for R-U₄O₉ was obtained by Fourier transformation of the neutron diffraction data obtained on the D4 instrument at the ILL. This PDF was refined, and we have proposed a description of the crystal structure of R-U₄O₉ in an area free from mechanical stress and chemical frustration. This description provides new insight into the interpretation of the transformation of U₄O₉ to U₃O₈, which would be driven by the shear transformation of crumpled sheets, associated with cuboctahedra, into regular U₃O₈ sheets.

ASSOCIATED CONTENT

b Supporting Information. X-ray crystallographic data in CIF format. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: Lionel.desgranges@cea.fr

ACKNOWLEDGMENT

The authors are indebted to the program MATAV funded by CEA, which supported this study. We also thank the technical staff and radioprotection personnel of the ILL for their contributions to the neutron diffraction work.

REFERENCES

- (1) Ferry, C.; Poinssot, C.; Cappelaere, C.; Desgranges, L.; Jegou, C.; Miserque, F.; Piron, J. P.; Roudil, D.; Gras, J. M. *J. Nucl. Mater.* 2006, **352**, 246.
- (2) McEachern, R. J.; Taylor, P. J. *Nucl. Mater.* 1998, **254**, 87.
- (3) Allen, G. C.; Holmes, N. R. J. *Nucl. Mater.* 1995, **223**, 231.
- (4) Willis, B. T. M. *J. Phys.* 1964, **25**, 431.
- (5) Loopstra, B. O. *Acta Crystallogr.* 1964, **17**, 651.
- (6) Bevan, D. J. M.; Grey, I. E.; Willis, B. T. M. *J. Solid State Chem.* 1986, **61**, 1.
- (7) Cooper, R. I.; Willis, B. T. M. *Acta Crystallogr.* 2004, **A60**, 322–325.
- (8) Desgranges, L.; Baldinozzi, G.; Rousseau, G.; Niepce, J.-C.; Calvarin, G. *Inorg. Chem.* 2009, **48**, 7585–7592.
- (9) Murray, A. D.; Willis, B. T. M. *J. Solid State Chem.* 1990, **84**, 52–57.
- (10) Geng, H. Y.; Che, Y.; Kaneta, Y.; Kinoshita, M. *Phys. Rev. B* 2008, **77**, 180101.
- (11) Geng, H. Y.; Che, Y.; Kaneta, Y.; Kinoshita, M. *Appl. Phys. Lett.* 2008, **93**, 201903.
- (12) Andersson, D. A.; Lezema, J.; Uberuaga, B. P.; Deo, C.; Conradson, S. D. *Phys. Rev. B* 2009, **79**, 024110.
- (13) Yakub, E.; Ronchi, C.; Staicu, D. *J. Nucl. Mater.* 2009, **389**, 119–126.
- (14) Conradson, S. D.; et al. *Inorg. Chem.* 2004, **43**, 6922–6935.
- (15) Garrido, F.; et al. *Inorg. Chem.* 2006, **45**, 8408–8413.
- (16) Fischer, H. E.; Cuello, G. J.; Palleau, P.; Feltn, D.; Barnes, A. C.; Badyal, Y. S.; Simonson, J. M. *Appl. Phys. A: Mater. Sci. Process.* 2002, **74**, S160–S162.

- (17) Farrow, C. L.; et al. *J. Phys.: Condens. Matter* 2007, 19, 335219.
- (18) Belbeoch, B.; Boivineau, J. C.; Perio, P. *J. Phys. Chem. Solids* 1967, 28, 1267–1275.
- (19) Burns, P. C. . In *Structural chemistry of inorganic actinide compounds*; Krivovichev, S. V., Burns, P. C., Tananaev, I. G., Eds.; Elsevier: New York, 2007.
- (20) Manara, D.; Renker, B. *J. Nucl. Mater.* 2003, 321, 233–237.
- (21) He, H.; Shoesmith, D. *Phys. Chem. Chem. Phys.* 2010, 12, 8108–8117.
- (22) Desgranges, L.; Palanchar, H.; Gamaléri, M.; Micha, J. S.; Optasanu, V.; Raceanu, L.; Montesin, T.; Creton, N. *J. Nucl. Mater.* 2010, 407, 44–47.