



# Diverse origin of P[19] rotaviruses in children with acute diarrhea in Taiwan: detection of novel lineages of the G3, G5, and G9 VP7 genes

Krisztian Banyai

## ► To cite this version:

Krisztian Banyai. Diverse origin of P[19] rotaviruses in children with acute diarrhea in Taiwan: detection of novel lineages of the G3, G5, and G9 VP7 genes. *Journal of Medical Virology*, 2011, 83 (7), pp.1279. 10.1002/jmv.22052 . hal-00640448

**HAL Id: hal-00640448**

**<https://hal.science/hal-00640448>**

Submitted on 12 Nov 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Diverse origin of P[19] rotaviruses in children with acute diarrhea in Taiwan: detection of novel lineages of the G3, G5, and G9 VP7 genes**

|                               |                                                                                                             |
|-------------------------------|-------------------------------------------------------------------------------------------------------------|
| Journal:                      | <i>Journal of Medical Virology</i>                                                                          |
| Manuscript ID:                | JMV-10-2139.R1                                                                                              |
| Wiley - Manuscript type:      | Research Article                                                                                            |
| Date Submitted by the Author: | 04-Jan-2011                                                                                                 |
| Complete List of Authors:     | Banyai, Krisztian; Baranya County Institute of State Public Health Service, Regional Laboratory of Virology |
| Keywords:                     | phylogenetic analysis, VP7, VP4, VP6, NSP4                                                                  |
|                               |                                                                                                             |

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

To *Journal of Medical Virology*

Diverse origin of P[19] rotaviruses in children with acute diarrhea in Taiwan: detection of novel lineages of the G3, G5, and G9 VP7 genes

Fang-Tzy Wu<sup>1,2</sup>, Krisztián Bányai<sup>3</sup>, Jason C. Huang<sup>2</sup>, Ho-Sheng Wu<sup>1,4\*</sup>, Feng-Yee Chang<sup>1</sup>, Jyh-Yuan Yang<sup>1</sup>, Chao Agnes Hsiung<sup>5</sup>, Yhu-Chering Huang<sup>6</sup>, Jen-Shiou Lin<sup>7</sup>, Kao-Pin Hwang<sup>8,9</sup>, Baoming Jiang<sup>10</sup>, and Jon R. Gentsch<sup>10</sup>

<sup>1</sup> Center for Research and Diagnostics, Centers for Disease Control, Department of Health, Taiwan

<sup>2</sup> Department of Biotechnology and Laboratory Science in Medicine, National Yang-Ming University, Taipei, Taiwan

<sup>3</sup> Veterinary Medical Research Institute, Budapest, Hungary

<sup>4</sup> School of Medical Laboratory Science and Biotechnology, Taipei Medical University, Taipei, Taiwan.

<sup>5</sup> Institute of Population Health Sciences, National Health Research Institutes, Zhunan, Taiwan

<sup>6</sup> Division of Pediatric Infectious Disease, Chang Gung Children's Hospital, Chang Gung University College of Medicine, Taoyuan, Taiwan

<sup>7</sup> Division of Pediatric Infectious Disease and Department of Laboratory Medicine, Changhua Christian Hospital, Changhua, Taiwan

<sup>8</sup> Division of Pediatric Infectious Disease, Department of Pediatrics, Chang Gung Memorial Hospital, Kaohsiung Medical Center, Chang Gung University College of Medicine, Kaohsiung, Taiwan

<sup>9</sup> School of Medicine, Children's Hospital, China Medical University & Hospitals, Taichung, Taiwan

<sup>10</sup> Division of Viral Diseases, Centers for Disease Control and Prevention, Atlanta, USA

\*Corresponding author. Tel: 886 2 26531377; Fax: 886 2 27837779. Email:[wuhs@cdc.gov.tw](mailto:wuhs@cdc.gov.tw)

Running head: Novel P[19] rotavirus strains in Taiwanese children

Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the CDC.

For Peer Review

**Abstract**

We previously reported the detection of genotype P[19] rotavirus strains from children hospitalized with acute dehydrating diarrhea during a 5-year surveillance period in Taiwan. The characterization of 5 P[19] strains (0.4% of all typed), including 3 G3P[19], a novel G5P[19] and a unique G9P[19] genotype is described in this study. Phylogenetic analysis of the VP4, VP7, VP6, and NSP4 genes was performed, which demonstrated novel lineages for respective genotypes of the VP4 and the VP7 genes. The sequence similarities of the P[19] VP4 gene among Taiwanese human strains was higher (nt, 91.5%-96.2%; aa, 93.7%-97.6%) than to other P[19] strains (nt, 83.5%-86.6%; aa, 89.4%-94.1%) from different regions of the world. The VP7 gene of the 3 G3P[19] Taiwanese strains shared up to 93.4% nt and 97.5% aa identity to each other but had lower similarity to reference strain sequences available in GenBank (nt, < 90.1%; aa, < 95.6%). Similarly, the VP7 gene of the novel G5P[19] strain was only moderately related to the VP7 gene of reference G5 strains (nt, 82.2%-87.3%; aa, 87.0%-93.1%), while the VP7 gene of the single G9P[19] strain was genetically distinct from other known human and animal G9 rotavirus strains (nt, ≤92.0%; aa, ≤95.7%). Together, these findings suggest that the Taiwanese P[19] strains originated by independent interspecies transmission events. Synchronized surveillance of human and animal rotaviruses in Taiwan should identify possible hosts of these uncommon human rotavirus strains.

**Key words**

phylogenetic analysis, VP7, VP4, VP6, NSP4

**Introduction**

Group A rotaviruses are the leading cause of severe dehydrating gastroenteritis in young children worldwide. Globally, rotavirus infections account for an estimated 611,000 deaths each year among children < 5 years of age [Parashar et al., 2006]. Rotavirus particles are

icosahedral, 70-75 nm in diameter and contain 11 genomic segments of double-stranded RNA. The two outer-layer proteins VP7 and VP4 form the basis of the dual classification system of group A rotavirus into G and P types [Estes and Kapikian, 2007]. Of the 24 G and 31 P genotypes currently known, at least 11 G types and 12 P types, in at least 70 different antigen combinations have been identified in humans [Matthijnssens et al., 2009]. Of these, 5 are globally common (G1P[8], G2P[4], G3P[8], G4P[8], and G9P[8]), while others are unusual combinations of these common antigen types and may be only regionally important. Rare strains include mainly those antigen specificities that originate from periodic occurrence of animal-to-human interspecies transmission events. Examples for such strains include G5, G6, G8, and G11 VP7 genes or P[3], P[7], P[11], and P[19] VP4 genes, as well as rare genetic variants of common antigen types, such as G3 VP7 genes of porcine, feline, or lapine origin or genetically diverse porcine-like P[6] or feline-like P[9] VP4 genes [De Grazia et al., 2010; Bányai et al., 2003, 2009a; 2009b, 2009c; Matthijnssens et al., 2006, 2009; Esona et al., 2009; Martella et al., 2006, 2008; Khamrin et al., 2006; Chitambar et al., 2009; Steyer et al., 2008].

Based on sequence similarities, recently a new classification system has been proposed, which designates independent genotypes for each genome segment [Matthijnssens et al., 2008a, 2008b]. This system is being appreciably adopted by the scientific community and several research groups initiated partial or whole genome based strain characterization studies to extend information on features of circulating rotavirus strains. For example, among the 9 non-serotype genes, the VP6 and NSP4 genes have been studied extensively due to their recognized relative genetic linkage and their potential use in determining the original host species for unusual strains with suspected heterologous origin [De Grazia et al., 2007, 2008, 2009; Chitambar et al., 2009; Bányai et al., 2009; Khamrin et al., 2009; Martella et al., 2008; Li et al., 2008; Araujo et al., 2007; Khamrin et al., 2006, 2007; Ghosh et al., 2007; Iturriza-Gomara et al., 2003].

1  
2  
3 90 In Taiwan, rotavirus strain surveillance started in 2001 [Chen et al., 2005]. Studies  
4  
5 91 have identified the globally common strains, although the predominant antigen types seem to  
6  
7 92 change over time. We have recently reported strain prevalence data over a 3-year period from  
8  
9  
10 93 3 sentinel hospitals located in northern, central and southern Taiwan [Wu et al., 2009]. That  
11  
12 94 study utilized a nucleotide sequence-based genotyping assay, in which type common PCR  
13  
14 95 primers were employed to amplify full-length or partial outer capsid genes. In total 829  
15  
16 96 strains were genotyped. Among these strains we identified G1P[8] strains to be predominant,  
17  
18 97 followed by G3P[8], G9P[8] and G2P[4] strains as additional medically important strains,  
19  
20 98 while 11 further antigen combinations represented minority strains. Unexpectedly, among  
21  
22 99 these minority combinations we detected 2 P[19] strains associated with a G3 and a G9 VP7  
23  
24  
25 100 gene. In subsequent extended surveillance, we detected 3 additional P[19] strains in 2008 and  
26  
27  
28 101 2009 seasons; 2 associated with G3 and 1 with G5 VP7 gene. In an attempt to better  
29  
30 102 understand the origin and relationship of these strains to one another and the few hitherto  
31  
32 103 characterized human and animal P[19] strains, we determined and analyzed the full length or  
33  
34 104 nearly full length VP7 genes, the variable VP8\* fragment of the VP4 gene, and the genotype  
35  
36 105 diversity and potential genetic linkage among the NSP4 and the VP6 genes.  
37  
38  
39  
40  
41  
42

43 **Materials and Methods**

44 *Case definition and data collection*

45  
46 108 From January 2005 to December 2009, a total of 6,346 stool specimens were collected from  
47  
48 109 children less than 5 years old who were hospitalized for treatment of acute gastroenteritis  
49  
50 110 (AGE) in 3 sentinel hospitals located in northern, central and southern Taiwan. AGE was  
51  
52 111 defined as three or more episodes of watery diarrhoea or looser-than-normal stool in the 24 h  
53  
54 112 before presentation.  
55  
56  
57  
58  
59  
60 114

### 115 *Laboratory testing*

116 Stool specimens were collected and screened for rotavirus VP6 antigen using an EIA kit  
117 (RIDASCREEN® Rotavirus, R-Biopharm AG, Germany) at each hospital. Rotavirus EIA-  
118 positive specimens were shipped to Rotavirus Reference Laboratory at Taiwan CDC for G-  
119 and P-genotyping.

### 121 *RT-PCR and Polyacrylamide Gel Electrophoresis*

122 Viral RNA was extracted from 10% (wt/v) fecal supernatants by MagNA Pure LC DNA  
123 isolation kit (Roche Diagnostics GmbH, Germany) according to the manufacturer's  
124 instructions. The extracted RNAs were used as the template for RT-PCR with random primers  
125 [Wu et al., 2009]. Viral VP4, VP6, VP7 and NSP4 genes were amplified with primers sets  
126 Con3/Con2, JRG7/JRG8 or GEN-VP6F/GEN-VP6R, Beg9-End9, and JRG30/JRG31 as  
127 described [Gentsch et al, 1992; Luan et al, 2009; Matthijnsens et al, 2006; Gouvea et al, 1990;  
128 Esona et al; 2009]. Viral RNA samples were also analyzed by a 10% polyacrylamide gel and  
129 silver staining according to the manufacturer's protocol (Bio-Rad, Hercules, CA, USA).

### 131 *Sequencing and phylogenetic analysis*

132 PCR amplicons were directly sequenced with primers sets Con3/Con2, JRG7/JRG8 or GEN-  
133 VP6F/GEN-VP6R, Beg9-End9, and JRG30/JRG31 on a ABI 3130 sequencer (Applied  
134 Biosystems, CA, US). Rotavirus genotypes were determined using the RotaC software [Maes  
135 et al., 2009]. Multiple nucleotide sequences were aligned and the phylogenetic analysis was  
136 performed using MEGA 4.0 software [Tamura et al., 2007].

## 137 **Results**

139 *Genetic analysis of the VP4 genes.* A 831 bp fragment of the VP4 gene (corresponding to nt  
140 34-864, strain Mc345) of 5 Taiwanese P[19] strains was sequenced. Along a shorter stretch

1  
2  
3 141 (750 bp and 250 aa), these strains shared 91.5%-96.2% (nt) and 93.7-97.6% (aa) identities  
4  
5 142 among each other and a lower similarity to 16 other P[19] strains detected in mainland Asia  
6  
7 143 (range, nt, 83.5-86.6%; aa, 89.4-94.1%). In our phylogenetic analysis, both nt (Fig. 1A) and  
8  
9 144 aa (not shown) trees demonstrated that Taiwanese P[19] strains fell into a discrete genetic  
10  
11 145 lineage. An aa alignment of the VP8\* fragment revealed 5 aa substitutions in the VP8\*  
12  
13 146 fragment of the protein, which were unique for the Taiwanese P[19] strains (Fig. 1B).  
14  
15  
16  
17  
18  
19

20 148 *Genetic analysis of the VP7 genes.* As determined by nucleotide sequence based genotyping,  
21  
22 149 the 5 Taiwanese P[19] strains exhibited 3 different VP7 specificities: G3 (03-98s185, 07-  
23  
24 150 94s126, 07-97s684), G5 (04-97s51), and G9 (07-96s1118) (Fig. 2).  
25  
26

27 151 The full or nearly full length coding region was determined for the G3 strains. These 3  
28  
29 152 strains shared 92.8% - 93.4% nt (and 96.3% - 97.5% aa) similarities along a 972 bp long  
30  
31 153 fragment to each other, and a lesser similarity to 183 reference G3 strains (nt, < 90.1%; aa, <  
32  
33 154 95.6%). They are most closely related to porcine isolates (e.g., CP-1). Along the antigenic  
34  
35 155 region these 3 Taiwanese strains shared high similarities, although 1 to 4 aa substitutions were  
36  
37 156 seen (Fig. 3). Phylogenetic analysis found that these 3 strains clustered on a separate branch,  
38  
39 157 distinct from other strains suggesting they represent a new lineage of the G3 VP7 gene (Fig.  
40  
41 158 2).  
42  
43  
44  
45

46 159 Identification of a G5P[19] strain in our sample collection was unexpected, as this  
47  
48 160 antigen combination was not identified previously in humans. Sequence analysis of a 828 bp  
49  
50 161 stretch of the VP7 gene of strain 04-97s51 revealed moderate sequence identity to 98  
51  
52 162 reference G5 strains of both animal and human origin (range: nt, 82.2-87.3%; aa, 87.0-  
53  
54 163 93.1%). Similar divergence was seen in the neutralization epitopes between 04-97s51 and  
55  
56 164 selected G5 strains. Overall, these similarity ranges together with results of phylogenetic  
57  
58  
59  
60

analysis suggested that this Taiwanese G5 strain fell into a new lineage of the G5 VP7 gene (Fig. 2).

A fifth Taiwanese P[19] strain contained a G9 VP7 gene. Because this specificity was seen in both human and porcine strains in South-East Asia, it was interesting to see whether the Taiwanese isolate clustered with these human or animal rotavirus strains. Of note, the 07-96s1118 strain displayed limited sequence similarity with other G9 strains detected in Taiwan or other parts of the world. Overall, this Taiwanese G9 strain shared  $\leq 92.0\%$  nt (and  $\leq 95.7\%$  aa) identity to any known G9 VP7 gene along the full coding region, including both human and porcine strains, and shared particularly lower similarity to the globally emerging variant of G9 VP7 gene (nt,  $< 90\%$ ). The closest relatives were among the unique US OM46-like G9 strains that were detected during 1997-1998 in Omaha (nt, 92%; aa, 95.1%). The neutralization epitopes of this novel Taiwanese G9 strain also suggested extensive divergence from other G9 strains. All these data demonstrated that 07-96s1118 is not only highly divergent from the globally emerging variant of G9 VP7 gene, but also divergent from all other G9 VP7 genes identified in surveillance studies over the last 30 years (Fig. 2, 3).

*Genetic analysis of the VP6 and NSP4 genes.* The full or partial coding regions of the VP6 and NSP4 genes were determined to further characterize the 5 identified Taiwanese P[19] strains. For the VP6 gene, the RotaC [Maes et al., 2009] online genotyping software that was developed to predict rotavirus A genotype specificities identified 2 VP6 genotypes among Taiwanese P[19] strains (Fig. 4). Four strains were classified as genotype I1, a common human VP6 specificity, while 07-96s1118, the unique Taiwanese G9P[19] strain shared only moderate sequence identity to other genotypes. The automated genotyping platform recommended validating the VP6 gene specificity of this strain through the RCWG, whose opinion was that this gene falls within the newly described I12 VP6 genotype. This genotype

1  
2  
3 190 is currently represented by 2 strains, the reference human Bangladeshi G11P[25] strain,  
4  
5 191 KTM368, and the Taiwanese G9P[19] strain, 07-96s1118. Phylogenetic analysis suggested  
6  
7  
8 192 this genotype is intermediate between genotype I1 ( $\leq 85\%$ ) and I5 VP6 ( $\leq 86\%$ ) genotypes and  
9  
10 193 even its representative strains share only marginal nt identities to one another (87.6% nt). The  
11  
12 194 other 4 P[19] strains shared ~90-98% nt identities among each other, and were most closely  
13  
14  
15 195 related to 2 porcine strains (i.e., 04-97s51-like P[19] strains vs. CN86, 97.8-97.9% nt  
16  
17 196 identities and 03-98s185 vs. JL94, 91.1% nt identity).

18  
19  
20 197 RotaC clustered all five Taiwanese P[19] strains into a single E1 NSP4 genotype with  
21  
22 198  $\leq 93.4\%$  identity to reference strains (Fig. 4). The P[19] strains shared 93.0%-96.0% nt  
23  
24 199 similarities among one another. Among the reference strains, the rare human-porcine  
25  
26 200 reassortant strains (e.g., LL4260) were most closely related to these Taiwanese P[19] strains  
27  
28  
29 201 (range: nt, 91.7%-93.4%).

30  
31 202 Three of the 5 Taiwanese P[19] rotavirus strains with adequate amount of RNA were  
32  
33  
34 203 further examined for their RNA profiles by polyacrylamide gel electrophoresis (Fig. 5). The 2  
35  
36 204 strains with G5P[19] and G9P[19] genotypes had long RNA electropherotypes similar to that  
37  
38 205 of the reference Wa strain, whereas one G3P[19] strain, 03-98s185, had a super-short  
39  
40 206 electropherotype as indicated by its rearranged gene segment 11.

41  
42  
43 207

44  
45  
46 208 **Discussion**

47  
48 209 This study reports new findings of unusual P[19] rotavirus strains from our sentinel hospital  
49  
50 210 based-surveillance during 2004 to 2009. Among 1183 strains G and P typed, there were 5  
51  
52 211 isolates (0.4% of total) belonging to genotype P[19]. These isolates were identified from  
53  
54 212 specimens collected from different years, one in 2005 (0.8%), one in 2007 (0.3%), two in  
55  
56 213 2008 (1%) and one in 2009 (0.6%). Thus, these rare strains were circulating among humans or  
57  
58 214 alternatively in an animal host in Taiwan during most of the surveillance period.

There have been a few reports on rotavirus P[19] strain detection among animals and humans since 1990, beginning with the first report of the isolation and characterization of P[19] prototype strain 4F in 1994 [Burke et al., 1994]. Several other studies document detection of the virus in humans from Thailand [Urasawa et al., 1992], India [Varghese et al., 2004] and Vietnam [Nguyen et al., 2008]. In the present study, phylogenetic analysis revealed that the Taiwanese strains were genetically closely related to each other but were distinct from human or animal strains from other countries. For example, in the VP8\* fragment of VP4 there were 5 conserved aa substitutions found in Taiwanese P[19] strains compared to human or animal P[19] strains from other countries. In addition, 4 other aa sites, positions 15, 44, 73 and 254, are more closely related to the prototype 4F strain than human or animal P[19] strains isolated from other Asian countries.

The most prevalent VP7 genotypes in Taiwan were G1 (48.3%), G3 (24.5%), G9 (14.1%), G2 (8.0%) and some rare genotypes like G4, G5 and G8 [Wu et al., 2009]. In most studies globally, G1 represented approximately 50-65% of the strains, followed by G3, G4, and G2 [Gentsch et al., 2005; Santos and Hoshino, 2005]. In addition, an increasing number of rare genotypes have been described recently, many of which have strong genetic relationships with animal rotaviruses [Matthijnssens et al., 2009].

Among the G3 strains, we found 3 with P[19] specificity, one each from 2005, 2008 and 2009. The nucleotide sequences of these 3 strains showed only moderate similarity between them (only 92.8% - 93.4%), and less similarity to 183 reference G3 strains. The most closely related strain was a porcine isolate, CP-1 which clustered on a distinct branch by phylogenetic analysis. Demonstration that there were several amino-acid substitutions in the antigenic sites of these Taiwan G3 strains supports the proposal that they represent a new G3 lineage.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Human G5 rotaviruses were first reported in combination with genotype P[8] as a regionally common strain in Brazil during 2004 [Alfieri et al., 1996]. Unusual human G5P[6] rotaviruses were identified from AGE children under 5 in China and Vietnam [Ahmed et al., 2007; Li et al., 2008] and more recently, a G5P[7] strain new to humans, has been detected in Cameroon [Esona et al., 2009]. However, G5 strains had not previously been reported in Taiwan. In addition, this is the first report on detection and molecular characterization of a G5P[19] strain. Molecular features of the G5 VP7 gene of our Taiwanese strain belonged to a phylogenetic lineage distinct from those represented by other strains.

Unlike the G5P[19] strain, the single Taiwanese G9P[19] strain is not the first representative of this particular antigen combination [Okada et al., 2000; Das et al., 2004; Mukherjee et al., 2010], however, it carried a G9 VP7 gene highly divergent from other currently known G9 VP7 genes, indicating that introduction of this particular gene into the human population in Taiwan was an independent evolutionary event from those events that occurred in other parts of the world during the past 3 decades. In a historical perspective, the DC706-like G9P[8] strains detected exclusively on the eastern coast of the USA represents the first documented evidence for human infection with G9 strains and could have caused a minor local outbreak [Cao et al., 2008]. Independent introduction associated with a different G9 VP7 gene lineage (WI61-like G9P[8]) was seen northeast from the site where DC706-like strains were detected during 1983 and this strain seemed to have some capacity to spread into other areas as evidenced by detections of closely related strains in subsequent years in Japan [Clark et al., 1987; Green et al., 1989]. A novel strain, Mc345-like G9P[19], was identified in Thailand in 1989 and its descendants were detected subsequently in this region [Urasawa et al., 1992; Okada et al., 2000; Das et al., 2004; Mukherjee et al., 2010]. Another novel strain, 116E-like G9P[11], appeared during early 1990s in India in asymptomatic neonates [Das et al., 1993; Gentsch et al., 1993]. The first description of the globally spread variant of G9

strains was also first detected in South Asia and genetically it was most closely related to Mc345-like strains. This variant of G9 strains that subsequently spread worldwide to become the 5<sup>th</sup> most common strain during a 10-year period are additional examples of introductions of G9 VP7 into human populations [Gentsch et al., 2005; Matthijnsens et al., 2009]. Additional emergences seen in central region of the USA during mid-1990s by a divergent G9P[8] strain (OM46-like) [Laird et al., 2003], and other G9 strains (97'SZ37 and T203) with divergent VP7 genes were reported in far-east Asia [Xu et al., GenBank record; Dong et al., 2008]. While to date only the lineage III G9 strains have been proven to possess significant epidemiologic potential, the data on a new G9 VP7 gene variant from Taiwan clearly demonstrates that interspecies transmission events are more frequent and the G9 VP7 gene is genetically more divergent than previously thought based on a limited number of gene sequences.

Overall, our findings demonstrate that the P[19] strains circulating in Taiwan are different from those in other Asian countries. The distinct evolutionary patterns of these strains may be explained by Taiwan's geographic separation from the rest of Asia by the Pacific Ocean and consequently minimal genetic contact with rotavirus strains of suspect animal reservoirs from other countries. Since P[19] strains may have originated in porcine species, it will be important to conduct surveillance and determine whether similar strains are also present among pigs in Taiwan.

## Acknowledgements

This study was financially supported in part by research grant of DOH97-DC-1102 and "National Research Program for Genome Medicine" (supported grant: 94-0324-19-F-01-00-00, 95-0324-19-F01-00-00-00-35, 96-0324-01-F-01) from Centers for Disease Control, Department of Health, Executive Yuan, Taiwan. K. B. was supported by the Hungarian

Research Fund (OTKA, PD76364). We thank Chen-Fu Yang for discussion and Shy-Yuan Liang for assistance in performing data management and molecular analysis, Yuhuan Wang for performing viral RNA analysis, and Jelle Matthijnsens for helpful discussion.

References

Ahmed K, Anh DD, Nakagomi O. 2007. Rotavirus G5P[6] in child with diarrhea, Vietnam. *Emerg Infect Dis* 13:1232-1235.

Alfieri AA, Leite JP, Nakagomi O, Kaga E, Woods PA, Glass RI, Gentsch JR. 1996. Characterization of human rotavirus genotype P[8]G5 from Brazil by probe-hybridization and sequence. *Arch Virol* 141:2353-2364

Araújo IT, Heinemann MB, Mascarenhas JD, Assis RM, Fialho AM, Leite JP. 2007. Molecular analysis of the NSP4 and VP6 genes of rotavirus strains recovered from hospitalized children in Rio de Janeiro, Brazil. *J Med Microbiol* 56:854-859.

Bányai K, Gentsch JR, Glass RI, Szucs G. 2003. Detection of human rotavirus serotype G6 in Hungary. *Epidemiol Infect* 130:107-112.

Bányai K, Bogdán A, Domonkos G, Kisfali P, Molnár P, Tóth A, Melegh B, Martella V, Gentsch JR, Szucs G. 2009a. Genetic diversity and zoonotic potential of human rotavirus strains, 2003-2006, Hungary. *J Med Virol* 81:362-370.

Bányai K, Esona MD, Mijatovic S, Kerin TK, Pedreira C, Mercado J, Balmaseda A, Perez MC, Patel MM, Gentsch JR. 2009b. Zoonotic bovine rotavirus strain in a diarrheic child, Nicaragua. *J Clin Virol* 46:391-393.

Bányai K, Esona MD, Kerin TK, Hull JJ, Mijatovic S, Vascenez N, Torres C, de Filippis AM, Foytich KR, Gentsch JR. 2009c. Molecular characterization of a rare, human-porcine reassortant rotavirus strain, G11P[6], from Ecuador. *Arch Virol* 154:1823-1829.

- 314 Burke B, McCrae MA, Desselberger U. 1994. Sequence analysis of two porcine rotaviruses  
315 differing in growth in vitro and in pathogenicity: distinct VP4 sequences and conservation  
316 of NS53, VP6 and VP7 genes. *J Gen Virol* 75:2205-2212.
- 317 Cao D, Santos N, Jones RW, Tatsumi M, Gentsch JR, Hoshino Y. 2008. The VP7 genes of  
318 two G9 rotaviruses isolated in 1980 from diarrheal stool samples collected in Washington,  
319 DC, are unique molecularly and serotypically. *J Virol* 82:4175-4179.
- 320 Chen KT, Chen PY, Tang RB, Huang YF, Lee PI, Yang JY, Chen HY, Bresee, J,  
321 Hummelman E, Glass R. 2005. Sentinel hospital surveillance for rotavirus diarrhea in  
322 Taiwan, 2001-2003. *J Infect Dis* 192 Suppl 1:S44-S48.
- 323 Chitambar SD, Arora R, Chhabra P. 2009. Molecular characterization of a rare G1P[19]  
324 rotavirus strain from India: evidence of reassortment between human and porcine  
325 rotavirus strains. *J Med Microbiol* 58:1611-1615.
- 326 Clark HF, Hoshino Y, Bell LM, Groff J, Hess G, Bachman P, Offit PA. 1987. Rotavirus  
327 isolate WI61 representing a presumptive new human serotype. *J Clin Microbiol* 25:1757-  
328 1762.
- 329 Das BK, Gentsch JR, Hoshino Y, Ishida S, Nakagomi O, Bhan MK, Kumar R, Glass RI.  
330 1993. Characterization of the G serotype and genogroup of New Delhi newborn rotavirus  
331 strain 116E. *Virology* 197:99-107.
- 332 Das S, Varghese V, Chaudhuri S, Barman P, Kojima K, Dutta P, Bhattacharya SK, Krishnan  
333 T, Kobayashi N, Naik TN. 2004. Genetic variability of human rotavirus strains isolated  
334 from Eastern and Northern India. *J Med Virol* 72:156-161.
- 335 De Grazia S, Martella V, Giammanco GM, Gòmara MI, Ramirez S, Cascio A, Colomba C,  
336 Arista S. 2007. Canine-origin G3P[3] rotavirus strain in child with acute gastroenteritis.  
337 *Emerg Infect Dis* 13:1091-1093.

- 338 De Grazia S, Giammanco GM, Martella V, Ramirez S, Colomba C, Cascio A, Arista S. 2008.  
339 Rare AU-1-like G3P[9] human rotaviruses with a Kun-like NSP4 gene detected in  
340 children with diarrhea in Italy. *J Clin Microbiol* 46:357-360.
- 341 De Grazia S, Martella V, Colomba C, Cascio A, Arista S, Giammanco GM. 2009. Genetic  
342 characterization of G3 rotaviruses detected in Italian children in the years 1993-2005. *J*  
343 *Med Virol* 81:2089-2095.
- 344 De Grazia S, Giammanco GM, Potgieter CA, Matthijnssens J, Banyai K, Platia MA, Colomba  
345 C, Martella V. 2010. Unusual assortment of segments in 2 rare human rotavirus genomes.  
346 *Emerg Infect Dis.* 16:859-862.
- 347 Dong HJ, Qian Y, Zhang Y, Zhao LQ, Zhu RN, Chen DM, Liu LY. 2009. Analysis on VP7  
348 and VP4 genes of human rotavirus G9 identified from children with diarrhea in Beijing,  
349 from 2007 to 2008. *Zhonghua Liu Xing Bing Xue Za Zhi* 30:1179-1183.
- 350 Esona MD, Geyer A, Banyai K, Page N, Aminu M, Armah GE, Hull J, Steele DA, Glass RI,  
351 Gentsch JR. 2009. Novel human rotavirus genotype G5P[7] from child with diarrhea,  
352 Cameroon. *Emerg Infect Dis* 15:83-86.
- 353 Estes MK, Kapikian AZ. 2007. Rotaviruses. In *Fields Virology*, 5th edn, vol. 2, pp. 1917–  
354 1974. Edited by D. M. Knipe, P. M. Howley, D. E. Griffin, R. A. Lamb, M. A. Martin, B.  
355 Roizman & S. E. Straus. Philadelphia: Lippincott Williams & Wilkins/Wolters Kluwer.
- 356 Gouvea V, Glass RI, Woods P, Taniguchi K, Clark HF, Forrester B, Fang ZY. Polymerase  
357 chain reaction amplification and typing of rotavirus nucleic acid from stool specimens.  
358 1990. *J Clin Microbiol* 28: 275-282.
- 359 Gentsch JR, Glass RI, Woods P, Gouvea V, Gorziglia M, Flores J, Das BK, Bhan MK.  
360 Identification of group A rotavirus gene 4 types by polymerase chain reaction. 1992. *J*  
361 *Clin Microbiol* 30: 1365-1373.

- 362 Gentsch JR, Das BK, Jiang B, Bhan MK, Glass RI. 1993. Similarity of the VP4 protein of  
363 human rotavirus strain 116E to that of the bovine B223 strain. *Virology* 194:424-430.
- 364 Gentsch JR, Laird AR, Bielfelt B, Griffin DD, Banyai K, Ramachandran M, Jain V, Cunliffe  
365 NA, Nakagomi O, Kirkwood CD, Fischer TK, Parashar UD, Bresee JS, Jiang B, Glass RI.  
366 2005. Serotype diversity and reassortment between human and animal rotavirus strains:  
367 implications for rotavirus vaccine programs. *J Infect Dis* 192 Suppl 1:S146-S159.
- 368 Ghosh S, Varghese V, Samajdar S, Bhattacharya SK, Kobayashi N, Naik TN. 2007. Evidence  
369 for independent segregation of the VP6- and NSP4- encoding genes in porcine group A  
370 rotavirus G6P[13] strains. *Arch Virol* 152:423-429.
- 371 Green KY, Hoshino Y, Ikegami N. 1989. Sequence analysis of the gene encoding the  
372 serotype-specific glycoprotein (VP7) of two new human rotavirus serotypes. *Virology*  
373 168:429-433.
- 374 Khamrin P, Maneekarn N, Peerakome S, Yagyu F, Okitsu S, Ushijima H. 2006. Molecular  
375 characterization of a rare G3P[3] human rotavirus reassortant strain reveals evidence for  
376 multiple human-animal interspecies transmissions. *J Med Virol* 78:986-994.
- 377 Khamrin P, Maneekarn N, Peerakome S, Tonusin S, Phan TG, Okitsu S, Ushijima H. 2007.  
378 Molecular characterization of rare G3P[9] rotavirus strains isolated from children  
379 hospitalized with acute gastroenteritis. *J Med Virol* 79:843-851.
- 380 Khamrin P, Maneekarn N, Peerakome S, Malasao R, Thongprachum A, Chan-It W,  
381 Mizuguchi M, Okitsu S, Ushijima H. 2009. Molecular characterization of VP4, VP6, VP7,  
382 NSP4, and NSP5/6 genes identifies an unusual G3P[10] human rotavirus strain. *J Med*  
383 *Virol* 81:176-182.
- 384 Laird AR, Gentsch JR, Nakagomi T, Nakagomi O, Glass RI. 2003. Characterization of  
385 serotype G9 rotavirus strains isolated in the United States and India from 1993 to 2001. *J*  
386 *Clin Microbiol* 41:3100-3111.

- Li DD, Duan ZJ, Zhang Q, Liu N, Xie ZP, Jiang B, Steele D, Jiang X, Wang ZS, Fang ZY. 2008. Molecular characterization of unusual human G5P[6] rotaviruses identified in China. *J Clin Virol* 42:141-148.
- Luan le T, Trang NV, Phuong NM, Nguyen HT, Ngo HT, Nguyen HT, Tran HB, Dang HN, Dang AD, Gentsch JR, Wang Y, Esona MD, Glass RI, Steele AD, Kilgore PE, Nguyen MV, Jiang B, Nguyen HD. Development and characterization of candidate rotavirus vaccine strains derived from children with diarrhoea in Vietnam. 2009. *Vaccine* 27S: F130-138.
- Maes P, Matthijnssens J, Rahman M, Van Ranst M. 2009. RotaC: a web-based tool for the complete genome classification of group A rotaviruses. *BMC Microbiol* 9:238.
- Martella V, Bányai K, Ciarlet M, Iturriza-Gómara M, Lorusso E, De Grazia S, Arista S, Decaro N, Elia G, Cavalli A, Corrente M, Lavazza A, Baselga R, Buonavoglia C. 2006. Relationships among porcine and human P[6] rotaviruses: evidence that the different human P[6] lineages have originated from multiple interspecies transmission events. *Virology* 344:509-519.
- Martella V, Colombrita D, Lorusso E, Draghin E, Fiorentini S, De Grazia S, Bányai K, Ciarlet M, Caruso A, Buonavoglia C. 2008. Detection of a porcine-like rotavirus in a child with enteritis in Italy. *J Clin Microbiol* 46:3501-3507.
- Matthijnssens J, Rahman M, Martella V, Xuelei Y, De Vos S, De Leener K, Ciarlet M, Buonavoglia C, Van Ranst M. 2006. Full genomic analysis of human rotavirus strain B4106 and lapine rotavirus strain 30/96 provides evidence for interspecies transmission. *J Virol* 80:3801-3810.
- Matthijnssens J, Ciarlet M, Rahman M, Attoui H, Bányai K, Estes MK, Gentsch JR, Iturriza-Gómara M, Kirkwood CD, Martella V, Mertens PP, Nakagomi O, Patton JT, Ruggeri FM, Saif LJ, Santos N, Steyer A, Taniguchi K, Desselberger U, Van Ranst M. 2008a.

- 412 Recommendations for the classification of group A rotaviruses using all 11 genomic  
413 RNA segments. *Arch Virol* 153:1621-1629.
- 414 Matthijnssens J, Ciarlet M, Heiman E, Arijs I, Delbeke T, McDonald SM, Palombo EA,  
415 Iturriza-Gómara M, Maes P, Patton JT, Rahman M, Van Ranst M. Full genome-based  
416 classification of rotaviruses reveals a common origin between human Wa-Like and  
417 porcine rotavirus strains and human DS-1-like and bovine rotavirus strains. 2008b. *J Virol*  
418 82:3204-3219.
- 419 Matthijnssens J, Bilcke J, Ciarlet M, Martella V, Bányai K, Rahman M, Zeller M, Beutels P,  
420 Van Damme P, Van Ranst M. 2009a. Rotavirus disease and vaccination: impact on  
421 genotype diversity. *Future Microbiol* 4:1303-1316.
- 422 Matthijnssens J, Potgieter CA, Ciarlet M, Parreño V, Martella V, Bányai K, Garaicoechea L,  
423 Palombo EA, Novo L, Zeller M, Arista S, Gerna G, Rahman M, Van Ranst M. 2009b. Are  
424 human P[14] rotavirus strains the result of interspecies transmissions from sheep or other  
425 ungulates that belong to the mammalian order Artiodactyla? *J Virol* 83:2917-2929.
- 426 Mukherjee A, Chattopadhyay S, Bagchi P, Dutta D, Singh NB, Arora R, Parashar UD,  
427 Gentsch JR, Chawla-Sarkar M. 2010. Surveillance and molecular characterization of  
428 rotavirus strains circulating in Manipur, north-eastern India: increasing prevalence of  
429 emerging G12 strains. *Infect Genet Evol* 10:311-320.
- 430 Nguyen TA, Hoang LP, Pham LD, Hoang KT, Okitsu S, Mizuguchi M, Ushijima H. 2008.  
431 Use of sequence analysis of the VP4 gene to classify recent Vietnamese rotavirus isolates.  
432 *Clin Microbiol Infect* 14:235-241.
- 433 Okada J, Urasawa T, Kobayashi N, Taniguchi K, Hasegawa A, Mise K, Urasawa S. 2000.  
434 New P serotype of group A human rotavirus closely related to that of a porcine rotavirus.  
435 *J Med Virol* 60:63-69.

- 1  
2  
3 436 Parashar UD, Gibson CJ, Breese JS, Glass RI. 2006. Rotavirus and severe childhood diarrhea.  
4  
5 437 *Emerg Infect Dis* 12:304-306.  
6  
7  
8 438 Santos N, Hoshino Y. 2005. Global distribution of rotavirus serotypes/genotypes and its  
9  
10 439 implication for the development and implementation of an effective rotavirus vaccine.  
11  
12 440 *Rev Med Virol* 15:29-56.  
13  
14  
15 441 Steyer A, Poljsak-Prijatelj M, Barlic-Maganja D, Marin J. 2008. Human, porcine and bovine  
16  
17 442 rotaviruses in Slovenia: evidence of interspecies transmission and genome reassortment. *J*  
18  
19 443 *Gen Virol* 89:1690-1698.  
20  
21  
22 444 Tamura K, Dudley J, Nei M, Kumar S. 2007. MEGA4: Molecular Evolutionary Genetics  
23  
24 445 Analysis (MEGA) software version 4.0. *Mol Biol Evol* 24:1596-1599.  
25  
26  
27 446 Urasawa S, Hasegawa A, Urasawa T, Taniguchi K, Wakasugi F, Suzuki H, Inouye S,  
28  
29 447 Pongprot B, Supawadee J, Suprasert S, et al. 1992. Antigenic and genetic analyses of  
30  
31 448 human rotaviruses in Chiang Mai, Thailand: evidence for a close relationship between  
32  
33 449 human and animal rotaviruses. *J Infect Dis* 166:227-234.  
34  
35  
36 450 Varghese V, Das S, Singh NB, Kojima K, Bhattacharya SK, Krishnan T, Kobayashi N, Naik  
37  
38 451 TN. 2004. Molecular characterization of a human rotavirus reveals porcine characteristics  
39  
40 452 in most of the genes including VP6 and NSP4. *Arch Virol* 149:155-172.  
41  
42  
43 453 Wu FT, Liang SY, Tsao KC, Huang CG, Lin CY, Lin JS, Su CY, Eng HL, Yang JY, Chen PJ,  
44  
45 454 Yang CF. 2009. Hospital-based surveillance and molecular epidemiology of rotavirus  
46  
47 455 infection in Taiwan, 2005-2007. *Vaccine* 27 Suppl 5:F50-F54.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure legends**

Fig. 1. A – Nucleotide sequence based phylogenetic trees of the P[19] VP4 gene. The arrows indicate the Taiwanese G3P[19], G5P[19], and G9P[19] strains. Bootstrap values >90% are indicated. Scale bars are proportional to the phylogenetic distance. Proposed lineage designation of the P[19] VP4 gene is indicated with I, II, and III, respectively. B – Amino acid alignment of the VP8\* domain of VP4 gene. Lineage specific substitutions characteristic to Taiwanese strains are highlighted. Identical residues are shown with dashed line.

Fig. 2. Nucleotide sequence based phylogenetic trees of the G3, G5 and G9 VP7 genes. The arrows indicate the Taiwanese G3P[19], G5P[19], and G9P[19] strains. Strain designated 03-98sP50 is a G5P[6] Taiwanese strain. Bootstrap values >90% are indicated. Scale bars are proportional to the phylogenetic distance.

Fig. 3. Primary structure of the predicted VP7 antigenic sites A, B, C, and F of Taiwanese and reference G3, G5, and G9 rotavirus strains. Dashed line indicates amino acid residue identities in the alignment.

Fig. 4. Nucleotide sequence based phylogenetic trees of the VP6 and the NSP4 genes of Taiwanese P[19] strains (the arrows indicate these strains). Bootstrap values >80% are indicated. Scale bars are proportional to the phylogenetic distance.

Fig. 5. RNA profiles of P[19] rotavirus strains from children with diarrhea in Taiwan. Viral RNA was analyzed by polyacrylamide gel electrophoresis and silver staining. Lanes 1-5,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

480 strains Wa, DS-1. 04-97s51, 07-96s1118, and 03-98s185, respectively. Segments of the  
481 genome from the reference strain Wa are shown on the left.

For Peer Review

Fig. 1A.





Fig. 1B.







Fig. 3.


| ANTIGENIC REGION: | A (87-101)      | B (143-152) | C (208-221)    | F (235-245) |
|-------------------|-----------------|-------------|----------------|-------------|
| G3 VP7            |                 |             |                |             |
| 07-94s126         | AEAAREINDNSWKDT | KYDATLQLDM  | LTDTNTFFEEVATA | HKLSVTNTCT  |
| 07-97s684         | T-----          | ---A---     | -----          | -----MD---  |
| 03-98s185         | -----           | ---A---     | -----          | -----       |
| CMP039            | T---TQ-----     | ---A---     | -----          | -----       |
| CP-1              | T---TQ-----     | -----       | -----          | -----       |
| T108              | T---T---D----   | -----       | -----          | ---N-----   |
| P                 | T---T-----      | -----       | -----          | ---N-----   |
| Cat2              | T---T-----      | -----       | ---S-----      | ---N-----   |
| A131              | T---T-----      | -----       | -----I-N       | -----       |
| RRV               | T---T-----      | -----       | ---A-----      | ---D---A--- |
| J63               | T---T-----      | ---V-----   | -A--AT-----    | ---D---T--- |
| Erv105            | T---T-----      | ---V-----   | ---T-----      | ---D---A--- |
| JE29              | T---VA-----     | ---NE-----  | ---T-----      | ---D---S--- |
| HCR3A             | T---T-----      | ---A-----   | ---VS-----T    | ---D---T--- |
| 30/96             | T---T-----      | ---A-----   | ---NAA-----    | ---D---A--- |
| G5 VP7            |                 |             |                |             |
| 04-97s51          | NEAATEIADNKWTET | KYDANLQLDM  | TTTDINSFEVVANA | HKLDVTNTCT  |
| C134              | -----Q-----     | -----       | L---T---T---S- | -----       |
| LL4260            | -----           | -----       | L---T---I---   | -----S---   |
| A34               | -----T-----     | ---G-----   | S-----T---     | -----S---   |
| KJ75              | -----T-----     | ---G-----   | S---D---T---   | -----S---   |
| H-1               | -----T-----     | ---G-----   | S-----T--ST    | -----S---   |
| 6784/2000/ARN     | -----S-----     | -----       | S---TS---T---  | -----       |
| 10/07/Ire         | -----           | -----       | S---T---T---   | -----       |
| 2F/05/Ire         | -----T-----     | -----       | S---T---M---   | -----       |
| IAL28             | -----D---D-     | ---I-----   | L---TS---T--ST | -----S---   |
| 344/04-1          | -----T-----     | ---G-----   | S---K---T---   | -----       |
| A46               | -----T-----     | ---G-----   | S-----I---     | -----       |
| 134/04-15         | -----A-----     | ---E-----   | L-----T---     | -----       |
| G9 VP7            |                 |             |                |             |
| 07-96s1118        | TEASTQIGDTEWKNT | KHDSLELDM   | TTTNPATFEEVAKN | HKLDVTSSCT  |
| 99-8F1542         | -----           | T-----      | ---A-----TS    | -----NT--   |
| 978237            | -----D-         | -Y-----     | -----AS        | -----NT--   |
| CMH274            | -----D-         | -Y-----     | I---T-----TS   | -----NT--   |
| 116E              | I-----G-        | -YN-----    | ---T-----T-    | -----NT--   |
| F45               | A-----D-        | -Y-----     | ---T-----AS    | -----TT--   |
| OM46              | -----D-         | -Y-----     | ---T-----AS    | -----NT--   |
| G2275             | -----D-         | -Y-----     | ---T-----AS    | -----NT--   |
| JP35-7            | -----P-----     | -Y-----     | ---T-----TG    | -----NT--   |
| A2                | -----           | -Y-----     | I---K-----T-   | -----NT--   |
| US1206            | -----D-         | -Y-----     | I---T-----TS   | -----NT--   |

Fig. 4.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



VP6 gene



NSP4 gene

Fig. 5.

