



HAL
open science

On the Finsler structure of the Teichmüller metric and Thurston's asymmetric metric

Athanase Papadopoulos, Weixu Su

► **To cite this version:**

Athanase Papadopoulos, Weixu Su. On the Finsler structure of the Teichmüller metric and Thurston's asymmetric metric. *Expositiones Mathematicae*, 2015, 33, pp.30-47. 10.1016/j.exmath.2013.12.007 . hal-00641949

HAL Id: hal-00641949

<https://hal.science/hal-00641949>

Submitted on 17 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE FINSLER STRUCTURE OF THE TEICHMÜLLER METRIC AND THURSTON'S ASYMMETRIC METRIC

A. PAPADOPOULOS AND W. SU

ABSTRACT. We highlight several analogies between the Finsler (infinitesimal) properties of Teichmüller's metric and Thurston's asymmetric metric on Teichmüller space. Thurston defined his asymmetric metric in analogy with Teichmüller's metric, as a solution to an extremal problem, which consists, in the case of the asymmetric metric, of finding the best Lipschitz maps in the homotopy class of homeomorphisms between two hyperbolic surface. (In the Teichmüller metric case, one searches for the best quasiconformal map between two conformal surfaces.) It turns out also that some properties of Thurston's asymmetric metric can be used to get new insight into Teichmüller's metric. In this direction, in analogy with Thurston's formula for the Finsler norm of a vector for the asymmetric metric that uses the hyperbolic length function, we give a new formula for the Finsler norm of a vector for the Teichmüller metric that uses the extremal length function. We also describe an embedding of projective measured foliation space in the cotangent space to Teichmüller space whose image is the boundary of the dual of the unit ball in the tangent space representing vectors of norm one for the Finsler structure associated to the Teichmüller metric.

AMS Mathematics Subject Classification: 32G15; 30F60; 57M50; 57N05.

Keywords: Teichmüller space; Thurston's asymmetric metric; Teichmüller metric; Finsler norm.

1. INTRODUCTION

Let $S = S_{g,n}$ be an oriented surface of genus $g \geq 0$ with $n \geq 0$ punctures. We assume that $3g - 3 + n > 0$. The Teichmüller space $\mathcal{T}(S)$ of S is the space of complex structures (or, equivalently, of complete and finite-area hyperbolic structures) on S up to equivalence, where two such structures X and Y are considered as equivalent if there is a conformal map (respectively, an isometry) $h : (S, X) \rightarrow (S, Y)$ which is homotopic to the identity map of S .

There are several natural metrics on $\mathcal{T}(S)$; some of them are Riemannian and others are only Finsler. Some metrics on Teichmüller space (e.g. the Weil-Petersson metric, and the Teichmüller metric) are strongly related to the complex structure of that space. A Finsler metric is a length metrics where the distance between two points is defined by minimizing lengths of peicewise C^1 paths joining them and where the length of a path is computed by integrating norms of tangent vectors. The fact that a metric is Finsler but not Riemannian means that the norm function on each tangent space is not associated to a scalar product. In this paper, we consider the Teichmüller metric and Thurston's asymmetric metric. Both metrics are Finsler, and in the case of Thurston's asymmetric metric, the Finsler norm not only is non-Riemannian but it is even not symmetric. These metrics are defined in terms of distances between geometric structures on the surface (conformal structures, in the case of the Techmüller metric, and hyperbolic structures, in the case of Thurston's asymmetric metric). The question of comparing the various metrics

Date: November 17, 2011.

on $\mathcal{T}(S)$ is a natural one, and it was also suggested by Thurston in [32]; see also [23].

We shall recall below the definition of Thurston's asymmetric metric. Thurston formulated several ideas concerning this asymmetric metric, and we shall review some of them below. He proved some of the major results and he outlined some other results in his paper [32]. After Thurston's paper was circulated, questions on the limiting behavior of stretch lines and of anti-stretch lines were considered, see [20], [23], [30] and [31]. Stretch lines are some special geodesics of that metric, and anti-stretch lines are stretch lines traversed in the opposite directions; we note that since the metric is not symmetric, a geodesic traversed in the opposite direction is not necessarily a geodesic. There were also generalizations of Thurston's asymmetric metric to the case of surfaces with boundary, see [14] and [15]. A renewal of interest in this metric has emerged recently, see e.g. the papers [33], [13] and [18]. It became also clear that among the known metrics on Teichmüller space, the Thurston metric is the one that has an interesting analogue on outer space, see [5]. Thus, there are very good reasons to study Thurston's asymmetric metric. Some analogies and some differences between Thurston's asymmetric metric and Teichmüller's metric have been described in the paper [23]. In the present paper, we point out new analogies that concern the Finsler character of these metrics.

Given a set X , a function $d : X \rightarrow X$ is said to be a *weak metric* on X if it satisfies all the axioms of a distance function except the symmetry axiom. The weak metric d is said to be *asymmetric* if it is strictly weak, that is if there exist two points x and y in X such that $d(x, y) \neq d(y, x)$.

A related notion is that of a *weak norm* on a vector space E . This is a function $p : E \rightarrow \mathbb{R}$ that is nonnegative, convex and positively homogeneous. In other words, p satisfies the following:

- (1) $p(x) \geq 0$ for all $x \in E$;
- (2) $p(\lambda x) = \lambda p(x)$ for all $x \in E$ and for all $\lambda \geq 0$;
- (3) $p(x_1 + x_2) \leq p(x_1) + p(x_2)$ for all $x_1, x_2 \in E$.

In this paper, to simplify terminology, we shall use in general the term *norm* to denote a weak norm and the terms *metric* to denote a weak metric.

Let X and Y be now two complete hyperbolic metrics on S . In the paper [32], Thurston defined an asymmetric metric d_L on $\mathcal{T}(S)$ by setting

$$(1) \quad d_L(X, Y) = \inf_f \log L_f(X, Y),$$

where the infimum is taken over all homeomorphisms $f : X \rightarrow Y$ homotopic to the identity map of S , and where $L_f(X, Y)$ is the Lipschitz constant of f , that is,

$$\text{Lip}(f) = \sup_{x \neq y \in S} \frac{d_Y(f(x), f(y))}{d_X(x, y)}.$$

We shall call this weak metric Thurston's asymmetric metric or, for short, Thurston's metric.

In the same paper, Thurston proved that there is a (non-necessarily unique) extremal Lipschitz homeomorphism that realizes the infimum in (1). He also proved that we have the following formula for the asymmetric metric:

$$(2) \quad d_L(X, Y) = \log \sup_{\gamma} \frac{\ell_Y(\gamma)}{\ell_X(\gamma)},$$

where $\ell_X(\gamma)$ denotes the hyperbolic length of γ with respect to the metric X and γ ranges over all essential simple closed curves on S .

Furthermore, Thurston proved that the asymmetric metric defined in (1) is Finsler, that is, it is a length metric which is defined by integrating a weak norm

on the tangent bundle of $\mathcal{T}(S)$ along paths in Teichmüller space, and taking the minimum lengths over all peicewise C^1 -paths. Thurston gave an explicit formula for the weak norm of a tangent vector V at a point X in $\mathcal{T}(S)$, namely,

$$(3) \quad \|V\|_L = \sup_{\lambda \in \mathcal{ML}} \frac{d\ell_\lambda(V)}{\ell_\lambda(X)}.$$

Here, $\mathcal{ML} = \mathcal{ML}(S)$ is the space of measured laminations on S , $\ell_\lambda : \mathcal{T}(S) \rightarrow \mathbb{R}$ is the length function on Teichmüller space associated to the measured lamination λ and $d\ell_\lambda$ is the differential of the function $\ell_\lambda(X)$ at the point $X \in \mathcal{T}(S)$.

For X in $\mathcal{T}(S)$ and λ in \mathcal{ML} , we shall use the notation $\ell_\lambda(X)$ or $\ell_X(\lambda)$ to denote the X -length of λ , depending on whether we consider the length function as a function on Teichmüller space or on measured lamination space.

We shall present an analogue of Formula (3) for the Teichmüller metric that is expressed in terms of extremal length, namely, we show that the Finsler weak norm associated to the Teichmüller distance is given by the following formula:

$$(4) \quad \|V\|_E = \sup_{\lambda \in \mathcal{MF}} \frac{d\text{Ext}_\lambda^{1/2}(V)}{\text{Ext}_\lambda^{1/2}(X)}.$$

Here, X is a conformal structure on S , considered as a point in Teichmüller space, V is a tangent vector at the point X , $\mathcal{MF} = \mathcal{MF}(S)$ is the space of measured laminations on S , $\text{Ext}_\lambda : \mathcal{T} \rightarrow \mathbb{R}$ is the extremal length function associated to the measured foliation λ and $d\text{Ext}_\lambda$ is the differential of that function at X .

There is a more geometric version of (4) in which the tangent vector V is interpreted as a Beltrami differential (see Corollary 3.7 below).

The following table summarizes some analogies between notions and results associated to Thurston's asymmetric metric and those associated to the Teichmüller metric.

	Thurston's metric	Teichmüller metric
(i)	Stretch maps	Teichmüller extremal maps
(ii)	Stretch lines	Teichmüller lines
(iii)	$d_L(X, Y) = \log \inf_f L(f)$	$d_T(X, Y) = \log \inf_f K(f)$
(iv)	$d_L(X, Y) = \log \sup_\gamma \frac{\ell_\gamma(Y)}{\ell_\gamma(X)}$	$d_T(X, Y) = \log \sup_\gamma \frac{\text{Ext}_\gamma^{1/2}(Y)}{\text{Ext}_\gamma^{1/2}(X)}$
(v)	$\ V\ _L = \sup_{\lambda \in \mathcal{ML}} \frac{d\ell_\lambda(V)}{\ell_\lambda(X)}$	$\ V\ _T = \sup_{\lambda \in \mathcal{MF}} \frac{d\text{Ext}_\lambda^{1/2}(V)}{\text{Ext}_\lambda^{1/2}(X)}$
(vi)	Thurston cataclysm coordinates $X \in \mathcal{T}(S) \mapsto F_\mu(X) \in \mathcal{MF}(\mu)$	The homeomorphism $X \in \mathcal{T}(S) \mapsto F_h(\Phi_F(X)) \in \mathcal{MF}(F)$
(vii)	$\ell_\lambda(X) = i(\lambda, F_{\lambda^*}(X))$	$\text{Ext}_\lambda(X) = i(\lambda, F_h(\Phi_\lambda(X)))$
(viii)	$d\ell_\gamma(\mu) = \frac{2}{\pi} \text{Re} \langle \Theta_\alpha, \mu \rangle$	$d\text{Ext}_\lambda(\mu) = -2 \text{Re} \langle \Phi_\lambda, \mu \rangle$
(ix)	The horofunction boundary is Thurston's boundary	The horofunction boundary is Gardiner-Masur's boundary
(x)	$\mathcal{L}_X(\lambda) = \frac{\ell_X(\lambda)}{L_X}$ for $\lambda \in \mathcal{ML}$	$\mathcal{E}_X(\lambda) = \frac{\text{Ext}_X^{1/2}(\lambda)}{K_X^{1/2}}$ for $\lambda \in \mathcal{ML}$

Most of the entries in this table are well know, some of them are known but need explanation, and some of the them are new and proved in this paper. We now make a few comments on all the entries.

(i) Stretch maps arise from the extremal problem of finding the best Lipschitz constant of maps homotopic to the identity between two hyperbolic structures on a surface S , in much the same way as Teichmüller maps arise from the extremal problem of finding the best quasiconformal constant of maps homotopic to the identity between two complex structures on S .

(ii) Stretch lines are geodesics for Thurston's metric. A stretch line is determined by a pair (μ, F) where μ is a complete lamination (not necessarily measured) and F a measured foliation transverse to μ . Thurston proved that any two points in Teichmüller space can be joined by a geodesic which is a concatenation of stretch lines, but in general such a geodesic is not unique. Furthermore, there exist geodesics for Thurston's metric that are not concatenations of stretch lines. This contrasts with Teichmüller's theorem establishing that existence and uniqueness of geodesics joining any two distinct points, cf. [28], [29] and Ahlfor's survey [3].

(iii) The left hand side is Thurston's definition of Thurston's metric, the infimum is over all homeomorphisms f homotopic to the identity and $L(f)$ is the Lipschitz constant of such a homeomorphism. The right hand side is the definition of the Teichmüller metric, the infimum is over all quasiconformal homeomorphisms f homotopic to the identity and $K(f)$ is the dilatation of such a homeomorphism.

(iv) The left hand side is another expression (also due to Thurston) of Thurston's metric, and the right hand side is Kerckhoff's formula for the Teichmüller metric.

(v) Here, V is a tangent vector to Teichmüller space at a point X . The left hand side formula is due to Thurston [32]; it is the infinitesimal form of Thurston's metric. The right hand side is an infinitesimal form of the Teichmüller metric, and it is proved below (Theorem 3.6).

(vi) On the left hand side, μ is a complete geodesic lamination and the range of the map, $\mathcal{MF}(\mu)$, is the subspace of \mathcal{MF} of equivalence classes of measured foliations that are transverse to μ . On the right hand side, F is a measured foliation, and the range of the map, $\mathcal{MF}(F)$, is the subspace of \mathcal{MF} consisting of equivalence classes of measured foliations that are transverse to F . The left hand side map is a homeomorphism defined in Thurston's paper [32]. The right hand side map is a homeomorphism that arises from the fact that a pair F_1, F_2 of measured foliations determines a unique point $X \in \mathcal{T}(S)$ and a unique quadratic differential Φ on X such that F_1 and F_2 are measure equivalent to the vertical and horizontal foliations of Φ . Here, $F_h(\Phi_F(X))$ is the (equivalence class of the) horizontal foliation of the quadratic differential on the Riemann surface X having F as vertical foliation.

(vii) The left hand side is an expression of the length of a complete lamination λ as the geometric intersection with the horocyclic foliation associated to a completion λ^* of λ . The formula is proved in [20] for the case where λ is complete. The case where λ is not complete follows easily from the geometric arguments used in that proof. The right hand side formula is due to Kerckhoff, see Lemma 3.3 below.

(viii) In the left hand side formula, μ is a Beltrami differential. The formula was given by Gardiner [7], and in this formula α is (the homotopy class of) a simple closed geodesic and Θ_α is the Poincaré series of α , that is, a quadratic differential on the hyperbolic surface X defined by

$$\Theta_\alpha = \sum_{B \in \langle A \rangle \setminus \Gamma} B^* \left(\frac{dz}{z} \right)^2$$

where $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a Fuchsian group associated to X acting on the upper half-plane \mathbb{H}^2 and $A(z) = e^{\ell_\alpha} z$ is the deck transformation corresponding to the simple closed geodesic α . The right hand side formula is called Gardiner's extremal length variational formula (see Lemma 3.2 below).

(ix) Walsh showed in [33] that the horofunction boundary of Thurston's metric is canonically identified with Thurston's boundary. Liu and Su showed in [17] that the horofunction boundary of the Teichmüller metric is canonically identified with Gardiner-Masur's boundary.

(x) Here, we choose a basepoint $X_0 \in \mathcal{T}(S)$ and we fix a complete geodesic lamination μ on S . For each $X \in \mathcal{T}(S)$, L_X is the Lipschitz constant of the extremal Lipschitz map between X_0 and X and K_X is the quasiconformal dilatation of the extremal quasiconformal map between X_0 and X .

We illustrate the use of the functions \mathcal{L}_X and \mathcal{E}_X . We use Thurston's homeomorphism $\phi_\mu : \mathcal{T}(S) \rightarrow MF(\mu)$ that maps each $X \in \mathcal{T}(S)$ to the horocyclic foliation $F_\mu(X)$. The definition of this homeomorphism is recalled in §2 below. We denote the projective class of a measured foliation F by $[F]$. By a result in [20], a sequence $(X_k)_{k \geq 1}$ in $\mathcal{T}(S)$ converges to a limit $[F] \in \mathcal{PMF}$ if and only if $F_\mu(X_k)$ tends to infinity and $[F_\mu(X_k)]$ converges to $[F]$. We prove the following:

Proposition 1.1. *If a sequence $(X_k)_{k \geq 1}$ in $\mathcal{T}(S)$ converges to a limit $[F] \in \mathcal{PMF}$, then $\mathcal{L}_{X_k}(\cdot)$ converges to $[F]$ in the following sense:*

(\star) *up to a subsequence, $\mathcal{L}_{X_k}(\cdot)$ converges to a positive multiple of $i(F, \cdot)$ uniformly on any compact subset of \mathcal{ML} .*

Proof. Assume that a sequence $(X_k)_{k \geq 1}$ converges to $[F] \in \mathcal{PMF}$. Let $L_{X_k} = L_k$ and $F_\mu(X_k) = F_k$. From Thurston's theory [6], there exists a sequence $(c_k)_{k \geq 1}$ of positive numbers such that $c_k \rightarrow 0$ and

$$(5) \quad c_k i(F_k, \lambda) \rightarrow i(F, \lambda)$$

for each $\lambda \in \mathcal{ML}$.

By the Fundamental Lemma in [20], there is a uniform constant $C > 0$ such that

$$(6) \quad i(F_k, \lambda) \leq \ell_\lambda(X_k) \leq i(F_k, \lambda) + C.$$

Note that

$$c_k \ell_\lambda(X_k) \leq c_k L_k \ell_\lambda(X_0)$$

and for each L_k , there is a measured lamination λ_k with $\ell_{\lambda_k}(X_0) = 1$ such that $L_k = \ell_{\lambda_k}(X_k)$. As a result, we have

$$(7) \quad \frac{c_k \ell_\lambda(X_k)}{\ell_\lambda(X_0)} \leq c_k L_k \leq c_k \ell_{\lambda_k}(X_k).$$

From (5), (6), (7), it follows that $c_k L_k$ is uniformly bounded from above and uniformly bounded below away from zero. It follows that, up to a subsequence, $\frac{\ell_\lambda(X_k)}{L_k}$ converges to a positive multiple of $i(F, \lambda)$ for each $\lambda \in \mathcal{ML}$. Since pointed convergence of $\frac{\ell_\lambda(X_k)}{L_k}$, $\lambda \in \mathcal{ML}$ is equivalent to uniform convergence on compact subsets of \mathcal{ML} , property (\star) holds. □

Concerning the right hand side formula in (x), a similar result was obtained by Miyachi in [19] :

Proposition 1.2. *If a sequence $(X_k)_{k \geq 1}$ in $\mathcal{T}(S)$ converges to a limit P in Gardiner-Masur's boundary, then $\mathcal{E}_{X_k}(\cdot)$ converges to some function $\mathcal{E}_P(\cdot)$ in the following sense: Up to a subsequence, $\mathcal{E}_{X_k}(\cdot)$ converges to a positive multiple of $\mathcal{E}_P(\cdot)$ uniformly on any compact subset of \mathcal{ML} .*

2. LIPSCHITZ NORM

This section contains results of Thurston from his paper [32] that we will use later in this paper. We have provided proofs because at times Thurston's proofs in [32] are considered as sketchy.

A geodesic lamination μ on a hyperbolic surface X is said to be *complete* if its complementary regions are all isometric to ideal triangles. (We note that we are dealing with laminations μ that are not necessarily measured, except if specified.) Associated with (X, μ) is a measured foliation $F_\mu(X)$, called the horocyclic foliation, satisfying the following three properties:

- (i) $F_\mu(X)$ intersects μ transversely, and in each cusp of an ideal triangle in the complement of μ , the leaves of the foliation are pieces of horocycles that make right angles with the boundary of the triangle;
- (ii) on the leaves of μ , the transverse measure for $F_\mu(X)$ agrees with arclength;
- (iii) there is a nonfoliated region at the centre of each ideal triangle of $S \setminus \mu$ whose boundary consists of three pieces of horocycles that are pairwise tangent (see Figure 1).




FIGURE 1. The horocyclic foliation of an ideal triangle.

We denote by $\mathcal{MF}(\mu)$ the space of measured foliations that are transverse to μ . Thurston [32] proved the following fundamental result.

Theorem 2.1. *The map $\phi_\mu : \mathcal{T}(S) \rightarrow \mathcal{MF}(\mu)$ defined by $X \mapsto F_\mu(X)$ is a homeomorphism.*

The *stretch line* directed by μ and passing through $X \in \mathcal{T}(S)$ is the curve

$$\mathbb{R} \ni t \mapsto X_t = \phi_\mu^{-1}(e^t F_\mu(X)).$$

We call a segment of a stretch line a *stretch path*.

Suppose that μ is the support of a measured geodesic lamination λ . Then, for any two points X_s, X_t , $s \leq t$ on the stretch line, their Lipschitz distance $d_L(X_s, X_t)$ is equal to $t - s$, and this distance is realized by

$$\log \frac{\ell_\lambda(X_t)}{\ell_\lambda(X_s)}.$$

We denote by \mathcal{ML} the space of measured geodesic laminations on X and we let $\mathcal{ML}_1 = \{\lambda \in \mathcal{ML} \mid \ell_\lambda(X) = 1\}$. We may identify \mathcal{ML}_1 with \mathcal{PL} , the space of projective measured laminations.

Thurston [32] introduced a Finsler structure on $\mathcal{T}(S)$ by defining the Finsler norm of a tangent vector $V \in T_X \mathcal{T}(S)$ by the following formula :

$$(8) \quad \|V\|_L = \sup_{\lambda \in \mathcal{ML}} \frac{d\ell_\lambda(V)}{\ell_\lambda(X)}.$$

Note that we may write

$$\|V\|_L = \sup_{\lambda \in \mathcal{ML}_1} d\ell_\lambda(V).$$

Lemma 2.2. *Let $\Gamma_\mu(t)$ be a stretch line through X with $\Gamma_\mu(0) = X$ and $\dot{\Gamma}_\mu(0) = V_0$. For any measured lamination λ supported by μ (λ may be not unique), we have*

$$\|V_0\|_L = \frac{d\ell_\lambda(V_0)}{\ell_\lambda(X)}.$$

Proof. By multiplying the transverse measure by a constant, we may assume that $\ell_\lambda(X) = 1$. The stretch map from X to $\Gamma_\lambda(t)$ is a Lipschitz map with Lipschitz constant e^t , and the hyperbolic length of λ in $\Gamma_\lambda(t)$ is equal to e^t . As a result, for any measured lamination $\gamma \in \mathcal{ML}_1$,

$$\ell_\gamma(\Gamma_\lambda(t)) \leq e^t.$$

It follows that

$$\sup_{\gamma \in \mathcal{ML}_1} d\ell_\gamma(V_0) = d\ell_{\lambda_0}(V_0).$$

□

Theorem 2.3 (Thurston [32] p. 20). $\|\cdot\|_L$ is the infinitesimal norm of Thurston's asymmetric distance d_L .

Proof. Recall that the Finsler norm $\|\cdot\|_L$ induces an (asymmetric) distance:

$$d(X, Y) = \inf_{\Gamma} \int_0^1 \|\dot{\Gamma}\|_L,$$

where the infimum is taken over all piecewise C^1 curves $\Gamma : [0, 1] \rightarrow \mathcal{T}$ in Teichmüller space joining X to Y .

By compactness of $\mathcal{PM}\mathcal{L}$, there exists an element $\lambda \in \mathcal{ML}$ such that the distance $d_L(X, Y)$ is attained at λ , that is,

$$d_L(X, Y) = \log \frac{\ell_\lambda(Y)}{\ell_\lambda(X)}.$$

For any piecewise C^1 path $\Gamma : [0, 1] \rightarrow \mathcal{T}(S)$ satisfying $\Gamma(0) = X$ and $\Gamma(1) = Y$, we have

$$\begin{aligned} d_L(X, Y) &= \log \frac{\ell_\lambda(Y)}{\ell_\lambda(X)} \\ &= \int_0^1 d \log \ell_\lambda(\dot{\Gamma}(t)) \\ &\leq \int_0^1 \|\dot{\Gamma}(t)\|_L. \end{aligned}$$

Therefore, $d_L(X, Y) \leq d(X, Y)$.

To prove the reverse inequality, we use a result proved by Thurston in [32], namely, that any two points in Teichmüller space can be connected by a finite concatenation of stretch paths, each of which stretches along some common measured geodesic lamination λ .

As a result, for any distinct X and Y in $\mathcal{T}(S)$, we can assume that there exists a path Γ connecting X and Y , with Γ being a concatenation $\Gamma_1 * \cdots * \Gamma_n$ of stretch paths. Up to reparametrization, we may assume that each Γ_i is defined on $[0, 1]$. Furthermore, by a result of Thurston, we may assume that each stretch path Γ_i is directed by a complete geodesic lamination μ_i that contains a common measured lamination λ , which is the *maximally stretched lamintation* from X to Y . This is a consequence of Theorem 8.2 in Thurston [32], in which Thurston shows that there

is a unique maximal ratio-maximizing chain recurrent lamination which contains all other ratio-maximizing chain recurrent laminations. For our purposes, we take λ to be the maximal (with respect to inclusion) measured lamination contained in $\mu(X, Y)$.

Along each Γ_i , by Lemma 2.2, we have

$$\sup_{\nu \in \mathcal{ML}} \frac{d\ell_\nu(\dot{\Gamma}_i)}{\ell_\nu} = \frac{d\ell_\lambda(\dot{\Gamma}_i)}{\ell_\lambda}.$$

Now we have

$$\begin{aligned} \int_\Gamma \|\dot{\Gamma}\|_L &= \sum_{i=1}^n \int_{\Gamma_i} \|\dot{\Gamma}\|_L \\ &= \sum_{i=1}^n \int_{\Gamma_i} \frac{d\ell_\lambda(\dot{\Gamma}_i)}{\ell_\lambda} \\ &= \sum_{i=1}^n \log \frac{\ell_\lambda(\Gamma_i(1))}{\ell_\lambda(\Gamma_i(0))} \\ &= \log \frac{\ell_\lambda(Y)}{\ell_\lambda(X)}. \end{aligned}$$

For the last equality, note that we have $\Gamma_i(1) = \Gamma_{i+1}(0)$ for each i and $\Gamma_1(0) = X, \Gamma_n(1) = Y$. It follows that $d(X, Y) \leq d_L(X, Y)$. \square

3. TEICHMÜLLER NORM

Extremal length is an important tool in the study of the Teichmüller metric. The notion is due to Ahlfors and Beurling [2], see also [1]. We briefly recall the definition. Given a Riemann surface X , a *conformal metric* σ on X is a metric that is locally of the form $\sigma(z)|dz|$ where z is a local holomorphic parameter and $\sigma(z) \geq 0$ is a Borel measurable function in the local chart. We define the σ -area of X by

$$A(\sigma) = \int_X \sigma^2(z)|dz|^2.$$

Given a homotopy class of simple closed curves α , its σ -length is defined by

$$L_\sigma(\alpha) = \inf_{\alpha'} \int_{\alpha'} \sigma(z)|dz|,$$

where the infimum is taken over all essential simple closed curves α' in the homotopy class α . (Note that it follows from the invariance property of the expression $\sigma(z)|dz|$ that the two integrals $A(\sigma)$ and $L_\sigma(\alpha)$ are well defined, that is, they can be computed in the holomorphic local coordinates and the value obtained does not depend on the choice of the coordinates, see [1].)

With the above notation, we define the extremal length of α on X by

$$\text{Ext}_\alpha(X) = \sup_\sigma \frac{L_\sigma^2(\alpha)}{A(\sigma)},$$

where $\sigma(z)|dz|$ ranges over all conformal metrics on X satisfying $0 < A(\sigma) < \infty$.

By a result of Kerckhoff [12], there is a unique continuous extension of the extremal length function to the space of measured foliations \mathcal{MF} , with $\text{Ext}_{a\gamma}(X) = a^2 \text{Ext}_\gamma(X)$ for $a > 0$ and γ a homotopy class of simple closed curves on X . Recall that a tangent vector to Teichmüller space at a point X can be represented by a Beltrami differential $\mu = \mu(z) \frac{d\bar{z}}{dz}$ [11]. (Note that in this section we use the letter μ to denote a Beltrami differential since this is the traditional notation, although

in other sections the same letter is used to denote a lamination. Hopefully, there will be no confusion.) Using the extremal length function, we define a norm on the tangent space at X by setting:

$$(9) \quad \|\mu\|_E = \sup_{\lambda \in \mathcal{MF}} \frac{d\text{Ext}_\lambda^{1/2}(\mu)}{\text{Ext}_\lambda^{1/2}(X)}.$$

The following theorem is due to Hubbard-Masur [9]; we refer to Kerckhoff [12] for a short proof.

Theorem 3.1. *For any Riemann surface X and for any measured foliation λ on X , there is exactly one quadratic differential, denoted by Φ_λ , whose vertical measured foliation is measure-equivalent to λ .*

Lemma 3.2 (Gardiner [8]). *The extremal length function Ext_λ is differentiable and we have the following formula, called the “first variational formula”:*

$$(10) \quad d\text{Ext}_\lambda(\mu) = -2\text{Re} \langle \Phi_\lambda, \mu \rangle,$$

where $\langle \Phi_\lambda, \mu \rangle$ is the natural pairing

$$\langle \Phi_\lambda, \mu \rangle = \iint_X \Phi_\lambda(z) \mu(z) dx dy.$$

The following observation is due to Kerckhoff [12]:

Lemma 3.3. *With the above notation,*

$$(11) \quad \text{Ext}_\lambda(X) = \iint_X |\Phi_\lambda(z)| dx dy.$$

Proof. The proof we give here is due to Ivanov [10]. We include it for completeness.

By continuity and the density of weighted simple closed curve in \mathcal{MF} , it suffices to prove (11) for the case where $\mu = a\gamma \in \mathcal{MF}$, where γ is (the homotopy class of) a simple closed curve and $a > 0$.

Let Φ be the one-cylinder Strebel differential on X determined by $a\gamma$. The complement of the vertical critical leaves of Φ is a cylinder foliated by circles isotopic to γ . Let us also set $\rho = |\Phi|^{1/2}|dz|$. Then ρ is a flat metric on S , with a finite number of singular points, which are conical singularities. Measured in the flat metric ρ , the circumference and height of the cylinder are equal to $L_\rho(\gamma)$ and a respectively. By a theorem of Jenkins-Strebel [27], the extremal length $\text{Ext}_X(\gamma)$ of γ is equal to

$$\text{Ext}_\gamma(X) = \frac{L_\rho(\gamma)}{a},$$

where $\rho = |\Phi|^{1/2}|dz|$.

The area $A(\rho) = \iint_X |\Phi_\lambda(z)| dx dy$ of the cylinder is equal to $aL_\rho(\gamma)$. As a result,

$$\text{Ext}_{a\gamma}(X) = a^2 \text{Ext}_\gamma(X) = aL_\rho(\gamma) = A(\rho).$$

□

The next result follows from (10) and (11).

Proposition 3.4. *The norm $\|\mu\|_E$ satisfies*

$$(12) \quad \|\mu\|_E = \sup_{\lambda \in \mathcal{MF}} \frac{-\text{Re} \langle \Phi_\lambda, \mu \rangle}{\iint_X |\Phi_\lambda(z)| dx dy} = \sup_{\|\Phi\|=1} \text{Re} \langle \Phi, \mu \rangle,$$

where Φ varies over all holomorphic quadratic differentials Φ .

Note that we have $\|\Phi\| = \int_X |\Phi(z)| dx dy$.

Now we consider the Teichmüller metric on $\mathcal{T}(S)$. We recall that it is defined by

$$d_T(X, Y) := \frac{1}{2} \inf_f \log K(f)$$

where $f : X \rightarrow Y$ is a quasi-conformal map homotopic to the identity map of S and

$$K(f) = \sup_{x \in X} K_x(f) \geq 1$$

is the quasi-conformal dilatation of f (the sup here denotes essential supremum), with

$$K_x(f) = \frac{|f_z(x)| + |f_{\bar{z}}(x)|}{|f_z(x)| - |f_{\bar{z}}(x)|}$$

being the pointwise quasiconformal dilatation at the point $x \in X$ with local conformal coordinate z .

Teichmüller's theorem states that given any $X, Y \in \mathcal{T}(S)$, there exists a unique quasi-conformal map $f : X \rightarrow Y$, called the Teichmüller map, such that

$$d_T(X, Y) = \frac{1}{2} \log K(f).$$

The Beltrami coefficient $\mu := \frac{\bar{\partial}f}{\partial f}$ is of the form $\mu = k \frac{\bar{\Phi}}{|\Phi|}$ for some quadratic differential Φ on X and some constant k with $0 \leq k < 1$. In some natural coordinates given by Φ on X and for some associated quadratic differential Φ' on Y , the Teichmüller map f is given by $f(x+iy) = K^{1/2}x + iK^{-1/2}y$, where $K = K(f) = \frac{1+k}{1-k}$.

It is known that the Teichmüller metric is a Finsler metric and that between any two points in $\mathcal{T}(S)$ there is exactly one geodesic. A geodesic ray with initial point X is given by the one-parameter family of Riemann surfaces $\{X_t\}_{t \geq 0}$, where there is a holomorphic quadratic differential Φ on X and a family of Teichmüller maps $f_t : X \rightarrow X_t$, with initial Beltrami differential $\mu(f_t) = \frac{e^{2t} - 1}{e^{2t} + 1} \frac{\bar{\Phi}}{|\Phi|}$. Here $\mu(f_t)$ is chosen such that the geodesic ray has unit speed, that is, $d_T(X_s, X_t) = t - s$ for all $s \leq t$. We also recall the following formula due to Kerckhoff [12].

Theorem 3.5. *Let X, Y be any two points in $\mathcal{T}(S)$. Then*

$$d_T(X, Y) = \frac{1}{2} \log \sup_{\lambda} \frac{\text{Ext}_{\lambda}(Y)}{\text{Ext}_{\lambda}(X)},$$

where λ ranges over elements in \mathcal{MF} .

Now we can prove the following:

Theorem 3.6. *The metric induced by the norm $\|\cdot\|_E$ defined in (9) is the Teichmüller metric.*

Proof. Denote by d_E the length metric on $\mathcal{T}(S)$ induced by the norm $\|\mu\|_E$. For any X, Y in $\mathcal{T}(S)$, by Kerckhoff's formula, the Teichmüller distance is realized by

$$d_T(X, Y) = \frac{1}{2} \log \frac{\text{Ext}_{\lambda}(Y)}{\text{Ext}_{\lambda}(X)}$$

for some measured foliation λ . An argument similar to the one in the first part of the proof of Theorem 2.3 shows that $d_T(X, Y) \leq d(X, Y)$.

For the converse, let $\Gamma(t)$ be a Teichmüller geodesic connecting X and Y . We parametrize Γ with unit speed, and define it on the interval $[0, T]$ with $T = d_T(X, Y)$ with $\Gamma(0) = X$ and $\Gamma(T) = Y$.

For each t in $[0, T]$, the Teichmüller map between $X = \Gamma(0)$ and $\Gamma(t)$ has quasi-conformal dilatation e^{2t} . It follows from the geometric definition of a quasiconformal map that for any λ in \mathcal{MF} , we have

$$\text{Ext}_\lambda(\Gamma(t)) \leq e^{2t} \text{Ext}_\lambda(\Gamma(0)).$$

Taking square roots, we get

$$\text{Ext}_\lambda^{1/2}(\Gamma(t)) \leq e^t \text{Ext}_\lambda^{1/2}(\Gamma(0))$$

and then

$$\begin{aligned} \lim_{t \rightarrow 0^+} \frac{\text{Ext}_\lambda^{1/2}(\Gamma(t)) - \text{Ext}_\lambda^{1/2}(\Gamma(0))}{t \text{Ext}_\lambda^{1/2}(\Gamma(t))} &\leq \lim_{t \rightarrow 0^+} \frac{e^t \text{Ext}_\lambda^{1/2}(\Gamma(0)) - \text{Ext}_\lambda^{1/2}(\Gamma(0))}{t \text{Ext}_\lambda^{1/2}(\Gamma(t))} \\ &= \lim_{t \rightarrow 0^+} \frac{e^t - 1}{t} \\ &= 1. \end{aligned}$$

As a result, $d \log \text{Ext}_\lambda^{1/2}(\dot{\Gamma}(0)) \leq 1$ and then

$$d \log \text{Ext}_\lambda^{1/2}(\dot{\Gamma}(t)) \leq 1$$

for all $t \in \mathbb{R}$.

Considering the integral of the norm $\|\cdot\|_E$ along $\Gamma(t)$, we have

$$\int_0^T \|\dot{\Gamma}(t)\|_E \leq T = d_T(X, Y).$$

This proves that $d(X, Y) \leq d_T(X, Y)$. \square

Combining Proposition 3.4 and Theorem 3.6, we have the following corollary.

Corollary 3.7. *The Teichmüller Finsler norm, denoted by $\|\mu\|_T$, is given by*

$$\|\mu\|_T = \sup_{\|\Phi\|=1} \text{Re} \langle \Phi, \mu \rangle,$$

where Φ varies over all holomorphic quadratic differentials Φ .

The above result was already known and can be obtained by using the famous Reich-Strebel inequality [26]. The result means that the Teichmüller norm $\|\cdot\|_T$ is dual to the L^1 -norm

$$\|\Phi\| = \iint_X |\Phi(z)| dx dy$$

on the cotangent space.

4. CONVEX EMBEDDING OF MEASURE FOLIATION SPACE USING EXTREMAL LENGTH

Following a usual trend, we shall call the boundary of a convex body a *convex sphere*. Note that the boundary of a convex body is always homeomorphic to a sphere.

Thurston [32] proved that for any $X \in \mathcal{J}(S)$, the function $d\ell : \mathcal{ML}_1 \rightarrow T^*_X \mathcal{J}(S)$ defined by $\lambda \mapsto d\ell_\lambda$ embeds \mathcal{ML}_1 as a convex sphere in the cotangent space of Teichmüller space at X containing the origin. This embedding is the dual of the boundary of the unit ball in the tangent space representing vectors of norm one for the Finsler structure associated to Thurston's metric. We prove an analogous result for the extremal length function.

Theorem 4.1. *Given a point X in $\mathcal{J}(S)$, let $\mathcal{PF} = \mathcal{MF}_1 = \{\lambda \in \mathcal{MF} \mid \text{Ext}_\lambda(X) = 1\}$. Then, the function $\mathcal{PF} \rightarrow \mathbb{T}^*_X\mathcal{J}(S)$ defined by $\lambda \mapsto d\text{Ext}_\lambda$ embeds \mathcal{PF} as a convex sphere in $\mathbb{T}^*_X\mathcal{J}(S)$ containing the origin. This sphere is the boundary of the dual of the unit ball in the tangent space representing vectors of norm one for the Finsler structure associated to the Teichmüller metric.*

Proof. We fix a point X in $\mathcal{J}(S)$. By Gardiner's formula,

$$d\text{Ext}_\lambda(\mu) = -2\text{Re} \langle \Phi_\lambda, \mu \rangle$$

(see Theorem 3.1 for the notation). Kerckhoff proved that if a sequence λ_n in \mathcal{PF} converges to λ then $\|\Phi_{\lambda_n} - \Phi_\lambda\| \rightarrow 0$ (see Page 34-35 of Kerckhoff's paper [12], where he proved the result for the case where each λ_n is a simple closed curve. Since the subset of weighted simple closed curves is dense in \mathcal{MF} , the result we need follows). It follows that the map $d\text{Ext}$ is continuous.

Moreover, since the pairing between quadratic differentials and harmonic Beltrami differentials is nondegenerate [11], if $d\text{Ext}_\lambda = d\text{Ext}_{\lambda'}$, then $\Phi_\lambda = \Phi_{\lambda'}$. This means that $\lambda = \lambda'$. Therefore, the map $d\text{Ext}$ is injective.

Let $d\text{Ext}(\mathcal{PF})$ be the image of \mathcal{PF} under the map $d\text{Ext}$, that is,

$$d\text{Ext}(\mathcal{PF}) = \{d\text{Ext}_\lambda \in \mathbb{T}^*_X\mathcal{J}(S) \mid \lambda \in \mathcal{PF}\}.$$

Since \mathcal{PF} is homeomorphic to a sphere, by invariance of domain, $d\text{Ext}(\mathcal{PF})$ is open and $d\text{Ext}$ is a homeomorphism between \mathcal{PF} and $d\text{Ext}(\mathcal{PF})$.

For any $\lambda \in \mathcal{PF}$, consider the Beltrami differential

$$V_\lambda = -\frac{\overline{\Phi_\lambda}}{2|\Phi_\lambda|}.$$

It is a tangent vector in $\mathbb{T}_X\mathcal{J}(S)$. By Gardiner's formula,

$$\begin{aligned} d\text{Ext}_\lambda(V_\lambda) &= -2\text{Re} \langle \Phi_\lambda, -\frac{\overline{\Phi_\lambda}}{2|\Phi_\lambda|} \rangle \\ &= \iint_X |\Phi_\lambda| dx dy = \text{Ext}_\lambda(X) = 1. \end{aligned}$$

As a result, the derivative $d\text{Ext}_\lambda$ in the direction V_λ is 1. Using Hölder's inequality, for any $\gamma \in \mathcal{PF}$, we have

$$\begin{aligned} d\text{Ext}_\gamma(V_\lambda) &= \iint \Phi_\gamma \frac{\overline{\Phi_\lambda}}{|\Phi_\lambda|} dx dy \\ &\leq \iint_X |\Phi_\gamma| dx dy = \text{Ext}_\gamma(X) = 1. \end{aligned}$$

Moreover, equality holds if and only if $\gamma = \lambda$. Therefore, the derivative $d\text{Ext}_\gamma(V_\lambda)$ of any other $\gamma \in \mathcal{PF}$ is strictly less than 1. Denote by $C(d\text{Ext}(\mathcal{PF}))$ the convex hull of $d\text{Ext}(\mathcal{PF})$ in $\mathbb{T}^*_X\mathcal{J}(S)$. Thus, V_λ defines a non-constant linear functional on the cotangent space which attains its maximal value on $C(d\text{Ext}(\mathcal{PF}))$ at $d\text{Ext}_\lambda$. As a result, $d\text{Ext}_\lambda$ is an extreme point of the convex set $C(d\text{Ext}(\mathcal{PF}))$.

We now use Thurston's argument in his proof of Theorem 5.1 of [32]. The set of extreme points of any convex set is a sphere of some dimension. Since the dimension of $d\text{Ext}(\mathcal{PF})$ is one less than the dimension of $\mathbb{T}^*_X\mathcal{J}(S)$, the set of extremal points of $C(d\text{Ext}(\mathcal{PF}))$ coincides with $d\text{Ext}(\mathcal{PF})$. As a result, $d\text{Ext}(\mathcal{PF})$ is a convex sphere. To see that the convex set $C(d\text{Ext}(\mathcal{PF}))$ contains the origin in its interior, note that since this convex hull has a nonempty interior, there must be at least one line through the origin of $\mathbb{T}^*_X\mathcal{J}(S)$ which intersects $d\text{Ext}(\mathcal{PF})$ in at least two points. For each of these points, there is a linear functional which attains its positive maximum value there. It follows that 0 must separate the two points, so 0 is a convex combination of them. \square

5. COMPARISON BETWEEN THE LIPSCHITZ AND TEICHMÜLLER NORMS

From an inequality called Wolpert’s inequality [34], we have $d_L \leq 2d_T$. (In fact, the inequality is contained in Sorvali’s paper [25], and it was rediscovered by Wolpert.) Choi and Rafi [4] proved that the two metrics are quasi-isometric in any thick part of the Teichmüller space, but that there are sequences $(X_n), (Y_n)$ in the thin part, with $d_L(X_n, Y_n) \rightarrow 0$.

It is interesting to compare the Lipschitz norm and the Teichmüller norm. In particular, we ask the following:

Question 5.1. Determine a function $C(\epsilon)$ such that

$$\|V\|_L \leq \|V\|_T \leq C(\epsilon)\|V\|_L$$

for any $V \in T_X\mathcal{T}(S)$ and for all $X \in \mathcal{T}_\epsilon(S)$, the ϵ -thick part of $\mathcal{T}(S)$.

Since $\|V\|_L$ (respectively $\|V\|_T$) is given by the logarithmic derivative of the hyperbolic (respectively extremal) length function of measured laminations, a further study of the relation between hyperbolic and extremal length may give the answer to the above question.

Question 5.2. Consider the length-spectrum metric d_{ls} on $\mathcal{T}(S)$. This can be defined as a symmetrization of Thurston’s metric¹, by the formula

$$\begin{aligned} d_{ls}(X, Y) &= \max\{\log \sup_{\gamma} \frac{\ell_Y(\gamma)}{\ell_X(\gamma)}, \log \sup_{\gamma} \frac{\ell_X(\gamma)}{\ell_Y(\gamma)}\} \\ &= \max\{d_L(X, Y), d_L(Y, X)\}. \end{aligned}$$

Is the length-spectrum metric d_{ls} a Finsler metric ? If yes, give a formula for the infinitesimal norm of a vector on Teichmüller space with respect to this Finsler structure.

There are comparisons between the length-spectrum metric and the Teichmüller metric. It was shown by Liu and Su [16] that the divergence of d_{ls} and d_T is only caused by the action of the mapping class group. In fact, they showed that d_{ls} and d_T are “almost” isometric on the moduli space and that the two metrics on the moduli space determine the same asymptotic cone.

Question 5.3. One can also seek for results analogous to those presented here for the weak metric L^* dual to Thurston’s metric, that is, the weak metric on Teichmüller space defined by

$$L^*(X, Y) = L(Y, X),$$

as well as for the length spectrum metric d_{ls} on the same space.

REFERENCES

- [1] L. V. Ahlfors, Conformal invariants: topics in geometric function theory. McGraw-Hill, New York (1973).
- [2] L. V. Ahlfors and A. Beurling, Conformal invariants and function-theoretic null-sets, Acta Math. 83 (1950), 101–129.
- [3] L. V. Ahlfors, Quasiconformal mappings, Teichmüller spaces, and Kleinian groups. Proc. int. Congr. Math., Helsinki 1978, Vol. 1, 71-84 (1980). Collected Works Volume II, pp. 485–498.
- [4] Y-E. Choi and K. Rafi, Comparison between Teichmüller and Lipschitz metrics, J. London Math. Soc. (2) 76 (2007), 739–756.
- [5] M. Culler and K. Vogtmann, Moduli of graphs and automorphisms of free groups. Invent. Math., 84(1), p. 91-119, 1986.

¹Note that historically the length spectrum metric was not introduced as a symmetrization. It was first defined by Sorvali [24], before Thurston introduced his asymmetric metric.

- [6] A. Fathi, F. Laudenbach and V. Poénaru, Travaux de Thurston sur les surfaces (second edition), Astérisque, vol. 66–67.
- [7] F. Gardiner, Schiffer’s interior variation and quasiconformal mapping, *Duke Math. J.* 42 (1975), 371–380.
- [8] F. Gardiner, Measured foliations and the minimal norm property for quadratic differentials, *Acta. Math.* 152 (1984), 57–76.
- [9] J. Hubbard and H. Masur, Quadratic differentials and foliations, *Acta. Math.* 142 (1979), 221–274.
- [10] N. V. Ivanov, Isometries of Teichmüller spaces from the point of view of Mostow rigidity, *Amer. Math. Soc. Transl. Ser. 2*, 202 (2001), 131–149.
- [11] Y. Iwayoshi and M. Taniguchi, *An Introduction to Teichmüller Spaces*. Springer 1992.
- [12] S. Kerckhoff, The asymptotic geometry of Teichmüller space, *Topology* 19 (1980), 23–41.
- [13] A. Lenzhen, K. Rafi and J. Tao, Bounded combinatorics and the Lipschitz metric on Teichmüller space, to appear in *Geometriae Dedicata*.
- [14] L. Liu, A. Papadopoulos, W. Su and G. Théret, On length spectrum metrics and weak metrics on Teichmüller spaces of surfaces with boundary, *Ann. Acad. Sci. Fenn., Math.* 35, No. 1, 255–274 (2010).
- [15] L. Liu, A. Papadopoulos, W. Su and G. Théret, Length spectra and the Teichmüller metric for surfaces with boundary, *Monatshefte für Mathematik*, 2010, vol. 161, no3, pp. 295–311
- [16] L. Liu and W. Su, Almost-isometry between Teichmüller metric and length-spectra metric on moduli space, arXiv:1012.1921, to appear in *Bulletin of the L. M. S.*
- [17] L. Liu and W. Su, The horofunction compactification of Teichmüller space, to appear in the *Handbook of Teichmüller theory*, Vol. IV, European Mathematical Society, 2012.
- [18] L. Liu, A. Papadopoulos, W. Su and G. Théret, On the classification of mapping class actions on Thurston’s asymmetric metric, preprint, 2011, submitted.
- [19] H. Miyachi, Teichmüller rays and the Gardiner-Masur boundary of Teichmüller space, *Geom. Dedicata* 137 (2008), 113–141.
- [20] A. Papadopoulos, On Thurston’s boundary of Teichmüller space and the extension of earthquakes, *Topology and its Application* 41 (1991), 147–177.
- [21] A. Papadopoulos and G. Théret, On Teichmüller’s metric and Thurston’s asymmetric metric on Teichmüller space, in : A. Papadopoulos (ed.), *Handbook of Teichmüller theory. Volume I*. Zürich, European Mathematical Society. IRMA Lectures in Mathematics and Theoretical Physics 11, 111-204 (2007).
- [22] A. Papadopoulos and G. Théret, On Teichmüller’s metric and Thurston’s asymmetric metric on Teichmüller space, in: *Handbook of Teichmüller theory, Volume I*, European Mathematical Society, Zürich, 2007, pp. 111-204.
- [23] A. Papadopoulos and G. Théret, Shift coordinates, stretch lines and polyhedral structures for Teichmüller space. *Monatsh. Math.* 153, No. 4, 309-346 (2008).
- [24] T. Sorvali, The boundary mapping induced by an isomorphism of covering groups, *Ann. Acad. Sci. Fenn. Math.* 526 (1972), 1–31.
- [25] T. Sorvali, On the dilatation of isomorphisms between covering groups, *Ann. Acad. Sci. Fenn. AI*, 551 15p. (1973).
- [26] E. Reich and K. Strebel, Extremal quasiconformal mappings with given boundary values, *Contributions to Analysis* (L. V. Ahlfors et al., eds.), Academic Press, New York, 1974, pp. 375–391.
- [27] K. Strebel, Quadratic differentials, *Ergebnisse der Math. (3)*, Vol.5 Spring-Verlag, Berlin, 1984.
- [28] O. Teichmüller, Extremale quasikonforme Abbildungen und quadratische Differentiale. *Abh. Preuss. Akad. Wiss., Math.-Naturw. Kl.* 22, pp. 1-197 (1940), *Collected Works*, Springer Verlag, 337-531.
- [29] Teichmüller, Bestimmung der extremalen quasikonformen Abbildungen bei geschlossenen orientierten Riemannschen Flächen. *Abh. Preuss. Akad. Wiss., Math.-Naturw. Kl.* 1943, No.4, 42 p. (1943).
- [30] G. Théret, On the negative convergence of Thurston’s stretch lines towards the boundary of Teichmüller space, *Ann. Acad. Sci. Fenn., Math.* 32, No. 2, 381-408 (2007).
- [31] G. Théret, On elementary antistretch lines, *Geom. Dedicata* 136, pp. 79-93 (2008).
- [32] W. P. Thurston, Minimal stretch maps between hyperbolic surfaces, 1986 preprint, arxiv:math/9801039v1.
- [33] C. Walsh, The horoboundary and isometry groups of Thurston’s Lipschitz metric, to appear in the *Handbook of Teichmüller theory*, Vol. IV, European Mathematical Society, 2012..
- [34] S. Wolpert, The length spectra as moduli for compact Riemann surfaces, *Ann. of Math.* 109 (1979), 323–351.

ATHANASE PAPADOPOULOS, UNIVERSITÉ DE STRASBOURG AND CNRS, 7 RUE RENÉ DESCARTES,
67084 STRASBOURG CEDEX, FRANCE

E-mail address: athanase.papadopoulos@math.unistra.fr

WEIXU SU, DEPARTMENT OF MATHEMATICS, FUDAN UNIVERSITY, 200433, SHANGHAI, P. R.
CHINA, AND UNIVERSITÉ DE STRASBOURG AND CNRS, 7 RUE RENÉ DESCARTES, 67084 STRAS-
BOURG CEDEX, FRANCE

E-mail address: suweixu@gmail.com