



**HAL**  
open science

## Low microzooplankton grazing rates in the Arctic Ocean during a *Phaeocystis pouchetii* bloom (Summer 2007): fact or artifact of the dilution technique?

Albert Calbet, Enric Saiz, Rodrigo Almeda, Juan Ignacio Movilla, Miguel Alcaraz

### ► To cite this version:

Albert Calbet, Enric Saiz, Rodrigo Almeda, Juan Ignacio Movilla, Miguel Alcaraz. Low microzooplankton grazing rates in the Arctic Ocean during a *Phaeocystis pouchetii* bloom (Summer 2007): fact or artifact of the dilution technique?. *Journal of Plankton Research*, 2010, 10.1093/plankt/FBQ142 . hal-00642440

**HAL Id: hal-00642440**

**<https://hal.science/hal-00642440>**

Submitted on 18 Nov 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Low microzooplankton grazing rates in the Arctic Ocean during a *Phaeocystis pouchetii* bloom (Summer 2007): fact or artifact of the dilution technique?**

Journal:	<i>Journal of Plankton Research</i>
Manuscript ID:	JPR-2010-197.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	01-Oct-2010
Complete List of Authors:	Calbet, Albert; Institut de Ciències del Mar (CSIC), Biologia Marina i Oceanografia Saiz, Enric; Institut de Ciències del Mar (CSIC), Biologia Marina i Oceanografia Almeda, Rodrigo; Institut de Ciències del Mar (CSIC), Biologia Marina i Oceanografia Movilla, Juan Ignacio; Institut de Ciències del Mar (CSIC), Biologia Marina i Oceanografia Alcaraz, Miguel; Institut de Ciències del Mar (CSIC), Biologia Marina i Oceanografia
Keywords:	Arctic Ocean, Microzooplankton, Dilution technique, Food web

SCHOLARONE™  
Manuscripts

1  
2  
3  
4 **Low microzooplankton grazing rates in the Arctic Ocean during a *Phaeocystis***  
5 ***pouchetii* bloom (Summer 2007): fact or artifact of the dilution technique?**  
6  
7  
8  
9

10  
11 Albert Calbet\*, Enric Saiz, Rodrigo Almeda, Juan Ignacio Movilla, and Miquel  
12 Alcaraz  
13

14  
15  
16 Institut de Ciències del Mar, CSIC. Pg. Marítim de la Barceloneta 37-49. 08003.  
17  
18 Barcelona, Spain.  
19

20  
21 \*Email: [acalbet@icm.csic.es](mailto:acalbet@icm.csic.es)  
22

23  
24 Phone: +34 932309500  
25

26  
27 FAX: +34 932309555  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract**

We studied the structure and dynamics of the microbial community of Arctic waters during July 2007 using a microzooplankton grazing dilution approach. The sampling covered a latitudinal transect along the East Greenland Sea, and a series of stations in the high Arctic (up to 80° 50' N), west and north of the Svalbard Islands. A main feature of the area was the presence of *Phaeocystis pouchetii*, which formed dense blooms. Despite the considerable biomass of microzooplankton (mostly large ciliates and dinoflagellates), their grazing impact on phytoplankton, assessed as total chlorophyll *a*, was significant in only 6 out of 16 experiments, which resulted in 8% of the standing stock being consumed on average. Overall, phytoplankton instantaneous growth rates were very low and even negative at times (range: -0.24 to 0.14; average: -0.04 for total chlorophyll), which could not be attributed to nutrient limitation nor the estimated microzooplankton grazing. We present 3 nonexclusive explanations for this fact: 1) we were facing a senescent community in which many organisms were dying either as a result of virus infections or for other natural causes, as corroborated by parallel estimates of natural cell mortality using membrane permeability probes; 2) the widespread and abundant *P. pouchetii* was probably deterring grazing and adversely affecting the entire planktonic community at the time of the study; and 3) the dilution technique failed to give a real estimate of grazing (i.e. either non significant or positive slopes), likely as a consequence of trophic cascades (decline of major grazers in the more concentrated treatments) combined with saturated-feeding responses. This last point calls for special attention when intending to use the dilution technique in productive environments, where grazing may be saturated.

**Keywords:** Arctic Ocean; Microzooplankton; *Phaeocystis pouchetii*; Dilution technique, Food web.

**Running title:** Food web interactions in the Arctic Ocean

## Introduction

The fast melting of Arctic ice, a consequence of global warming (Cavalieri *et al.*, 2003; Johannessen *et al.*, 2004), has profound ecological consequences on Arctic wildlife, including polar bears, walruses, and ringed seals, which are threatened by the reduction of sea-ice cover. Not so evident are the changes that the inhabitants of the waters underneath and surrounding the Polar ice cap will suffer under progressive melting conditions. One of the key groups in marine food webs, which are potentially sensitive to these Arctic scenario changes, are the microzooplankton.

Microzooplankton are of paramount importance in controlling primary production (Calbet and Landry, 2004). Regrettably, their role in the high Arctic still remains uncertain and needs to be determined to fully understand and predict the consequences of the changes this ecosystem will experience in the future. For instance, the relevance of the trophic impact of this group in the Arctic Ocean food web has mostly been studied either in bays or in relatively low latitude sites (Paranjape, 1987; Gifford *et al.*, 1995; Olson and Strom, 2002; Verity *et al.*, 2002; Strom *et al.*, 2008), or it has been derived indirectly (Levinsen *et al.*, 1999; Rysgaard *et al.*, 1999; Levinsen and Nielsen, 2002). Most of these studies indicate a strong control of primary production by microzooplankton grazing. However, recent work in high Arctic waters by Sherr *et al.* (Sherr *et al.*, 2009) questions such strong control, likely due, according to the authors, to a strong top-down impact of copepods on microzooplankton (Levinsen and Nielsen, 2002; Campbell *et al.*, 2009).

For this reason, we undertook a study of the microbial interactions in Arctic waters during the melting season. Our research coincided with a bloom of *P. pouchetii* (Lasternas and Agustí, in press), which will add further value to our results because, despite being a successful species in Arctic waters (Schoemann *et al.*, 2005), the very few data dealing with the impact of microzooplankton feeding on these algae have resulted in contradictory conclusions. Weisse and Scheffel-Moser (Weisse and Scheffel-Moser, 1990) measured microzooplankton grazing in a *P. cf. pouchetii* bloom in the North Sea using the dilution technique (Landry and Hassett, 1982)

1  
2  
3 finding grazing loss rates from 0.037 to 0.174 h<sup>-1</sup>, grazing rates increasing in the  
4 course of the bloom and exceeding phytoplankton growth rates at the end. Gifford *et*  
5 *al.* (Gifford *et al.*, 1995), also using the dilution technique, did not detect any grazing  
6 on phytoplankton in the high-latitude North Atlantic Ocean during a *P. pouchetii*  
7 bloom, but they obtained increased microzooplankton grazing as the bloom declined.  
8 Archer *et al.* (Archer *et al.*, 2000) measured microzooplankton grazing impact under  
9 relatively low abundance of *P. pouchetii* in 3 fjords of the northern Norway, and  
10 obtained significant grazing rates on total phytoplankton and on fluorescently labeled  
11 algae of similar size to *P. pouchetii*. On the other hand, Wolfe *et al.* (2000) only found  
12 significant grazing on chlorophyll and DMSP when *P. pouchetii* cells were in poor  
13 condition in the Labrador Sea. In this regard, the ability of microzooplankton to ingest  
14 *Phaeocystis* spp. has been widely discussed in the literature (see reviews by Whipple  
15 *et al.*, 2005, and Nejstgaard *et al.*, 2007). It is usually accepted that microzooplankton  
16 exerts their pressure mostly on single cells; however, there is evidence that *Noctiluca*  
17 *scintillans* (Weisse *et al.*, 1994; Jakobsen and Tang, 2002) and *Gyrodinium cf. spirale*  
18 (Stelfox-Widdicombe *et al.*, 2004) are able to ingest small colonies.

## 45 Method

46  
47 This study was part of a multidisciplinary project (ATOS: POL2006-00550/CTM),  
48 and took place on board the research vessel BIO Hespérides from July 1 to 24, 2007.  
49 The cruise departed Reykjavik (Iceland), and sampling started northward across the  
50 Greenland Sea (Table 1, Fig. 1). In Arctic waters, we sampled a series of stations in  
51 the vicinity of the ice-edge, alternating between several stations free of ice and two  
52 coastal stations near the Svalbard Islands. During the study, we reached a historical  
53 minimum of Arctic ice cover (Zhang *et al.* 2008), allowing samples to be taken from  
54 areas up to 80° 50' N. The stations lasted 24 h, starting with profiles for the  
55  
56  
57  
58  
59  
60

1  
2  
3 measurement of salinity, temperature, and fluorescence during the early morning  
4 using a Seabird CTD911, followed by collection of water samples for the  
5 determination of chlorophyll *a* concentration (hereafter Chl *a*) with 12 l Niskin bottles  
6 fitted to a rosette during the ascending CTD casts.  
7  
8  
9

10  
11 At each station we conducted standard grazing dilution experiments (Landry  
12 and Hassett, 1982) to assess the microzooplankton grazing impact on primary  
13 producers and on other components of the food web (see below). This technique  
14 consists of the sequential dilution of natural water with filtered seawater to obtain a  
15 gradient of net grazing impact on phytoplankton. The water for these experiments was  
16 collected at the fluorescence maximum (Table 1) using a rosette equipped with 12-L  
17 Niskin bottles, according to the fluorescence profile. Once on deck, the water was  
18 gravity-filtered through a Pall Acropak 0.8/0.2 500 capsule (previously flushed,  
19 including tubing, with diluted HCl and thoroughly rinsed with deionised water), and  
20 then 2 replicate bottles (2.3-L acid washed polycarbonate) for each of the dilution  
21 treatments were filled with the corresponding required amount of filtered seawater.  
22 Afterwards, we added natural non-filtered seawater from the selected depth to the  
23 bottles to generate experimental water percentages of 13, 27, 50, 73, and 100%. At  
24 some stations, the presence of a dense bloom of the haptophyte *Phaeocystis pouchetii*  
25 made it impossible to efficiently filter the water through the Acropak capsules;  
26 therefore, filtered seawater originated from below the bloom depth.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 To guarantee the homogeneity of the natural water poured from different Niskin  
42 bottles filled at the same depth, we used a 20-L intermediary carboy in which the  
43 water was gently mixed by its own flow. All the process was carried out under dim  
44 light conditions to avoid cell light-damage. To promote constant and saturated  
45 phytoplankton growth in the dilution series, each bottle received added nutrients (10  
46  $\mu\text{M}$   $\text{NH}_4\text{Cl}$  and 0.7  $\mu\text{M}$   $\text{Na}_2\text{HPO}_4$ ). In addition, four 100% (i.e., not diluted) bottles  
47 were prepared without nutrients to assess the natural growth of the algae. Two of  
48 these latter bottles were sacrificed for initial samples. Because we did not screen the  
49 water used for the dilution series to avoid damaging delicate microzooplankton, the  
50 experimental suspension may have contained some mesozooplankton. We examined  
51 the bottles by eye to observe the presence of large copepods, and the very few times  
52 this occurred did not result in inconsistent results in the dilution series.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

All bottles were incubated on deck in a large (600 L), dark incubator with open-circuit water running from a 5-m depth at a temperature about the same as *in situ*. The natural sunlight was dimmed with appropriate dark plastic mesh to mimic the light intensity at the fluorescence maximum. We gently mixed the bottles by repeatedly turning them upside down and moving them around the incubator at least three times per day. After 27-32 h, we finished the incubations and took samples for quantification of total and > 5- $\mu\text{m}$  Chl *a* concentration. To get further insight into the actual trophic interactions during dilution experiments and to detect possible artifacts (Dolan *et al.*, 2000; Agis *et al.*, 2007; Modigh and Franzè, 2009), we additionally took samples for the determination of nano- and microplankton from the 2 initials and in one of the replicates per dilution level, and preserved them with Acidic Lugol's solution (2% final concentration). To avoid damaging the delicate cells we first added the fixative and then gently siphoned the water sample directly into the sample bottle

For total Chl *a*, we filtered 50-250 ml of water (depending on station and dilution level) under low vacuum pressure (< 100 mm Hg) through Whatmann glass fiber filters (GF/F, 25 mm diameter). For the > 5- $\mu\text{m}$  fraction, we filtered 100-300-ml samples through 5- $\mu\text{m}$  pore-size polycarbonate Osmonics Inc. filters (25 mm diameter). After filtration, the filters were stored frozen at -20°C until fluorometric analysis of acetone extracts, with and without acidification (Parsons *et al.*, 1984) on a Turner Designs Fluorometer.

Lugol-preserved samples were processed in the laboratory. We concentrated the most diluted treatments (13%, 25% and 50%) by first settling the entire bottle for 72 h and gently siphoning off 50-75% of the supernatant water without re-suspending the sample. Then, for all the samples, 100 ml of the concentrate was settled in Utermöhl chambers for at least 48 h prior and counted under the microscope. The whole chamber, or a fraction of it for the smallest and more abundant organisms, was counted under an inverted microscope (XSB-1A) at 100, 250, and 400X magnification, depending on the group. Fifty to one-hundred cells per group were sized, adjusted to their closest geometric shape, and converted into carbon using the equations of Menden-Deuer and Lessard (Menden-Deuer and Lessard, 2000). We did not use any correction factor to compensate for ciliate losses due to fixation, as previously suggested by Broglio *et al.* (Broglio *et al.*, 2004), and Calbet and Saiz



(Calbet and Saiz, 2005), because recent research has revealed that such corrections should apply to many other planktonic groups, not only ciliates (Zarauz and Irigoien, 2008), and universal factors have not been developed yet. Because plankton were preserved with acidic Lugol's solution, no distinction between strict heterotrophs and auto/mixotrophs was made for flagellates, ciliates, and some dinoflagellates. However, the groups identified to species level were classified trophically according to the literature.

Instantaneous growth rates in dilution grazing experiments were derived from net growth in the un-amended bottles (no nutrients added) adding the mortality by microzooplankton grazing from dilution experiments when the latter was significant. We should note that in the cases where the grazing mortality was not significant the instantaneous growth rate of prey would be equivalent to the net growth rate in the un-amended bottles. Therefore, we may be underestimating the actual instantaneous growth rate of the prey because some mortality may still occur in the bottles, even if not measured by the dilution method. All statistical tests were conducted with JMP 7.0 statistical software.

## Results

### *Plankton biomass and distribution*

We present a summary of the different fractions of Chl *a* in Table 1 and the contribution of micro- and nanoplankton to total plankton community biomass at the sampled stations in Table 2 (a detailed summary of the species composition is presented in the Supplementary Material online). The main feature of the data for the Arctic stations is the almost ubiquitous presence of the haptophyte *Phaeocystis pouchetii*, which formed dense blooms (Table 2, Fig. 2) and whose distribution was directly correlated with total Chl *a* ( $r=0.76$ ,  $p<0.01$ ), and inversely correlated with temperature ( $r = 0.77$ ;  $p < 0.05$ ). Following *P. pouchetii*, diatoms (mostly chain forming species  $> 20 \mu\text{m}$  in total length of the genera *Chaetoceros*, *Fragilariopsis*, *Nitzschia*, *Rhizosolenia*, *Thalassiothrix*, and *Thalassiosira*) were the second biomass grouping within the phytoplankton, being dominant in the Greenland Sea and at warmer stations. Regarding micrograzers, the important contribution of large ciliates

1  
2  
3 (mostly Tintinnida, *Strombidium* spp., and the mixotroph *Laboea* spp.) was noticeable  
4 in the Greenland Sea and Svalbard Islands coastal station (Table 2). Actually, ciliates  
5 were the overall major contributors to microzooplankton during the study and were  
6 substantial components of the total plankton biomass (ca. 25%), whereas  
7 heterotrophic dinoflagellates (considering 50% of unidentified dinoflagellates as  
8 heterotrophic; Lessard and Swift, 1985; Sherr and Sherr, 2007) represented, on  
9 average, 12% of the total carbon biomass (Table 2). In relation to the relevance of  
10 heterotrophs, it is interesting to note that the quotient between heterotrophic and  
11 autotrophic carbon (indicative of the trophic characteristics of the system) was > 1 in  
12 the Greenland Sea and Svalbard Islands coastal stations (Fig. 3).  
13  
14  
15  
16  
17  
18  
19  
20  
21

22 Despite the high heterotrophic biomass, we did not find significant  
23 correlations between any of the size fractions of Chl *a*, large heterotrophic  
24 dinoflagellates (e.g., *Gyrodinium* spp., *Protoperidinium* spp., and *Katodinium* spp.),  
25 and ciliates. However, large mixotrophic dinoflagellates (*Dinophysis* spp., *Ceratium*  
26 spp., *Gonyaulax* spp., and *Amphidinium* spp.) were positively correlated with diatoms  
27 ( $r = 0.85$ ;  $p < 0.05$ ). Correlations between groups do not necessarily mean causality,  
28 but they can indicate that an association exists. Likewise, unidentified > 20- $\mu\text{m}$   
29 dinoflagellates were also positively correlated with *P. pouchetti* biomass ( $r = 0.72$ ;  $p <$   
30  $0.05$ ). However, when excluded from the analysis station 43 (the station with the  
31 maximum biomass of *P. pouchetii*) this relationship is strongly affected and it  
32 becomes non-significant.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 We calculated the C:Chl *a* ratios using the Chl *a* data and the autotrophic  
44 biomass (in carbon), obtained by cell counting and conversion to carbon using  
45 literature equations (Table 2). For unidentified dinoflagellates we assumed 50% of  
46 the organisms were phototrophs. Because our microscope counting technique, based  
47 on Lugol-preserved samples, does not allow for good resolution of the lower size-  
48 fractions, we can better estimate the C:Chl *a* ratio using only > 5- $\mu\text{m}$  Chl *a*. The  
49 values averaged 18 for the entire data-set, and ranged from 4 to 47.  
50  
51  
52  
53  
54  
55  
56  
57  
58

#### 59 *Dilution grazing experiments*

60

1  
2  
3 Table 3 shows the outcome of the dilution grazing experiments based on the different  
4 size-fractions of Chl *a*. Unanticipated results from these experiments include the  
5 following: i) the little, or even at times negative, phytoplankton instantaneous growth  
6 (range: -0.24 to 0.14; average -0.04 for total Chl *a*) and ii) the low microzooplankton  
7 grazing impact on primary producers. Significant microzooplankton grazing (i.e.,  
8 significant negative slopes in the dilution experiments) were found only in 6 out of 16  
9 experiments for total and <5- $\mu\text{m}$  Chl *a*, and no significant mortality was detected in  
10 the >5- $\mu\text{m}$  fraction at any of the stations. Overall, microzooplankton grazing cannot  
11 explain the observed phytoplankton negative net growth rates (especially in the >5-  
12  $\mu\text{m}$  size-fractions) and must be attributed to other causes (see discussion). However, it  
13 is interesting to note the inverse relationship between the phytoplankton net growth  
14 rates and Chl *a* concentration (Fig. 4) in the Arctic Ocean stations. This points  
15 towards a naturally-occurring, density-dependent mortality effect, probably linked to  
16 the presence of *Phaeocystis pouchetii*, although not significantly related to the  
17 occurrence of either this or any other planktonic group. Likewise, and related to net  
18 phytoplankton growth rates, there was no clear evidence of nutrient limitation at most  
19 of the stations (Fig. 5).

20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
The phytoplankton and microzooplankton composition analysis in the initial  
and final dilution experiments revealed a very complex food web scenario, with  
frequent negative growth rates both for autotrophs and heterotrophs, frequent positive  
slopes, and very few cases of significant microzooplankton grazing impact (Table 4).  
Overall, it is difficult to extract any clear interpretation or global pattern out of the  
dilution data for the different plankton groups. Nevertheless, we decided to present  
these data because negative dilution grazing results in the literature have seldom been  
discussed (Dolan and McKeon, 2005). As example, we show in Fig. 6 the dilution  
experiment plots for stations 4 and 33. It is interesting to note in these plots the  
positive slopes for some prey and the negative slopes (theoretically indicative of  
grazing) for some potential grazers of phytoplankton. It should be noted that the  
mortality rates of top predators (large ciliates and heterotrophic dinoflagellates) are  
not necessarily estimated well by the dilution method. However, we think the  
information about the changes of these groups in the incubation bottles is relevant and  
should be also presented in the results.

## Discussion

### *Community composition and microzooplankton grazing on phytoplankton*

A main characteristic that distinguishes our study from previous ones in Arctic waters is the peculiarity of the composition of the heterotrophic microbial community. Past works stressed the relevance of heterotrophic dinoflagellates in Arctic waters, especially when diatoms dominated the autotrophic community (Levinsen *et al.*, 1999; Rysgaard *et al.*, 1999; Sherr *et al.*, 1997; Sherr and Sherr, 2007; Sherr *et al.*, 2009). We found a population rich in ciliates, which accounted for ca. 2/3 of the total microzooplankton biomass (assuming 50% of the unidentified dinoflagellates were heterotrophs). This high contribution was especially evident in the Greenland Sea, where they completely dominated the microplankton. We suspect that in these waters we were facing a very unusual situation, likely the result of temporal imbalances in the structure of the community (e.g., the end of a phytoplankton bloom), as the biomass of producers seems unable to maintain such a high abundance of grazers.

Despite the relatively high biomass of microzooplankton, the grazing rates on phytoplankton (Chl *a*) were low. Certainly, trophic relationships between predators and prey do not necessarily have to be directly related to biomass, especially in areas where annual blooms are intense. If a phytoplankton bloom is senescent, grazers, even if abundant, may not graze phytoplankton cells in poor health (the end of the bloom situation). Similar inconsistencies between grazing impacts and grazer biomass have also been reported for Antarctic waters (Caron *et al.*, 2000). Regarding the Arctic Ocean, we find contrasting results for microzooplankton grazing impact in different areas. For instance, during the summer along the western coast of Greenland, Levinsen *et al.* (Levinsen *et al.*, 1999) studied the microzooplankton grazing impact by indirect methods and concluded that if this group had a purely autotrophic diet they could remove 362% of primary daily production. Nevertheless, the authors pointed out that cannibalism could likely reduce the actual impact of this group on phytoplankton. Also by indirect measurements, Rysgaard *et al.* (Rysgaard *et al.*, 1999) estimated that the combined grazing activity of ciliates and dinoflagellates in Young Sound (NE Greenland) would potentially remove only 14% of the annual

1  
2  
3 primary production, a value that contrasts strongly with the previous estimate.  
4  
5 Levinsen and Nielsen (Levinsen and Nielsen, 2002) found that potential  
6  
7 microzooplankton grazing could account for 32-55% of the primary production in  
8  
9 Disko Bay. These data, as well as the 40-114% primary production daily grazed in  
10  
11 Jones Sound (Paranjape, 1987), the 37-88% in Baffin Bay (Paranjape, 1987), the 64-  
12  
13 97% in the Barents Sea (Verity *et al.*, 2002), and the 2-293% (average 110 and 81%  
14  
15 of phytoplankton growth rates for > 10  $\mu\text{m}$  and < 10  $\mu\text{m}$  phytoplankton, respectively)  
16  
17 grazed in the southeast Bering Sea (Olson and Strom, 2002) are much higher than the  
18  
19 values observed in our study. Recent research in the high Western Arctic Ocean  
20  
21 (Sherr *et al.*, 2009), however, advocates a lower control of microzooplankton grazing  
22  
23 on primary producers (average  $22 \pm 26\%$ ). The study by Sherr *et al.* (Sherr *et al.*,  
24  
25 2009), even if in a different area, with lower average temperatures, and with a  
26  
27 community of phytoplankton not dominated by *Phaeocystis*, but by diatoms, is the  
28  
29 study that *a priori* seems more appropriate to contrast with ours; both were located in  
30  
31 open waters near the ice-edge zone, and both were conducted at high latitudes using  
32  
33 the same methodology. Our data agree with the results of the Sherr *et al.* (Sherr *et al.*,  
34  
35 2009) study, which does not show significant grazing in about half of the experiments  
36  
37 at the fluorescence maximum and contains total average grazing rates of  $< 0.1 \text{ d}^{-1}$ .  
38  
39 Similarly, in the Sherr *et al.* (Sherr *et al.*, 2009) study they found low and even  
40  
41 negative values for phytoplankton growth rates during summer. They attributed this  
42  
43 natural mortality to low light levels (samples were collected at the base of the  
44  
45 euphotic zone) and to post-bloom conditions (protist grazing rates were low because  
46  
47 the diatom blooms were senescent).

45  
46 Despite the overall low grazing rates, some associations between the  
47  
48 distributions of several organisms seem to be evident. For instance, mixotrophic  
49  
50 dinoflagellates were positively correlated with diatoms. Mixotrophy is widespread  
51  
52 among dinoflagellates, and it is not uncommon that these organisms contribute  
53  
54 significantly to the community grazing on phytoplankton (Stoecker, 1999; Stoecker *et*  
55  
56 *al.*, 2009), and particularly on diatoms (Yoo *et al.*, 2009). Moreover, the role of  
57  
58 mixotrophic dinoflagellates in the fate of primary producers is likely to have been  
59  
60 underestimated for several reasons. First, the presence of their own chloroplasts may  
mask the detection of prey inside the organisms, and second, the different feeding  
mechanisms displayed by this group (direct engulfment, tube-feeding, and pallium-

1  
2  
3 feeding; Hansen and Calado, 1999) make it quite difficult to correctly assess their  
4 contribution to total community grazing. Because *pallium*-feeding and tube-feeding  
5 do not leave evident remains of the preyed cell inside the predator and because the  
6 *pallium* and peduncles are not persistent structures, they are not easily quantified  
7 when microscopically observing preserved samples. Therefore, we are of the opinion  
8 that a predator-prey association between armoured mixotrophic dinoflagellates and  
9 diatoms is meaningful. Certainly, other microbial grazers, such as ciliates among  
10 others, can impact on diatoms (Aberle *et al.*, 2007), but we did not obtain proof of this  
11 behaviour in our study. On the other hand, the diatom-dinoflagellate relationship has  
12 been widely suggested in the literature (e.g., Sherr and Sherr, 2007; Saito *et al.*, 2006;  
13 Calbet, 2008), although seldom quantified in natural communities (Archer *et al.*,  
14 1996).

15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28 *Finding explanations for the low microzooplankton grazing impact, but the high net*  
29 *mortality rates of phytoplankton:*

30  
31  
32  
33 It is difficult to ascertain whether the low grazing found in our study is a general  
34 characteristic describing the system or if it was the result of some particular  
35 conditions. It is surprising, however, that we found many negative net phytoplankton  
36 growth rates (based on Chl *a* changes) not associated with microzooplankton grazing.  
37 The simplest explanation for this would be that the incubation light-level was not  
38 appropriate, the cells adjusting their Chl *a* contents to the new conditions. This could  
39 actually be the cause at some stations; however, the same pattern persists in cell  
40 counts in many others. Therefore, we have to find alternative hypotheses, which may  
41 be nonexclusive:

42  
43  
44  
45  
46  
47  
48  
49  
50 - *The natural mortality hypothesis*

51  
52  
53 Parallel to our study, Lasternas and Agustí (Lasternas and Agustí, in press) used a  
54 membrane permeability probe (Agustí and Sánchez, 2002) to estimate the natural  
55 mortality of the *P. pouchetii* community, and they observed that higher percentages of  
56 dead *P. pouchetii* cells (up to ca. 90% at some stations) were associated to both cold  
57 and less saline waters across the area studied. The mechanisms behind this mortality,  
58 not related to grazing rates, can be several. It could well be that we were facing a  
59  
60



1  
2  
3 senescent community at the end of the bloom. On the other hand, we cannot disregard  
4 viruses as playing a role in controlling the population of these and other protists and  
5 responsible for the frequent negative growth rates (Baudoux *et al.*, 2006; Jacobsen *et*  
6 *al.*, 2007). Unfortunately, our experimental set up was not adequate for virus-related  
7 mortality quantification.  
8  
9

10  
11  
12  
13 Given the contribution of *P. pouchetii* to the total phytoplankton biomass, the  
14 natural mortality of these algae could be driving the growth rates observed on the  
15 basis of Chl *a*. Corroborating this, we observed an inverse relationship between  
16 phytoplankton instantaneous growth rates and Chl *a* distribution. Although other  
17 groups of plankton could also have shown natural mortality rates, we do not have  
18 solid evidence of this.  
19  
20  
21  
22  
23

#### 24 25 - *The Phaeocystis pouchetii hypothesis*

26  
27  
28 As previously mentioned, our cruise coincided with high abundances of *Phaeocystis*  
29 *pouchetii*, which commonly blooms in these waters in July (Schoemann *et al.*, 2005).  
30 The peculiarities of the food web dominated by these algae make it difficult to extract  
31 general conclusions. We believe that our low grazing estimates (not significant at  
32 most of the stations) could be partially a consequence of the presence of *P. pouchetii*.  
33 We observed the presence of many colonies in the samples, although we have not  
34 been able to quantify the percentage of free cells and colonies in the Lugol preserved  
35 samples. However, it is quite likely this species either introduced variability into the  
36 samples (colonies can be heterogeneously distributed in the samples) and precluded  
37 the establishment of significant regressions based on Chl *a*, evaded grazing when in  
38 colonial form (Hansen *et al.*, 1994; Weisse *et al.*, 1994; Tang, 2003), or chemically  
39 deterred grazers (Barnard *et al.*, 1984; Nejstgaard *et al.*, 2007; van Rijssel *et al.*,  
40 2007). Certainly, *P. pouchetii* seems the most obvious candidate when seeking  
41 chemical deterrence of grazing. Similar to macroalgae and other phytoplankters,  
42 *Phaeocystis* spp. exude chemicals that can interfere with grazing activity (see review  
43 by van Rijssel *et al.*, 2007). Although the chemicals involved in this process have not  
44 yet been identified, it has been suggested that grazing-activated DMSP cleavage by *P.*  
45 *pouchetii* contributes to grazing deterrence (Wolfe *et al.*, 2000). Allelopathic  
46 interactions could also be responsible for the negative growth rates of a phytoplankton  
47 (and other protists) community, which apparently was not limited by nutrients.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Overall, we believe that the net transfer of energy to higher trophic levels in this ecosystem would be greatly diminished if the autotrophic community was dominated by *Phaeocystis*, provided that other groups not considered here (e.g., copepods) do not exert a strong impact on this alga. However, the low tolerance of *P. pouchetii* to relatively high temperatures (Schoemann *et al.*, 2005), as shown by the reduced presence at St. 27 where temperatures reached 7.5°C, indicates a limited relevance of this species under global warming scenarios. Certainly, this does not mean that other *Phaeocystis* species, such as *P. globosa*, which is adapted to warmer waters, cannot replace *P. pouchetii*, further diminishing the trophic efficiency of the system because *P. globosa* is seldom consumed by zooplankton (see review by Nejstgaard *et al.*, 2007).

#### -The dilution grazing artifact hypothesis

Chl *a* is a rough proxy for phytoplankton because it does not capture the complexity of this group, and its use in dilution experiments has been questioned because chlorophyll content per cell may change during the incubation (McManus, 1995). Moreover, the need for a close examination of the microzooplankton community during dilution experiments to detect and correct possible artifacts has also been discussed (Dolan *et al.*, 2000; Agis *et al.*, 2007; Modigh and Franzè, 2009). To shed light on this point and to deepen our understanding of the food web interactions in Arctic waters, we further examined changes in the entire microbial community during the dilution experiments. We did not make any attempt to compare the rates obtained based on Chl *a*, with those derived from cell counts and their corresponding conversion to carbon because the uncertainties associated when depicting a trophic role (autotrophy vs heterotrophy) to unidentified dinoflagellates and nanoflagellates. Besides, the contribution of mixotrophic species at some stations (e.g. the mixotrophic ciliate *Laboea* sp. represented most of the planktonic biomass at station St. 3; Supplementary Material online) precluded any comparison.

When opening the planktonic black box in the dilution experimental bottles we faced unanticipated results suggesting a complex and intricate food web, in which choosing the major microbial grazers of phytoplankton was not so straightforward. A clear example of a puzzling response was the occurrence of positive regression slopes between the net growth rates of certain groups against the dilution factor. Positive slopes for heterotrophs and mixotrophs, even if sometimes the result of complicated



1  
2  
3 ecological interactions, can be easily interpreted as growth enhancement due to  
4 increased feeding in the more concentrated treatments. However, positive slopes of  
5 phytoplankton occur when the organism considered is adversely affected by the  
6 dilution treatment. The explanation for these particular responses is not easy because  
7 they can have different non-exclusive causes. For instance, they may either be the  
8 result of strong trophic cascade effects during the incubations (Calbet *et al.*, 2008),  
9 chemical grazing deterrence by the algae or other organisms, toxic effects of the  
10 filtered seawater (Landry, 1993), mixotrophs being important contributors of  
11 phytoplankton biomass (then, favoured in less diluted conditions), or of complex  
12 cycling of nutrients between internal and external pools (Landry, 1993), because  
13 nutrients would be taken up by smaller algae more efficiently and would become  
14 limiting for larger phytoplankton. We do not believe that the last three hypotheses  
15 apply to our experiments for the following reasons: i) any toxic effect would likely be  
16 persistently evident at all the groups and stations, ii) some of the groups of  
17 phytoplankton showing positive slopes, as far as we know, were not mixotrophic  
18 (e.g., diatoms), and iii) nutrients were supplied in excess. Therefore, either (or both)  
19 trophic cascades or grazing deterrence seem to be the most reasonable explanation. If  
20 *P. pouchetii* was responsible for the positive slopes found for diatoms and other  
21 groups, its effects would not be apparent at the stations where the haptophyte was not  
22 present (stations 2 and 3). While this was the case and thus supporting the feeding  
23 deterrence hypothesis, it did not fully demonstrate the hypothesis because the  
24 response was not directly related to the *P. pouchetii* concentration.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 We, therefore, contemplated the trophic cascade explanation for the positive  
44 slopes found in our (and others) study. It has been argued that changes of grazer  
45 abundance during dilution grazing incubation may result in results that are artifacts  
46 (Dolan *et al.*, 2000; Agis *et al.*, 2007; Modigh and Franzè, 2009). These changes  
47 usually involve a decrease in abundance in the most diluted treatments, the result of  
48 starvation. However, our experiments showed on many occasions the opposite, the  
49 grazers diminishing in the most concentrated treatments. This can be a consequence  
50 of predation from other microzooplankers, either protozoans (intraguild predation) or  
51 metazoans (e.g., copepod nauplii; not included in our sampling), during the  
52 incubations.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

It is relatively easy to mathematically simulate a dilution grazing experiment involving a grazer that reduces their abundance inversely related to the dilution level during the incubation. We can actually base our example on data from one of our experiments. For instance, we can use as an example the response of  $< 20 \mu\text{m}$  ciliates at Station 4, and the positive slope for nanoflagellates, one of their likely prey (Fig. 6). We assume nanoflagellates doubled their abundance in 24h, and that  $< 20 \mu\text{m}$  ciliates were the only group grazing on them. If ciliate feeding rates were linearly related to food concentration we will most likely obtain, after the incubation period, a negative slope for nanoflagellates when plotting net growth rates as function of dilution level, as predicted by dilutions (although it would not be a true estimate of the natural grazing rate on this group because grazers varied their concentrations in the experimental bottles). However, if feeding was saturated, we could easily mimic the results found in the experiments using a constant feeding rate of only 76 nanoflagellates consumed per ciliate per day (Fig. 7). This happens because the grazing pressure is in this case only proportional to the abundance of grazers, and we have a higher net growth rate of grazers ( $< 10 \mu\text{m}$  ciliates) in the more concentrated treatments. Moreover, varying the concentration of grazers, prey and the growth and grazing rates we can also obtain non significant from zero grazing estimates, which are not true rates, but artifacts of the method in very special situations. This mathematical exercise was not intended to correct our grazing rates, as suggested by Modigh and Franzè (Modigh and Franzè, 2009), because given the complexity of the food web we cannot anticipate the microzooplankton group responsible for most of the phytoplankton grazing impact. However, we can use our reasoning to prove that positive slopes (and non-significant slopes) are easily the result of a combination of trophic cascades during the incubations (the main grazers decreasing their abundance in the most concentrated treatments) with a saturated feeding responses. The picture complicates further if microzooplankton feeding behaviour changes with food concentration, as described by Teixeira and Figueras (Teixeira and Figueras, 2009), and if there is nutrient limitation during the incubations. Actually, and regarding this latter artifact, severe nutrient limitation during the dilution experiments inversely proportional to the dilution level will most likely favour fake negative slopes, exaggerating the grazing activity of microzooplankton because phytoplankton instantaneous growth rates will be higher in the most diluted treatments, where more nutrients per cell are available.

1  
2  
3  
4 In summary, the data presented here depict a planktonic Arctic community dominated  
5 by *P. pouchetii* and rich in microzooplankton, which at first sight did not seem to be exerting  
6 a strong control on a phytoplankton community in decline. However, several natural and  
7 causes as well as artifacts may have been playing important roles in some of our  
8 experiments, precluding clear grazing estimates in this very complex food web. Even though,  
9 it may seem disappointing to conclude that our rates might not be actual estimates, but bound  
10 for the lower grazing impact of the microzooplankton in the area, they point out the need for  
11 presenting negative results, when these are not consequence of evident mistakes or artifacts  
12 (Dolan and McKeon, 2005). Only with a whole picture of the existing data we will be able to  
13 extract solid conclusions on the dynamics of marine systems. Maybe in the future someone  
14 will find the way of extracting information from such results. Specifically regarding dilution  
15 grazing experiments, this study calls for special caution when applying the technique,  
16 originally developed for oligotrophic areas, to rich environments where saturated feeding  
17 responses may be common. In any case, as previously suggested (Dolan *et al.*, 2000; Agis *et*  
18 *al.*, 2007; Modigh and Franzè, 2009), it is evident we need a detailed examination of the  
19 grazer and prey dynamics during the incubations if we want to present trustable  
20 microzooplankton grazing estimates. By presenting data on counts-based rates we will  
21 enhance our resolution and avoid artifacts associated with chlorophyll analysis. However,  
22 these sorts of data involve a considerable amount of time and are highly dependent on the  
23 taxonomic skills of the researcher.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

- Aberle N., Lengfellner K., Sommer U. (2007) Spring bloom succession, grazing impact and herbivore selectivity of ciliate communities in response to winter warming. *Oecologia*, **150**, 668-681.
- Agis, M., Granda, A., Dolan J. R. (2007) A cautionary note: Examples of possible microbial community dynamics in dilution grazing experiments. *J. Exp. Mar. Biol. Ecol.*, **341**, 176-183.
- Agustí, S., Sánchez, C. M. (2002) Cell viability in natural phytoplankton communities quantified by a membrane permeability probe. *Limnol. Oceanogr.*, **47**, 818–828.
- Archer S. D., Leakey, R. J. G., Burkill, P. H., Sleigh M. A. (1996) Microbial dynamics in coastal waters of East Antarctica: herbivory by heterotrophic dinoflagellates. *Mar. Ecol. Prog. Ser.*, **139**, 239-255.
- Archer, S. D., Verity, P. G., Stefels, J. (2000) Impact of microzooplankton on the progression and fate of the spring bloom in fjords of northern Norway. *Aquat. Microb. Ecol.*, **22**, 27-41.
- Barnard, W. R., Andreae, M. O., Iverson, R. L. (1984) Dimethylsulde and *Phaeocystis poucheti* in the Southeastern Bering Sea. *Cont. Shelf. Res.*, **3**, 103–113.
- Baudoux, A- C., Noordeloos, A. A. M., Veldhuis, M. J. W., Brussaard, C. P. D. (2006) Virally induced mortality of *Phaeocystis globosa* during a spring bloom in temperate coastal waters. *Aquat. Microb. Ecol.*, **44**, 201-217.
- Broglio, E., Saiz, E., Calbet, A., Trepas, I., Alcaraz, M. (2004) Trophic impact and prey selection by crustacean zooplankton on the microbial communities of an oligotrophic coastal area (NW Mediterranean Sea). *Aquat. Microb. Ecol.*, **35**, 65-78.
- Calbet, A. (2008) The trophic roles of microzooplankton in marine systems. *Ices J. Mar. Science*, **65**, 325-331.
- Calbet, A., Landry, M. R. (2004) Phytoplankton growth, microzooplankton grazing, and carbon cycling in marine systems. *Limnol. Oceanogr.*, **49**, 51-57.

1  
2  
3 Calbet, A., Saiz, E. (2005) The ciliate-copepod link in marine ecosystems. *Aquat.*  
4 *Microb. Ecol.*, **38**, 157-167.

5  
6  
7  
8 Calbet, A., Trepát, I., Almeda, R., Saló, V., Saiz, E., Movilla, J. I., Alcaraz, M.,  
9 Yebra, L., Simó, R. (2008) Impacts of micro- and nanograzers on phytoplankton  
10 assessed by standard and size-fractionated dilutions. *Aquat. Microb. Ecol.*, **50**, 154-  
11 156.

12  
13  
14  
15  
16 Campbell, R. G., Sherr, E. B., Ashjian, C. J., Plourde, S., Sherr, B. F., Hill, V.,  
17 Stockwell, D. A. (2009) Mesozooplankton prey preference and grazing impact in the  
18 Western Arctic Ocean. *Deep-Sea Research II* **56**, 1274-1289.

19  
20  
21  
22 Caron, D. A., Dennett, M. R., Lonsdale, D. J., Moran, D. M., Shalapyonok, L. (2000)  
23 Microzooplankton herbivory in the Ross Sea, Antarctica. *Deep-Sea Res. II*, **47**, 3249-  
24 3272.

25  
26  
27  
28 Cavalieri, D. J., Parkinson, C. L., Vinnikov, K. Y. (2003) 30-Year satellite record  
29 reveals contrasting Arctic and Antarctic decadal sea ice variability. *Geophys. Res.*  
30 *Lett.*, **30**, DOI: 10.1029/2003GL018031.

31  
32  
33  
34  
35 Dolan, J. R., Gallegos, C. L., Moigis, A. (2000) Dilution effects on microzooplankton  
36 in dilution grazing experiments. *Mar. Ecol. Prog. Ser.*, **200**, 127-139.

37  
38  
39  
40 Dolan, J. R., McKeon, K. (2005) The reliability of grazing rate estimates from  
41 dilution experiments: Have we over-estimated rates of organic carbon consumption by  
42 microzooplankton? *Ocean Science*, **1**, 1-7.

43  
44  
45  
46 Gallegos, C. L. (1989) Microzooplankton grazing on phytoplankton in the Rhode  
47 River, Maryland: nonlinear feeding kinetics. *Mar. Ecol. Prog. Ser.*, **57**, 23-33.

48  
49  
50  
51 Gifford, D. J., Fessenden, L. M, Garrahan, P. R. and Martin, E. (1995) Grazing by  
52 microzooplankton and mesozooplankton in the high-latitude North Atlantic Ocean:  
53 Spring versus summer dynamics, *J. Geophys. Res.*, **100**, 6665–6675.

54  
55  
56  
57 Hansen, B., Verity, P., Falkenhaus, T., Tande, K. S., Norrbin, F. (1994) On the  
58 trophic fate of *Phaeocystis pouchetti* (Harriot). 5. Trophic relationships between  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Phaeocystis* and zooplankton: An assessment of methods and size dependence. *J. Plankton Res.*, **16**, 487-511.

Hansen, P. J., Calado, A. J. (1999) Phagotrophic mechanisms and prey selection in free-living dinoflagellates. *J. Euk. Microbiol.*, **46**, 282-289.

Jacobsen, A., Larsen, A., Martínez-Martínez, J., Verity, P. G., Frischer, M. E. (2007) Susceptibility of colonies and colonial cells of *Phaeocystis pouchetii* (Haptophyta) to viral infection *Aquat. Microb. Ecol.*, **48**, 105-112.

Johansson, M., Gorokhova, E., Larsson, U. (2004) Annual variability in ciliate community structure, potential prey and predators in the open northern Baltic Sea proper. *J. Plankton Res.* **26**, 67–80.

Landry, M.R. (1993) Estimating rates of growth and grazing mortality of phytoplankton by the dilution method. In: Kemp P. F., Sherr B. F., Sherr E. B., Cole J. L. (eds) Handbook of methods in aquatic microbial ecology. Lewis Publishers, Boca Raton, FL, p 715–722.

Landry, M. R., Hassett, R. P. (1982) Estimating the grazing impact of marine microzooplankton. *Mar. Biol.*, **67**, 283-288.

Lasternas, S., Agustí, S. Phytoplankton production and community structure during the arctic ice-melting record in summer 2007. In press, *Polar Biol.*

Lessard, E. J., Swift, E. (1985) Species-specific grazing rates of heterotrophic dinoflagellates in oceanic waters, measured with a dual radioisotope technique. *Mar. Biol.*, **87**, 289-296.

Levinsen, H., Nielsen, T. G. (2002) The trophic role of marine pelagic ciliates and heterotrophic dinoflagellates in arctic and temperate coastal ecosystems: A cross-latitude comparison. *Limnol. Oceanogr.*, **47**, 427-439.

Levinsen, H., Nielsen, T. G., Hansen, B. W. (1999) Plankton community structure and carbon cycling on the western coast of Greenland during the stratified summer situation. II. Heterotrophic dinoflagellates and ciliates. *Aquat. Microb. Ecol.*, **6**, 217–232.

1  
2  
3 Mcmanus G. B. (1995) Phytoplankton abundance and pigment changes during  
4 simulated in situ dilution experiments in estuarine waters: possible artifacts caused by  
5 algal light adaptation. *J. Plankton Res.*, **17**, 1705-1716.  
6  
7

8  
9  
10 Menden-Deuer S., Lessard E. J. (2000) Carbon to volume relationships for  
11 dinoflagellates, diatoms, and other protist plankton. *Limnol. Oceanogr.*, **45**, 569-579.  
12  
13

14 Modigh, M., Franzè, G. (2009) Changes in phytoplankton and microzooplankton  
15 populations during grazing experiments at a Mediterranean coastal site. *J. Plankton*  
16 *Res.*, **31**, 853–864.  
17  
18

19  
20 Nejtgaard, J., Tang, K., Steinke, M., Dutz, J., Koski, M., Antajan, E., Long J. (2007)  
21 Zooplankton grazing on *Phaeocystis*: a quantitative review and future challenges.  
22 *Biogeochemistry*, **83**, 147-172.  
23  
24

25  
26  
27 Olson, M. B., Strom, S. L. (2002) Phytoplankton growth, microzooplankton herbivory  
28 and community structure in the southeast Bering Sea: insight into the formation and  
29 temporal persistence of an *Emiliana huxleyi* bloom. *Deep Sea Res. II*, **49**, 5969-  
30 5990.  
31  
32

33  
34  
35 Paranjape, M. A. (1987) Grazing by microzooplankton in the eastern Canadian Arctic  
36 in summer 1983. *Mar. Ecol. Prog. Ser.*, **40**, 239-246.  
37  
38

39  
40 Parsons, T. R., Maita, Y., Lalli, C.M. (1984) A manual of chemical and biological  
41 methods for sea water analysis. Pergamon Press, Oxford.  
42  
43

44 Rysgaard, S., Nielsen, T. G., Hansen, B. W. (1999) Seasonal variation in nutrients,  
45 pelagic primary production and grazing in a high-Arctic coastal marine ecosystem,  
46 Young Sound, Northeast Greenland. *Mar. Ecol. Prog. Ser.*, **179**, 13-25.  
47  
48

49  
50 Saito, H., Ota, T., Suzuki, K., Nishioka, J., Tsuda, A. (2006) Role of heterotrophic  
51 dinoflagellate *Gyrodinium* sp. in the fate of an iron induced diatom bloom *Geophys.*  
52 *Res. Lett.* **33** L09602, doi, 10.1029/2005GL025366.  
53  
54

55  
56  
57 Schoemann, W., Becquevort, S., Stefels, J., Rousseau, W., Lancelot, C. (2005)  
58 *Phaeocystis* blooms in the global ocean and their controlling mechanisms: a review. *J.*  
59 *Sea Res.*, **53**, 43-66.  
60



- 1  
2  
3 Sherr, E. B., Sherr, B. F. (2007) Heterotrophic dinoflagellates: a significant  
4 component of microzooplankton biomass and major grazers of diatoms in the sea.  
5 *Mar. Ecol. Prog. Ser.*, **352**, 187-197.  
6  
7  
8  
9  
10 Sherr, E. B., Sherr, B. F., Fessenden, L. (1997) Heterotrophic protists in the Central  
11 Arctic Ocean. *Deep-Sea Res. II*, **44**, 1665-1682.  
12  
13  
14 Sherr, E. B., Sherr, B. F., Hartz, A. J. (2009) Microzooplankton grazing impact in the  
15 Western Arctic Ocean. *Deep-Sea Res. Part II*, **56**, 1264-1273.  
16  
17  
18  
19 Stelfox-Widdicombe, C.E., Archer, S.D., Burkill, P.H. and Barnes, P.M. (2004)  
20 Microzooplankton grazing in *Phaeocystis* and diatom-dominated waters in the  
21 southern North Sea in spring. *Journal of Sea Research*, **51**, (1), 37-51.  
22  
23  
24  
25 Stoecker, D.K. (1999) Mixotrophy among dinoflagellates. *J. Eukary. Microbiol.*, **46**,  
26 397-401.  
27  
28  
29  
30 Stoecker, D. K., Johnson, M. D., de Vargas, C., Not, F. (2009) Acquired phototrophy  
31 in aquatic protists. *Aquat. Microb. Ecol.*, **57**, 279-310.  
32  
33  
34  
35 Strom, S. L., Fredrickson, K. A. (2008) Intense stratification leads to phytoplankton  
36 nutrient limitation and reduced microzooplankton grazing in the southeastern Bering  
37 Sea. *Deep-Sea Res II*, **55**, 1761-1774.  
38  
39  
40  
41 Tang, K. W. (2003) Grazing and colony size development in *Phaeocystis globosa*  
42 (Prymnesiophyceae): the role of a chemical signal. *J. Plankton Res.*, **25**, 831-842.  
43  
44  
45  
46 Teixeira, I. G., Figueiras, F. C. (2009) Feeding behaviour and non-linear responses in  
47 dilution experiments in a coastal upwelling system. *Aquat. Microb. Ecol.*, **55**, 53-63.  
48  
49  
50  
51 van Rijssel, M., Alderkamp, A-C., Nejstgaard, J. C., Sazhin, A. F., Verity, P. G.  
52 (2007) Haemolytic activity of living *Phaeocystis pouchetii* during mesocosm blooms.  
53 Biogeo- chemistry. doi, 10.1007/s10533-007-9095-1.  
54  
55  
56  
57 Verity, P. G., Wassmann, P., Frischer, M. E., Howard Jones, M. H., Allen, A. E.  
58 (2002) Grazing of phytoplankton by microzooplankton in the Barents Sea during  
59 early summer. *J. Mar. Sys.*, **38**, 109-123.  
60



1  
2  
3 Weisse, T., Scheffel-Moser, U. (1990) Growth and grazing loss rates in single-celled  
4 *Phaeocystis* sp. (Prymnesiophyceae). *Marine Biology* **106**, 153– 158.  
5  
6

7  
8 Weisse, T., Tande, K., Verity, P., Hansen, F., Gieskes, W. (1994) The trophic  
9 significance of *Phaeocystis* blooms. *J. Mar. Syst.*, **5**, 67–79.  
10  
11

12 Whipple, S. J., Patten, B. P. and Verity, P. G. (2005) Life cycle of the marine alga  
13 *Phaeocystis*: a conceptual model to summarize literature and guide research. *J. Mar.*  
14 *Syst.* **57**, 83-110.  
15  
16

17  
18 Wolfe, G. V., Levasseur, M., Cantin, G., Michaud, S. (2000) DMSP and DMS  
19 dynamics and microzooplankton grazing in the Labrador Sea: application of the  
20 dilution technique. *Deep-Sea Res. I*, **47**, 2243-2264.  
21  
22

23  
24 Du Yoo, Y., Jeong H. J., Kim, M. S., Kang N. S., Song, J. Y., Shin, W., Kim, K. Y.,  
25 Lee, K. (2009) Feeding by phototrophic red-tide dinoflagellates on the ubiquitous  
26 marine diatom *Skeletonema costatum*. *Euk. Microbiol.*, **56**, 413-420.  
27  
28

29  
30 Zarauz, L., Irigoien, X. (2008) Effects of Lugol's fixation on the size structure of  
31 natural nano–microplankton samples, analyzed by means of an automatic counting  
32 method. *J. Plankton Res.*, **30**, 1297–1303.  
33  
34

35  
36 Zhang, J., Lindsay, R., Steele, M., Schweiger, A. (2008) What drove the dramatic  
37 retreat of Arctic sea ice during summer 2007? *Geophys. Res. Lett.*, **35**(11), L11505.  
38  
39  
40  
41  
42

#### 43 44 **Acknowledgements**

45  
46 This research was supported by the Spanish grants POL2006-0550/CTM to C. Duarte,  
47 CTM2009-08783 to A.C., CTM2006-12344-C02-01 to M. A, CTM2007-60052 to E.S., and  
48 grant 2005SGR-00219 from the Generalitat de Catalunya to the research group, and a PhD  
49 fellowship, BES-2005-7491 to RA. The authors wished to thank the crew of the BIO  
50 Hesperides and the special help of the technical staff of the UTM (D. Afonso, R. Ametller, J.  
51 Llinàs J. Pozo, J. Sanz, and X. Vidal) for their invaluable logistic help. Plankton  
52 composition was analyzed by Dr. M. Delgado. Five anonymous reviewers highly contributed  
53 to improve earlier versions of the ms.  
54  
55  
56  
57  
58  
59  
60

**TABLE 1.** Summary of the dates of sampling (July 2007), geographic position and area, depth of sampling (m) and *in situ* chlorophyll *a* concentration ( $\mu\text{g l}^{-1} \pm \text{SE}$ ) and temperature for the different stations sampled.

Station	Date	Latitude	Longitude	Area	Sampling depth	Total Chl <i>a</i>	> 5 $\mu\text{m}$ Chl <i>a</i>	Temp. ( $^{\circ}\text{C}$ )
2	2	70° 43.19 N	17° 07.70 W	Greenland Sea	20	0.48 $\pm$ 0.02	0.15 $\pm$ 0.00	-1.2
3	3	72° 57.21 N	12° 39.19 W	Greenland Sea	25	1.31 $\pm$ 0.21	0.79 $\pm$ 0.00	0
4	4	74° 53.89 N	7° 24.50 W	Greenland Sea	32	1.41 $\pm$ 0.02	0.49 $\pm$ 0.01	0.8
5	5	77° 23.29 N	1° 40.57 W	Arctic Ocean	15	6.67 $\pm$ 0.42	2.90 $\pm$ 0.13	4
6	6	78° 00.44 N	2° 29.94 E	Arctic Ocean	23	1.75 $\pm$ 0.03	1.09 $\pm$ 0.03	4
9	7	78° 43.72 N	2° 58.51 E	Arctic Ocean	15	2.97 $\pm$ 0.07	1.48 $\pm$ 0.03	5
12	8	79° 30.83 N	7° 29.74 E	Arctic Ocean	20	5.21 $\pm$ 0.33	2.57 $\pm$ 0.07	5
15	9	80° 08.39 N	11° 19.54 E	Svalbard Coast	20	3.30 $\pm$ 0.13	1.81 $\pm$ 0.36	5
18	10	80° 26.90 N	15° 35.38 E	Svalbard Coast	35	2.67 $\pm$ 0.10	0.81 $\pm$ 0.08	5
20	12	80° 13.98 N	10° 10.97 E	Arctic Ocean	24	4.81 $\pm$ 0.20	2.31 $\pm$ 0.07	5
23	13	79° 22.16 N	6° 49.39 E	Arctic Ocean	17	8.97 $\pm$ 0.63	5.01 $\pm$ 0.08	5
27	15	79° 52.71 N	8° 36.44 E	Arctic Ocean	30	1.81 $\pm$ 0.09	1.38 $\pm$ 0.07	7.5
33	17	80° 23.46 N	12° 25.98 E	Arctic Ocean	25	1.97 $\pm$ 0.01	0.66 $\pm$ 0.00	5
39	19	80° 49.96 N	13° 12.82 E	Arctic Ocean	39	0.77 $\pm$ 0.02	0.28 $\pm$ 0.03	5
43	22	80° 25.29 N	7° 57.57 E	Arctic Ocean	20	8.97 $\pm$ 0.21	2.28 $\pm$ 0.12	-1
46	23	79° 59.15 N	3° 39.63 E	Arctic Ocean	16	4.94 $\pm$ 0.07	0.73 $\pm$ 0.06	-1.5

**Table 2.** In situ biomass in  $\mu\text{gC L}^{-1}$  of the different protist groups considered at the stations sampled. Numbers between parentheses are SE of 2 replicates. The quotient “total carbon /  $> 5 \mu\text{m Chl } a$ ” are also shown.

Station	Diatoms	Nanofl	P. pouchetii	< 20 $\mu\text{m}$ dinofl.	> 20 $\mu\text{m}$ dinofl.	Mixo. dinofl.	Het. dinofl.	< 20 $\mu\text{m}$ ciliat	> 20 $\mu\text{m}$ ciliat	Total biomass	C / > 5 $\mu\text{m}$ Chl a
2	1.25 (0.07)	0.36 (0.06)	0.00 (0.00)	0.51 (0.08)	0.24 (0.03)	0.24 (0.02)	0.28 (0.07)	4.48 (0.46)	34.91 (4.88)	42.26	14.03
3	1.55 (0.21)	3.12 (0.05)	0.00 (0.00)	0.84 (0.02)	0.22 (0.03)	0.12 (0.01)	0.08 (0.00)	4.74 (0.33)	50.53 (4.00)	61.18	4.75
4	1.26 (0.15)	0.60 (0.01)	0.83 (0.12)	0.77 (0.12)	0.57 (0.01)	0.72 (0.08)	0.43 (0.08)	12.2 (0.00)	20.90 (2.31)	38.31	7.73
5	1.77 (0.47)	0.51 (0.12)	2.81 (0.32)	14.1 (3.40)	0.26 (0.06)	1.76 (0.28)	0.47 (0.13)	0.66 (0.19)	11.49 (3.03)	33.83	4.75
6	1.53 (0.05)	3.78 (0.81)	12.2 (2.30)	10.3 (4.38)	0.87 (0.23)	1.20 (0.27)	0.69 (0.13)	3.70 (0.11)	6.08 (1.66)	40.42	20.64
9	1.79 (0.11)	1.02 (0.09)	1.59 (0.10)	1.93 (0.32)	0.34 (0.03)	0.72 (0.06)	0.15 (0.02)	0.40 (0.11)	0.83 (0.06)	8.77	3.88
12	0.81 (0.03)	1.81 (0.05)	5.90 (0.82)	1.49 (0.04)	0.75 (0.11)	0.11 (0.04)	0.17 (0.01)	1.91 (0.15)	0.94 (0.27)	13.88	3.44
15	2.97 (0.48)	3.82 (0.12)	13.0 (1.27)	5.56 (0.14)	3.09 (0.74)	0.99 (0.05)	1.53 (0.24)	2.45 (0.33)	15.09 (6.67)	48.50	12.80
18	0.18 (0.02)	4.01 (1.87)	1.19 (0.71)	6.14 (0.72)	5.24 (0.28)	0.27 (0.05)	3.33 (0.67)	2.12 (0.15)	59.9 (7.61)	82.38	11.46
20	4.24 (0.36)	3.43 (0.22)	14.8 (2.40)	3.99 (0.07)	3.89 (0.73)	0.98 (0.08)	0.57 (0.17)	6.59 (0.42)	4.52 (0.50)	43.02	10.39
23	67.8 (5.19)	3.10 (0.39)	24.8 (2.63)	5.00 (0.28)	1.79 (0.53)	3.65 (1.20)	0.65 (0.11)	5.73 (0.21)	7.12 (0.81)	119.58	20.18
27	25.1 (1.21)	1.36 (0.10)	0.09 (0.01)	5.49 (0.34)	1.91 (0.10)	2.55 (0.23)	0.51 (0.04)	0.81 (0.24)	1.74 (0.38)	39.59	23.22
33	6.47 (0.38)	2.33 (0.28)	7.89 (4.22)	3.36 (0.47)	2.70 (0.53)	1.05 (0.17)	0.70 (0.05)	4.56 (0.13)	5.60 (0.63)	34.67	29.68
39	6.73	1.61	1.79	2.10	2.50	0.25	0.34	1.45	1.02	17.78	42.16

	(1.05)	(0.15)	(0.04)	(0.08)	(0.39)	(0.03)	(0.04)	(0.25)	(0.20)		
43	6.09	1.15	49.0	1.96	9.73	1.37	1.08	2.77	4.77	77.93	27.63
	(0.78)	(0.20)	(7.40)	(0.20)	(2.02)	(0.06)	(0.16)	(0.27)	(0.73)		
46	3.67	2.52	25.7	1.65	2.33	1.39	1.02	5.94	3.43	47.61	46.53
	(0.18)	(0.06)	(9.41)	(0.39)	(0.34)	(0.21)	(0.02)	(0.57)	(0.12)		

---

For Peer Review

**Table 3.** Dilution grazing experiments. Phytoplankton instantaneous growth rates without nutrients ( $\mu \pm \text{SE}$ ;  $\text{d}^{-1}$ ) and mortality rates ( $m \pm \text{SE}$ ;  $\text{d}^{-1}$ ) for total and the 2 fractions of chlorophyll *a* at the different stations. %SS correspond to the calculated impact on the phytoplankton standing stock. Determination coefficient for the regression analysis is also provided. SE error for  $\mu$  was obtained using the equation  $(\text{SE}_k^2 + \text{SE}_m^2)^{1/2}$  where  $\text{SE}_k$  is the SE of the average of the 100% un-amended bottles and  $\text{SE}_m$  is the standard error associated to the slope of the regression equation (when significant).

Station	$\mu$ (TOTAL)	$m$ (TOTAL)	$r^2$ (TOTAL)	$\mu$ (< 5 $\mu\text{m}$ )	$m$ (< 5 $\mu\text{m}$ )	$r^2$ (< 5 $\mu\text{m}$ )	$\mu$ (> 5 $\mu\text{m}$ )	$m$ (> 5 $\mu\text{m}$ )	%SS
2	0.12 $\pm$ 0.029	ns	–	0.10 $\pm$ 0.035	ns	–	0.17 $\pm$ 0.14	ns	0.0
3	0.07 $\pm$ 0.11	ns	–	-0.05 $\pm$ 0.019	ns	–	0.13 $\pm$ 0.18	ns	0.0
4	-0.23 $\pm$ 0.11	-0.31 $\pm$ 0.11	0.50	-0.39 $\pm$ 0.25	-0.44 $\pm$ 0.19	0.45	-0.22 $\pm$ 0.13	ns	23.9
5	-0.11 $\pm$ 0.077	-0.13 $\pm$ 0.039	0.59	0.00 $\pm$ 0.21	-0.34 $\pm$ 0.056	0.84	-0.15 $\pm$ 0.069	ns	11.6
6	-0.05 $\pm$ 0.010	ns	–	0.29 $\pm$ 0.11	-0.30 $\pm$ 0.086*	0.67	-0.03 $\pm$ 0.050	ns	0.0
9	-0.11 $\pm$ 0.046	ns	–	-0.33 $\pm$ 0.026	ns	–	0.07 $\pm$ 0.090	ns	0.0
12	0.15 $\pm$ 0.113	-0.31 $\pm$ 0.11*	0.73	-0.25 $\pm$ 0.026	ns	–	-0.08 $\pm$ 0.022	ns	28.6
15	0.09 $\pm$ 0.030	ns	–	-0.16 $\pm$ 0.24	ns	–	0.23 $\pm$ 0.079	ns	0.0
18	-0.06 $\pm$ 0.059	ns	–	-0.11 $\pm$ 0.00	ns	–	0.19 $\pm$ 0.036	ns	0.0
20	-0.24 $\pm$ 0.015	ns	–	-0.29 $\pm$ 0.031	ns	–	-0.19 $\pm$ 0.060	ns	0.0
23	-0.24 $\pm$ 0.00	ns	–	-0.48 $\pm$ 0.15	ns	–	-0.09 $\pm$ 0.07	ns	0.0
27	-0.04 $\pm$ 0.043	ns	–	-0.04 $\pm$ 0.16	ns	–	-0.04 $\pm$ 0.007	ns	0.0
33	0.14 $\pm$ 0.081	-0.24 $\pm$ 0.073	0.57	0.00 $\pm$ 0.11	-0.30 $\pm$ 0.11	0.48	-0.06 $\pm$ 0.041	ns	22.9
39	-0.04 $\pm$ 0.022	ns	–	-0.05 $\pm$ 0.002	ns	–	-0.03 $\pm$ 0.061	ns	0.0
43	0.01 $\pm$ 0.152	-0.35 $\pm$ 0.15	0.41	-0.05 $\pm$ 0.19	-0.46 $\pm$ 0.19**	0.45	0.17 $\pm$ 0.069	ns	29.7
46	0.01 $\pm$ 0.030	-0.16 $\pm$ 0.027	0.83	-0.02 $\pm$ 0.43	-0.25 $\pm$ 0.058	0.70	0.34 $\pm$ 0.70	ns	14.9

1 ns: not significant ( $p > 0.05$ ); \* Grazing saturation: 3-point method used (Gallegos 1989); \*\* One outlier removed

**Table 4.** Dilution grazing experiments. Instantaneous growth rates without nutrients ( $\mu$ ;  $d^{-1}$ ) and mortality rates ( $m$ ;  $d^{-1}$ ) for the different planktonic groups considered. Determination coefficient for the regression analysis is also provided. Positive  $m$  values indicate positive slope.

Plank group		St 2	St 3	St 4	St 5	St 6	St 9	St 12	St 15	St 18	St 20	St 23	St 27	St 33	St 39	St 43	St 46
Diatoms	$\mu$	-0.10	0.83	-0.21	-0.41	0.02	0.82	1.49	0.24	-0.38	0.21	-0.03	-0.36	-0.20	-0.08	0.05	0.17
	$m$	ns	ns	ns	ns	ns	1.16	ns	1.08	ns	ns	ns	ns	0.68	ns	ns	0.72
	$r^2$						0.91		0.86					0.92	0.66		0.93
Nanoflagellates	$\mu$	-0.08	0.69	-0.22	0.58	0.08	0.99	0.87	0.62	-0.06	-0.42	-0.06	-0.62	0.62	0.12	-0.18	-0.61
	$m$	ns	ns	1.19	ns	-0.33	ns	ns	1.59	ns	ns	ns	ns	1.38	0.95	0.66	1.05
	$r^2$			0.91		0.96			0.96					0.94	0.81	0.77	0.79
<i>P. pouchetii</i>	$\mu$	nd	nd	-0.47	0.95	-3.97	1.20	0.87	-0.02	0.44	-0.37	0.04	0.11	-0.62	-0.21	-0.25	0.19
	$m$			ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	1.19
	$r^2$																0.93
< 20 $\mu\text{m}$ dinofl.	$\mu$	-0.29	-0.17	-0.48	-1.39	-0.77	0.83	0.32	-0.49	-0.19	-0.52	0.01	-0.78	-0.23	-0.55	-0.35	-0.29
	$m$	ns	ns	1.14		0.94	0.48	1.28	ns	ns	ns	ns	ns	ns	ns	0.93	ns
	$r^2$			0.72		0.86	0.79	0.96								0.77	
> 20 $\mu\text{m}$ dinofl.	$\mu$	0.96	1.53	-0.79	-0.37	-0.30	0.31	1.19	-0.48	0.02	0.26	0.52	0.32	-0.23	-0.02	-0.38	-0.50
	$m$	-0.53	-0.71	ns	ns	ns	ns	ns	ns	ns	ns	2.08	ns	ns	ns	ns	ns
	$r^2$	0.78	0.88									0.88					
Mixo. dinofl.	$\mu$	0.26	0.10	0.32	-0.14	-0.61	-0.04	0.63	-0.17	0.06	1.30	0.12	-0.75	0.21	0.24	-0.03	0.03
	$m$	ns	ns	-0.65	ns	ns	ns	ns	ns	ns	-0.97	ns	ns	ns	ns	0.47	ns
	$r^2$			0.72							0.90					0.87	
Het. dinofl.	$\mu$	1.17	0.28	0.29	-1.04	-0.22	1.88	1.02	-0.34	-0.01	0.19	0.47	0.05	0.12	0.23	0.54	-0.06
	$m$	-0.58	ns	-1.39	ns	ns	ns	ns	ns	ns	ns	1.00	ns	-0.60	ns	ns	ns
	$r^2$	0.79		0.90								0.76		0.80			

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

< 20 μm ciliates	μ	-0.12	0.25	0.70	1.21	-0.61	1.03	0.95	-0.54	0.15	-0.30	0.50	0.10	-0.16	0.13	0.12	-0.16
	<i>m</i>	ns	ns	-0.79	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
	<i>r</i> <sup>2</sup>			0.69													
> 20 μm ciliates	μ	-0.08	-0.17	0.03	-0.71	0.26	-1.05	0.17	-1.71	-0.45	-0.44	-0.73	-1.26	-0.14	0.06	-0.20	-1.07
	<i>m</i>	ns	ns	-0.70	ns	ns	ns	ns	0.55	ns	ns	ns	ns	ns	ns	ns	ns
	<i>r</i> <sup>2</sup>			0.74					0.84								

ns: not significant regression equation (p > 0.05); nd: not determined because not enough cells;

## Figure legends

**Figure 1.** Map of the surveyed area indicating the sampling stations.

**Figure 2.** *Phaeocystis pouchetii* biomass as related to latitude.

**Figure 3.** The quotient heterotrophic carbon / autotrophic carbon as a function of latitude. Greenland Sea and Coastal stations are indicated, the rest of stations correspond to Arctic Ocean open waters.


**Figure 4.** Relationship between instantaneous phytoplankton growth rates in the un-amended (without added nutrients) bottles ( $d^{-1}$ ) and chlorophyll *a* concentration ( $\mu g$  Chl *a*  $l^{-1}$ ).

**Figure 5.** Comparison of instantaneous phytoplankton growth rates (from total Chl *a* analysis) in bottles amended (with added nutrients) and un-amended. The discontinuous line represents the 1:1.

**Figure 6.** Example of grazing dilution plots for the microbial components of the planktonic community of stations 4 (left) and 33 (right).

**Figure 7.** Simulated outcome of a dilution grazing experiment using the abundance and growth rates of  $< 20 \mu m$  ciliates in St. 4 as grazers and the abundance of nanoflagellates as prey. See text for further details.





183x228mm (300 x 300 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Fig. 2

209x297mm (200 x 200 DPI)






Fig. 4

209x297mm (200 x 200 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Fig. 5

209x297mm (200 x 200 DPI)




Fig. 6

209x297mm (200 x 200 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





Fig. 7


209x297mm (200 x 200 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51



<http://mc.manuscriptcentral.com/jplankt>

km








Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7

**Web Annex 1.** Species composition and abundance of the plankton at the different stations sa

Station	2	2	3	3	4
	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>
<b>Diatoms</b>					
<i>Chaetoceros spp.</i>	1224		526		80959
<i>Corethron hystrix</i>	0		0		11497
<i>Eucampia zodiacus</i>	0		0		1770
<i>Fragilariopsis spp.</i>	280		20		20
<i>Nitzschia spp.</i>	0		20		60
<i>Rhizosolenia spp.</i>	10		10		60
<i>Skeletonema spp.</i>	0		0		130
<i>Thalassiosira spp.</i>	955		445		20
Other centric diatoms	220		20		10
Pennated diatoms	0		0		0
<b>Flagellates</b>					
<i>Phaeocystis pouchetii</i>	0		0		0
<i>Choanoflagellidea</i>	0		0		212940
<i>Dictyocha sp.</i>	35		5		10
Other nanoflagellates	29250		4500		10
<b>Dinoflagellates</b>					
<i>Amphidinium spp.</i>	65		35		35
<i>Ceratium spp.</i>	10		0		5
<i>Dinophysis spp.</i>	0		0		0
<i>Gyrodinium spp.</i>	85		15		45
<i>Katodinium glaucum</i>	70		30		15
<i>Micracanthodinium claytoni</i>	20		20		95
<i>Prorocentrum balticum</i>	70		10		0
<i>Protooperidinium bipes</i>	85		15		20
<i>Protooperidinium spp.</i>	25		5		10
<i>Torodinium robustum</i>	50		10		10
Dinoflagellates < 20 µm	1644		274		2719.5
Dinoflagellates > 20 µm	479.5		68.5		490
<b>Ciliates</b>					
<i>Didinium sp.</i>	80		0		15
<i>Laboea sp.</i>	545		65		5
<i>Mesodinium spp.</i>	55		35		5
<i>Strombidium spp.</i>	505		65		680
Tintinnids	0		0		5
< 20 µm ciliates	3699		548		5
> 20 µm ciliates	420		70		240

mpled. SE corresponde to the Standard Error of 2 replicate samples. Sizes correspo

	<b>4</b>	<b>5</b>	<b>5</b>	<b>6</b>	<b>6</b>	<b>9</b>
SE	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>	
	180	1630	530	2875	465	50
	0	0	0	0	0	0
	0	35	15	0	0	3600
	130	10545	1110	1330	10	1820
	225	41850	15750	15300	900	28575
	0	0	0	0	0	0
	0	0	0	0	0	0
	90	10405	1325	6930	550	10195
	0	30	10	125	5	5
	5	12375	3825	2025	1125	9450
	30758	1138785	129735	4947228	934092	357492
	8649	0	0	0	0	0
	20	75	25	35	5	35
	2366	262353	60543	227757	49011	170097
	10	15	5	240	80	10
	10	0	0	5	5	0
	5	10	10	5	5	5
	130	155	55	705	95	50
	5	20	10	370	130	0
	10	30	20	20	0	15
	35	7425	1125	4050	900	2925
	5	888	222	20	10	230
	5	25	5	40	10	5
	25	35	15	55	5	5
	388.5	45675	11025	33525	14175	10800
	20	550	110	1835	455	335
	5	20	0	15	5	5
	50	0	0	0	0	0
	0	0	0	0	0	0
	304.5	65	15	95	5	0
	10	0	0	0	0	0
	301	686.5	203.5	6555	215	205
	100.5	530	140	330	90	130



standard Error of 2 replicate samples. Sizes correspond to equivalent spherical diam

	<b>9</b>	<b>12</b>	<b>12</b>	<b>15</b>	<b>15</b>	<b>18</b>
SE	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>	
9	30	3115	805	1845	325	5
10	0	55	5	110	60	0
11	450	10	10	30	30	0
12	760	370	50	1080	260	135
13	2475	9450	450	27000	5850	170
14	0	0	0	0	0	0
15	0	0	0	1575	675	0
16	195	1770	410	15935	675	849.5
17	5	10	10	242	232	5
18	1350	900	450	3150	900	479.5
19						
20						
21						
22						
23	23064	1326180	184512	3992955	389205	268119
24	0	0	0	49011	14415	0
25	25	10	0	5	5	35
26	14415	194553	4641	224874	23064	605430
27						
28						
29						
30	10	60	20	120	20	390
31	0	0	0	0	0	0
32	5	0	0	15	5	0
33	10	145	15	790	170	935
34	0	120	0	120	20	685
35	5	20	10	80	20	10
36	225	110	50	3825	225	479.5
37	30	40	10	105	25	890.5
38	5	5	5	70	40	10
39	5	15	5	10	10	5
40						
41	1800	8325	225	18000	450	98022
42	45	435	75	1695	405	2429
43						
44						
45						
46	5	5	5	10	0	10
47	0	0	0	45	5	1400
48	0	0	0	0	0	0
49	0	65	15	145	35	35
50	0	0	0	5	5	5
51	55	1040	70	2555	325	1465
52	10	140	40	810	370	760
53						
54						
55						
56						
57						
58						
59						
60						

	<b>18</b>	<b>20</b>	<b>20</b>	<b>23</b>	<b>23</b>	<b>27</b>
SE	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>	
5	9960	180	149858	4990	27281.5	
0	440	40	1600	280	810	
0	25	25	60	40	25	
15	1280	540	5148.5	901.5	1030	
90	57825	4275	380556	46128	533355	
0	0	0	0	0	10	
0	1575	225	2475	225	0	
315.5	10530	1310	79785	2585	12135	
5	10	10	21190	2240	8365	
68.5	4725	1125	6300	2250	450	
158565	2485146	403620	5420040	576600	20181	
0	0	0	0	0	0	
5	15	5	10	10	0	
282534	804186	50586	323880	41262	149916	
110	230	10	140	20	80	
0	0	0	0	0	0	
0	0	0	0	0	0	
167	195	65	445	35	210	
274	220	80	460	100	190	
0	10	10	10	10	30	
68.5	2475	225	14175	4725	9900	
68.5	100	40	70	30	140	
0	60	10	35	5	45	
5	120	40	120	20	60	
11532	24750	450	20025	1125	25425	
117	1065	165	990	290	950	
0	0	0	10	0	20	
130	0	0	15	5	5	
0	0	0	0	0	0	
25	180	0	120	30	65	
5	0	0	0	0	0	
132	4650	310	2636.5	72.5	400	
140	720	80	1000	120	250	

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

	<b>27</b>	<b>33</b>	<b>33</b>	<b>39</b>	<b>39</b>	<b>43</b>
SE	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>	SE	Cells L <sup>-1</sup>	
	7434.5	2610	190	2345	675	1350
	150	220	40	75	15	0
	5	5	5	15	5	0
	50	450	110	300	20	890
	72075	25650	2250	9675	225	9000
	10	25	15	10	10	0
	0	0	0	0	0	0
	1195	11000	1750	4760	340	15885
	225	1660	250	2410	470	255
	0	4275	225	2475	225	0
	2883	1588533	850485	360375	8649	8014740
	0	0	0	0	0	900
	0	60	20	20	10	285
	11532	194553	24189	154599	13965	207801
	20	140	20	65	15	90
	0	0	0	0	0	5
	0	10	10	0	0	35
	50	215	25	180	30	150
	10	410	30	110	10	320
	20	25	15	20	0	75
	900	2250	450	45	5	3825
	20	70	50	10	0	120
	5	65	5	25	15	55
	0	160	20	15	5	45
	1575	16200	2250	16875	675	15075
	20	1745	345	1040	150	3340
	0	0	0	0	0	30
	5	40	0	0	0	110
	0	0	0	0	0	120
	25	310	80	155	35	225
	0	15	15	0	0	0
	120	1905	145	725	115	1055
	50	520	80	150	30	280

	<b>43</b>	<b>46</b>	<b>46</b>	
SE	Cells L <sup>-1</sup>	SE	verage cell size (ESD; μm)	
9	260	350	250	21.2
10	0	0	0	44.2
11	0	0	0	16
12	430	240	240	8.6
13	2250	5175	675	4.2
14	0	0	0	36
15	0	0	0	12.4
16	1035	11475	225	18
17	65	725	205	24.24
18	0	450	0	9.6
19				
20				
21				
22				
23	1210860	4050615	1484745	3.8
24	900	1350	1350	4
25	65	630	110	24.4
26	34821	284559	5991	4.04
27				
28				
29				
30	20	300	20	13.8
31	5	0	0	52.2
32	5	15	5	32.8
33	20	370	30	32.4
34	60	370	10	15
35	25	260	20	20
36	225	3825	675	19.4
37	20	50	20	18.8
38	5	35	5	36.6
39	15	40	10	27.6
40				
41	1575	12375	2925	
42	680	1510	200	
43				
44				
45	10	45	5	28.8
46	20	20	0	71
47	20	20	20	21
48	95	440	100	33
49	0	5	5	57.2
50	55	2735	195	
51	40	400	20	
52				
53				
54				
55				
56				
57				
58				
59				
60				