



HAL
open science

The yeast ABC transporter Pdr18 (ORF YNR070w) controls plasma membrane sterol composition, playing a role in multidrug resistance

Tânia Rodrigues Cabrito, Miguel Cacho Teixeira, Ashutosh Singh, Rajendra Prasad, Isabel Sá-Correia

► To cite this version:

Tânia Rodrigues Cabrito, Miguel Cacho Teixeira, Ashutosh Singh, Rajendra Prasad, Isabel Sá-Correia. The yeast ABC transporter Pdr18 (ORF YNR070w) controls plasma membrane sterol composition, playing a role in multidrug resistance. *Biochemical Journal*, 2011, 440 (2), pp.195-202. 10.1042/BJ20110876 . hal-00642839

HAL Id: hal-00642839

<https://hal.science/hal-00642839>

Submitted on 19 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **The yeast ABC transporter Pdr18 (ORF YNR070w) controls plasma membrane sterol**
2 **composition, playing a role in multidrug resistance**

3 Tânia R Cabrito^{1,2}, Miguel C Teixeira^{1,2}, Ashutosh Singh³, Rajendra Prasad³ and Isabel
4 Sá-Correia^{1,2*}

5 ¹IBB - Institute for Biotechnology and Bioengineering, Centre for Biological and
6 Chemical Engineering, Instituto Superior Técnico, 1049-001 Lisboa, Portugal.

7 ²Department of Bioengineering, Instituto Superior Técnico, Technical University of
8 Lisbon, 1049-001 Lisboa, Portugal.

9 ³ Membrane Biology Laboratory, School of Life Sciences, Jawaharlal Nehru University,
10 New Delhi 110067, India.

11
12
13
14 Corresponding author: Prof. Isabel Sá-Correia, IBB - Institute for Biotechnology and
15 Bioengineering, Biological Sciences Research Group, Centro de Engenharia Biológica e
16 Química, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal; e-mail
17 isacorreia@ist.utl.pt; Tel. +351-218417682; Fax +351-218419199.

18
19
20 **Short title:** The MDR transporter Pdr18 controls plasma membrane sterol composition

21
22 **Keywords:** multidrug resistance, ergosterol homeostasis, *Saccharomyces cerevisiae*,
23 pleiotropic drug resistance transporters, herbicide resistance, PDR network
24

25

26 **ABSTRACT**

27

28 The action of multidrug efflux pumps in multidrug resistance (MDR) acquisition
29 has been proposed to partially depend on the transport of physiological substrates
30 which may indirectly affect drug partition and transport across cell membranes.

31

32 In this work, *PDR18* gene (ORF *YNR070w*), encoding a putative pleiotropic drug
33 resistance (PDR) transporter of the ATP-binding cassette superfamily, was found to
34 mediate plasma membrane sterol incorporation in yeast. Pdr18 physiological role is
35 demonstrated to affect plasma membrane potential and is proposed to underlie its
36 action as a MDR determinant, conferring resistance to the herbicide 2,4-D. The action
37 of Pdr18 in yeast tolerance to 2,4-D, which was found to contribute to reduce [¹⁴C]-2,4-
38 D intracellular accumulation, may be indirect, given the observation that 2,4-D
39 exposure deeply affects the sterol plasma membrane composition, this effect being
40 much stronger in $\Delta pdr18$ background. *PDR18* activation under 2,4-D stress is regulated
41 by the transcription factors Nrg1, controlling carbon source availability and stress
42 response, and, less significantly, Yap1, involved in oxidative stress and MDR, and Pdr3,
43 a key regulator of the yeast PDR network, consistent with a broad role in stress
44 defence. Altogether, our results suggest that Pdr18 plays a role in plasma membrane
sterol incorporation, this physiological trait contributing to a MDR phenotype.

44

THIS IS NOT THE VERSION OF RECORD - see doi:10.1042/BJ20110876

Accepted Manuscript

45

46 **INTRODUCTION**

47 Multidrug resistance (MDR), the ability to acquire simultaneous resistance
48 to unrelated chemical compounds, is a widespread phenomenon that often
49 results from the activity of multidrug efflux pumps of the ATP-Binding
50 Cassette (ABC) Superfamily and of the Major Facilitator Superfamily (MFS). These
51 transporters are proposed to actively extrude or compartmentalize a wide range of
52 chemically and structurally disparate drugs and other xenobiotics, thus
53 providing protection from these compounds. However, the prevalence and
54 apparent redundancy of so many MDR transporters, which protect the cell
55 against toxic compounds that are not present in its natural environment, has
56 led to speculation concerning a natural physiological function [1-3].

57 In recent years, a physiological role for many of these transporters in the model
58 eukaryote *Saccharomyces cerevisiae* has been proposed. Among MFS-MDR
59 transporters, for example, Tpo1-4 and Qdr3 were suggested to mediate the
60 export of polyamines [4, 5], while Qdr2 was proposed to catalyse K⁺ influx [6]. These
61 results suggest that the chemoprotection role of these MDR transporters may come,
62 at least in some cases, as a result of their influence on plasma membrane
63 potential and/or ΔpH control, which in turn can alter the partitioning and
64 accumulation of drugs [2]. On the other hand, most
65 of the yeast pleiotropic drug resistance (PDR) transporters, of the ABC
66 superfamily, characterized as multidrug resistance determinants, have been
67 associated to the control of lipid incorporation into cell membranes [3]. Interestingly,
68 Pdr5 and its homologs from *Candida albicans*, Cdr1 and Cdr2, were shown to function
69 as phospholipid translocators [7]. Pdr10 and Pdr15 were also implicated in membrane
70 lipid organization [8, 9] and Aus1 and Pdr11 were found to mediate nonvesicular
71 movement of plasma membrane sterol to the endoplasmic reticulum in *S. cerevisiae*,
72 facilitating exogenous sterol uptake [10]. The deletion of yet another *S. cerevisiae*
73 transporter encoding gene, *PDR16*, which is regulated by the PDR network
74 transcription factor Pdr1, also leads to changes in yeast plasma membrane sterol
75 composition [11]. Altogether, these results have been suggested to imply that the
76 multidrug resistance phenotype conferred by PDR transporters may occur, to some
77 extent, due to the changes they impose in the composition of all membranes, which in
78 turn may affect drug partition or transport. This hypothesis is consistent with the fact
79 that *ERG2*, *ERG4* and *ERG5* genes, encoding the final enzymes of the ergosterol
80 biosynthetic pathway, were found to be determinants of multidrug resistance [12].

81 In this study, the functional analysis of *PDR18* (ORF *YNR070w*), encoding a
82 putative yeast PDR transporter of the plasma membrane, was undertaken. Based on
83 the fact that *PDR18* is up-regulated (1.6 fold) in yeast cells challenged by the herbicide
84 2,4-dichlorophenoxyacetic acid (2,4-D), as previously revealed by microarray analysis
85 [13], *PDR18* gene was found to confer yeast resistance to the auxin-like herbicides 2,4-
86 D and MCPA, among other xenobiotic compounds. Our previous studies on the yeast
87 adaptive response mechanisms to the herbicide 2,4-D (reviewed in [14]) indicate that
88 the expression of the drug:H⁺ antiporter Tpo1, of the ABC transporter Pdr5, and of the
89 *Arabidopsis thaliana* Tpo1 homolog At5g13750, confers 2,4-D resistance in *S.*
90 *cerevisiae* and leads to a lower intracellular accumulation of the herbicide [15, 16]. In
91 this work, the role and regulation of *PDR18* expression in yeast resistance to 2,4-D is

92 scrutinized. Given the previous implication of PDR transporters in phospholipid and
93 ergosterol, the role of Pdr18 in yeast plasma membrane lipid composition was studied.
94 *PDR18* expression was found to affect plasma membrane potential and sterol
95 composition, allowing us to propose Pdr18 as mediator of non-vesicular ergosterol
96 transport into the plasma membrane, this physiological role being, at least partially,
97 responsible for the observed multidrug resistance phenotype.
98
99

100 **EXPERIMENTAL**

101 **Strains, plasmids and growth conditions.** The parental *Saccharomyces cerevisiae* strain
102 BY4741 (*MATa*, *his3Δ1*, *leu2Δ0*, *met15Δ0*, *ura3Δ0*) and the derived single deletion
103 mutants *Δpdr18*, *Δnrg1*, *Δyap1* and *Δpdr3* used in this study, were obtained from the
104 EUROSCARF collection. Plasmids pRS416_ *PDR18* (*EUROSCARF*), expressing the *PDR18*
105 gene from its natural promoter, pRS416_ *PDR18Δp* and the cloning vector pRS416
106 (*EUROSCARF*) were also used. Plasmid pRS416_ *PDR18Δp* was obtained through site-
107 directed mutagenesis of pRS416_ *PDR18* using the QuikChange® XL Site-Directed
108 Mutagenesis Kit (Stratagene), and exhibit a substitution of 2 nucleotides in the
109 putative Nrg1 binding site in the *PDR18* promoter region. The pair of mutagenic
110 oligonucleotides used for this procedure were 5'
111 GTTCGTTTCATATGATTGCTTAGGTACCATTC 3' and its complementary sequence 5'
112 GAATGGTACCTAAGCAATATGAAACGAAAC 3', in which the replacement nucleotides
113 are underlined.

114 Yeast cells were cultivated at 30 °C with 250 rpm agitation in minimal growth
115 medium MM4, pH 3.5 (adjusted with HCl), containing (per litre): 1.7 g yeast nitrogen
116 base without amino acids or (NH₄)₂SO₄ (Difco), 20 g glucose (Merck), 2.65 g (NH₄)₂SO₄
117 (Merck), 20 mg L-histidine (Merck), 20 mg L-methionine (Merck), 60 mg L-leucine
118 (Sigma) and 20 mg L-uracil (Sigma). Solid medium, pH 4.0 (adjusted with HCl), was
119 prepared by adding 20 gL⁻¹ agar (Iberagar). Cells harbouring pRS416 or derived
120 plasmids were grown in the same media conditions and without uracil (MM4-U) to
121 assure plasmid segregational stability.

122 **Susceptibility assays.** The susceptibility of the parental strain BY4741 and *Δpdr18*
123 deletion mutant to toxic concentrations of 2,4D was assessed by comparing their
124 growth curves or growth in spot assays in MM4 medium supplemented or not with
125 inhibitory concentrations of 2,4-D (0.45mM in liquid medium and 1 - 2.5mM in solid
126 medium). Cell suspensions used for the spot assays were prepared as described
127 previously [5]. Besides 2,4-D, other chemical stress inducers (obtained from Sigma)
128 were tested in the specified concentration ranges: the herbicides MCPA (2-methyl-4-
129 chlorophenoxyacetic acid) (1 – 1.5 mM) and barban (0.08 – 0.1 mM); 2,4-
130 dichlorophenol, the 2,4-D degradation intermediate 2,4-DCP (0.5 – 1 mM); the
131 fungicides benomyl (17 – 35 mgL⁻¹) and mancozeb (1 – 1.2 mgL⁻¹); and salts of several
132 toxic metal cations, namely, CdSO₄ (0.025 – 0.04 mM), CuSO₄ (0.5 – 1 mM), MnSO₄ (5 –
133 10 mM), ZnSO₄ (5 – 10 mM), Al₂(SO₄)₃ (0.5 – 1 mM), TiCl₃ (0.1 – 0.5 mM), CoSO₄ (1.5 –
134 2 mM) and PbSO₄ (1.5 – 2 mM).

135 ***PDR18* gene expression assays.** The changes registered in the transcript levels from
136 the *PDR18* gene in BY4741 and in derived mutants *Δpdr18*, *Δnrg1*, *Δyap1* and *Δpdr3*,
137 upon yeast exposure to 0mM, 0.3mM or 0.45mM of 2,4-D were assessed by real time
138 RT-PCR. RNA extraction from yeast cells were carried out as described previously [15].
139 The used RT-PCR protocol followed the manufacturer's instructions and was described
140 elsewhere [5]. Primers for the amplification of the *PDR18* and *ACT1* cDNA were
141 designed using Primer Express Software (Applied Biosystems) and are 3'-
142 TTGGCAAGCCGGATCTGT-5', 3'- CCACGCGATTGGGAAT -5' and 3'-
143 CTCCACCACTGCTGAAAGAGAA -5', 3'- CCAAGGCGACGTAACATAGTTTT -5', respectively.
144 The RT-PCR reaction was carried out using a thermal cycler block (7500 Real-Time PCR
145 System - Applied Biosystems). The *ACT1* mRNA level was used as the internal control.

146 The relative values obtained for unstressed conditions were set as 1 and the remaining
147 values are relative to that value.

148 **[¹⁴C]-2,4-D accumulation assays.** The intracellular accumulation ratio of [¹⁴C]-labelled
149 2,4-D in the parental strain BY4741 and the derived deletion mutant $\Delta pdr18$ was
150 assessed as described previously [16].

151 **Plasma membrane potential ($\Delta\Psi$) estimation.** To estimate the differences in BY4741
152 and derived mutant $\Delta pdr18$ plasma membrane potential two methods were used: the
153 [¹⁴C]-Methylamine uptake assay [17] and the 3-3'-dihexyloxacarbocyanine iodide
154 (DiOC₆(3)) accumulation assay [18].

155 The uptake of [¹⁴C]-methylamine in the parental strain BY4741 and the mutant
156 strain $\Delta pdr18$ was monitored as described previously [5, 17].

157 To estimate DiOC₆(3) fluorescence, cells were harvested as described above,
158 resuspended in MES-glucose buffer (10 mM MES, 0.1 mM MgCl₂ and 20 gL⁻¹ glucose,
159 pH 6), supplemented with DiOC₆(3) (Molecular probes) at a final concentration of
160 0.25nM and incubated in the dark for 30 min at 30°C with orbital agitation. After
161 centrifugation, cells were immediately observed with a Zeiss Axioplan microscope
162 equipped with adequate epifluorescence filters (BP450-490 and LP520). And
163 fluorescence emission was collected with a CCD camera (Cool SNAPFX, Roper Scientific
164 Photometrics). Bright-field images for determination of $\Delta\Psi$ were obtained
165 concurrently and recorded at 1 min intervals, each experiment being finished within 15
166 min. The images were analysed using MetaMorph 3.5. The fluorescence images were
167 background corrected using dark-current images. The intensity values were calculated
168 for a minimum of 80 cells per experiment. Individual cells were selected using regions
169 of interest obtained from bright-field images recorded before or after the experiment.
170 The value of fluorescence intensity emitted by each cell was obtained pixel-by-pixel in
171 the region of interest. Fluorescence levels given by the software were expressed as a
172 percentage.

173 **Plasma membrane sterol composition assessment** Total cell membranes (CMs) were
174 extracted and prepared from yeast cells grown in MM4 medium (pH 3.5) and
175 harvested in the exponential growth phase. For studying the effect of 2,4-D,
176 exponential cells grown in MM4 medium were transferred to fresh medium
177 supplemented with 0.45mM of 2,4-D, and grown for 1h at 30 °C. Cells were harvested
178 by centrifugation, resuspended in homogenization buffer containing 50 mM Tris pH
179 7.5, 2.5 mM EDTA and a protease inhibitor cocktail (1 mM PMSF, 1 µg/ml leupeptin,
180 pepstatin A and aprotinin) and broken by vortexing with glass beads (Glaperlon 0.40-
181 0.60 mm). The CM were recovered by centrifugation at 1000 g to remove unbroken
182 cells and finally the CM were pelleted by ultracentrifugation at 25000 rpm for 1 hour.
183 The CM were resuspended in a buffer containing 20 mM Tris pH 7.5, 150 mM NaCl,
184 20% glycerol and protease inhibitors at the concentrations mentioned above. Plasma
185 membrane (PM) fractions were obtained from CM fractions by sucrose gradient
186 centrifugation as described by Monk *et al.* (1991) [19]. The obtained CM and PM
187 protein concentrations, measured by the BCA test, ranged between 8 – 19 µg/µl and 3
188 – 9 µg/µl respectively. Equal amounts of PMs were used for lipid extraction from each
189 of the yeast strains using the method described by Bligh and Dyer (1959) with slight
190 modifications [20]. Sterols were extracted and analyzed using the method described
191 previously with slight modifications [21]. The saponified lipids were re-extracted using
192 5 ml hexane and vortexed several times. 1 ml of water was added to separate the

193 organic phase, which was then dried in N₂ at 60°C. The extracted sterols were
194 derivatized using 100 µl N,O-Bis(trimethylsilyl) trifluoroacetamide with trimethyl-
195 chlorosilane BSTFA/TMCS (Sigma) at 80°C for 1 hr in N₂. The derivatized sterols were
196 then analyzed using GCMS (Schimadzu QP2010 Plus, Japan) and aDB5-MS column 60 m
197 X 0.2 mm, film thickness 0.20 µm. The carrier gas was helium with a flow rate of
198 1ml/min and a pressure of 80.8 kPa. Initial column temperature of 120°C was held for
199 1 min. then programmed at 120°C to 250°C at 5°C/min where it was held for 30 min. 1
200 µl injection was made using a split ratio of 10. Injection temperature was 300°C. The
201 total ion mass spectra were recorded in the mass range m/z 40-650 at the scan rate of
202 1 s/scan. The interface and detector temperature was 300°C. Peak identification was
203 based on relative retention time and total ion mass spectral comparison with an
204 external standard. The sterol standards were obtained from Sigma-Aldrich.

205
206

THIS IS NOT THE VERSION OF RECORD - see doi:10.1042/BJ20110876

Accepted Manuscript

207

208 **RESULTS**

209 **The ABC transporter *Pdr18*, encoded by ORF YNR070w, confers yeast**
210 **resistance to 2,4-D and other chemical stresses.** The susceptibility towards 2,4-D
211 imposed stress of the single deletion mutant $\Delta pdr18$ was found to be higher compared
212 to the parental strain, based on spot assays and liquid growth (Fig. 1). In the absence
213 of *Pdr18*, yeast cells become susceptible to 2,4-D, even when supplemented in
214 concentrations that hardly affect wild-type viability (Fig. 1A). For higher concentrations
215 of the herbicide, the expression of this transporter becomes essential for survival
216 under stress (Fig. 1A). *PDR18* deletion was also found to lead to a longer 2,4-D-induced
217 lag-phase, and also a reduced value of final biomass concentration (Fig. 1B). The
218 expression of *PDR18* from a centromeric plasmid was found to rescue the 2,4-D
219 susceptibility phenotype of $\Delta pdr18$ cells, to levels comparable to the parental strain,
220 while no effect was detected in the control cells harbouring the corresponding cloning
221 vector (Fig. 1C). The ability of this gene expression to confer resistance to other
222 pesticides and chemical compounds of agro-economical importance was further
223 analyzed and *PDR18* was also found to be a determinant of yeast resistance to the
224 herbicides MCPA and barban, to the 2,4-D degradation intermediate 2,4-
225 dichlorophenol (2,4-DCP), to the agricultural fungicide mancozeb, and to the metal
226 cations Zn^{2+} , Mn^{2+} , Cu^{2+} and Cd^{2+} (Fig. 2). No protection was conferred by *PDR18*
227 expression towards benomyl, Co^{2+} , Pb^{2+} , Al^{3+} and Tl^{3+} (data not shown).

228 ***PDR18* transcription is activated in response to 2,4-D imposed stress in a *Nrg1*-**
229 **, *Yap1*- and *Pdr3*-dependent manner.** A strong increase in the transcript levels of
230 *Pdr18* (~8 fold) was registered after one hour of exposure of an un-adapted *S.*
231 *cerevisiae* cell population to 0.3 mM of 2,4-D. This strong but transient increase was
232 followed by a rapid decrease of transcript levels to basal values as cells adapt to
233 growth in the presence of the herbicide. When exposed to a higher concentration of
234 2,4-D, 0.45 mM, *PDR18* transcriptional activation reached a maximum of up to 5-fold
235 after 4h of stress exposure, correlating with the longer duration of the lag-phase
236 imposed by this higher 2,4-D concentration tested (Fig 3). The fact that a higher
237 herbicide concentration apparently leads to a lower activation of *PDR18* may be due to
238 the fact that 0.45mM of 2,4-D induces viability loss, thus reducing the number of cells
239 in the studied population with the ability to generate a stress response [22].

240 The YEASTRACT database (www.yeasttract.com) [23, 24], was used to guide the
241 analysis of the transcriptional control underlying *PDR18* activation. Five transcription
242 factors were identified as documented *PDR18* regulators previously proven to bind to
243 its promoter region, and, thus, were selected as candidates for, directly or indirectly,
244 controlling 2,4-D-induced *PDR18* up-regulation (Fig. 4A). Of these, only *Nrg1*, *Pdr3* and
245 *Yap1* were examined in this study, because mutants deleted for the other two genes
246 are either unviable ($\Delta rap1$) or exhibits marked growth defects ($\Delta swi4$) (SGD,
247 *Saccharomyces* Genome Database – www.yeastgenome.com). The three tested
248 mutant strains, devoid of *PDR3*, *YAP1* or *NRG1*, exhibit nearly identical *PDR18* mRNA
249 basal levels, as registered in unstressed parental strain cells. However, the *PDR18* up-
250 regulation detected in the wild type cells after 4h of incubation with 2,4-D is abrogated
251 in $\Delta nrg1$ mutant or reduced in $\Delta yap1$ and $\Delta pdr3$ mutants (Fig. 4B), suggesting that
252 both *Pdr3* and *Yap1* transcription factors are necessary to assure full *PDR18* activation.

253 Since Nrg1 has been described as a transcriptional repressor, its action as an
254 activator of *PDR18* was hypothesized to be indirect. Nonetheless, this transcription
255 factor was previously demonstrated, through genome-wide screenings [25, 26], to
256 bind to the *PDR18* promoter region. Furthermore, according to the YEASTRACT
257 database, a potential Nrg1 binding site can be found in the *PDR18* promoter at position
258 -567. To evaluate whether or not the action of Nrg1 on *PDR18* transcriptional up-
259 regulation might be direct, site-directed mutagenesis was used to abrogate the
260 putative Nrg1 binding site found in the pRS416_*PDR18* expression plasmid. Both
261 pRS416_*PDR18* and pRS416_*PDR18* Δ p plasmids were transformed into Δ *pdr18* cells, so
262 that the genomic expression of *PDR18* from its natural promoter would not be
263 accounted for. RT-PCR was used to measure the *PDR18* transcript levels in these cells,
264 upon exposure to 0.45mM of 2,4-D. The abrogation of the Nrg1 binding site was seen
265 to have only a moderate effect on herbicide-induced *PDR18* up-regulation (Fig. 4C)
266 when compared to the full effect observed upon Nrg1 deletion. Altogether, these
267 results suggest that the action of Nrg1 on *PDR18* expression appears to be indirect in
268 this case.

269 **Role of *PDR18* expression in 2,4-D intracellular accumulation.** Given the
270 presumed role of Pdr18 as a plasma membrane multidrug resistance transporter that
271 confers resistance to yeast cells against 2,4-D imposed stress, the effect of *PDR18*
272 expression in the intracellular accumulation of [¹⁴C]-2,4-D was assessed. The
273 accumulation of [¹⁴C]-2,4-D in non-adapted yeast cells suddenly exposed to the
274 presence of 0.3 mM 2,4-D (at pH 3.5), which induces a mild growth inhibition in both
275 the parental strain and Δ *pdr18* cells (data not shown), is 2.5 fold higher in cells devoid
276 of *PDR18* than in parental cells (Fig. 5). This result strongly suggests that Pdr18 activity
277 increases yeast resistance towards 2,4-D by reducing the accumulation of the 2,4-D
278 anion within yeast cells, presumably by catalyzing the direct extrusion of the herbicide.

279 ***PDR18* deletion causes changes in yeast plasma membrane sterol
280 composition.** In the absence of 2,4-D supplementation, upon disruption of *PDR18*, a
281 nearly 2-fold accumulation of squalene and lanosterol, the precursors of ergosterol
282 biosynthetic pathway, and a 1,5 fold reduction of ergostatetraenol and ergosterol
283 content, the end products of the ergosterol biosynthetic pathways, in the plasma
284 membrane were detected (Fig. 6). Upon episomal complementation of *PDR18*, there
285 was a partial complementation of the Δ *pdr18* plasma membrane sterol composition
286 phenotype (Fig. 6).

287 The levels of ergosterol in the plasma membrane of *S. cerevisiae* BY4741
288 exposed to 2,4-D were found to decrease 1.5 fold compared with unstressed
289 conditions, while the levels of squalene increased 5.5 fold. This effect is exacerbated in
290 the absence of *PDR18* gene. Indeed, in Δ *pdr18* cells 2,4-D exposure leads to a 4.3 fold
291 increase in squalene and a 5.2 fold decrease in ergosterol plasma membrane
292 concentrations (Fig. 6). Remarkably, under 2,4-D challenge the relative abundance of
293 squalene becomes higher than the relative abundance of ergosterol in both wild type
294 and Δ *pdr18* backgrounds and the concentration of the other sterols detected in the
295 plasma membrane of unstressed yeast cells become undetectable (Fig. 6).

296 **Lower plasma membrane potential is observed in Δ *pdr18* cells.** The role of
297 *PDR18* expression in the maintenance of yeast membrane potential was also analyzed.
298 Yeast plasma membrane potential was first estimated based on the uptake of
299 methylammonium, a non-metabolizable ammonium analogue, whose influx is strongly

300 dependent on the maintenance of the transmembrane potential [27]. The deletion of
301 *PDR18* was found to decrease in around 60% the level of methylammonium uptake in
302 yeast cells (Fig. 7A). Consistent with these results, the fluorescence intensity levels of
303 cells loaded with the DiOC₆(3) probe, whose accumulation inside yeast cells is
304 dependent on the plasma membrane potential [18], is three fold higher in wild-type
305 cells than in $\Delta pdr18$ cells (Fig. 7B). Both methods indicate a strong depletion of the
306 plasma membrane potential in the absence of *PDR18*.
307

308

309 **DISCUSSION**

310 This study provides the first functional report on the uncharacterized yeast PDR
311 transporter, Pdr18, encoded by ORF *YNR070w*. Although this gene was not previously
312 characterized, a microarray analysis from our group showed that *PDR18* is up-
313 regulated in yeast cells exposed to inhibitory concentrations of the herbicide 2,4-D
314 [13]. Guided by this preliminary result, the current study provides evidence showing
315 that *PDR18* is a determinant of yeast resistance to 2,4-D, to MCPA, another auxin like
316 herbicide, and to several other unrelated chemical stresses, including, barban, an
317 herbicide of the carbanilate family, mancozeb, an agricultural fungicide and the soil
318 contaminant metals cadmium, copper, manganese and zinc. This study provides
319 evidence showing that *PDR18* gene expression decreases the intracellular
320 accumulation of radiolabelled 2,4-D. The intracellular accumulation pattern observed
321 for $\Delta pdr18$ deletion mutant, compared with the wild-type strain, is similar to the one
322 observed previously for $\Delta tpo1$ deletion mutant [16]. Interestingly, the *PDR18* homolog
323 in the plant model *Arabidopsis thaliana*, *AtPDR9*, was seen to confer 2,4-D resistance
324 in plant, also contributing to decreased 2,4-D accumulation in plant roots [28].

325 During 2,4-D induced lag-phase period, preceding exponential growth
326 resumption under herbicide stress, *PDR18* transcript levels were shown to increase
327 transiently. This fact, together with the reduction of the duration of the lag phase
328 induced by 2,4-D due to *PDR18* expression, indicates that the role of Pdr18 is
329 preponderant during the period of adaptation to the herbicide, while the effect over
330 the inhibition of specific growth exert by the herbicide is not significant. Consistent
331 with a broad role in stress defence, the transcriptional up-regulation of *PDR18* was
332 found to be partially reduced in mutants with either *PDR3* or *YAP1* genes deleted and
333 completely abolished in a mutant devoid of *NRG1*. The partial effect of Pdr3 in *PDR18*
334 activation resembles the effect described before exerted by Pdr3 over the
335 transcriptional up-regulation of *TPO1* under 2,4-D stress [15] and places *PDR18* within
336 the yeast PDR network. At the same time, the role of Yap1, the major regulator of *S.*
337 *cerevisiae* oxidative stress response, in 2,4-D induced *PDR18* up-regulation may
338 correlate with the observation that this herbicide exerts a pro-oxidant action in yeast
339 [22]. Moreover, Yap1 also plays a role in the control of multidrug resistance, regulating
340 the expression of at least ten other MDR proteins: the ABC drug efflux pumps Snq2,
341 Ycf1 and Pdr5 and the drug-H⁺ antiporters Flr1, Tpo1, Tpo2, Tpo4, Azr1, Yhk8 and Qdr3
342 (www.yeasttract.com) [5, 23, 24]. On the other hand, in the absence of *NRG1* the 2,4-D-
343 induced transcriptional up-regulation is completely abrogated. Consistent with the
344 notion that Nrg1 acts as a transcriptional repressor, the action of Nrg1 in this case is
345 proposed to be indirect, based on the fact that the abrogation of the Nrg1 binding site
346 in the *PDR18* promoter led only to a slight change in its herbicide-dependent
347 transcriptional up-regulation. The action of Nrg1 is likely to occur through the
348 regulation of other genes, possibly other transcription factors. Interestingly, the
349 transcript levels of the *NRG1* gene suffer a 5-fold increase in yeast cells exposed for 15
350 min to 0.3mM of 2,4-D, as described in a previous microarray analysis [13]. The same
351 global analysis suggests that yeast cells challenged with toxic concentrations of 2,4-D
352 experience a state of glucose and energy limitation, despite the saturating
353 concentrations of this preferential carbon source in the surrounding medium, which
354 could account for an Nrg1-mediated response [13].

355 Most significantly, in this work the deletion of *PDR18* in *S. cerevisiae* cells was
356 found to lead to an accumulation of the precursors of ergosterol biosynthetic pathway,
357 squalene and lanosterol, and to a decrease in the content of ergostatetraenol and
358 ergosterol, the end products of the ergosterol biosynthetic pathway, in yeast plasma
359 membrane. In the same conditions, no changes in the phospholipid composition of the
360 yeast plasma membrane were registered upon *PDR18* deletion (results not shown).
361 Although the exact role of Pdr18 in sterol homeostasis requires clarification, Pdr18 is
362 proposed to play a direct role in the incorporation of ergosterol in the plasma
363 membrane as part of the non-vesicular ER to plasma membrane ergosterol transport
364 mechanism [10]. Both in mammalian and yeast cells, newly synthesized
365 cholesterol/ergosterol has been shown to be transported from the endoplasmic
366 reticulum to the plasma membrane via two mechanisms: one dependent on vesicular
367 transport and the other dependent on ATP, but independent of vesicular transport
368 [29]. However, no specific transporter has been implicated so far in the mediation of
369 this non-vesicular ergosterol movement. Our results suggest that Pdr18 may
370 contribute to this important physiological function. Given this proposed physiological
371 role, the observed apparent inhibitory effect of *PDR18* deletion on sterol biosynthesis
372 could result from probing local sterol concentrations, thus influencing the activity of
373 ergosterol synthesizing enzymes, as suggested for Pdr16 [11].

374 The lipid composition of a cellular membrane has profound effects on its
375 biophysical properties that may affect a membrane's fusibility, including intrinsic
376 curvature, thickness, stiffness, and permeability [30-32]. Unlike intracellular
377 membranes, the yeast plasma membrane is highly enriched in ergosterol. In various
378 plant models, ergosterol induces changes in membrane potential [33, 34] and
379 modifications of H⁺ fluxes across the membranes [33, 35, 36], among other effects. A
380 low level of ergosterol leads to disruption of the membrane lateral order [37], which
381 results in membrane fluidization, compromising the physiological membrane potential.
382 Consistent with the depletion of ergosterol in the plasma membrane of $\Delta pdr18$ cells,
383 *PDR18* expression was also found to be essential in the maintenance of yeast plasma
384 membrane potential. Two probes were used to assess the differences between wild-
385 type and $\Delta pdr18$ plasma membrane potential to rule out the hypothesis that the
386 observed variation might result from the direct action of Pdr18 in the excretion of one
387 of the selected probes.

388 The action of Pdr18 in 2,4-D resistance can be explained in light of its
389 contribution to sterol homeostasis. It is interesting to see that exposure to the
390 herbicide 2,4-D leads to several changes in membrane sterol composition similar to
391 those caused by *PDR18* deletion, including a decrease in ergosterol and an increase in
392 squalene relative concentrations. These changes occurring under 2,4-D stress indicate
393 a possible action of the herbicide as an inhibitor of ergosterol biosynthesis or transport
394 into the plasma membrane and are consistent with the requirement for *PDR18*
395 expression and the observed *PDR18* up-regulation registered in this study.
396 Furthermore, in the absence of *PDR18* the effect of 2,4-D in the plasma membrane
397 sterol content is even more pronounced than in wild-type cells. Such a reduced
398 ergosterol content in $\Delta pdr18$ cells is likely to increase the permeability of the plasma
399 membrane towards 2,4-D and to affect the active export of 2,4-D to the outer medium,
400 through dedicated transporters, eventually including Tpo1, Pdr5 [15] and Pdr18 itself,

401 consistent with the observed increase accumulation of 2,4-D in yeast cells devoid of
402 *PDR18*.

403 Altogether, based on the results presented in this paper a physiological role for
404 Pdr18 in the control of sterol homeostasis, specifically, in maintaining ergosterol
405 physiological levels in the plasma membrane is proposed. Pdr18 role as a multidrug
406 resistance determinant is suggested to derive, at least partially, from its physiological
407 role, which is expected to affect drug partition and transport across cell membranes.
408 These results are expected to increase current knowledge on the action of this family
409 of transporters with impact in the design of strategies to deal with MDR. Given the
410 particular role of Pdr18 in pesticide resistance, these results may also guide the design
411 of new pesticide resistant crops of agro-economic interest.
412

413

414 **FUNDING**

415

416 This work was supported by FEDER and “Fundação para a Ciência e a
417 Tecnologia” (FCT) (Contracts PTDC/AGR-AAM/102967/2008 and PTDC/BIA-
418 MIC/72577/2006 and PhD grant SFRH/BD/40885/2007 to TRC). This work is also
419 supported in part by grants to R.P. from Department of Biotechnology
420 (BT/PR9100/Med/29/03/2007, BT/PR9563/BRB/10/567/2007,
421 BT/PR11158/BRB/10/640/2008, BT/PR13641/MED/29/175/2010). We acknowledge
422 Advanced Instrumentation Research Facility (AIRF), JNU for providing GC-MS.

423

424

THIS IS NOT THE VERSION OF RECORD - see doi:10.1042/BJ20110876

Accepted Manuscript

425

426 **REFERENCES**

- 427 1 Roepe, P. D., Wei, L. Y., Hoffman, M. M. and Fritz, F. (1996) Altered drug translocation
428 mediated by the MDR protein: direct, indirect, or both? *J. Bioenerg. Biomembr.* **28**, 541-555
- 429 2 Sá-Correia, I., Santos, S., Teixeira, M., Cabrito, T. and Mira, N. (2009) Drug:H⁺
430 antiporters in chemical stress response in yeast. *Trends Microbiol.* **17**, 22-31
- 431 3 Jungwirth, H. and Kuchler, K. (2006) Yeast ABC transporters-- a tale of sex, stress, drugs
432 and aging. *FEBS Lett.* **580**, 1131-1138
- 433 4 Tomitori, H., Kashiwagi, K., Asakawa, T., Kakinuma, Y., Michael, A. J. and Igarashi, K.
434 (2001) Multiple polyamine transport systems on the vacuolar membrane in yeast. *Biochem. J.*
435 **353**, 681-688
- 436 5 Teixeira, M. C., Cabrito, T. R., Hanif, Z. M., Vargas, R. C., Tenreiro, S. and Sá-Correia, I.
437 (2011) Yeast response and tolerance to polyamine toxicity involving the drug:H⁺ antiporter
438 Qdr3 and the transcription factors Yap1 and Gcn4. *Microbiology.* **157**, 945-956
- 439 6 Vargas, R. C., Garcia-Salcedo, R., Tenreiro, S., Teixeira, M. C., Fernandes, A. R., Ramos,
440 J. and Sá-Correia, I. (2007) *Saccharomyces cerevisiae* Multidrug Resistance Transporter Qdr2 Is
441 Implicated in Potassium Uptake, Providing a Physiological Advantage to Quinidine-Stressed
442 Cells. *Eukaryot. Cell.* **6**, 134-142
- 443 7 Smriti, Krishnamurthy, S., Dixit, B. L., Gupta, C. M., Milewski, S. and Prasad, R. (2002)
444 ABC transporters Cdr1p, Cdr2p and Cdr3p of a human pathogen *Candida albicans* are general
445 phospholipid translocators. *Yeast.* **19**, 303-318
- 446 8 Schuller, C., Mamnun, Y. M., Wolfger, H., Rockwell, N., Thorner, J. and Kuchler, K.
447 (2007) Membrane-active compounds activate the transcription factors Pdr1 and Pdr3
448 connecting pleiotropic drug resistance and membrane lipid homeostasis in *saccharomyces*
449 *cerevisiae*. *Mol. Biol. Cell.* **18**, 4932-4944
- 450 9 Rockwell, N. C., Wolfger, H., Kuchler, K. and Thorner, J. (2009) ABC transporter Pdr10
451 regulates the membrane microenvironment of Pdr12 in *Saccharomyces cerevisiae*. *J. Membr.*
452 *Biol.* **229**, 27-52
- 453 10 Li, Y. and Prinz, W. A. (2004) ATP-binding cassette (ABC) transporters mediate
454 nonvesicular, raft-modulated sterol movement from the plasma membrane to the
455 endoplasmic reticulum. *J. Biol. Chem.* **279**, 45226-45234
- 456 11 van den Hazel, H. B., Pichler, H., do Valle Matta, M. A., Leitner, E., Goffeau, A. and
457 Daum, G. (1999) *PDR16* and *PDR17*, two homologous genes of *Saccharomyces cerevisiae*,
458 affect lipid biosynthesis and resistance to multiple drugs. *J. Biol. Chem.* **274**, 1934-1941
- 459 12 Hillenmeyer, M. E., Fung, E., Wildenhain, J., Pierce, S. E., Hoon, S., Lee, W., Proctor, M.,
460 St Onge, R. P., Tyers, M., Koller, D., Altman, R. B., Davis, R. W., Nislow, C. and Giaever, G.
461 (2008) The chemical genomic portrait of yeast: uncovering a phenotype for all genes. *Science.*
462 **320**, 362-365
- 463 13 Teixeira, M. C., Fernandes, A. R., Mira, N. P., Becker, J. D. and Sá-Correia, I. (2006) Early
464 transcriptional response of *Saccharomyces cerevisiae* to stress imposed by the herbicide 2,4-
465 dichlorophenoxyacetic acid. *FEMS Yeast Res.* **6**, 230-248
- 466 14 Teixeira, M. C., Duque, P. and Sá-Correia, I. (2007) Environmental genomics:
467 mechanistic insights into toxicity of and resistance to the herbicide 2,4-D. *Trends Biotechnol.*
468 **25**, 363-370
- 469 15 Teixeira, M. C. and Sá-Correia, I. (2002) *Saccharomyces cerevisiae* resistance to
470 chlorinated phenoxyacetic acid herbicides involves Pdr1p-mediated transcriptional activation
471 of *TPO1* and *PDR5* genes. *Biochem. Biophys. Res. Commun.* **292**, 530-537
- 472 16 Cabrito, T. R., Teixeira, M. C., Duarte, A. A., Duque, P. and Sá-Correia, I. (2009)
473 Heterologous expression of a Tpo1 homolog from *Arabidopsis thaliana* confers resistance to
474 the herbicide 2,4-D and other chemical stresses in yeast. *Appl. Microbiol. Biotechnol.* **84**, 927-
475 936

- 476 17 Hoeberichts, F. A., Perez-Valle, J., Montesinos, C., Mulet, J. M., Planes, M. D., Hueso,
477 G., Yenush, L., Sharma, S. C. and Serrano, R. (2010) The role of K(+) and H(+) transport systems
478 during glucose- and H₂O₂-induced cell death in *Saccharomyces cerevisiae*. *Yeast*. **27**, 713-725
- 479 18 Madrid, R., Gomez, M. J., Ramos, J. and Rodriguez-Navarro, A. (1998) Ectopic
480 potassium uptake in *trk1 trk2* mutants of *Saccharomyces cerevisiae* correlates with a highly
481 hyperpolarized membrane potential. *J. Biol. Chem.* **273**, 14838-14844
- 482 19 Monk, B. C., Kurtz, M. B., Marrinan, J. A. and Perlin, D. S. (1991) Cloning and
483 characterization of the plasma membrane H(+)-ATPase from *Candida albicans*. *J. Bacteriol.* **173**,
484 6826-6836
- 485 20 Bligh, E. G. and Dyer, W. J. (1959) A rapid method of total lipid extraction and
486 purification. *Can. J. Biochem. Physiol.* **37**, 911-917
- 487 21 Sharma, M., Dhamgaye, S., Singh, A. and Prasad, R. (in press) Lipidome analysis reveals
488 antifungal polyphenol curcumin affects membrane lipid homeostasis: targets $\Delta 5,6$ desaturase
489 *ERG3* of *Candida albicans*. *Frontiers in Bioscience*
- 490 22 Teixeira, M. C., Telo, J. P., Duarte, N. F. and Sá-Correia, I. (2004) The herbicide 2,4-
491 dichlorophenoxyacetic acid induces the generation of free-radicals and associated oxidative
492 stress responses in yeast. *Biochem. Biophys. Res. Commun.* **324**, 1101-1107
- 493 23 Abdulrehman, D., Monteiro, P. T., Teixeira, M. C., Mira, N. P., Lourenco, A. B., dos
494 Santos, S. C., Cabrito, T. R., Francisco, A. P., Madeira, S. C., Aires, R. S., Oliveira, A. L., Sá-
495 Correia, I. and Freitas, A. T. (2011) YEASTRACT: providing a programmatic access to curated
496 transcriptional regulatory associations in *Saccharomyces cerevisiae* through a web services
497 interface. *Nucleic. Acids Res.* **39**, D136-140
- 498 24 Teixeira, M. C., Monteiro, P., Jain, P., Tenreiro, S., Fernandes, A. R., Mira, N. P.,
499 Alenquer, M., Freitas, A. T., Oliveira, A. L. and Sá-Correia, I. (2006) The YEASTRACT database: a
500 tool for the analysis of transcription regulatory associations in *Saccharomyces cerevisiae*.
501 *Nucleic. Acids Res.* **34**, D446-451
- 502 25 Harbison, C. T., Gordon, D. B., Lee, T. I., Rinaldi, N. J., Macisaac, K. D., Danford, T. W.,
503 Hannett, N. M., Tagne, J. B., Reynolds, D. B., Yoo, J., Jennings, E. G., Zeitlinger, J., Pokholok, D.
504 K., Kellis, M., Rolfe, P. A., Takusagawa, K. T., Lander, E. S., Gifford, D. K., Fraenkel, E. and Young,
505 R. A. (2004) Transcriptional regulatory code of a eukaryotic genome. *Nature*. **431**, 99-104
- 506 26 Lee, T. I., Rinaldi, N. J., Robert, F., Odom, D. T., Bar-Joseph, Z., Gerber, G. K., Hannett,
507 N. M., Harbison, C. T., Thompson, C. M., Simon, I., Zeitlinger, J., Jennings, E. G., Murray, H. L.,
508 Gordon, D. B., Ren, B., Wyrick, J. J., Tagne, J. B., Volkert, T. L., Fraenkel, E., Gifford, D. K. and
509 Young, R. A. (2002) Transcriptional regulatory networks in *Saccharomyces cerevisiae*. *Science*.
510 **298**, 799-804
- 511 27 van de Mortel, J. B., Mulders, D., Korthout, H., Theuvenet, A. P. and Borst-Pauwels, G.
512 W. (1988) Transient hyperpolarization of yeast by glucose and ethanol. *Biochim. Biophys. Acta*.
513 **936**, 421-428
- 514 28 Ito, H. and Gray, W. M. (2006) A gain-of-function mutation in the *Arabidopsis*
515 pleiotropic drug resistance transporter *PDR9* confers resistance to auxinic herbicides. *Plant*
516 *Physiol.* **142**, 63-74
- 517 29 Sullivan, D. P., Ohvo-Rekila, H., Baumann, N. A., Beh, C. T. and Menon, A. K. (2006)
518 Sterol trafficking between the endoplasmic reticulum and plasma membrane in yeast.
519 *Biochem. Soc. Trans.* **34**, 356-358
- 520 30 Hancock, J. F. (2006) Lipid rafts: contentious only from simplistic standpoints. *Nat. Rev.*
521 *Mol. Cell Biol.* **7**, 456-462
- 522 31 Mukherjee, S. and Maxfield, F. R. (2004) Membrane domains. *Annu. Rev. Cell Dev. Biol.*
523 **20**, 839-866
- 524 32 Dupont, S., Beney, L., Ferreira, T. and Gervais, P. (2011) Nature of sterols affects
525 plasma membrane behavior and yeast survival during dehydration. *Biochim. Biophys. Acta*.
526 **1808**, 1520-1528

- 527 33 Amborabe, B. E., Rossard, S., Perault, J. M. and Roblin, G. (2003) Specific perception of
528 ergosterol by plant cells. *C. R. Biol.* **326**, 363-370
- 529 34 Rossard, S., Luini, E., Perault, J. M., Bonmort, J. and Roblin, G. (2006) Early changes in
530 membrane permeability, production of oxidative burst and modification of PAL activity
531 induced by ergosterol in cotyledons of *Mimosa pudica*. *J. Exp. Bot.* **57**, 1245-1252
- 532 35 Granado, J., Felix, G. and Boller, T. (1995) Perception of Fungal Sterols in Plants
533 (Subnanomolar Concentrations of Ergosterol Elicit Extracellular Alkalinization in Tomato Cells).
534 *Plant Physiol.* **107**, 485-490
- 535 36 Kasparovsky, T., Blein, J. P. and Mikes, V. (2004) Ergosterol elicits oxidative burst in
536 tobacco cells via phospholipase A2 and protein kinase C signal pathway. *Plant Physiol.*
537 *Biochem.* **42**, 429-435
- 538 37 Tierney, K. J., Block, D. E. and Longo, M. L. (2005) Elasticity and phase behavior of DPPC
539 membrane modulated by cholesterol, ergosterol, and ethanol. *Biophys. J.* **89**, 2481-2493
- 540
541
542

543 **FIGURE LEGENDS**

544

545 Figure 1 – Comparison of the susceptibility to 2,4-D of *S. cerevisiae* BY4741 and the derived
546 deletion mutant $\Delta pdr18$, through cultivation in liquid medium (B) or spot assays (A and C). A:
547 Spot assays were carried as described in the M&M section. The cell suspensions used to
548 prepare the spots in lanes b) and c) were 1:5 and 1:25 serial dilutions, respectively, of the
549 suspensions with an $OD_{600nm}=0.05\pm 0.005$ spotted in lane a). Pictures were taken after 2 days
550 of incubation. B: Growth curves of wild-type (wt) (■, □) and $\Delta pdr18$ (▲, △) cells in MM4 liquid
551 medium, pH 3.5, (■, ▲) or in this medium supplemented with 0.45mM of 2,4-D (□, △). Cells of
552 the inocula were grown in the absence of the herbicide and the growth curves are
553 representative of at least three independent experiments. C: Spot assays were carried as
554 described in the M&M section. The cell suspensions used to prepare the spots in lanes b) and
555 c) were 1:5 and 1:25 serial dilutions, respectively, of the suspensions with an
556 $OD_{600nm}=0.05\pm 0.005$ spotted in lane a). Pictures were taken after 4 days of incubation.

557

558 Figure 2 - Susceptibility to the herbicides 2,4-D, MCPA and barban, to the 2,4-DCP, to the
559 agricultural fungicide mancozeb, and to the metal ions Zn^{2+} , Mn^{2+} , Cu^{2+} and Cd^{2+} induced stress
560 of deletion mutant $\Delta pdr18$ compared to the wt, through spot assays. Cells used for the spot
561 assays were prepared as described in the M&M section. The cell suspensions used to prepare
562 the spots in lanes b) and c) were 1:5 and 1:25 serial dilutions, respectively, of the suspensions
563 with an $OD_{600nm}=0.05\pm 0.005$ spotted in lane a). Pictures were taken after 2 or 3 days of
564 incubation.

565

566 Figure 3 – *PDR18* transcript levels in yeast cells exposed to 2,4-D imposed stress. A:
567 Comparison of the susceptibility to 2,4-D induced stress of *S. cerevisiae* parental strain BY4741
568 exposed to 0 (◆), 0.3 (■) or 0.45mM (▲) of 2,4-D through cultivation in MM4 liquid medium,
569 pH 3.5. Cells used to prepare the inocula were previously grown in the absence of 2,4-D until
570 mid-exponential phase. Growth curves are representative of at least three independent
571 growth experiments. B: Comparison of the relative transcript values of *PDR18* mRNA/*ACT1*
572 mRNA, in cells of parental BY4741 during the period of adaptation to 2,4-D by RT-PCR. The
573 *PDR18* mRNA value for the control conditions (0h, unsupplemented medium) was set as 1 and
574 the remaining values were relative values. Values are the mean of at least three independent
575 experiments and error bars indicate the standard deviation.

576

577 Figure 4 – A: Representation of the putative regulatory network controlling *PDR18*
578 transcription, according to the information in the YEASTRACT database (www.yeasttract.com).
579 B: Relative values of *PDR18* mRNA in wild type strain (wt) and $\Delta nrg1$, $\Delta pdr3$ and $\Delta yap1$ mutant
580 cells before and four hours following an yeast cell population exposure to 0.45 mM of 2,4-D.
581 The mRNA relative value for the wild type strain immediately before exposure to the herbicide
582 (control) was set as 1. Values are means of at least three independent experiments and error
583 bars indicate the standard deviation. C: Relative values of *PDR18* mRNA in $\Delta pdr18$ cells
584 transformed with pRS416_*PDR18*, pRS416_*PDR18* Δp or the corresponding empty vector
585 before and four hours following an yeast cell population exposure to 0.45 mM of 2,4-D. The
586 mRNA relative value for the $\Delta pdr18$ strain, harboring the pRS416_*PDR18* plasmid,
587 immediately before exposure to the herbicide was set as 1. Values are means of at least three
588 independent experiments and error bars indicate the standard deviation.

589

590 Figure 5 - Comparison of [^{14}C]-2,4-D accumulation in non-adapted cells of *S. cerevisiae* BY4741
591 (■) and the derived deletion mutant $\Delta pdr18$ (▲), during cultivation for 30min in MM4 liquid
592 medium (pH 3.5) supplemented with 0.3 mM of cold 2,4-D (Sigma) and 0.5 μM of [^{14}C]-2,4-D.

593 The accumulation (2,4-D* intra/2,4-D* extra) values are the means of at least three
594 independent experiments, error bars indicating standard deviation.

595

596 Figure 6 – Comparison of the relative abundance of sterol content in yeast plasma membrane
597 of the *S. cerevisiae* BY4741 and the derived deletion mutant $\Delta pdr18$ harboring the *PDR18*
598 expression plasmid or the corresponding empty vector, both grown under control conditions
599 or after 1h of exposure to 0.45 mM of 2,4-D, measured by GCMS. Values are means of at least
600 three independent experiments and error bars indicate the standard deviation.

601

602 Figure 7 – *PDR18* gene expression is important to maintain plasma membrane potential in
603 yeast cells. A: Time course accumulation of [14 C]-Methylammonium was followed during
604 incubation of *S. cerevisiae* BY4741 (■) and the derived deletion mutant $\Delta pdr18$ (▲), at 30°C, in
605 growth MM4 liquid medium, pH 3.5, supplemented with the radiolabelled ammonium
606 analogue. Relative levels of [14 C]-Methylammonium, assessed as described in M&M, are
607 average \pm standard deviation of at least three independent experiments. B: Comparison of the
608 average membrane potential using the fluorescent probe DiCO₆(3). Values of membrane
609 potential are set as the percentage of the value obtained for wild-type cells and result of at
610 least three independent experiments.

611

612

613

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

