



HAL
open science

Is it possible to construct an asymptotically efficient estimator of the proportion of true null hypotheses in a multiple testing setup?

van Hanh Nguyen, Catherine Matias

► To cite this version:

van Hanh Nguyen, Catherine Matias. Is it possible to construct an asymptotically efficient estimator of the proportion of true null hypotheses in a multiple testing setup?. 2011. hal-00647082v2

HAL Id: hal-00647082

<https://hal.science/hal-00647082v2>

Preprint submitted on 16 May 2012 (v2), last revised 8 Jan 2013 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Is it possible to construct an asymptotically efficient estimator of the proportion of true null hypotheses in a multiple testing setup?

Van Hanh Nguyen^{1,2} and Catherine Matias²

May 16, 2012

1. Laboratoire de Mathématiques d'Orsay, Université Paris Sud, UMR CNRS 8628, Bâtiment 425, 91 405 Orsay Cedex, France. E-mail: nvanhanh@genopole.cnrs.fr
2. Laboratoire Statistique et Génome, Université d'Évry Val d'Essonne, UMR CNRS 8071- USC INRA, 23 bvd de France, 91 037 Évry, France. E-mail: catherine.matias@genopole.cnrs.fr

Abstract

One important problem in the multiple testing context is the estimation of the proportion θ of true null hypotheses. This proportion appears in a semiparametric mixture model with two components: a uniform distribution on the interval $[0, 1]$ and a nonparametric density f . A large number of estimators of this proportion exist under different identifiability assumptions but their rate of convergence or asymptotic efficiency has only been partly studied. We shall focus here on two different categories of identifiability assumptions previously introduced in the literature: in the first case, f vanishes on a set with positive Lebesgue measure (and a subcase is obtained when this set is an interval) and in the second case, the set of points where f vanishes has a null Lebesgue measure. We first compute a lower bound on the local asymptotic minimax (LAM) quadratic risk of any estimator under the first case. To our knowledge, it has not been investigated whether the parametric rate of convergence may be achieved by a consistent estimator of the proportion θ in this semiparametric setup. Thus, we study an estimator previously proposed by [Celisse and Robin \(2010\)](#), and improve the results concerning its consistency by establishing its almost sure convergence as well as \sqrt{n} -consistency, under the assumption that f vanishes on an interval. As a consequence, it is natural to discuss the existence of asymptotically efficient estimators of the proportion θ in the sense of a convolution theorem. In the first case, we conjecture that no \sqrt{n} -consistent estimator is efficient. In the second case, we prove that the efficient information matrix for estimating θ is zero. Hence in this case, the LAM quadratic risk is not finite and there is no regular estimator of the proportion θ . These results are illustrated on simulations.

Key words and phrases: Asymptotic efficiency; efficient score, false discovery rate; information bound; multiple testing; p -values; semiparametric model.

1 Introduction

The problem of estimating the proportion θ of true null hypotheses is of interest in situation where several thousands of (independent) hypotheses can be tested simultaneously. One of the typical applications in which multiple testing problems occur is estimating the proportion of genes that are not differentially expressed in deoxyribonucleic acid (DNA) microarray experiments (see for instance [Dudoit and van der Laan, 2008](#)). Among other application domains, we mention astrophysics ([Meinshausen and Rice, 2006](#)) or neuroimaging ([Turkheimer et al., 2001](#)). A reliable estimate of θ is important when one wants to control multiple error rates, such as the false discovery rate (FDR) introduced by [Benjamini and Hochberg \(1995\)](#). In this work, we shall discuss asymptotic properties of estimators of the true proportion of null hypotheses. We stress that the asymptotic framework is particularly relevant in the above mentioned contexts where the number of tested hypotheses is huge.

In many recent articles (such as [Broberg, 2005](#); [Celisse and Robin, 2010](#); [Efron, 2004](#); [Efron et al., 2001](#); [Genovese and Wasserman, 2004](#), etc), a two-component mixture density is used to model the behavior of p -values X_1, X_2, \dots, X_n associated with n independent tested hypotheses. More precisely, assume the test statistics are independent and identically distributed (iid) with a continuous distribution under the corresponding null hypotheses, then the p -values X_1, X_2, \dots, X_n are iid and follow the uniform distribution $\mathcal{U}([0, 1])$ on interval $[0, 1]$ under the null hypotheses. The density g of p -values is modeled by a two-component mixture with following expression

$$\forall x \in [0, 1], \quad g(x) = \theta + (1 - \theta)f(x), \quad (1)$$

where $\theta \in [0, 1]$ is the unknown proportion of true null hypotheses and f denotes the density function of p -values generated under the alternative (false null hypotheses).

Starting from this model, two different points of view may be adopted. The first one consists in adding assumptions to the parameter set, so that the parameters (θ, f) become identifiable. It is easy to see that the parameters (θ, f) of model (1) are identifiable if and only if the infimum of density f on $[0, 1]$ is zero. This is a rather strong assumption, nonetheless it is unavoidable to ensure identifiability. Another point of view consists in not requiring identifiability of the model parameters. In such a case, model (1) is equivalent to

$$\forall x \in [0, 1], \quad g(x) = \bar{\theta} + (1 - \bar{\theta})\bar{f}(x), \quad \text{where } \bar{f} = \frac{f - c_f}{1 - c_f}, c_f = \inf_{[0,1]} f \text{ and } \bar{\theta} = \theta + (1 - \theta)c_f, \quad (2)$$

where now $(\bar{\theta}, \bar{f})$ is always identifiable from the distribution g . Thus, when it is not assumed that f vanishes on the set $[0, 1]$, we can only recover the parameter $\bar{\theta}$ instead of θ . In the following, we discuss the *best possible* properties of any estimator of $\bar{\theta}$ in general, under different assumptions on f (or equivalently \bar{f}). With some abuse of notation, we may still refer to $\bar{\theta}$ as *the proportion of true null hypotheses*, although it does not correspond exactly to this quantity when the infimum of f is not assumed to be zero.

Let us come back to the different cases that may occur with respect to the behavior of f and its infimum. First, one may assume the existence of $x_0 \in [0, 1]$ with $f(x_0) = \inf_{[0,1]} f$ or equivalently $\bar{f}(x_0) = 0$. Put in other words, density f may or not attain its infimum

on the set $[0, 1]$. Note that as soon as f is assumed to be continuous (a rather mild assumption), such a point x_0 will indeed exist. However, from an estimation perspective, this assumption is very weak and it is hopeless to obtain a reliable estimate of $\bar{\theta}$ without additional assumptions. Thus, in the following, we shall classify the setups in two different categories: models assuming that f attains its infimum on a set of points with positive measure and models where this set of points may have zero measure, and where regularity or monotonicity assumptions are added on f .

One of the most common assumptions made on density f in the literature is that it attains its infimum on an interval of the form $[\lambda, 1]$. This assumption underlies many estimation procedures further discussed below, such as variants of Efron's estimator. However, the fact that on real data, density f may not be minimum in the neighborhood of $x = 1$ has previously been noticed in the literature (see [Pounds and Cheng, 2006](#), and the references therein). As a consequence, authors such as [Celisse and Robin \(2010\)](#) propose a milder identifiability assumption that f attains its infimum on an interval $[\lambda^*, \mu^*]$ with $\mu^* \leq 1$. This assumption is more general than the previous one and contains $\mu^* = 1$ as a particular case. Nonetheless, we shall remark that when $\mu^* < 1$, such an assumption makes little sense in the specific context of modeling the distribution of p -values. Indeed, the high values of g (or f) near $x = 1$ that may be observed come from a misspecification of the distribution of the test statistic under the null or the alternative hypotheses. For instance, the test statistics may in fact be discrete, or the performed test is one-sided while some statistics follow the *untested alternative* ([Pounds and Cheng, 2006](#)). In this case, rather than generalizing assumptions on the semiparametric mixture model, one should reconsider the use of this mixture model itself (in the context of modeling the distribution of p -values). Indeed, it is then inappropriate to assume that the p -values corresponding to the null hypotheses follow a uniform distribution.

Let us remark that when identifiability of (θ, f) in model (1) is required, assuming f vanishes on a subset with positive measure is a rather strong assumption. Indeed, it implies that the support density of the underlying statistic T under the alternative and the null are different. More precisely, if the rejection region has the form $\{T \geq c\}$, then it is easy to see that the density f of the p -values under the alternative satisfies

$$\forall x \in [0, 1], \quad f(x) = \frac{\phi_1}{\phi_0} (\Phi_0^{-1}(1 - x)), \quad (3)$$

where ϕ_ϵ and Φ_ϵ are the density and cumulative distribution function (cdf) of T under hypothesis H_ϵ , respectively, and $\epsilon \in \{0, 1\}$. Then, assuming that the likelihood ratio ϕ_1/ϕ_0 is increasing (at least for x larger than some $A \in \mathbb{R}$ which is a reasonable assumption for a rejection region of the form $\{T \geq c\}$), density f vanishes on some interval if and only if the support of density ϕ_1 is strictly included in the support of ϕ_0 . In other words, the density of T should vanish on some interval $(-\infty, a_1]$ under the alternative and on some interval $(-\infty, a_0]$ under the null hypothesis, with $a_0, a_1 \in \mathbb{R}$ satisfying $a_0 < a_1$. However, when identifiability of (θ, f) is not required and one aims only at estimating the less informative parameter $\bar{\theta}$, assuming that f attains its minimum value on a subset with positive measure is a milder assumption. Indeed, this happens for instance when testing the nullity of the location parameter in a Laplace model (see for instance [Donoho and Jin, 2004](#), Section 5.2). More precisely, assume that the tests statistics T_1, T_2, \dots, T_n are iid with density $\phi_0(t) = (1/2) \exp\{-|t|\}$ under the null hypothesis and $\phi_1(t) = (1/2) \exp\{-|t - \mu|\}$ under

the alternative, so that we want to test $H_0 : \mu = 0$ versus a one-sided alternative $H_1 : \mu > 0$. In this case, the rejection region has the form $\{T \geq c\}$ and according to (3), density f attains its infimum on an interval of the form $[\mu, 1]$ if and only if the ratio ϕ_1/ϕ_0 does the same on an interval of the form $(-\infty, M]$. In the Laplace location model we are considering, this is exactly the case as for any $t < 0 < \mu$ we obtain that the ratio $(\phi_1/\phi_0)(t)$ equals its minimum value $e^{-\mu}$.

Nonetheless, to avoid assuming that f attains its minimum value on a whole interval while preserving some hope to obtain a consistent estimator of $\bar{\theta}$, some authors proposed to rather assume that density f attains its minimum value at point $x = 1$. They further assume either that f is decreasing such as in the work of Langaas et al. (2005); or that f is regular near $x = 1$, such as in the work of Neuvial (2010).

Let us now discuss the different estimators proposed in the literature, starting with those assuming (implicitly or not) that f attains its minimum value on a whole interval. In the following, we stress the fact that it is completely equivalent to estimate θ assuming that f vanishes (at some point, or in some interval) or to estimate only $\bar{\theta}$ (and making assumptions on the set of points where f attains its infimum). First, Schweder and Spjøtvoll (1982) suggested a procedure to estimate $\bar{\theta}$, that has been later used by Storey (2002). This estimator depends on an unspecified parameter $\lambda \in [0, 1)$ and is equal to the proportion of p -values larger than this threshold λ divided by $1 - \lambda$. It is thus consistent only if f attains its minimum value on the interval $[\lambda, 1]$ (an assumption not made in the article by Schweder and Spjøtvoll (1982) nor the one by Storey (2002)). Note that even if such an assumption were made, it would not solve the problem of choosing λ such that f attains its infimum on $[\lambda, 1]$. Adapting this procedure in order to end up with an estimate of the positive FDR (pFDR), Storey (2002) proposes a bootstrap strategy to pick λ . More precisely, his procedure minimizes the mean-squared error for estimating the pFDR. Note that Genovese and Wasserman (2004) established that, for fixed value λ such that the cdf F of f satisfies $F(\lambda) < 1$, Storey's estimator converges at parametric rate and is asymptotically normal, but is also asymptotically biased (thus it does not converge to $\bar{\theta}$ at parametric rate). Some other choices of λ are, for instance, based on break point estimation (Turkheimer et al., 2001) or spline smoothing (Storey and Tibshirani, 2003). Recently, Celisse and Robin (2010) proposed an estimator of θ , relying on a histogram estimate of g , under the assumption that f vanishes on an interval $[\lambda^*, \mu^*]$. This estimator is shown to converge in probability to θ . We want to stress here an important difference between Storey's and Celisse and Robin's estimators. Indeed, both are constructed using nonparametric estimates \hat{g} of the density g and then estimate θ (or $\bar{\theta}$ in Storey's case) relying on the value of \hat{g} on a specific interval. However, contrarily to Storey's procedure, the estimator proposed by Celisse and Robin automatically selects an interval where g is identically equal to $\bar{\theta}$.

Other estimators of $\bar{\theta}$ are based on regularity or monotonicity assumptions made on f or equivalently on g , combined with the assumption that the infimum of g is attained at $x = 1$ (thus we have $\bar{\theta} = g(1)$). These estimators rely on nonparametric estimates of g and appear to inherit nonparametric rates of convergence. Langaas et al. (2005) derive estimators based on nonparametric maximum likelihood estimation of the p -value density, in two setups: decreasing and convex decreasing densities f . We mention that no theoretical properties of these estimators are given. Hengartner and Stark (1995) propose a very general finite sample confidence envelope for a monotone density. Relying on this result

and assuming moreover that cdf G is concave and that g is Lipschitz in a neighborhood of $x = 1$, [Genovese and Wasserman \(2004\)](#) construct an estimator converging to $g(1) = \bar{\theta}$ at rate $(\log n)^{1/3}n^{-1/3}$. Under some regularity assumptions on f near $x = 1$, [Neuviel \(2010\)](#) established that by letting $\lambda \rightarrow 1$, [Storey's](#) estimator may be turned into a consistent estimator of $\bar{\theta}$, with a nonparametric rate of convergence equal to $n^{-k/(2k+1)}\eta_n$, where $\eta_n \rightarrow +\infty$ and k controls the regularity of f near $x = 1$.

We mention that [Meinshausen and Bühlmann \(2005\)](#) discuss probabilistic upper bounds for the proportion of true null hypotheses, which are valid under general and unknown dependence structures between the test statistics. We also mention that, in a very general and nonparametric setup, [Swanepoel \(1999\)](#) proposes a two-step estimator of the minimum of an unknown density h based on the distribution of the spacings between observations. Assuming that in some neighborhood of the value at which the density h achieves its minimum, h has a bounded second derivative h'' satisfying a Lipschitz condition, [Swanepoel](#) establishes that for any $\delta > 0$, there exists an estimator of this infimum converging at rate $(\log n)^\delta n^{-2/5}$ to the true minimum. However, this procedure assumes that the minimum is achieved in the interior of the support of h . As discussed above, such an assumption is not realistic in the context of estimation of the true proportion of null hypotheses, unless the minimum is achieved in a whole interval of the form $[\lambda, 1]$.

Finally, note that we do not discuss here estimators of the proportion of non null effects in Gaussian mixtures such as in [Cai and Jin \(2010\)](#); [Jin \(2008\)](#); [Jin and Cai \(2007\)](#), a related but although different problem as the one we study.

Despite a large number of different estimation procedures of the proportion of true null hypotheses proposed in the literature, very few results on their convergence properties exist. In particular, none of the above mentioned estimators is shown to achieve a parametric rate of convergence towards $\bar{\theta}$. Thus, natural open questions are whether it is possible to construct an estimator converging at parametric rate; whether there exists an *asymptotically efficient* estimator and what is the value of an *optimal* variance. Here, asymptotic efficiency stands in the sense of a convolution theorem (see [van der Vaart, 1998](#), Chapter 25, for more details on efficiency theory for semiparametric models). An estimator is said to be *regular* if it converges in distribution at parametric rate. According to a convolution theorem (see Theorem 25.20 in [van der Vaart, 1998](#)), regular estimators converge in distribution to the convolution of a Gaussian random variable with some (minimal) variance and another random variable. Thus, an estimator is asymptotically efficient if among regular estimators, it achieves the *best* limit -namely the Gaussian distribution with minimal variance. In fact in our context, at least two different cases occur, whether density f is assumed to attain its infimum on a set with either positive or null Lebesgue measure. We thus consider the following semi-parametric model

$$\mathcal{P} = \{p_{\theta,f} : [0, 1] \rightarrow \mathbb{R}^+; p_{\theta,f} := \theta + (1 - \theta)f, \theta \in (0, 1), \\ f : [0, 1] \rightarrow \mathbb{R}^+ \text{ density with } \inf_{[0,1]} f = 0\},$$

and remind once again the reader about the fact that this model is completely equivalent to model (2). In other words, when it is not assumed that f vanishes on $[0, 1]$ then the following discussion remains valid replacing (θ, f) with $(\bar{\theta}, \bar{f})$. In this semiparametric setup, we aim at estimating the finite-dimensional parameter θ and consider f as a nuisance parameter.

The main aim of our work is discussing the existence of an asymptotically efficient estimator of θ under the two categories of identifiability assumptions mentioned above.

We shall thus consider two semiparametric submodels of \mathcal{P} , respectively denoted by \mathcal{P}_δ and \mathcal{P}_0 and defined in the following sections. Model \mathcal{P}_δ assumes that the subset of points where f is zero has a positive Lebesgue measure whereas in model \mathcal{P}_0 , it has a null measure. We first compute a lower bound for the local asymptotic minimax (LAM) quadratic risk for estimating θ in model \mathcal{P}_δ and then look for the existence of a \sqrt{n} -consistent estimator of θ in this model. We prove that this \sqrt{n} -consistency is achieved by the estimator previously proposed by [Celisse and Robin \(2010\)](#). By doing that, we improve the results known on the convergence properties of this estimator. Indeed, we recall that [Celisse and Robin \(2010\)](#) established only convergence in probability of their estimator and did not establish that it achieves a parametric rate of convergence. Then, looking for an efficient estimator among the regular ones (namely among those converging at parametric rate), we conjecture that no \sqrt{n} -consistent estimator is efficient in this model. In a second part, we prove that the efficient information matrix for estimating θ in model \mathcal{P}_0 is zero. This implies both that the LAM quadratic risk at parametric rate is not finite and that if there exists a \sqrt{n} -consistent estimator in this model, it can not have finite asymptotic variance. We stress that this does not necessarily imply that \sqrt{n} -consistent estimators do not exist in model \mathcal{P}_0 . However, the only rates of convergence obtained until now in this case are nonparametric ones.

The article is organized as follows. Section 2 is devoted to the easiest case (from an estimation perspective) where it is assumed that f vanishes on a set with positive Lebesgue measure. In Section 2.1, we compute a lower bound of the LAM quadratic risk for estimating θ in model \mathcal{P}_δ and give conditions under which an estimator is asymptotically efficient in the sense of a convolution theorem. Since the former lower bound for the LAM quadratic risk is positive, it is natural to look for an estimator converging at parametric rate, and then for an efficient one. Thus in Section 2.2, we establish the consistency properties (almost sure convergence and \sqrt{n} -consistency) of the estimator previously proposed by [Celisse and Robin \(2010\)](#). In Section 2.3, we recall a general method (one-step estimators) of construction of efficient estimators relying on first step \sqrt{n} -consistent ones. Section 2.4 further discusses the implications of semiparametric theory on the existence of efficient estimators of θ in model \mathcal{P}_δ . In particular, it is conjectured that no efficient estimator exists in this setup. In a second part (Section 3), we turn to the more difficult case where f is only assumed to vanish on a set with null Lebesgue measure. In this case, we prove that the efficient information matrix for estimating θ is zero. Hence in this case, the LAM quadratic risk is not finite and there is no regular estimator of the proportion θ . Some simulations are given as an illustration in Section 4. Finally, some technical proofs have been postponed to Appendix A.

2 Asymptotically efficient estimators of θ when f vanishes on a set with positive Lebesgue measure

2.1 A lower bound on the LAM quadratic risk

For any fixed unknown positive parameter $\delta > 0$, we introduce the set of densities $\mathcal{F}_\delta = \{\text{densities } f \text{ on } [0, 1]; \mu(I_f) = \delta\}$, where μ denotes Lebesgue measure and $I_f = \{x \in$

$[0, 1]; f(x) = 0\}$. We now consider a semiparametric submodel of \mathcal{P} defined by

$$\mathcal{P}_\delta = \left\{ p_{\theta,f} \in \mathcal{P}; p_{\theta,f} = \frac{d\mathbb{P}_{\theta,f}}{d\mu} = \theta + (1 - \theta)f, \theta \in (0, 1), f \in \mathcal{F}_\delta \right\}.$$

In this model, we assume that the density f vanishes on a set whose Lebesgue measure is exactly $\delta > 0$. We aim at estimating the parameter $\psi(\mathbb{P}_{\theta,f}) = \theta$ and consider f as a nuisance parameter. We shall calculate a lower bound on the LAM quadratic risk of any estimator of θ and we also present conditions under which an estimator is asymptotically efficient in the sense of a convolution theorem.

Let us start by recalling some results from semiparametric theory (we refer to Chapter 25 in [van der Vaart, 1998](#), for more details on this subject). We consider a tangent set, denoted by $\dot{\mathcal{P}}_\delta$, of the model \mathcal{P}_δ at $\mathbb{P}_{\theta,f}$ (with respect to the two parameters (θ, f)) while $\dot{l}_{\theta,f}$ is the ordinary score function for θ in the model in which f is fixed and $\dot{\mathcal{P}}_{f,\delta}$ is a tangent set for the nuisance parameter f in the model where θ is fixed. We consider the efficient score function $\tilde{l}_{\theta,f}$ for estimating θ and let $\tilde{\psi}_{\theta,f}$ be the efficient influence function relative to the tangent set $\dot{\mathcal{P}}_\delta$. For any function h , let us denote $\mathbb{P}_{\theta,f}h = \int h d\mathbb{P}_{\theta,f}$. We now consider the path

$$f_t(x) = \frac{k(th_0(x))f(x)}{\int k(th_0(u))f(u)du} = c(t)k(th_0(x))f(x), \quad (4)$$

where $k(u) = 2(1 + e^{-2u})^{-1}$ and $[c(t)]^{-1} = \int k(th_0(u))f(u)du$. By using Lemma 1.8 in [van der Vaart \(2002\)](#), we obtain a tangent set for f given by

$$\dot{\mathcal{P}}_{f,\delta} = \left\{ h \in \mathbb{L}^2(\mathbb{P}_{\theta,f}); \exists h_0 \in \mathbb{L}^1(\mathbb{P}_{\theta,f}), h = \frac{(1 - \theta)fh_0}{\theta + (1 - \theta)f} \text{ and } \int fh_0 = 0 \right\}.$$

We remark that if $\dot{l}_{\theta,f}$ is the ordinary score function for θ in the model in which f is fixed, then for every $a \in \mathbb{R}$ and for every $h \in \dot{\mathcal{P}}_{f,\delta}$, we have $a\dot{l}_{\theta,f} + h$ is a score function for (θ, f) corresponding to the path $t \mapsto \mathbb{P}_{\theta+ta, f_t}$. Hence, the linear span

$$\dot{\mathcal{P}}_\delta = \text{lin}(\dot{l}_{\theta,f} + \dot{\mathcal{P}}_{f,\delta}) = \{\alpha\dot{l}_{\theta,f} + \beta h; (\alpha, \beta) \in \mathbb{R}^2, h \in \dot{\mathcal{P}}_{f,\delta}\}$$

is a tangent set for (θ, f) at $\mathbb{P}_{\theta,f}$. This set is a linear subspace of Hilbert space $\mathbb{L}^2(\mathbb{P}_{\theta,f})$ with infinite dimension. For every score function g in the tangent set $\dot{\mathcal{P}}_\delta$, we write $P_{t,g}$ for a submodel with score function g along which the function $\psi : \mathbb{P}_{\theta,f} \rightarrow \theta$ is differentiable. An estimator sequence T_n is called regular at $\mathbb{P}_{\theta,f}$ for estimating $\psi(\mathbb{P}_{\theta,f})$ (relative to the tangent set $\dot{\mathcal{P}}_\delta$) if there exists a probability measure L such that

$$\sqrt{n}(T_n - \psi(P_{1/\sqrt{n},g})) \xrightarrow[P_{1/\sqrt{n},g}]{d} L, \text{ for every } g \in \dot{\mathcal{P}}_\delta.$$

According to a convolution theorem (Theorem 25.20 in [van der Vaart, 1998](#)), this limit distribution writes as the convolution between the Gaussian distribution $N(0, P(\tilde{\psi}_P^2))$ and another distribution. Thus we shall say that an estimator sequence is asymptotically efficient at P (relative to the tangent set $\dot{\mathcal{P}}_\delta$) if it is regular at P with limit distribution $L = N(0, P(\tilde{\psi}_P^2))$, in other words it is the best regular estimator. We shall call LAM quadratic risk of an estimator sequence T_n (relative to the tangent set $\dot{\mathcal{P}}_\delta$) the quantity

$$\sup_E \liminf_{n \rightarrow \infty} \sup_{g \in E} \mathbb{E}_{P_{1/\sqrt{n},g}} [\sqrt{n}(T_n - \psi(P_{1/\sqrt{n},g}))]^2,$$

where the first supremum is taken over all finite subsets E of the tangent set $\dot{\mathcal{P}}_\delta$. According to LAM theorem (see Theorem 25.21 in [van der Vaart, 1998](#)), this quantity is lower bounded by the minimal variance $P(\tilde{\psi}_P^2)$.

In the following lemma, we shall calculate the efficient information matrix and the efficient influence function in order to establish conditions under which an estimator is asymptotically efficient.

Lemma 1. *The efficient information matrix $\tilde{I}_{\theta,f}$ for θ and the efficient influence function $\tilde{\psi}_{\theta,f}$ relative to the tangent set $\dot{\mathcal{P}}_\delta$ are respectively given by*

$$\tilde{I}_{\theta,f} = \frac{\delta}{\theta(1-\theta\delta)} \text{ and } \tilde{\psi}_{\theta,f}(x) = \frac{1}{\delta} \mathbf{1}_{\{f(x)=0\}} - \theta.$$

By using Theorem 25.21 and Lemma 25.23 in [van der Vaart \(1998\)](#), an immediate consequence of this lemma is expressed (without any proof) in the following theorem.

Theorem 1. *i) For any estimator sequence T_n we have,*

$$\sup_E \liminf_{n \rightarrow \infty} \sup_{g \in E} \mathbb{E}_{P_{1/\sqrt{n},g}} [\sqrt{n}(T_n - \psi(P_{1/\sqrt{n},g}))]^2 \geq \theta \left(\frac{1}{\delta} - \theta \right),$$

where the first supremum is taken over all finite subsets E of the tangent set $\dot{\mathcal{P}}_\delta$.

ii) A sequence of estimators $\hat{\theta}_n$ is asymptotically efficient in the sense of a convolution theorem (best regular estimator) if and only if it satisfies

$$\hat{\theta}_n = \frac{1}{n} \sum_{i=1}^n \frac{1}{\delta} \mathbf{1}_{\{f(X_i)=0\}} + o_{\mathbb{P}_{\theta,f}}(n^{-1/2}). \quad (5)$$

Remark 1. *i) Note that for fixed parameter value λ such that $G(\lambda) < 1$, [Storey's](#) estimator $\hat{\theta}^{\text{Storey}}(\lambda)$ satisfies*

$$\sqrt{n} \left(\hat{\theta}^{\text{Storey}}(\lambda) - \frac{1-G(\lambda)}{1-\lambda} \right) \xrightarrow[n \rightarrow \infty]{d} \mathcal{N} \left(0, \frac{G(\lambda)(1-G(\lambda))}{(1-\lambda)^2} \right)$$

(see for instance [Genovese and Wasserman, 2004](#)). In particular, if we assume that f vanishes on $[\lambda, 1]$ then we obtain that $G(\lambda) = 1 - \theta(1 - \lambda)$ and $\hat{\theta}^{\text{Storey}}(\lambda)$ becomes a \sqrt{n} -consistent estimate of θ , which is moreover asymptotically distributed, with asymptotic variance

$$\theta \left(\frac{1}{1-\lambda} - \theta \right).$$

In this sense, the oracle version of [Storey's](#) estimator that picks $\lambda = 1 - \delta$ (namely choosing λ as the smallest value such that f vanishes on $[\lambda, 1]$) is asymptotically efficient. Note also that $\hat{\theta}^{\text{Storey}}(\lambda)$ automatically satisfies (5).

ii) In the same way, an oracle version of [Celisse and Robin's](#) estimator that would know the true interval Λ^ and include the partition $I = (N, \lambda^*, \mu^*)$ in the collection of possible partitions \mathcal{I} would have asymptotically minimal variance and thus be efficient (see [Section 2.2](#) and [Remark 3](#) for more details).*

Proof of Lemma 1. First, we can easily compute the ordinary score function $\dot{l}_{\theta,f}$ for θ in the model in which f is fixed:

$$\dot{l}_{\theta,f}(x) = \frac{\partial}{\partial \theta} \log[\theta + (1 - \theta)f(x)] = \frac{1 - f(x)}{\theta + (1 - \theta)f(x)}. \quad (6)$$

Define $\Pi_{\theta,f}$ as the orthogonal projection onto the closure of the linear span of $\dot{\mathcal{P}}_{f,\delta}$ in $\mathbb{L}_2(\mathbb{P}_{\theta,f})$. We write

$$\dot{l}_{\theta,f} = \left(\frac{1 - f}{\theta + (1 - \theta)f} + \frac{\delta}{1 - \theta\delta} \right) \mathbf{1}_{\{f>0\}} + \frac{1}{\theta} \mathbf{1}_{\{f=0\}} - \frac{\delta}{1 - \theta\delta} \mathbf{1}_{\{f>0\}}.$$

We shall prove that

$$\Pi_{\theta,f} \dot{l}_{\theta,f}(x) = \left(\frac{1 - f(x)}{\theta + (1 - \theta)f(x)} + \frac{\delta}{1 - \theta\delta} \right) \mathbf{1}_{\{f(x)>0\}}, \quad (7)$$

and then the efficient score function for θ is

$$\tilde{l}_{\theta,f}(x) = \dot{l}_{\theta,f}(x) - \Pi_{\theta,f} \dot{l}_{\theta,f}(x) = \frac{1}{\theta} \mathbf{1}_{\{f(x)=0\}} - \frac{\delta}{1 - \theta\delta} \mathbf{1}_{\{f(x)>0\}}.$$

In fact, we can write

$$\left(\frac{1 - f}{\theta + (1 - \theta)f} + \frac{\delta}{1 - \theta\delta} \right) \mathbf{1}_{\{f>0\}} = \frac{(1 - \theta)fh_0}{\theta + (1 - \theta)f},$$

where

$$h_0(x) = \left(\frac{1 - f(x)}{(1 - \theta)f(x)} + \frac{\delta}{1 - \theta\delta} \times \frac{\theta + (1 - \theta)f(x)}{(1 - \theta)f(x)} \right) \mathbf{1}_{\{f(x)>0\}}.$$

Let us denote by I_f^c the complement of I_f in $[0, 1]$. It is not difficult to examine the condition $\int fh_0 = 0$. Indeed,

$$\begin{aligned} \int_0^1 f(x)h_0(x)dx &= \int_{I_f^c} \left(\frac{1 - f(x)}{(1 - \theta)} + \frac{\delta}{1 - \theta\delta} \times \frac{\theta + (1 - \theta)f(x)}{(1 - \theta)} \right) dx \\ &= \frac{1}{1 - \theta} \left[\int_{I_f^c} dx - \int_{I_f^c} f(x)dx + \frac{\delta}{1 - \theta\delta} \int_{I_f^c} (\theta + (1 - \theta)f(x))dx \right] \\ &= \frac{1}{1 - \theta} \left[1 - \int_{I_f} dx - \int_0^1 f(x)dx + \frac{\delta}{1 - \theta\delta} (1 - \int_{I_f} \theta dx) \right] \\ &= \frac{1}{1 - \theta} \left[-\delta + \frac{\delta}{1 - \theta\delta} (1 - \theta\delta) \right] = 0, \end{aligned}$$

and hence

$$\left(\frac{1 - f}{\theta + (1 - \theta)f} + \frac{\delta}{1 - \theta\delta} \right) \mathbf{1}_{\{f>0\}} \in \dot{\mathcal{P}}_{f,\delta}.$$

Therefore, it is necessary to prove that

$$\frac{1}{\theta} \mathbf{1}_{\{f=0\}} - \frac{\delta}{1 - \theta\delta} \mathbf{1}_{\{f>0\}} = \frac{1}{\theta(1 - \theta\delta)} \mathbf{1}_{\{f=0\}} - \frac{\delta}{1 - \theta\delta} \perp \dot{\mathcal{P}}_{f,\delta},$$

where \perp means orthogonality in $\mathbb{L}^2(\mathbb{P}_{\theta,f})$. In fact, for every score function

$$h = \frac{(1-\theta)fh_0}{\theta + (1-\theta)f} \in \dot{\mathcal{P}}_{f,\delta} \text{ with } \int fh_0 = 0,$$

we have

$$\begin{aligned} & \int_0^1 \left[\frac{1}{\theta(1-\theta\delta)} \mathbf{1}_{\{f(x)=0\}} - \frac{\delta}{1-\theta\delta} \right] h(x) d\mathbb{P}_{\theta,f}(x) \\ = & \int_0^1 \left[\frac{1}{\theta(1-\theta\delta)} \mathbf{1}_{\{f(x)=0\}} - \frac{\delta}{1-\theta\delta} \right] \frac{(1-\theta)f(x)h_0(x)}{\theta + (1-\theta)f(x)} (\theta + (1-\theta)f(x)) dx \\ = & \frac{1-\theta}{\theta(1-\theta\delta)} \int_0^1 f(x)h_0(x) \mathbf{1}_{\{f(x)=0\}} dx - \frac{(1-\theta)\delta}{1-\theta\delta} \int_0^1 f(x)h_0(x) dx = 0. \end{aligned}$$

This establishes (7). Let us now calculate the efficient information matrix

$$\begin{aligned} \tilde{I}_{\theta,f} &= \mathbb{P}_{\theta,f}(\tilde{l}_{\theta,f}^2) \\ &= \int_0^1 \left(\frac{1}{\theta^2} \mathbf{1}_{\{f(x)=0\}} + \frac{\delta^2}{(1-\theta\delta)^2} \mathbf{1}_{\{f(x)>0\}} \right) (\theta + (1-\theta)f(x)) dx \\ &= \frac{\delta}{\theta} + \frac{\delta^2}{(1-\theta\delta)^2} (1-\theta\delta) = \frac{\delta}{\theta(1-\theta\delta)} > 0. \end{aligned}$$

Using Lemma 25.25 in [van der Vaart \(1998\)](#), we remark that the functional $\psi(\mathbb{P}_{\theta,f}) = \theta$ is differentiable at $\mathbb{P}_{\theta,f}$ relative to the tangent set $\dot{\mathcal{P}}_\delta$ with efficient influence function given by

$$\begin{aligned} \tilde{\psi}_{\theta,f}(x) &= \tilde{I}_{\theta,f}^{-1} \tilde{l}_{\theta,f}(x) \\ &= \frac{\theta(1-\theta\delta)}{\delta} \left(\frac{1}{\theta} \mathbf{1}_{\{f(x)=0\}} - \frac{\delta}{1-\theta\delta} \mathbf{1}_{\{f(x)>0\}} \right) \\ &= \frac{1-\theta\delta}{\delta} \mathbf{1}_{\{f(x)=0\}} - \theta \mathbf{1}_{\{f(x)>0\}} \\ &= \frac{1}{\delta} \mathbf{1}_{\{f(x)=0\}} - \theta. \end{aligned}$$

We can thus conclude that a lower bound on the minimax quadratic risk for estimating θ is $\mathbb{P}_{\theta,f}(\tilde{\psi}_{\theta,f}^2) = \tilde{I}_{\theta,f}^{-1} = \theta(\delta^{-1} - \theta)$. \square

2.2 A \sqrt{n} -consistent estimator of θ

In this section, we shall introduce the estimator of the parameter θ proposed by [Celisse and Robin \(2010\)](#). In particular, we specify some assumptions that establish its almost sure convergence and \sqrt{n} -consistency. Throughout this section, we shall always assume that the density f belongs to $\mathbb{L}^2([0, 1])$.

Assumption 1. *Density f is null on an interval $[\lambda^*, \mu^*] \subset (0, 1]$ (with unknown values λ^* and μ^*) and f is monotone outside the interval $[\lambda^*, \mu^*]$.*

For example, f is decreasing on $[0, \lambda^*]$ and increasing on $[\mu^*, 1]$. This assumption is stronger than Assumption A' in [Celisse and Robin \(2010\)](#), the latter not being sufficient to

establish *the result they claim* (see the proof of Lemma 4 for more details). The monotonicity part of our assumption is not necessary and we shall explain what is exactly required and how we use the previous assumption in the proof of Lemma 4. Under Assumption 1, the true parameter θ is equal to $g(x)$ for all x in $[\lambda^*, \mu^*]$. Note that the case where we impose $\mu^* = 1$ is included in this setting. The general idea underlying the estimator's construction is the following one. Let us consider a nonparametric density estimator of g . For example, let \hat{g}_I be a histogram estimator corresponding to a partition $I = (I_k)_{1, \dots, D}$ of $[0, 1]$, defined by

$$\hat{g}_I(x) = \sum_{k=1}^D \frac{n_k}{n|I_k|} \mathbf{1}_{I_k}(x),$$

where $n_k = \text{card}\{i : X_i \in I_k\}$ is the number of observations in I_k , $\mathbf{1}_{I_k}$ is the indicator function of I_k and $|I_k|$ is the width of interval I_k . We estimate θ by the value of \hat{g}_I on an interval \hat{I} which is the closest as possible to the interval $[\lambda^*, \mu^*]$.

Let us now recall more precisely the procedure for estimating θ that is presented in [Celisse and Robin \(2010\)](#). For a given integer N , define \mathcal{I}_N as the set of partitions of $[0, 1]$ such that for some integers k, l with $2 \leq k + 2 \leq l \leq N$, the first k intervals and the last $N - l$ ones are regular of width $1/N$, namely

$$\mathcal{I}_N = \left\{ I = (I_i)_i : \forall i \neq k + 1, |I_i| = \frac{1}{N}, |I_{k+1}| = \frac{l - k}{N}, 2 \leq k + 2 \leq l \leq N \right\}.$$

Then for two given integers $m_{\min} < m_{\max}$, denote by \mathcal{I} the collection of considered partitions, defined by

$$\mathcal{I} = \bigcup_{m_{\min} \leq m \leq m_{\max}} \mathcal{I}_{2^m}. \quad (8)$$

Every partition I in \mathcal{I} is characterized by a triplet $(N = 2^m, \lambda = k/N, \mu = l/N)$ and the quality of the histogram estimator \hat{g}_I is measured by its quadratic risk. So in this sense, the *oracle estimator* $\hat{g}_{\hat{I}}$ is obtained through

$$\hat{I} = \underset{I \in \mathcal{I}}{\text{argmin}} \mathbb{E}[\|g - \hat{g}_I\|_2^2] = \underset{I \in \mathcal{I}}{\text{argmin}} R(I), \text{ where } R(I) = \mathbb{E} \left[\|\hat{g}_I\|_2^2 - 2 \int_0^1 \hat{g}_I(x)g(x)dx \right].$$

However, for every partition I , the quantity $R(I)$ depends on g which is unknown. Thus \hat{I} is an oracle and not an estimator. It is then natural to replace $R(I)$ by an estimator. In [Celisse and Robin \(2008, 2010\)](#), the authors use leave-p-out (LPO) estimator of $R(I)$ with $p \in \{1, \dots, n - 1\}$, whose expression is given by (see [Celisse and Robin, 2008](#), Theorem 2.1)

$$\hat{R}_p(I) = \frac{2n - p}{(n - 1)(n - p)} \sum_k \frac{n_k}{n|I_k|} - \frac{n(n - p + 1)}{(n - 1)(n - p)} \sum_k \frac{1}{|I_k|} \left(\frac{n_k}{n}\right)^2. \quad (9)$$

The best theoretical value of p is the one that minimizes the mean-square error (MSE) of $\hat{R}_p(I)$, namely

$$p^*(I) = \underset{p \in \{1, \dots, n-1\}}{\text{argmin}} \text{MSE}(p, I) = \underset{p \in \{1, \dots, n-1\}}{\text{argmin}} \mathbb{E} \left[(\hat{R}_p(I) - R(I))^2 \right].$$

It clearly appears that $\text{MSE}(p, I)$ has the form of a function $\Phi(p, I, \alpha)$ (see [Celisse and Robin, 2008](#), Proposition 2.1) depending on the unknown vector $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_D)$ with

$\alpha_k = \mathbb{P}(X_1 \in I_k)$. A natural idea would be to replace the α_k s in $\Phi(p, I, \alpha)$ by their empirical counterparts $\hat{\alpha}_k = n_k/n$ and an estimator of $p^*(I)$ is therefore given by

$$\hat{p}(I) = \underset{p \in \{1, \dots, n-1\}}{\operatorname{argmin}} \widehat{MSE}(p, I) = \underset{p \in \{1, \dots, n-1\}}{\operatorname{argmin}} \Phi(p, I, \hat{\alpha}).$$

The exact calculation of $\hat{p}(I)$ may be found in Theorem 3.1 from [Celisse and Robin \(2008\)](#). Hence, the procedure for estimating θ is the following one

1. For each partition $I \in \mathcal{I}$, define $\hat{p}(I) = \underset{p \in \{1, \dots, n-1\}}{\operatorname{argmin}} \widehat{MSE}(p, I)$,
2. Choose $\hat{I} = (\hat{N}, \hat{\lambda}, \hat{\mu}) \in \underset{I \in \mathcal{I}}{\operatorname{argmin}} \hat{R}_{\hat{p}(I)}(I)$ such that the width of the interval $[\hat{\lambda}, \hat{\mu}]$ is maximum,
3. Estimate θ by $\hat{\theta}_n = \operatorname{card}\{i : X_i \in [\hat{\lambda}, \hat{\mu}]\} / [n(\hat{\mu} - \hat{\lambda})]$.

Remark 2. In our procedure, we consider the set of natural partitions defined by (8), while [Celisse and Robin \(2010\)](#) use the one defined by

$$\mathcal{I} = \bigcup_{N_{\min} \leq N \leq N_{\max}} \mathcal{I}_N.$$

This change is natural for lowering the complexity of the algorithm and has no consequences on the theoretical properties of the estimator. In particular, if we assume the function f vanishes on an interval $[1 - \delta, 1]$, then the complexity of the algorithm is simpler when we consider the following set of partitions

$$\mathcal{I} = \bigcup_{m_{\min} \leq m \leq m_{\max}} \mathcal{I}_{2^m},$$

where

$$\mathcal{I}_N = \left\{ I^{(k)} = (I_i)_{i=1, \dots, k+1} : \forall i \leq k, |I_i| = \frac{1}{N}, |I_{k+1}| = \frac{N-k}{N}, 1 \leq k \leq N-2 \right\}.$$

Let us now study the almost sure convergence and \sqrt{n} -consistency of this estimator. First, we present some lemmas whose proofs have been postponed to Appendix A. The two first ones are proved for all partitions of $[0, 1]$ without relying on Assumption 1, while in Lemma 4 we assume that the function f satisfies Assumption 1. This hypothesis is stronger than Assumption A' in [Celisse and Robin \(2010\)](#), the latter not being sufficient to establish their result (see discussion in the proof of Lemma 4). We mention that in their simulations, [Celisse and Robin \(2010\)](#) use a function f satisfying our assumption. In Lemma 3, we improve the results of convergence on LPO risk estimator established in Proposition 2.1 and Corollary 2.1 from [Celisse and Robin \(2010\)](#). Indeed, we prove its almost sure convergence and asymptotic normality.

For each partition I , let us denote by \mathcal{F}_I the vector space of piecewise constant functions built from the partition I and g_I the orthogonal projection of $g \in L^2([0, 1])$ onto \mathcal{F}_I . The mean square error of a histogram estimator \hat{g}_I can be written as the sum of a bias term and a variance term

$$\mathbb{E}[\|g - \hat{g}_I\|_2^2] = \|g - g_I\|_2^2 + \mathbb{E}[\|g_I - \hat{g}_I\|_2^2].$$

Lemma 2. Let $I = (I_k)_{k=1}^D$ be an arbitrary partition of $[0, 1]$. Then the variance term of the mean square error of a histogram estimator \hat{g}_I is bounded by C/n , where C is a positive constant. In other words,

$$\mathbb{E}[|g_I - \hat{g}_I|_2^2] = O\left(\frac{1}{n}\right).$$

Lemma 3. Let $I = (I_k)_{1,\dots,D}$ be an arbitrary partition of $[0, 1]$. Define $L(I) = \|g_I - g\|_2^2$ the bias term of the mean square error of a histogram estimator \hat{g}_I and $\hat{L}_p(I) = \hat{R}_p(I) + \|g\|_2^2$. Let $p \in \{1, 2, \dots, n-1\}$ such that $\lim_{n \rightarrow \infty} p/n < 1$. Then we have the following results

i) $\hat{L}_p(I) \xrightarrow[n \rightarrow \infty]{as} L(I)$

ii) $\sqrt{n}(\hat{L}_p(I) - L(I)) = \sqrt{n}(\hat{R}_p(I) - R(I)) + \frac{1}{\sqrt{n}}(s_{11} - s_{21}) \xrightarrow[n \rightarrow \infty]{d} \mathcal{N}(0, 4\sigma_I^2)$,

where \xrightarrow{d} means convergence in distribution and \xrightarrow{as} is almost sure convergence and

$$\sigma_I^2 = s_{32} - s_{21}^2 \text{ with } s_{ij} = \sum_k \frac{\alpha_k^i}{|I_k|^j}, \forall (i, j) \in \mathbb{N}^2.$$

Lemma 4. Let I, J be two partitions in \mathcal{I} , then I is called a subdivision of J and we denote $I \trianglelefteq J$, if $\mathcal{F}_J \subset \mathcal{F}_I$. Suppose that function f satisfies Assumption 1, let us consider m_{max} large enough such that $\mu^* - \lambda^* > 2^{1-m_{max}}$. Define $N = 2^{m_{max}}$ and $I^{(N)} = (N, \lambda_N, \mu_N) \in \mathcal{I}$ with $\lambda_N = \lceil N\lambda^* \rceil / N$, $\mu_N = \lfloor N\mu^* \rfloor / N$. Then for every partition $I \in \mathcal{I}$, we have

i) If I is a subdivision of $I^{(N)}$, then $L(I) = L(I^{(N)})$.

ii) If I is not a subdivision of $I^{(N)}$, then $L(I) > L(I^{(N)})$.

In the two following theorems, we shall improve the properties of estimator $\hat{\theta}_n$ with respect to those obtained in Theorem 2.1 in [Celisse and Robin \(2010\)](#). Indeed, the latter establish only a convergence in probability, while we shall prove the almost sure convergence and \sqrt{n} -consistency of $\hat{\theta}_n$.

Theorem 2. (Almost sure convergence of $\hat{\theta}_n$). Suppose that density f satisfies Assumption 1 and furthermore,

$$\forall I \in \mathcal{I}, \quad 8s_{11}s_{21} - 2s_{11}^2 + 8s_{32} - 10s_{21}^2 - 4s_{22} \neq 0, \quad s_{21} - s_{22} - s_{32} + 3s_{11} \neq 0. \quad (10)$$

Then for m_{max} large enough, we have $\hat{\theta}_n \xrightarrow[n \rightarrow \infty]{as} \theta$.

Proof. First, we remark that under conditions (10), [Celisse and Robin](#) prove in their Proposition 2.1 that

$$\frac{\hat{p}(I)}{n} \xrightarrow[n \rightarrow \infty]{as} l_\infty(I) \in [0, 1).$$

Denoting by $\Lambda^* = [\lambda^*, \mu^*]$ and $\hat{\Lambda} = [\hat{\lambda}, \hat{\mu}]$, we may write

$$\begin{aligned} \hat{\theta}_n &= \theta + (\hat{\theta}_n - \theta) \mathbf{1}_{\hat{\Lambda} \subseteq \Lambda^*} + (\hat{\theta}_n - \theta) \mathbf{1}_{\hat{\Lambda} \not\subseteq \Lambda^*} \\ &= \theta + \sum_{I=(2^m, \lambda, \mu): [\lambda, \mu] \subseteq \Lambda^*} \left[\frac{1}{n(\mu - \lambda)} \sum_{i=1}^n \mathbf{1}\{X_i \in [\lambda, \mu]\} - \theta \right] \mathbf{1}\{\hat{\lambda} = \lambda, \hat{\mu} = \mu\} \\ &\quad + (\hat{\theta}_n - \theta) \mathbf{1}_{\hat{\Lambda} \not\subseteq \Lambda^*}. \end{aligned} \quad (11)$$

For each partition $I = (2^m, \lambda, \mu)$ such that $[\lambda, \mu] \subseteq \Lambda^*$, applying the strong law of large numbers we get that

$$\frac{1}{n(\mu - \lambda)} \sum_{i=1}^n \mathbf{1}\{X_i \in [\lambda, \mu]\} \xrightarrow[n \rightarrow \infty]{as} \frac{\mathbb{P}(X_i \in [\lambda, \mu])}{\mu - \lambda} = \theta.$$

Since the cardinality $card(\mathcal{I})$ of \mathcal{I} is finite and does not depend on n , in order to finish the proof, it is sufficient to establish that

$$(\hat{\theta}_n - \theta) \mathbf{1}_{\hat{\Lambda} \not\subseteq \Lambda^*} \xrightarrow[n \rightarrow \infty]{as} 0.$$

In fact, if $\hat{\Lambda} \not\subseteq \Lambda^*$ then \hat{I} is not a subdivision of $I^{(N)}$. Using Lemma 4, we have $L(\hat{I}) > L(I^{(N)})$. Let

$$\gamma = \min_{I \not\subseteq I^{(N)}} L(I) - L(I^{(N)}) > 0, \quad (12)$$

where $I \not\subseteq I^{(N)}$ means that I is not a subdivision of $I^{(N)}$. We obtain that

$$\begin{aligned} |\hat{\theta}_n - \theta| \mathbf{1}_{\hat{\Lambda} \not\subseteq \Lambda^*} &\leq (N + \theta) \mathbf{1}\{L(\hat{I}) - L(I^{(N)}) \geq \gamma\} \leq \\ &(N + \theta) \mathbf{1}\{|\hat{L}_{\hat{p}(\hat{I})}(\hat{I}) - L(\hat{I})| + |\hat{L}_{\hat{p}(I^{(N)})}(I^{(N)}) - L(I^{(N)})| + \hat{L}_{\hat{p}(\hat{I})}(\hat{I}) - \hat{L}_{\hat{p}(I^{(N)})}(I^{(N)}) \geq \gamma\} \\ &\leq (N + \theta) \mathbf{1}\{2 \sup_{I \in \mathcal{I}} |\hat{L}_{\hat{p}(I)}(I) - L(I)| + \hat{L}_{\hat{p}(\hat{I})}(\hat{I}) - \hat{L}_{\hat{p}(I^{(N)})}(I^{(N)}) \geq \gamma\}. \end{aligned}$$

By definition of \hat{I} , we have $\hat{L}_{\hat{p}(\hat{I})}(\hat{I}) - \hat{L}_{\hat{p}(I^{(N)})}(I^{(N)}) \leq 0$, so that

$$\begin{aligned} |\hat{\theta}_n - \theta| \mathbf{1}_{\hat{\Lambda} \not\subseteq \Lambda^*} &\leq (N + \theta) \mathbf{1}\{\sup_{I \in \mathcal{I}} |\hat{L}_{\hat{p}(I)}(I) - L(I)| \geq \frac{\gamma}{2}\} \\ &\leq (N + \theta) \sum_{I \in \mathcal{I}} \mathbf{1}\{|\hat{L}_{\hat{p}(I)}(I) - L(I)| \geq \frac{\gamma}{2}\}. \end{aligned} \quad (13)$$

Since $\forall I \in \mathcal{I}$, we both have $\hat{L}_p(I) \xrightarrow[n \rightarrow \infty]{as} L(I)$ and $\hat{p}(I)/n \xrightarrow[n \rightarrow \infty]{as} l_\infty(I) \in [0, 1]$ as well as the fact that $\hat{R}_p(I)$ (given by (9)) is a continuous function of p/n , we obtain $\hat{L}_{\hat{p}(I)}(I) \xrightarrow[n \rightarrow \infty]{as} L(I)$. Therefore,

$$\mathbf{1}\{|\hat{L}_{\hat{p}(I)}(I) - L(I)| \geq \frac{\gamma}{2}\} \xrightarrow[n \rightarrow \infty]{as} 0.$$

Indeed, if $X_n \xrightarrow{as} X$ then $\forall \epsilon > 0$, we have $\mathbf{1}\{|X_n - X| \geq \epsilon\} \xrightarrow{as} 0$. It thus follows that $(\hat{\theta}_n - \theta) \mathbf{1}_{\hat{\Lambda} \not\subseteq \Lambda^*} \xrightarrow{as} 0$. We finally get that $\hat{\theta}_n \xrightarrow{as} \theta$. \square

Remark 3. As previously noted in ii) from Remark 1, an oracle version of *Celisse and Robin*'s estimator relying on the true interval Λ^* would have asymptotically minimal variance and thus be efficient. However, from an expression such as (11), it is not possible to ensure in general that $\sqrt{n}\hat{\theta}_n$ has asymptotic variance equal to $\theta(|\Lambda^*|^{-1} - \theta)$.

In the following, the expression $O_{\mathbb{P}}(1)$ denotes a sequence that is bounded in probability. A sequence estimator T_n of θ is said to be \sqrt{n} -consistent if $\sqrt{n}(T_n - \theta) = O_{\mathbb{P}}(1)$. We remark that if T_n is asymptotically normal then T_n is \sqrt{n} -consistent but the converse of

this statement is not true. In the following theorem, we shall prove the \sqrt{n} -consistency of $\hat{\theta}_n$ but we were not able to prove its asymptotically normality. However, according to Prohorov's theorem, there exists a subsequence of $\{\sqrt{n}(\hat{\theta}_n - \theta)\}_n$ that converges in distribution to some random variable Z .

Theorem 3. (\sqrt{n} -consistency of $\hat{\theta}_n$). *Suppose that density f satisfies the hypotheses of Theorem 2, then for m_{max} large enough, $\hat{\theta}_n$ is \sqrt{n} -consistent.*

Proof. We may write as in the proof of Theorem 2,

$$\begin{aligned} \sqrt{n}(\hat{\theta}_n - \theta) &= \sum_{I=(2^m, \lambda, \mu): [\lambda, \mu] \subseteq \Lambda^*} \sqrt{n} \left[\frac{1}{n(\mu - \lambda)} \sum_{i=1}^n \mathbf{1}\{X_i \in [\lambda, \mu]\} - \theta \right] \mathbf{1}_{\{\hat{\lambda}=\lambda, \hat{\mu}=\mu\}} \\ &\quad + \sqrt{n}(\hat{\theta}_n - \theta) \mathbf{1}_{\{\hat{\Lambda} \not\subseteq \Lambda^*\}}. \end{aligned}$$

For each partition $I = (2^m, \lambda, \mu)$ such that $[\lambda, \mu] \subset \Lambda^*$, by applying the central limit theorem, we get that

$$\sqrt{n} \left[\frac{1}{n(\mu - \lambda)} \sum_{i=1}^n \mathbf{1}_{X_i \in [\lambda, \mu]} - \theta \right] \xrightarrow[n \rightarrow \infty]{d} \mathcal{N} \left(0, \theta \left(\frac{1}{\mu - \lambda} - \theta \right) \right).$$

Hence, using again that $\text{card}(\mathcal{I})$ is finite,

$$\sum_{I=(2^m, \lambda, \mu): [\lambda, \mu] \subseteq \Lambda^*} \sqrt{n} \left[\frac{1}{n(\mu - \lambda)} \sum_{i=1}^n \mathbf{1}_{X_i \in [\lambda, \mu]} - \theta \right] \mathbf{1}_{\hat{\lambda}=\lambda, \hat{\mu}=\mu} = O_{\mathbb{P}}(1). \quad (14)$$

We shall now prove that $\sqrt{n}(\hat{\theta}_n - \theta) \mathbf{1}_{\hat{\Lambda} \not\subseteq \Lambda^*} \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0$. In fact, according to (13), for all $\epsilon > 0$, we have

$$\begin{aligned} \mathbb{P}(\sqrt{n}|\hat{\theta}_n - \theta| \mathbf{1}_{\hat{\Lambda} \not\subseteq \Lambda^*} > \epsilon) &\leq \mathbb{P}(\hat{\Lambda} \not\subseteq \Lambda^*) \\ &\leq \mathbb{P}(\sup_{I \in \mathcal{I}} |\hat{L}_{\hat{\rho}(I)}(I) - L(I)| \geq \frac{\gamma}{2}) \\ &\leq \sum_{I \in \mathcal{I}} \mathbb{P}(|\hat{L}_{\hat{\rho}(I)}(I) - L(I)| \geq \frac{\gamma}{2}) \xrightarrow[n \rightarrow \infty]{} 0, \end{aligned}$$

where γ is defined by (12). Therefore, $\sqrt{n}(\hat{\theta}_n - \theta) \mathbf{1}_{\hat{\Lambda} \not\subseteq \Lambda^*} = o_{\mathbb{P}}(1)$. We finally conclude that $\sqrt{n}(\hat{\theta}_n - \theta) = O_{\mathbb{P}}(1)$. \square

Note that the difficulty in establishing asymptotic normality of $\hat{\theta}_n$ comes from the fact that there are many different terms in the left-hand side of equation (14) and we did not succeed in proving that $\hat{I} = I^{(N)}$ a.s for large enough n .

2.3 General overview of one-step estimators

In this section, we introduce the one-step method to construct an asymptotically efficient estimator, relying on a \sqrt{n} -consistent one (see [van der Vaart, 1998](#), Section 25.8). Let $\hat{\theta}_n$ be a \sqrt{n} -consistent estimator of θ , then $\hat{\theta}_n$ can be discretized on grids of mesh width

$n^{-1/2}$. Suppose that we are given a sequence of estimators $\hat{l}_{n,\theta}(\cdot) = \hat{l}_{n,\theta}(\cdot; X_1, \dots, X_n)$ of the efficient score function $\tilde{l}_{\theta,f}$. Define with $m = \lfloor n/2 \rfloor$,

$$\hat{l}_{n,\theta,i}(\cdot) = \begin{cases} \hat{l}_{m,\theta}(\cdot; X_1, \dots, X_m) & \text{if } i > m, \\ \hat{l}_{n-m,\theta}(\cdot; X_{m+1}, \dots, X_n) & \text{if } i \leq m. \end{cases}$$

Thus, for X_i ranging through each of the two halves of the sample, we use an estimator $\hat{l}_{n,\theta,i}$ based on the other half of the sample. We assume that, for every deterministic sequence $\theta_n = \theta + O(n^{-1/2})$, we have

$$\sqrt{n} \mathbb{P}_{\theta_n, f} \hat{l}_{n,\theta_n} \xrightarrow[n \rightarrow \infty]{\mathbb{P}_{\theta, f}} 0, \quad (15)$$

$$\mathbb{P}_{\theta_n, f} \|\hat{l}_{n,\theta_n} - \tilde{l}_{\theta_n, f}\|^2 \xrightarrow[n \rightarrow \infty]{\mathbb{P}_{\theta, f}} 0, \quad (16)$$

$$\int \|\tilde{l}_{\theta_n, f} d\mathbb{P}_{\theta_n, f}^{1/2} - \tilde{l}_{\theta, f} d\mathbb{P}_{\theta, f}^{1/2}\|^2 \xrightarrow[n \rightarrow \infty]{} 0. \quad (17)$$

Then, the one-step estimator defined as

$$\tilde{\theta}_n = \hat{\theta}_n - \left(\sum_{i=1}^n \hat{l}_{n,\hat{\theta}_n,i}^2(X_i) \right)^{-1} \sum_{i=1}^n \hat{l}_{n,\hat{\theta}_n,i}(X_i),$$

is asymptotically efficient at (θ, f) (see [van der Vaart, 1998](#), Section 25.8). This estimator $\tilde{\theta}_n$ can be considered a one-step iteration of the Newton-Raphson algorithm for solving an approximation of the equation $\sum_i \tilde{l}_{\theta, f}(X_i) = 0$ with respect to θ , starting at the initial guess $\hat{\theta}_n$.

In the next section, we discuss a converse result on necessary conditions for existence of an asymptotically efficient estimator of θ and its implications in model \mathcal{P}_δ .

2.4 Existence of efficient estimators of θ in model \mathcal{P}_δ

Under condition (17), it is shown in Theorem 7.4 from [van der Vaart \(2002\)](#) that the existence of an asymptotically efficient sequence of estimators of θ implies the existence of a sequence of estimators $\hat{l}_{n,\theta}$ of $\tilde{l}_{\theta,f}$ satisfying (15) and (16). In our case, it is not difficult to prove that condition (17) holds. As a consequence, we obtain the following proposition.

Proposition 1. *The existence of an asymptotically efficient sequence of estimators of θ in model \mathcal{P}_δ is equivalent to the existence of a sequence of estimators $\hat{l}_{n,\theta}$ of the efficient score function $\tilde{l}_{\theta,f}$ satisfying (15) and (16).*

Proof. Let us first establish that condition (17) holds. In fact, with the notation $p_{\theta, f} =$

$\theta + (1 - \theta)f$, we have

$$\begin{aligned}
& \int \|\tilde{l}_{\theta_n, f} d\mathbb{P}_{\theta_n, f}^{1/2} - \tilde{l}_{\theta, f} d\mathbb{P}_{\theta, f}^{1/2}\|^2 = \int_0^1 \left(\tilde{l}_{\theta_n, f}(x) \sqrt{p_{\theta_n, f}(x)} - \tilde{l}_{\theta, f}(x) \sqrt{p_{\theta, f}(x)} \right)^2 dx \\
& \leq 2 \int_0^1 (\tilde{l}_{\theta_n, f}(x) - \tilde{l}_{\theta, f}(x))^2 p_{\theta_n, f}(x) dx + 2 \int_0^1 \tilde{l}_{\theta, f}^2(x) \left(\sqrt{p_{\theta_n, f}(x)} - \sqrt{p_{\theta, f}(x)} \right)^2 dx \\
& \leq 2 \int_0^1 \left[\frac{1}{\theta_n} - \frac{1}{\theta} + \left(\frac{1}{\theta(1 - \theta\delta)} - \frac{1}{\theta_n(1 - \theta_n\delta)} \right) \mathbf{1}_{\{f(x) > 0\}} \right]^2 p_{\theta_n, f}(x) dx \\
& \quad + 2 \int_0^1 \left[\frac{1}{\theta} - \frac{1}{\theta(1 - \theta\delta)} \mathbf{1}_{\{f(x) > 0\}} \right]^2 \frac{(\theta_n - \theta)^2 (1 - f(x))^2}{\left(\sqrt{p_{\theta_n, f}(x)} + \sqrt{p_{\theta, f}(x)} \right)^2} dx \\
& \leq 2 \int_0^1 (\theta_n - \theta)^2 \left[\frac{1}{\theta\theta_n} + \frac{\delta(\theta + \theta_n) + 1}{\theta\theta_n(1 - \theta\delta)(1 - \theta_n\delta)} \mathbf{1}_{\{f(x) > 0\}} \right]^2 p_{\theta_n, f}(x) dx \\
& \quad + 2 \int_0^1 (\theta_n - \theta)^2 \left[\frac{1}{\theta^2} + \frac{1}{\theta^2(1 - \theta)^2} \right] \frac{(1 - f(x))^2}{(\sqrt{\theta_n} + \sqrt{\theta})^2} dx \\
& \leq (\theta_n - \theta)^2 \left[\frac{C}{\theta^2} + \frac{C(1 + 2C\theta)}{\theta^2(1 - \theta)^2} \right]^2 + C(\theta_n - \theta)^2 \left[\frac{1}{\theta^3} + \frac{1}{\theta^3(1 - \theta)^2} \right] = O\left(\frac{1}{n}\right),
\end{aligned}$$

where C is some positive constant. Thus, according to Theorem 7.4 from [van der Vaart \(2002\)](#), the existence of an asymptotically efficient sequence of estimators of θ is equivalent to the existence of a sequence of estimators $\hat{l}_{n, \theta}$ satisfying (15) and (16). \square

The estimator $\hat{l}_{n, \theta}$ of the efficient score function $\tilde{l}_{\theta, f}$ must satisfy both a "no-bias" (15) and a consistency (16) condition. The consistency is usually easy to arrange, but the "no-bias" condition requires a convergence to zero of the bias at a rate faster than $1/\sqrt{n}$. In our model, the efficient score function $\tilde{l}_{\theta, f}$ is given by

$$\tilde{l}_{\theta, f}(x) = \frac{1}{\theta} - \frac{1}{\theta(1 - \theta\delta)} \mathbf{1}_{\{f(x) > 0\}},$$

so that we must estimate the set I_f and its measure δ in order to estimate $\tilde{l}_{\theta, f}$. To our knowledge, the rate of convergence of (the bias of) such level-sets estimators is nonparametric (see [Baillio et al., 2001](#); [Cadre, 2006](#); [Mason and Polonik, 2009](#)). Thus, it is likely that there does not exist an asymptotically efficient sequence of estimators of θ in model \mathcal{P}_δ .

3 Asymptotically efficient estimators of θ when f vanishes on a set with null Lebesgue measure

We now consider a second submodel of \mathcal{P} given by $\mathcal{P}_0 = \{p_{\theta, f} \in \mathcal{P} : \theta \in (0, 1), f \in \mathcal{F}_0\}$, where $\mathcal{F}_0 = \{\text{density } f \text{ on } [0, 1] \text{ such that } \mu(I_f) = 0\}$. This model is a limiting case of model \mathcal{P}_δ when $\delta \rightarrow 0$. It corresponds to a difficult case with respect to estimation in model (1), where density f vanishes only on a subset of null Lebesgue measure. We recall that estimators constructed under this setup (and assuming moreover either regularity or monotonicity properties on f) exhibit nonparametric rates of convergence. Besides, the

results of Theorem 1 suggest that no estimator of θ can have finite asymptotic variance at rate \sqrt{n} in this case. By using for instance the path given by Equation (4), we obtain a tangent set for f ,

$$\dot{\mathcal{P}}_{f,0} = \left\{ h \in \mathbb{L}^2(\mathbb{P}_{\theta,f}); \exists h_0 \in \mathbb{L}^1(\mathbb{P}_{\theta,f}), h = \frac{(1-\theta)fh_0}{\theta + (1-\theta)f} \text{ with } \int_0^1 fh_0 = 0 \right\}.$$

In the following lemma, we shall prove that the ordinary score $\dot{l}_{\theta,f}$ belongs to this tangent set $\dot{\mathcal{P}}_{f,0}$. So the efficient information matrix $\tilde{I}_{\theta,f}$ for θ is equal to 0 and we can prove that relative to the tangent set $\dot{\mathcal{P}}_0 = \text{lin}(\dot{l}_{\theta,f} + \dot{\mathcal{P}}_{f,0})$, there is no regular estimator sequence for θ or equivalently, the minimax quadratic risk is not finite.

Lemma 5. *The efficient information matrix $\tilde{I}_{\theta,f}$ for θ relative to the tangent set $\dot{\mathcal{P}}_0$ is equal to 0. Moreover, the functional $\psi(\mathbb{P}_{\theta,f}) = \theta$ is not differentiable at $\mathbb{P}_{\theta,f}$ relative to the tangent set $\dot{\mathcal{P}}_0$.*

Proof. First, we prove that the ordinary score function $\dot{l}_{\theta,f}$ for the parameter θ belongs to this tangent set $\dot{\mathcal{P}}_{f,0}$. Indeed, let the function

$$h_0 = \frac{1-f}{(1-\theta)f} \mathbf{1}_{\{f>0\}}.$$

Since $\mu(I_f) = 0$, we have $\int fh_0 = 0$ and according to (6), we obtain

$$\dot{l}_{\theta,f} = \frac{(1-\theta)fh_0}{\theta + (1-\theta)f}, \quad \mu - \text{almost everywhere,}$$

so that $\dot{l}_{\theta,f} \in \dot{\mathcal{P}}_{f,0}$. Then, the efficient information matrix $\tilde{I}_{\theta,f}$ is equal to 0. Now, we show that the functional $\psi(\mathbb{P}_{\theta,f}) = \theta$ is not differentiable at $\mathbb{P}_{\theta,f}$ relative to the tangent set $\dot{\mathcal{P}}_0 = \text{lin}(\dot{l}_{\theta,f} + \dot{\mathcal{P}}_{f,0}) = \dot{\mathcal{P}}_{f,0}$. In fact, if this were true, there would exist a function $\tilde{\psi}_{\theta,f}$ such that

$$a = \left. \frac{\partial}{\partial t} \psi(\mathbb{P}_{\theta+ta, f_t}) \right|_{t=0} = \langle \tilde{\psi}_{\theta,f}, a\dot{l}_{\theta,f} + h \rangle, \quad \forall a \in \mathbb{R}, h \in \dot{\mathcal{P}}_{f,0},$$

where $\langle \cdot, \cdot \rangle$ denotes scalar product in $\mathbb{L}^2(\mathbb{P}_{\theta,f})$. Choosing $h = \dot{l}_{\theta,f}$, we obtain $a = (a+1)\langle \tilde{\psi}_{\theta,f}, \dot{l}_{\theta,f} \rangle$ for every value $a \in \mathbb{R}$, which is impossible. \square

Theorem 4. *For any estimator sequence T_n we have,*

$$\sup_E \liminf_{n \rightarrow \infty} \sup_{g \in E} \mathbb{E}_{P_{1/\sqrt{n},g}} [\sqrt{n}(T_n - \psi(P_{1/\sqrt{n},g}))]^2 = +\infty,$$

where the first supremum is taken over all finite subsets E of the tangent set $\dot{\mathcal{P}}_0$.

Proof. We first remark that the tangent set $\dot{\mathcal{P}}_0$ is a linear subspace of $\mathbb{L}^2(\mathbb{P}_{\theta,f})$ with infinite dimension. So we can choose an orthonormal basis $\{h_i\}_{i=1}^{\infty}$ of $\dot{\mathcal{P}}_0$ such that for every m , $\dot{l}_{\theta,f} \notin \dot{\mathcal{P}}_{0,m} := \text{lin}(h_1, h_2, \dots, h_m)$. We thus have

$$\begin{aligned} \sup_E \liminf_{n \rightarrow \infty} \sup_{g \in E} \mathbb{E}_{P_{1/\sqrt{n},g}} [\sqrt{n}(T_n - \psi(P_{1/\sqrt{n},g}))]^2 \\ \geq \sup_F \liminf_{n \rightarrow \infty} \sup_{g \in F} \mathbb{E}_{P_{1/\sqrt{n},g}} [\sqrt{n}(T_n - \psi(P_{1/\sqrt{n},g}))]^2, \end{aligned}$$

where E and F range through all finite subsets of the tangent sets $\dot{\mathcal{P}}_0$ and $\text{lin}(\dot{l}_{\theta,f} + \dot{\mathcal{P}}_{0,m})$, respectively. The efficient score function for θ corresponding to the tangent set $\text{lin}(\dot{l}_{\theta,f} + \dot{\mathcal{P}}_{0,m})$ is

$$\tilde{l}_{\theta,f,m} = \dot{l}_{\theta,f} - \sum_{i=1}^m \langle \dot{l}_{\theta,f}, h_i \rangle h_i \neq 0,$$

where $\langle \cdot, \cdot \rangle$ stands for the scalar product in $\mathbb{L}^2(\mathbb{P}_{\theta,f})$. Moreover, the efficient information matrix $\tilde{I}_{\theta,f,m} = \mathbb{P}_{\theta,f}(\tilde{l}_{\theta,f,m}^2)$ is nonsingular. Using Lemma 25.25 from [van der Vaart \(1998\)](#), we remark that the functional $\psi(\mathbb{P}_{\theta,f}) = \theta$ is differentiable at $\mathbb{P}_{\theta,f}$ relative to the tangent set $\text{lin}(\dot{l}_{\theta,f} + \dot{\mathcal{P}}_{0,m})$ with efficient influence function $\tilde{\psi}_{\theta,f,m} = \tilde{I}_{\theta,f,m}^{-1} \tilde{l}_{\theta,f,m}$. So we can apply Theorem 25.21 from [van der Vaart \(1998\)](#) to obtain that

$$\sup_F \liminf_{n \rightarrow \infty} \sup_{g \in F} \mathbb{E}_{P_{1/\sqrt{n},g}} [\sqrt{n}(T_n - \psi(P_{1/\sqrt{n},g}))]^2 \geq \tilde{I}_{\theta,f,m}^{-1}.$$

Since $\tilde{I}_{\theta,f,m} \xrightarrow{m \rightarrow \infty} 0$, we have

$$\sup_E \liminf_{n \rightarrow \infty} \sup_{g \in E} \mathbb{E}_{P_{1/\sqrt{n},g}} [\sqrt{n}(T_n - \psi(P_{1/\sqrt{n},g}))]^2 \geq \tilde{I}_{\theta,f,m}^{-1} \xrightarrow{m \rightarrow \infty} +\infty.$$

□

Remark 4. Using Theorem 2 in [Chamberlain \(1986\)](#), we can conclude that there is no regular estimator sequence for θ relative to the tangent set $\dot{\mathcal{P}}_0$. This result implies that if there exists a \sqrt{n} -consistent estimator in model \mathcal{P}_0 , it can not have finite asymptotic variance. It does not imply that \sqrt{n} -consistent estimators do not exist in model \mathcal{P}_0 , namely, we could have $\sqrt{n}(\hat{\theta} - \theta) = O_{\mathbb{P}}(1)$ for some estimator $\hat{\theta}$ but then $\text{Var}(\sqrt{n}\hat{\theta}) \rightarrow +\infty$. However, we note that the only rates of convergence obtained until now in this case are nonparametric ones.

4 Simulations

In this section, we give some illustrations of the previous results on some simulated experiments and explore the non asymptotic performances of the estimators previously discussed. We choose to study the rates of convergence of [Celisse and Robin \(2010\)](#)'s estimator and a one-step estimator of $\bar{\theta}$, the latter being constructed from the former. For each estimator $\hat{\theta}_n$ of $\bar{\theta}$, we compare the quantity $n\mathbb{E}[(\hat{\theta}_n - \bar{\theta})^2]$ with the optimal variance $\bar{\theta}(\delta^{-1} - \bar{\theta})$ when this bound exists. Equivalently, we compare the logarithm of mean squared error (MSE) of each estimator $\hat{\theta}_n$ with $-\log(n) + \log(\bar{\theta}(\delta^{-1} - \bar{\theta}))$. When the alternative density f is constant on a set with null Lebesgue measure, we only compare the slope of the line induced by $\log(MSE)$ with the parametric rate -1 . We choose to investigate the behaviour of these two different estimators of $\bar{\theta}$ under two different models. The first one corresponds to the double exponential model already mentioned in the introduction. The second model we consider was introduced in [Celisse and Robin \(2010\)](#). In each case, we simulated data with sample size $n \in \{1000; 2000; 3000; 3500; 4000; 4500; 5000\}$ and perform $R = 100$ repetitions.

In the first model, the p -value corresponding to the statistic T is defined by $p\text{-value}(T) = 1 - \Phi_0(T)$, where

$$\Phi_0(t) = \begin{cases} \frac{1}{2} \exp(t) & \text{if } t \leq 0, \\ 1 - \frac{1}{2} \exp(-t) & \text{if } t > 0, \end{cases}$$

and the density function of the p -values under the alternative is given by

$$f(x) = \begin{cases} \exp(\mu) & \text{if } 0 \leq x \leq \frac{1}{2} \exp(-\mu), \\ \exp(-\mu)/(4x^2) & \text{if } \frac{1}{2} \exp(-\mu) \leq x \leq \frac{1}{2}, \\ \exp(-\mu) & \text{if } \frac{1}{2} \leq x \leq 1. \end{cases}$$

As already mentioned in the introduction, in this case we can only estimate $\bar{\theta} = \theta + (1 - \theta)e^{-\mu}$. In the second model, we simulate p -values under the alternative with distribution

$$f(x) = \frac{s}{\lambda} \left(1 - \frac{x}{\lambda}\right)^{s-1} \mathbf{1}_{[0,\lambda]}(x),$$

where $\lambda \in (0, 1)$ and $s > 1$. In this case, the infimum of f is zero and we exactly recover θ . Note that density f is always decreasing, convex when $s \geq 2$ and concave when $s \in (1, 2]$. As an illustration, we represent the densities of the statistics and corresponding p -values in Figure 1 for the first model and of the p -values in Figure 2 for the second model. The parameter values chosen for the experiments are described in Table 1.




Figure 1: Density functions of the statistics and p -values of the first model with $\theta = 0.7$ and $\mu = 2$.

In order to obtain a one-step estimator of $\bar{\theta}$, starting from [Celisse and Robin \(2010\)](#)'s preliminary estimator, we estimate the efficient score function $\tilde{l}_{\theta,f}$ by

$$\hat{l}_{n,\theta}(x; X_1, \dots, X_n) = \frac{1}{\bar{\theta}} - \frac{1}{\bar{\theta}(1 - \bar{\theta}\hat{\delta}_n(\bar{\theta}))} \mathbf{1}_{\{\hat{g}_n(x) > \bar{\theta}\}},$$

where \hat{g}_n is a kernel density estimator of the function g and $\hat{\delta}_n(\bar{\theta}) = 1 - \mu(\{x : \hat{g}_n(x) > \bar{\theta}\})$. This may be done only when the alternative density f is constant (or vanishes) on an




Figure 2: Density function of the p -values of the second model. Left: $s = 1.3, \lambda = 0.45$ and $\theta = 0.86$; Right: $s = 4, \lambda = 1$ and $\theta = 0.7$.

	Model 1 (μ, θ)	Model 2 (s, λ, θ)
(a)	(2, 0.7)	(4, 0.75, 0.9)
(b)	(2, 0.85)	(1.3, 0.45, 0.86)
(c)	($\sqrt{13}$, 0.7)	(4, 1, 0.7)
(d)	(5, 0.87)	(1.4, 1, 0.86)

Table 1: Parameter values in each model.

interval. Thus, in cases (c) and (d) from model 2, we do not have a one-step estimator. The results are presented in Figures 3 (first model) and 4 (second model).

According to Figures 3 and 4, the rates of convergence of the estimators are both non-parametric. In some cases, the rate of [Celisse and Robin \(2010\)](#)'s estimator is nearly parametric (green line with approximately the same slope as the red line) but with a variance always larger than the optimal bound (y -intercept larger for the green line than for the red one). Moreover, the rate of convergence of the one-step estimator is always slower. This may be due to the non-parametric rate of convergence of the kernel density estimator of g used in the estimation of the efficient score function. Thus, it seems that in these cases, the one-step method does not even keep the parametric rate of convergence of the preliminary estimator it relies on.

Acknowledgments The authors are grateful to Cyril Dalmasso, Elisabeth Gassiat and Pierre Neuvial for fruitful discussions concerning this work.




Figure 3: Logarithm of the mean squared error as a function of $\log(n)$ and corresponding linear regression for [Celisse and Robin \(2010\)](#)'s estimator (\bullet and green line, respectively) and the one-step estimator (\circ and blue line, respectively) in the first model, for different parameter values ((a) top left; (b) top right; (c) bottom left; (d) bottom right). Red line represents the line $y = -\log(n) + \log(\bar{\theta}(\delta^{-1}) - \bar{\theta})$.




Figure 4: Logarithm of the mean squared error as a function of $\log(n)$ and corresponding linear regression for [Celisse and Robin \(2010\)](#)'s estimator (\bullet and green line, respectively) and the one-step estimator (\circ and blue line, respectively, upper cases only) in the second model, for different parameter values ((a) top left; (b) top right; (c) bottom left; (d) bottom right). Red line represents the line $y = -\log(n) + \log(\theta(\delta^{-1} - \theta))$ (upper cases) and the line $y = -\log(n) + c$ for some well chosen constant c (bottom cases).

A Appendix. Proofs of technical lemmas

A.1 Proof of Lemma 1

Note that [Celisse and Robin \(2010\)](#) prove that $\mathbb{E}[\|g - \hat{g}_I\|_2^2] \xrightarrow{n \rightarrow \infty} 0$, while we further establish that it is $O(1/n)$. By a simple bias-variance decomposition, we may write

$$\mathbb{E}[\|g_I - \hat{g}_I\|_2^2] = \mathbb{E}[\|g - \hat{g}_I\|_2^2] - \|g_I - g\|_2^2.$$

As for the bias term, it is easy to show that

$$\begin{aligned} \|g - g_I\|_2^2 &= \inf_{h \in \mathcal{F}_I} \|g - h\|_2^2 \\ &= \inf_{(a_k)_k \in \mathbb{R}} \left[\|g\|_2^2 - 2 \int_0^1 \left(\sum_k a_k \mathbf{1}_{I_k}(x) \right) g(x) dx + \int_0^1 \left(\sum_k a_k \mathbf{1}_{I_k}(x) \right)^2 dx \right] \\ &= \inf_{(a_k)_k \in \mathbb{R}} \left[\|g\|_2^2 - 2 \sum_k a_k \alpha_k + \sum_k a_k^2 |I_k| \right] \\ &= \|g\|_2^2 - \sum_k \frac{\alpha_k^2}{|I_k|} = \|g\|_2^2 - s_{21}. \end{aligned} \tag{18}$$

Let us now calculate the mean square error of \hat{g}_I

$$\begin{aligned} \mathbb{E}[\|g - \hat{g}_I\|_2^2] &= \|g\|_2^2 + \mathbb{E} \left[\|\hat{g}_I\|_2^2 - 2 \int_0^1 \hat{g}_I(x) g(x) dx \right] \\ &= \|g\|_2^2 + \mathbb{E} \left[\int_0^1 \left(\sum_k \frac{n_k}{n|I_k|} \mathbf{1}_{I_k}(x) \right)^2 dx - 2 \int_0^1 \sum_k \frac{n_k}{n|I_k|} \mathbf{1}_{I_k}(x) g(x) dx \right] \\ &= \|g\|_2^2 + \mathbb{E} \left[\sum_k \frac{n_k^2}{n^2|I_k|} - 2 \sum_k \frac{n_k \alpha_k}{n|I_k|} \right]. \end{aligned}$$

Since n_k follows a Binomial distribution $\mathcal{B}(n, \alpha_k)$, we have

$$\mathbb{E}[n_k] = n\alpha_k \text{ and } \mathbb{E}[n_k^2] = n^2\alpha_k^2 + n\alpha_k(1 - \alpha_k).$$

Therefore,

$$\begin{aligned} \mathbb{E}[\|g - \hat{g}_I\|_2^2] &= \|g\|_2^2 + \sum_k \frac{n^2\alpha_k^2 + n\alpha_k(1 - \alpha_k)}{n^2|I_k|} - 2 \sum_k \frac{n\alpha_k^2}{n|I_k|} \\ &= \|g\|_2^2 - s_{21} + \frac{1}{n}(s_{11} - s_{21}). \end{aligned} \tag{19}$$

Using (18) and (19), we obtain the desired result, namely

$$\mathbb{E}[\|g_I - \hat{g}_I\|_2^2] = \mathbb{E}[\|g - \hat{g}_I\|_2^2] - \|g_I - g\|_2^2 = \frac{1}{n}(s_{11} - s_{21}) = O\left(\frac{1}{n}\right).$$

A.2 Proof of Lemma 2

i) Since

$$\lim_{n \rightarrow \infty} \frac{p}{n} < 1 \text{ and } \frac{n_k}{n} \xrightarrow[n \rightarrow \infty]{as} \alpha_k, \text{ for all } k,$$

we obtain that

$$\begin{aligned} \hat{L}_p(I) &= \|g\|_2^2 + \frac{2n-p}{(n-1)(n-p)} \sum_k \frac{n_k}{n|I_k|} - \frac{n(n-p+1)}{(n-1)(n-p)} \sum_k \frac{1}{|I_k|} \left(\frac{n_k}{n}\right)^2 \\ &\xrightarrow[n \rightarrow \infty]{as} \|g\|_2^2 - \sum_k \frac{\alpha_k^2}{|I_k|} = \|g\|_2^2 - s_{21} = \|g_I - g\|_2^2 = L(I). \end{aligned}$$

ii) By definition of $R(I)$ and using (19), we have

$$R(I) = \mathbb{E}[\|g - \hat{g}_I\|_2^2] - \|g\|_2^2 = -s_{21} + \frac{1}{n}(s_{11} - s_{21}).$$

This gives that

$$\begin{aligned} \sqrt{n}[\hat{R}_p(I) - R(I)] &= \sqrt{n} \left[\frac{2n-p}{(n-1)(n-p)} \sum_k \frac{n_k}{n|I_k|} - \frac{n(n-p+1)}{(n-1)(n-p)} \sum_k \frac{1}{|I_k|} \left(\frac{n_k}{n}\right)^2 \right. \\ &\quad \left. + s_{21} - \frac{1}{n}(s_{11} - s_{21}) \right] \\ &= \frac{2n-p}{(n-1)(n-p)} \sum_k \frac{1}{|I_k|} \left[\sqrt{n} \left(\frac{n_k}{n} - \alpha_k\right) \right] + \frac{(2n-p)\sqrt{n}}{(n-1)(n-p)} s_{11} \\ &\quad - \frac{n(n-p+1)}{\sqrt{n}(n-1)(n-p)} \sum_k \frac{1}{|I_k|} \left[\sqrt{n} \left(\frac{n_k}{n} - \alpha_k\right) \right]^2 - \frac{(2n-p)\sqrt{n}}{(n-1)(n-p)} s_{21} \\ &\quad - \frac{2n(n-p+1)}{(n-1)(n-p)} \sum_k \frac{\alpha_k}{|I_k|} \left[\sqrt{n} \left(\frac{n_k}{n} - \alpha_k\right) \right] - \frac{1}{\sqrt{n}}(s_{11} - s_{21}) \\ &= T_1 - \frac{2n(n-p+1)}{(n-1)(n-p)} \sum_k \frac{\alpha_k}{|I_k|} \left[\sqrt{n} \left(\frac{n_k}{n} - \alpha_k\right) \right]. \end{aligned} \quad (20)$$

Then, using the central limit theorem and the continuity of the function $x \mapsto x^2$, we have

$$\sqrt{n} \left(\frac{n_k}{n} - \alpha_k\right) \xrightarrow[n \rightarrow \infty]{d} \mathcal{N}(0, \alpha_k(1 - \alpha_k)),$$

$$\left[\sqrt{n} \left(\frac{n_k}{n} - \alpha_k\right) \right]^2 \xrightarrow[n \rightarrow \infty]{d} Z_k^2 \text{ with } Z_k \sim \mathcal{N}(0, \alpha_k(1 - \alpha_k)).$$

It thus follows that $T_1 = o_{\mathbb{P}}(1)$. We now consider the remaining term in (20). We have

$$\begin{aligned} \sum_k \frac{\alpha_k}{|I_k|} \left[\sqrt{n} \left(\frac{n_k}{n} - \alpha_k\right) \right] &= \frac{1}{\sqrt{n}} \sum_k \frac{\alpha_k}{|I_k|} n_k - \sqrt{n} \sum_k \frac{\alpha_k^2}{|I_k|} \\ &= \frac{1}{\sqrt{n}} \sum_k \frac{\alpha_k}{|I_k|} \left(\sum_{i=1}^n \mathbf{1}_{X_i \in I_k} \right) - \sqrt{n} s_{21} \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left(\sum_k \frac{\alpha_k}{|I_k|} \mathbf{1}_{X_i \in I_k} - s_{21} \right). \end{aligned}$$

Let us denote

$$Y_i = \sum_k \frac{\alpha_k}{|I_k|} \mathbf{1}_{X_i \in I_k} - s_{21}.$$

Then the random variables Y_1, Y_2, \dots, Y_n are iid centered with variance

$$\sigma_I^2 = \mathbb{E}(Y_1^2) = \mathbb{E}\left(\sum_k \frac{\alpha_k^2}{|I_k|^2} \mathbf{1}_{X_1 \in I_k} - 2s_{21} \sum_k \frac{\alpha_k}{|I_k|} \mathbf{1}_{X_1 \in I_k} + s_{21}^2\right) = s_{32} - s_{21}^2.$$

By the central limit theorem, we obtain

$$\sum_k \frac{\alpha_k}{|I_k|} \left[\sqrt{n} \left(\frac{n_k}{n} - \alpha_k\right)\right] \xrightarrow[n \rightarrow \infty]{d} \mathcal{N}(0, \sigma_I^2).$$

Combining this with (20) implies that

$$\sqrt{n}[\hat{R}_p(I) - R(I)] \xrightarrow[n \rightarrow \infty]{d} \mathcal{N}(0, 4\sigma_I^2).$$

It is easy to calculate that

$$\sqrt{n}(\hat{L}_p(I) - L(I)) = \sqrt{n}(\hat{R}_p(I) - R(I)) + \frac{1}{\sqrt{n}}(s_{11} - s_{21}).$$


Hence, we have

$$\sqrt{n}[\hat{L}_p(I) - L(I)] \xrightarrow[n \rightarrow \infty]{d} \mathcal{N}(0, 4\sigma_I^2),$$

which completes the proof.


A.3 Proof of Lemma 3

i) If I is a subdivision of $I^{(N)}$, then $I = (N, \lambda, \mu)$ with $[\lambda, \mu] \subset [\lambda^*, \mu^*]$. For example, we may have the following situation



Since g is constant on the interval $[\lambda^*, \mu^*] \supset [\lambda_N, \mu_N] \supset [\lambda, \mu]$, we have $g_I = g_{I^{(N)}} = g$ on the interval $[\lambda_N, \mu_N]$. This implies that $\|g_I - g\|_2^2 = \|g_{I^{(N)}} - g\|_2^2$.

ii) If $I = (2^m, \lambda, \mu)$ is not a subdivision of $I^{(N)}$, then there are two cases to consider: If $m = m_{max}$ then $[\lambda, \mu] \not\subset [\lambda_N, \mu_N]$. For example, we may have



Since $g_I = g_{I^{(N)}} = g$ on the interval $[\lambda_N, \mu_N]$ and the two partitions I and $I^{(N)}$ restricted to the interval $[\lambda, \mu]^c \cap [\lambda_N, \mu_N]^c$ are the same, we thus have

$$\|g_I - g\|_{2, [\lambda, \mu]^c}^2 = \|g_{I^{(N)}} - g\|_{2, [\lambda, \mu]^c}^2,$$


so that

$$\|g_I - g\|_2^2 - \|g_{I^{(N)}} - g\|_2^2 = \|g_I - g\|_{2, [\lambda, \mu]}^2 - \|g_{I^{(N)}} - g\|_{2, [\lambda, \mu]}^2.$$

Using the monotonicity of f on the intervals $[0, \lambda^*]$ and $[\mu^*, 1]$, we get that

$$\|g_I - g\|_{2, [\lambda, \mu]}^2 > \|g_{I^{(N)}} - g\|_{2, [\lambda, \mu]}^2, \text{ which implies that } L(I) > L(I^{(N)}).$$

If $m < m_{max}$, we may have for example




As before, we may show that

$$\|g_I - g\|_2^2 - \|g_{I^{(N)}} - g\|_2^2 \geq \|g_I - g\|_{2, [\lambda, \mu]^c}^2 - \|g_{I^{(N)}} - g\|_{2, [\lambda, \mu]^c}^2 > 0,$$

which completes the proof.

We remark that the assumptions in Lemma 2.1 or Theorem 2.1 in [Celisse and Robin \(2010\)](#) are not sufficient to show these results. In fact, the assumption "g is non-constant outside Λ^* " is not sufficient to imply that $\|g - g_{I^{(N)}}\|_2^2 < \|g - g_{\hat{I}}\|_2^2$ in the case where \hat{I} is not a subdivision of $I^{(N)}$. For example, let us consider the following situation



We may then calculate that

$$\|g - g_{\hat{I}}\|_2^2 - \|g - g_{I^{(N)}}\|_2^2 = (c - a)(\alpha_1 - \alpha)^2 + (b - c)(\alpha_2 - \alpha)^2,$$

where

$$\alpha = \frac{1}{b - a} \int_a^b g(x) dx, \quad \alpha_1 = \frac{1}{c - a} \int_a^c g(x) dx, \quad \alpha_2 = \frac{1}{b - c} \int_c^b g(x) dx.$$

So that if the function g satisfies $\alpha = \alpha_1 = \alpha_2$ (and g is non-constant outside Λ^*) then $\|g - g_{I^{(N)}}\|_2^2 = \|g - g_{\hat{I}}\|_2^2$.

References

- Baíllo, A., J. A. Cuesta-Albertos, and A. Cuevas (2001). Convergence rates in nonparametric estimation of level sets. *Statist. Probab. Lett.* 53(1), 27–35.
- Benjamini, Y. and Y. Hochberg (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. Roy. Statist. Soc. Ser. B* 57(1), 289–300.
- Broberg, P. (2005). A comparative review of estimates of the proportion unchanged genes and the false discovery rate. *BMC Bioinformatics* 6(1), 199.
- Cadre, B. (2006). Kernel estimation of density level sets. *J. Multivariate Anal.* 97(4), 999–1023.
- Cai, T. T. and J. Jin (2010). Optimal rates of convergence for estimating the null density and proportion of nonnull effects in large-scale multiple testing. *Ann. Statist.* 38(1), 100–145.
- Celisse, A. and S. Robin (2008). Nonparametric density estimation by exact leave- p -out cross-validation. *Comput. Statist. Data Anal.* 52(5), 2350–2368.
- Celisse, A. and S. Robin (2010). A cross-validation based estimation of the proportion of true null hypotheses. *J. Statist. Plann. Inference* 140(11), 3132–3147.
- Chamberlain, G. (1986). Asymptotic efficiency in semiparametric models with censoring. *J. Econometrics* 32(2), 189–218.
- Donoho, D. and J. Jin (2004). Higher criticism for detecting sparse heterogeneous mixtures. *Ann. Statist.* 32(3), 962–994.
- Dudoit, S. and M. J. van der Laan (2008). *Multiple testing procedures with applications to genomics*. Springer Series in Statistics. New York: Springer.
- Efron, B. (2004). Large-scale simultaneous hypothesis testing: the choice of a null hypothesis. *J. Amer. Statist. Assoc.* 99(465), 96–104.
- Efron, B., R. Tibshirani, J. D. Storey, and V. Tusher (2001). Empirical Bayes analysis of a microarray experiment. *J. Amer. Statist. Assoc.* 96(456), 1151–1160.
- Genovese, C. and L. Wasserman (2004). A stochastic process approach to false discovery control. *Ann. Statist.* 32(3), 1035–1061.
- Hengartner, N. W. and P. B. Stark (1995). Finite-sample confidence envelopes for shape-restricted densities. *Ann. Statist.* 23(2), 525–550.
- Jin, J. (2008). Proportion of non-zero normal means: universal oracle equivalences and uniformly consistent estimators. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 70(3), 461–493.
- Jin, J. and T. Cai (2007). Estimating the null and the proportion of nonnull effects in large-scale multiple comparisons. *J. Amer. Statist. Assoc.* 102(478), 495–506.

- Langaas, M., B. H. Lindqvist, and E. Ferkingstad (2005). Estimating the proportion of true null hypotheses, with application to DNA microarray data. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 67(4), 555–572.
- Mason, D. M. and W. Polonik (2009). Asymptotic normality of plug-in level set estimates. *Ann. Appl. Probab.* 19(3), 1108–1142.
- Meinshausen, N. and P. Bühlmann (2005). Lower bounds for the number of false null hypotheses for multiple testing of associations under general dependence structures. *Biometrika* 92(4), 893–907.
- Meinshausen, N. and J. Rice (2006). Estimating the proportion of false null hypotheses among a large number of independently tested hypotheses. *Ann. Statist.* 34(1), 373–393.
- Neuvial, P. (2010). Intrinsic bounds and false discovery rate control in multiple testing problems. Technical report, arXiv:1003.0747.
- Pounds, S. and C. Cheng (2006). Robust estimation of the false discovery rate. *Bioinformatics* 22(16), 1979–1987.
- Schweder, T. and E. Spjøtvoll (1982). Plots of p-values to evaluate many tests simultaneously. *Biometrika* 69(3), 493–502.
- Storey, J. D. (2002). A direct approach to false discovery rates. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 64(3), 479–498.
- Storey, J. D. and R. Tibshirani (2003). Statistical significance for genomewide studies. *Proc. Natl. Acad. Sci. USA* 100(16), 9440–9445 (electronic).
- Swanepoel, J. W. H. (1999). The limiting behavior of a modified maximal symmetric $2s$ -spacing with applications. *Ann. Statist.* 27(1), 24–35.
- Turkheimer, F., C. Smith, and K. Schmidt (2001). Estimation of the number of true null hypotheses in multivariate analysis of neuroimaging data. *NeuroImage* 13(5), 920 – 930.
- van der Vaart, A. (2002). Semiparametric statistics. Bolthausen, Erwin et al., Lectures on probability theory and statistics. Ecole d’été de probabilités de Saint-Flour XXIX - 1999, Saint-Flour, France, July 8-24, 1999. Berlin: Springer. Lect. Notes Math. 1781, 331-457 (2002).
- van der Vaart, A. W. (1998). *Asymptotic statistics*, Volume 3 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge: Cambridge University Press.