



HAL
open science

Nationwide surveillance of 18 respiratory viruses in patients with influenza like illnesses: a pilot feasibility study in the French Sentinel Network

A. Falchi, Clement Turbelin, Laurent Andreoletti, Christophe Arena, Thierry Blanchon, Isabelle Bonmarin, Thomas Hanslik, Marianne Leruez, Xavier de Lamballerie, Fabrice Carrat

► To cite this version:

A. Falchi, Clement Turbelin, Laurent Andreoletti, Christophe Arena, Thierry Blanchon, et al.. Nationwide surveillance of 18 respiratory viruses in patients with influenza like illnesses: a pilot feasibility study in the French Sentinel Network. *Journal of Medical Virology*, 2011, 83 (8), pp.1451. 10.1002/jmv.22113. hal-00647401

HAL Id: hal-00647401

<https://hal.science/hal-00647401>

Submitted on 2 Dec 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Nationwide surveillance of 18 respiratory viruses in patients with influenza like illnesses: a pilot feasibility study in the French Sentinel Network

Journal:	<i>Journal of Medical Virology</i>
Manuscript ID:	JMV-10-2023.R3
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	09-Mar-2011
Complete List of Authors:	Falchi, A. turbelin, clement andreoletti, laurent arena, christophe blanchon, thierry bonmarin, isabelle hanslik, thomas leruez, marianne de Lamballerie, Xavier carrat, fabrice
Keywords:	respiratory viruses

SCHOLARONE™
Manuscripts

Table I: Viral aetiology of the ILI patients.

Viral aetiology (co-infections included)	Number	% of patients
Adenovirus	2	2.1
Bocavirus	2	2.1
Enterovirus	0	0
Human Coronavirus type 229E	9	9.4
Influenza virus type A subtype H1N1	0	0
Influenza virus type A subtype H1N1 2009	5	5.2
Influenza virus type A subtype H3N2	1	1
Influenza virus type B	1	1
Influenza virus type C	0	0
Metapneumovirus type A	11	13.5
Metapneumovirus type B	2	
Parainfluenza virus type 1	0	1
Parainfluenza virus type 2	1	
Parainfluenza virus type 3	0	
Parainfluenza virus type 4	0	
Respiratory Syncytial Virus type A	10	25
Respiratory Syncytial Virus type B	14	
Rhinovirus	3	3.1
Total number of detected virus	61	
Details of detected Co-infections		
Metapneumovirus A + Respiratory Syncytial Virus A+Respiratory Syncytial Virus B	1	1
Influenza B + Metapneumovirus A	1	1
Adenovirus + Human Coronavirus 229E	1	1
Bocavirus + Respiratory Syncytial Virus A	1	1
Number of positive patients	56	58.3 %
Total patients	96	



Aetiology of viral respiratory agents (RSV, hMPV and the other viruses detected) by age group.
266x186mm (96 x 96 DPI)

1
2 1 **Nationwide surveillance of 18 respiratory viruses in patients with influenza-like illnesses: a**
3 2 **pilot feasibility study in the French Sentinel Network**
4 3
5 4
6 4

7 5 Running title: Nationwide Surveillance of 18 Respiratory Viruses
8 6

9
10 7 Alessandra Falchi,^{1,2*} Clément Turbelin,^{1,2} Laurent Andreoletti,^{3,4} Christophe Arena,^{1,2} Thierry
11 8 Blanchon,^{1,2} Isabelle Bonmarin,⁵ Thomas Hanslik,^{1,6,7} Marianne Leruez-Ville,⁸ Xavier De
12 9 Lamballerie,^{9,10} Fabrice Carrat^{1,2,11}
13
14

15 10
16 11 *1.INSERM, UMR-S 707, F-75012, Paris, France;*

17 12 *2.UPMC Université Paris 06, UMR-S U707, F-75012, Paris, France;*

18 13 *3.Unité de Virologie Médicale et Moléculaire, Centre Hospitalier Universitaire, Reims, France;*

19 14 *4.IFR 53/EA-4303 (DAT/PPCIDH), Faculté de Médecine, Reims, France;*

20 15 *5.Institut de Veille Sanitaire, St Maurice, Paris, France;*

21 16 *6.Université Versailles Saint Quentin en Yvelines, F-78000, Versailles, France;*

22 17 *7.Assistance Publique Hôpitaux de Paris, Service de Médecine Interne, Hôpital Ambroise Paré, F-92100, Boulogne Billancourt, France ;*

23 18 *8.Unité de Virologie, Hôpital Necker-Enfants Malades, Université Paris Descartes, EA 36-20, Paris, France;*

24 19 *9.Université de la Méditerranée, Unité des Virus Emergents UMR190, Marseille, France;*

25 20 *10.Institut de Recherche pour le Développement, Marseille, France;*

26 21 *11.Assistance Publique-Hôpitaux de Paris, Hôpital Saint Antoine, Unité de Santé Publique, F-75012 Paris, France.*
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

23
24
25
26 23
27
28 24
29 25
30 26 Correspondence: Alessandra Falchi, réseau Sentinelles, UMR-S 707, Laboratory of Virology,
31 27 University of Corsica, 20250 Corte, France

32 28 Tel: + 33 0 495 45 06 77

33 29 Fax: + 33 1 44 73 84 54

34 30 E-mail: falchi@u707.jussieu.fr (A. Falchi)
35 31
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

The aim of the present study was to test the feasibility of integrating the diagnosis of 18 respiratory viruses into clinical surveillance of influenza-like illness using a PCR-DNA microarray detection assay.

The study took place in the French Sentinel Network, a nationwide surveillance network of General Practitioners (GPs) representative of French GPs in terms age, location and type of practice (urban/rural). Three virological laboratories also participated in the study. The study was planned for five weeks from January 25, 2010 to February 27, 2010.

A subset of 150 Sentinel GPs, located in mainland France, was enrolled to collect clinical data and nasopharyngeal samples from every first patient of the week having a medical visit for influenza-like illness defined as a sudden fever of 39°C or more with respiratory symptoms and myalgia. Sixty-three GPs (42%) collected 103 samples while 87 GPs (58%) did not. GPs did not differ with respect to their age, gender, urban/rural distribution or years of inscription in the Sentinel Network.

Patients included were of a similar age and had similar vaccination characteristics, but were more frequently men than influenza-like illness patients reported to the network during the study period. Sixty-one viruses were detected from 56 of 96 (58%) interpretable samples. The respiratory viruses detected most frequently were metapneumovirus and respiratory syncytial virus.

This study showed that virological diagnosis of 18 respiratory viruses can be combined with surveillance of clinical influenza-like illness in general practice. Although feasibility has not been demonstrated yet, it will be evaluated over the winter of 2010-2011.

Keywords: respiratory viruses, influenza-like illness, surveillance system, general practitioners

Abstract (word count: 252)

Text (word count: 2906)

1
2 72 **INTRODUCTION**

3
4 73 The spread of the new variant A/(H1N1) 2009 influenza virus in the Northern hemisphere at
5
6
7 74 an earlier time than usual for seasonal influenza in previous years has highlighted the limited
8
9 75 knowledge concerning the aetiology of acute respiratory illnesses outside of influenza outbreaks.

10
11 76 In early September 2009, while reports of influenza-like illness were increasing in medical practice-
12
13
14 77 based surveillance in France and other European countries, the detection of pandemic influenza
15
16 78 virus remained sporadic [Linde et al., 2009]. This finding was attributed to the circulation of other
17
18 79 respiratory viruses and, to a lesser extent, may have been associated with the increased propensity
19
20
21 80 of patients with influenza-like illness to seek medical advice due to increased anxiety in the
22
23 81 pandemic context.

24
25 82 This finding was also observed for influenza-like illness incidence rates reported by the
26
27
28 83 French Sentinel Network [Turbelin et al., 2009]. The Sentinel network is a continuous
29
30 84 epidemiological surveillance system based on General Practitioners (GPs) and has been operating
31
32
33 85 since 1984 in France. It consists of a sample of 1,319 GPs throughout France, of which 200 to 400
34
35 86 participate each week in continuous surveillance [Valleron et al., 1986]. The GPs participating in
36
37 87 the Sentinel Network are representative of the whole French GP population as regards to age,
38
39
40 88 location, type (rural/urban) and kind of practice (alone/pluridisciplinary) [Chauvin et al., 1995].

41
42 89 The main objective of the French Sentinel Network is to provide the national health
43
44
45 90 authorities with near to real-time information on several health-related events occurring in the
46
47 91 community, including influenza-like illness (for the full list of events, see <http://www.sentiweb.fr>).
48
49 92 These events are reported by Sentinel GPs through a dedicated website, aggregated at different
50
51
52 93 spatial and temporal levels and used to inform health authorities. In the Sentinel network, influenza-
53
54 94 like illness cases are defined on the basis of clinical signs, but no virological collection is carried
55
56 95 out routinely.

57
58
59 96 Given the need for improving virological surveillance after the first A/(H1N1)2009
60

1
2 97 pandemic wave, which spanned from September 6, 2009 to December 26, 2009 in France, the
3
4 98 feasibility of integrating the diagnosis of respiratory viruses into clinical surveillance of influenza-
5
6
7 99 like illness in the French Sentinel Network was tested. As a secondary objective, virological and
8
9 100 clinical data were analysed.
10
11
12 101

13 14 102 METHODS

15 16 103 **Study Design and Inclusion Criteria**

17
18 104 Three virological laboratories participated in the study. A subset of 150 volunteer GPs,
19
20
21 105 located in mainland France were recruited, on the basis of their participation in continuous
22
23 106 surveillance in the Sentinel Network.
24

25
26 107 The 150 GPs were distributed over all the departments (administrative units) of France. The
27
28 108 GPs participating to the present study did not differ from other GPs of the Sentinel Network with
29
30 109 respect to age (51.7 ± 8.1 years vs. 52.1 ± 7.8 years; $P=0.93$), gender (male) (79% vs. 83%; $P=0.60$)
31
32 110 urban/rural distribution (urban 75% vs. 74%; rural 25% vs. 26%; $P=0.68$) and years of inscription
33
34 111 in the Sentinel Network (10.4 ± 7.6 years vs. 11.1 ± 7.7 years; $P=0.27$).
35
36

37 112 The pilot study was carried out over five weeks between January 25, 2010 and February 27,
38
39
40 113 2010, after the end of the French A/(H1N1)2009 epidemic. The 150 enrolled volunteer GPs
41
42 114 received swabs and other study materials to include every first patient of the week having a medical
43
44 115 visit for influenza-like illness (Sentinel network case-definition: a sudden fever of 39°C or above,
45
46 116 respiratory symptoms and myalgia), with the additional criteria of the time between symptom onset
47
48
49 117 and the visit of less than 48 hours.
50

51 52 118 **Sample and patient data collection**

53
54 119 Sample collection was performed with Σ -Virocult® swabs (ELItech, France). The
55
56 120 nasopharyngeal swabs were sent by mail in 2 ml of viral transport medium within two days to one
57
58
59 121 of the three virological laboratories participating in the study. Detailed demographic and clinical
60

1
2 122 data (time of the onset of symptoms, reported symptoms, physical findings and influenza
3
4 123 vaccination status) were obtained from patients during the medical visit.
5

7 124 **Nucleic Acid Extraction and Microarray Detection**

8

9 125 Following RNA and DNA extraction from clinical specimens using either an automated
10
11 126 MagNA Pure LC with the Total Nucleic Acid High Performance kit (Roche Diagnostic, Meylan,
12
13 France) or an EZ1 Biorobot with the virus mini kit (Qiagen, Courtaboeuf, France), all samples were
14 127
15 tested for the simultaneous detection of a panel of 18 respiratory viral pathogens using a PCR-DNA
16 128
17 microarray detection assay (CLART® PneumoVir and FluaVir Version 3.0, Genomica, Madrid,
18 129
19 Spain). The CLART® PneumoVir allows the detection of 17 respiratory viruses distinguishing
20
21 130 between the seasonal influenza A subtypes A/H1N1 and A/H3N2. The CLART® FluaVir kit was
22
23 131 able to type and to distinguish the influenza virus A/H1N12009 from the A/H1N1 and A/H3N2
24
25 132 subtypes of the seasonal influenza A virus. All respiratory samples were tested simultaneously using
26
27 these two kits.
28 133
29
30 134

31
32
33 135 The combination of these two systems allows for the simultaneous detection of Adenovirus
34
35 136 (AV), Bocavirus (hBov), Coronavirus 229E (hCoV), Enterovirus (Echovirus) (EV), Influenza A
36
37 137 ((H1N1) 2009, H3N2, H1N1), Influenza B, Influenza C, MetaPneumoVirus (hMPVa and hMPVb),
38
39 Parainfluenza (PIV) (1, 2, 3 and 4), Respiratory Syncytial Virus (RSVa and RSVb) and Rhinovirus
40 138
41 (RV). The technique includes an internal control that tests the efficiency of the amplification
42 139
43 process and to detect the presence of inhibition products. The sensitivity and specificity of the
44
45 140 multiplex RT-PCR assay for the simultaneous detection of this panel of respiratory viral pathogens
46
47 141 have been previously reported. [Renois et al., 2010].
48
49 142

50
51
52 143 Surveillance virological results (number of samples, positive samples, type and subtype of
53
54 144 viruses isolated weekly) were available on the website the week following sample collection. Each
55
56 145 week, the GPs also received the virological results of her/his patients on a secure study website. The
57
58 interval between specimen collection and availability of the results for GPs was 10-14 days. Finally,
59 146
60

1
2 147 an e-mail survey in GPs who did not include any patients to describe the reasons for the lack of
3
4 148 sample collection was performed, and another survey was performed to evaluate the GPs'
5
6
7 149 satisfaction with the pilot study in those who included patients.
8

9 150 **Statistical Analysis**

10
11 151 Patients with at least one virus detected (positive patients) were compared to patients
12
13
14 152 with no virus detected (negative patients). Factors and signs associated with positivity for the
15
16 153 viruses that were most frequently isolated or with co-infections were studied. The Fisher exact test
17
18
19 154 was used to compare independent categorical variables and the Mann-Whitney test was used to
20
21 155 compare continuous variables. A logistic regression model was used for binomial outcomes, and
22
23 156 odds ratios (OR) and their 95% confidence intervals were estimated.
24
25
26 157

28 158 **RESULTS**

30 159 **Number of samples collected by GPs**

31
32
33 160 Sixty-three (42%) GPs collected 103 nasopharyngeal samples. Based on clinical data
34
35 161 reported by the GPs, all samples were collected in patients with influenza-like illness as defined by
36
37
38 162 the surveillance network. The maximum number of samples collected was during week 04 (from
39
40 163 January, 25 to January 29, 2010) (n=36). Thirty-nine (62%) GPs provided one sample, eleven
41
42 164 (17%) provided two samples, ten (16%) GPs provided three samples and three (11%) GPs provided
43
44
45 165 four samples.
46

47 166 **Comparison of GPs who collected samples and GPs who did not**

48
49 167 GPs who collected samples (n=63) did not differ from those who did not (n=87) with respect
50
51
52 168 to their age (51.8 ± 7.4 years vs. 51.7 ± 8.8 years; $P=0.93$), gender (male) (78% vs. 80%; $P=0.68$) and
53
54 169 years of inscription in the Sentinel Network (10.8 ± 7.2 years vs. 10.7 ± 7.4 years; $P=0.68$).
55

56 170 57 58 59 171 **GP satisfaction survey** 60

1
2 172 Sixty-three percent (40/63) of GPs responded to the satisfaction survey. Thirty-five out of 40
3
4 173 (87.5%) GPs were satisfied with the swab collection instructions and 38 (95%) found the dedicated
5
6
7 174 study website useful for following the development of the study and for consulting instructions.
8
9 175 Thirty-four (82.5%) GPs were satisfied with the feedback on the results.

11 176 **Survey in GPs who did not collect samples**

13
14 177 Eighty-seven GPs did not collect nasopharyngeal samples. Four items have been proposed
15
16 178 to explain the reasons for not participating (lack of any influenza-like illness cases during the study
17
18 179 period; lack of time; the study was too complex; declined to participate). Ninety-five percent
19
20
21 180 (83/87) of GPs declared that they did not collect any samples because of the lack of any influenza-
22
23 181 like illness case during the study period, but they reported themselves to be very interested in
24
25
26 182 participating in the study. Four GPs did not participate because of a lack of time and they declined
27
28 183 to participate in the following phase of the study.

30 184 **Characteristics of patients**

32
33 185 The median age of the 103 patients with influenza-like illness patients was 18 years [3-75],
34
35 186 of whom 58 (56%) were males. These patients did not differ from cases of influenza-like illness
36
37 187 reported by the Sentinel network during the study period (n=380) as regards age ($P=0.49$) or
38
39
40 188 influenza vaccination status ($P=0.46$). However, the proportion of men was higher in this sample
41
42 189 (56%) than in cases of influenza-like illness reported in the network (45% men) ($P=0.02$).

44
45 190 Of note, 50% of the included patients were children under the age of 18 which, compared to
46
47 191 the proportion of children in the general French population (22%), indicated that children were 3.5
48
49 192 times more likely than adults to be included in the sample of patients with an influenza-like illness.

51 193 **Virological findings**

53
54 194 Seven samples were excluded since the virological results were not interpretable by the
55
56 195 FluaVir (n=4) microarray or the PneumoVir microarray (n=3) because of undetectable internal
57
58
59 196 controls. Among the remaining 96 influenza-like illness cases, 61 viruses were detected in
60

nasopharyngeal samples from 56 (58%) patients: RSV in 24 (25%), hMPV in 13 (13.5%), hCoV 229E in nine (9.4%), Influenza A/H1N1 2009 in five (5.2%), RV in three (3.1%), hBoV and AV in two (2.1%) and Influenza B, Influenza A/H3N2 and PIV 2 in one (1.0%) (Table I). More than one virus was detected in four patients (Table I).

The age distribution (0–5; 6–18; 19–25; 26–44; 45–87 years) differed significantly between positive and negative patients, with positive patients being younger than negative patients (OR=0.98, IC 0.95–0.99; P=0.0226). All patients aged 0–5 years (n=12) were positive for one or more respiratory viruses, among whom eight had a positive diagnosis for RSVb infection (Fig. 1).

The most common clinical findings in patients infected with hMPV (n=13) and RSV (n=24) were cough (n=11 and n=24), rhinorrhea (n=11 and n=22), fatigue (n=11 and n=21) and myalgia (n=10 and n=16). Co-infected patients (Table I) had a median age of 15 years and did not differ clinically from mono-infected patients.

DISCUSSION

Despite the small number of samples included, the present pilot study showed that it was feasible to combine microarray detection of 18 respiratory virus infections with clinical surveillance of influenza-like illness in order to provide a more complete understanding of the epidemiology of these viruses in the community.

Combined community virological and clinical surveillance for influenza or RSV have been implemented in several countries. Notably, in France, the Groupes Régionaux d'Observation de la Grippe (GROGs) is an influenza-devoted surveillance system which has been operating for more than 25 years in which GPs or paediatricians perform virological diagnosis of influenza in patients with acute respiratory illness [Quénel et al., 1998]. Similar networks have been launched in other European countries, e.g. the UK, Switzerland, Italy and Spain (www.ecdc.europe.eu). But, to our knowledge, no surveillance system combines the simultaneous diagnosis of a large number of

1
2 222 respiratory viruses with clinical surveillance of influenza-like illness using a well-defined case
3
4 223 definition.
5

6
7 224 The simultaneous diagnosis of a large number of respiratory viruses associated with clinical
8
9 225 surveillance allows for extrapolation of the prevalence rates of each virus in patients matching the
10
11 226 clinical case definition to a nationwide level and facilitates the interpretation of surveillance data.
12

13
14 227 In addition, from a technical viewpoint, the microarray assay requires fewer human interventions
15
16 228 (up to 5 h of hands-on time can be saved compared to qRT-PCR) and allows for the testing of up to
17
18 229 24 samples per run (66 minutes of setup, excluding RNA extraction and RT-PCR) with the results
19
20
21 230 available on the next day [Frobert et al., 2011; Raymond et al., 2009]. Thus, the surveillance system
22
23 231 could be sustained during periods when the number of cases of influenza-like illness is much higher.
24

25
26 232 Although obtaining the results on the same day would be the ideal scenario, the ease of
27
28 233 implementation and higher throughput are important factors, especially in high-volume laboratories
29
30 234 [Raymond et al., 2009]. The present pilot study was not optimized to shorten the delay between
31
32
33 235 collection and delivering the results to the GPs, the reason why the actual delay was 10 to 14 days.
34
35 236 However, the results of the surveillance were made available the week following sample collection,
36
37
38 237 notably because results were aggregated over the preceding week.
39

40 238 The feasibility of the programme has been questioned, given the fact that only 63 of 150
41
42 239 GPs collected at least one sample. Although this has not been demonstrated yet, it will be evaluated
43
44
45 240 over the winter of 2010-2011. The small number of participants in this first season might easily be
46
47 241 explained by the case definition for the collection of specimens, which were rather stringent and
48
49 242 included elevated fever. Of note, 146 (97%) GPs have agreed to continue participating in the
50
51
52 243 programme during the winter of 2010-2011.
53

54 244 Among the 103 samples tested, the results of seven samples were not available because of
55
56 245 PCR-inhibited or uncertain results. In many cases, the failure in identification could have resulted
57
58
59 246 from a failure in the nucleic acid amplification step rather than from a failure of the microarray
60

1
2 247 [Townsend et al., 2006], even if technical training is necessary to avoid false positive results
3
4 248 because of insufficient washing or overexposure of the arrays [Frobert et al., 2011]. Excluding
5
6 249 these samples, 58% were positive for at least one virus. This proportion of positive diagnosis is
7
8
9 250 comparable to findings from other studies using qRT-PCR or a combination of qRT-PCR and
10
11 251 classical cell culture techniques, in which 41–61.8% of the samples were positive [Follin et al.,
12
13
14 252 2009; Heikkinen et al., 2008; Laguna-Torres et al., 2009; Nougairède et al., 2010].
15

16 253 It is noteworthy that the clinical criteria used for sample collection in those studies included
17
18 254 a broad spectrum of respiratory symptoms, while the criteria for influenza-like illness were rather
19
20
21 255 narrow/specific. Therefore, the number of positive specimens was not altered by the selection of
22
23 256 patients with a strict case definition. On the other hand, false-negative results may have occurred in
24
25
26 257 the adult population and could have been the consequence of insufficient quality of the sample
27
28 258 collected or of a lack of sensitivity of the laboratory techniques. Nevertheless, a recent study
29
30 259 validated the performance of the commercial kits used in the present study [Renois et al., 2010].
31
32

33 260 Negative results can also be due to other unknown or known respiratory pathogens (e.g.
34
35 261 Parechovirus, Polyomavirus KI and WU, Human coronavirus OC43, NL63 and KU1 or *M.*
36
37 262 *pneumoniae* and *C. pneumoniae*). This proportion of undiagnosed cases is in line with those found
38
39
40 263 in other surveys [Hasman et al., 2009; Laguna-Torres et al., 2009] and appeal for complementary
41
42 264 virological investigations using, for example, high throughput methods (e.g. pyrosequencing
43
44 265 methods [Deyde et al., 2009], mass spectrometry [Chen et al., 2011], high resolution melting
45
46
47 266 (HRM) techniques [Varillas et al., 2011] and Padlock Probes [Wu et al., 2009]).
48

49 267 In this investigation, it was found that age was associated with the chance of being positive,
50
51
52 268 with a rate of positivity of 100% in younger patients (0–5 years), in which RSV and hMPV were the
53
54 269 most frequently detected viruses. Similar findings were obtained in other settings and were
55
56 270 attributed to the developing immune status of young children and their vulnerability to infections
57
58
59 271 [Raboni et al., 2011].
60

1
2 272 Given the temperature criteria used for the influenza-like illness case-definition, children
3
4 273 (50%) were overrepresented in comparison with adults relative to the general population (22%).
5
6
7 274 This figure is well in line with current knowledge that fever is usually higher in children than in
8
9 275 adults [Feigin et al., 2009].
10

11
12 276 However, the age distribution of patients did not differ from the age distribution of reported
13
14 277 influenza-like illness cases in the network during the period. Therefore, a selection bias related to
15
16 278 age was excluded. Moreover, the Sentinel Network involves GPs and does not involve
17
18
19 279 paediatricians, which explains the low proportion of very young children (<2 years) in the sample.
20
21 280 A substantial proportion of adult subjects with RSV (11.5%) or hMPV (8.3%) were found in this
22
23 281 study. These findings were also consistent with other reports in which RSV was detected in 11–
24
25
26 282 22% of adult patients during medical visits [Falsey et al., 2006; Zambon et al., 2001] and hMPV in
27
28 283 2.2% to 10.6% in adults participants during a prospective follow-up [Walsh et al., 2008].
29

30 284 In the present study, there were no significant differences in the clinical characteristics of
31
32
33 285 patients infected with hMPV, RSV or other viral infections, but the number of patients in each
34
35 286 group was low and comparisons may have been underpowered.
36

37
38 287 As in previous studies [Heikkinen et al., 2008], the fact that co-infections were common was
39
40 288 confirmed. In this study, co-infections with hMPV occurred twice, with RSVa and Influenza B
41
42 289 respectively, but they were not associated with more severe signs than mono-infections, as reported
43
44
45 290 elsewhere [Frobert et al., 2011; Wolf et al., 2006], but here again the number of coinfecting events
46
47 291 was too low to reach a conclusion. In addition, because these RT-PCR DNA microarray detection
48
49 292 techniques are only qualitative, not quantitative, it was not possible to determine which virus was
50
51
52 293 predominant in concomitant respiratory viral infections.
53

54 294 In conclusion, although feasibility was not formally demonstrated (e.g. with respect to a set
55
56 295 of pre-specified criteria), this study showed that virological diagnosis of 18 respiratory viruses can
57
58
59 296 be combined with surveillance of clinical influenza-like illness in general practice. Such an
60

1
2 297 approach, if used routinely, could help to describe better the seasonal and community burden of
3
4 298 these respiratory viruses and to assess the clinical outcome of single and multiple viral infections.
5
6

7 299
8
9 300
10
11 **Acknowledgements:** This work was made possible thanks to the participating general
12 301 practitioners of the Sentinel Network: Andrieu S, Aubry C, Baert M, Baradel J, Barbary M,
13 302 Behar M, Benard C, Bertrand-Desvages G, Beylot JM, Bon C, Breton X, Collet P, Comelli Jy,
14 303 Damalix G, Dao B, Delanian Jp, Demure P, Descotes C, Dupont-Vidal J, Duvochel P, Ecochard
15 304 Cirica N, Em Ngourn M, Evellin F, Flamerion JM, Fondere M, Galesne-Herceg G, Galluet B,
16 305 Galopin G, Gueux P, Kleiber H, Kurth D, Lavot R, Le Duff N, Lefebure P, Leharle P, Lherbier C,
17 306 Maillard Jm, Mao G, Mathieu B, Mener E, Menu G, Meyrand B, Mullie B, Nogrel C, Nortier B,
18 307 Odoux X, Olaya E, Ostermann P, Pasquio O, Petit L, Pigache C, Ravinet O, Richard P, Rieu A
19 308 Ruillier F, Samzun JL, Saugues M, Schweckler P, Sebbag P, Semeraro JP, Soulie P, Virondaud M.
20 309 We thank the Institut de Veille Sanitaire (InVS, France) for providing funding and support for the
21 310 design and conduct of the study
22 311
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2 312 **Bibliography**
3
4 313
5
6 314 Chauvin P, Valleron AJ 1995. Attitude of French general practitioners to the public health
7
8 315 surveillance of communicable diseases. *Int J Epidemiol* 24:435-440.
9
10
11
12 316 Chen KF, Rothman RE, Ramachandran P, Blyn L, Sampath R, Ecker DJ, Valsamakis A, Gaydos CA
13
14 317 2011. Rapid identification viruses from nasal pharyngeal aspirates in acute viral respiratory
15
16 318 infections by RT-PCR and electrospray ionization mass spectrometry. *J Virol Methods* 173:60-66.
17
18
19
20 319 Deyde VM, Okomo-Adhiambo M, Sheu TG, Wallis TR, Fry A, Dharan N, Klimov AI, Gubareva LV
21
22 320 2009. Pyrosequencing as a tool to detect molecular markers of resistance to neuraminidase
23
24 321 inhibitors in seasonal influenza A viruses. *Antiviral Res* 81:16-24.
25
26
27
28 322 Falsey AR, Criddle MC, Walsh EE 2006. Detection of respiratory syncytial virus and human
29
30 323 metaPneumoVirus by reverse transcription polymerase chain reaction in adults with and without
31
32 324 respiratory illness. *J Clin Virol* 35:46-50.
33
34
35
36 325 Feigin RD, Cherry J, Demmler-Harrison GJ, Kaplan SL 2009. *Textbook Of Pediatric Infectious*
37
38 326 *Diseases*, 6th ed Expert Consult: Saunders, Inc 3856 p.
39
40
41
42 327 Follin P, Lindqvist A, Nyström K, Lindh MA 2009. A variety of respiratory viruses found in
43
44 328 symptomatic travellers returning from countries with ongoing spread of the new influenza. *Euro*
45
46 329 *Surveill* 14: 19242.
47
48
49
50 330 Frobert E, Escuret V, Javouhey E, Casalegno JS, Bouscambert-Duchamp M, Moulinier C, Gillet Y,
51
52 331 Lina B, Floret D, Morfin F 2011. Respiratory viruses in children admitted to hospital intensive care
53
54 332 units: evaluating the CLART® PneumoVir DNA array. *J Med Virol* 83:150-155.
55
56
57
58 333 Hasman H, Pachucki CT, Unal A, Nguyen D, Devlin T, Peeples ME, Kwilas SA 2009. Aetiology of
59
60 334 influenza-like illness in adults includes parainfluenzavirus type 4. *J Med Microbiol* 58:408-413.

- 1
2 335 Heikkinen T, Österback R, Peltola V, Jartti T, Vainionpää R 2008. Human metapneumoVirus
3
4 336 infections in children *Emerg Infect Dis* 14:101–106.
5
6
7 337 Laguna-Torres VA, Gómez J, Ocaña V, Aguilar P, Saldarriaga T, Chavez E, Perez J, Zamalloa H,
8
9 Forshey B, Paz I, Gomez E, Ore R, Chauca G, Ortiz E, Villaran M, Vilcarromero S, Rocha C,
10 338 Chinchá O, Jiménez G, Villanueva M, Pozo E, Aspajo J, Kochel T 2009. Influenza-like illness
11
12 339 sentinel surveillance in Peru *PLoS One* 4:6118.
13
14 340
15
16
17 341 Linde A, Rotzén-Ostlund M, Zwegberg-Wirgart B, Rubinova S, Brytting M 2009. Does viral
18
19 342 interference affect spread of influenza? *Euro Surveill* 14: 19354.
20
21
22 343 Nougairède A, Ninove L, Zandotti C, Salez N, Mantey K, Resseguier N, Gazin C, Raoult D, Charrel
23
24 344 RN, de Lamballerie X 2010. Novel virus influenza A (H1N1sw) in South-Eastern France, April–
25
26 August 2009 *PLoS One* 5:9214.
27 345
28
29
30 346 Quénel P, Dab W 1998 Influenza A and B epidemic criteria based on time-series analysis of health
31
32 347 services surveillance data. *Eur J Epidemiol* 14:275-285.
33
34
35 348 Raboni SM, Stela V, Cruz CR, França JB, Moreira S, Gonçalves L, Nogueira MB, Vidal LR,
36
37 349 Almeida SM, Debur MC, Carraro H Jr, Duarte Dos Santos CN 2011. Pandemic influenza A and
38
39 350 community respiratory viral infections in southern Brazil - laboratory diagnosis, epidemiology and
40
41 clinical outcome *J Clin Microbiol* [Epub ahead of print].
42 351
43
44
45 352 Raymond F, Carbonneau J, Boucher N, Robitaille L, Boisvert S, Wu WK, De Serres G, Boivin G,
46
47 353 Corbeil J 2009. Comparison of automated microarray detection with real-time PCR assays for
48
49 354 detection of respiratory viruses in specimens obtained from children. *J Clin Microbiol* 47:743–750
50
51
52 355 Renois F, Talmud D, Huguenin A, Moutte L, Strady C, Cousson J, Lévêque N, Andréoletti L 2010.
53
54 356 Rapid detection of respiratory tract viral infections and co-infections in patients with ILI by use of
55
56 RT-PCR DNA microarray systems *J Clin Microbiol* 48:3836-3842.
57 357
58
59
60 358 Townsend MB, Dawson ED, Mehlmann M, Smagala JA, Dankbar DM, Moore CL, Smith CB, Cox

- 1
2 359 NJ, Kuchta RD, Rowlen KL 2006. Experimental evaluation of the FluChip diagnostic microarray
3
4 360 for influenza virus surveillance J Clin Microbiol 44:2863-2871.
5
6
7 361 Turbelin C, Pelat C, Boëlle PY, Lévy-Bruhl D, Carrat F, Blanchon T, Hanslik T 2009. Early
8
9 362 estimates of 2009 pandemic influenza A(H1N1) virus activity in general practice in France:
10
11 363 incidence of influenza-like illness and age distribution of reported cases Euro Surveill 14:19341.
12
13
14 364 Valleron AJ, Bouvet E, Garnerin P, Ménarès J, Heard I, Letrait S, Lefauchaux J 1986. A computer
15
16 365 network for the surveillance of communicable diseases: the French experiment Am J Public Health
17
18 366 76:1289–1292.
19
20
21 367 Varillas D, Bermejo-Martin JF, Almansa R, Rojo S, Nogueira B, Eiros JM, Rico L, Iglesias V, de
22
23 368 Lejarazu RO 2011. A new method for detection of pandemic influenza virus using High Resolution
24
25 369 Melting analysis of the neuraminidase gene J Virol Methods 171:284-286.
26
27
28
29 370 Walsh EE, Peterson DR, Falsey AR 2008. Human MetaPneumoVirus infections in adults: another
30
31 371 piece of the puzzle Arch Intern Med 168:2489-2496.
32
33
34 372 Wolf DG, Greenberg D, Kalkstein D, Shemer-Avni Y, Givon-Lavi N, Saleh N, Goldberg MD,
35
36 373 Dagan R 2006. Comparison of human metaPneumoVirus, respiratory syncytial virus and influenza
37
38 374 A virus lower respiratory tract infections in hospitalized young children Pediatr Infect Dis J 25:320-
39
40 375 324.
41
42
43
44 376 Wu W, Tang YW 2009. Emerging molecular assays for detection and characterization of respiratory
45
46 377 viruses Clin Lab Med 29:673-693.
47
48
49 378 Zambon MC, Stockton JD, Clewley JD, Fleming DF 2001. Contribution of influenza and
50
51 379 respiratory syncytial virus to community cases of influenza-like illness: an observational study
52
53 380 Lancet 358:1410–1416.
54
55
56
57
58
59
60