



HAL
open science

Prediction and interpretation of the hydration entropies of monovalent cations and anions

Sheeba Jem Irudayam, Richard Henchman

► **To cite this version:**

Sheeba Jem Irudayam, Richard Henchman. Prediction and interpretation of the hydration entropies of monovalent cations and anions. *Molecular Physics*, 2010, pp.1. 10.1080/00268976.2010.532162 . hal-00647965


HAL Id: hal-00647965

<https://hal.science/hal-00647965>

Submitted on 4 Dec 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Prediction and interpretation of the hydration entropies of monovalent cations and anions

Journal:	<i>Molecular Physics</i>
Manuscript ID:	TMPH-2010-0326.R1
Manuscript Type:	Special Issue Paper -EIGHTH LIBLICE CONFERENCE on the Statistical Mechanics of Liquids
Date Submitted by the Author:	08-Oct-2010
Complete List of Authors:	Irudayam, Sheeba; The University of Manchester, Manchester Interdisciplinary Biocentre Henchman, Richard; The University of Manchester, Manchester Interdisciplinary Biocentre
Keywords:	solutions, free energy, statistical mechanics, water, ion
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
TMPH-2010-0326.R1.tex	

SCHOLARONE™
Manuscripts

Prediction and interpretation of the hydration entropies of monovalent cations and anions

Sheeba Jem Irudayam and Richard H. Henchman

Manchester Interdisciplinary Biocentre, The University of Manchester, 131 Princess Street, Manchester M1 7DN, United Kingdom and School of Chemistry, The University of Manchester, Oxford Road, Manchester M13 9PL, United Kingdom.

henchman@manchester.ac.uk

Abstract

The hydration entropy of the monovalent cations and anions are calculated directly from molecular dynamics simulations of the hydrated ion and bulk water using theory previously applied to the hydration of noble gases [Irudayam and Henchman, *J. Phys.: Cond. Matter*, 2010, 22, 284108]. Extensions are included to account for differential hydrogen-bonding of first-shell waters with themselves, the ion, and bulk water. The entropies, enthalpies and Gibbs energies agree reasonably with simulation and experiment when the effect of force field is taken into account. The anions' entropy losses are mostly vibrational and librational, consistent with their stronger hydration. The cations' entropy losses are mostly orientational which implies fewer hydrogen-bond arrangements because the cation substantially inhibits the ability of surrounding water molecules to accept hydrogen bonds. Owing to the many entropy terms and different decompositions, it is shown that the terms of kosmotropes and chaotropes must be appropriately applied so as not to lead to contradictions. It is also proposed that the number of hydrogen bond arrangements help explain the ordering in Hofmeister series of ions whereby anions increase this number but cations decrease it.

Introduction

Much knowledge has been accrued about the properties of ions in aqueous solutions.¹⁻⁶ However, one quantity for which our understanding is incomplete is the solution's entropy. We might be able to measure or calculate entropy changes for processes such as the hydration of ions but deeper insights come from knowing the entropy itself and how it is distributed between all the constituent molecules. This makes imperative the development of new methods with these capabilities. Concerning predictive approaches, a large number of methods have been developed or applied to calculate the related Gibbs energy of ion hydration which is easier to quantify than entropy because it connects directly with the equilibrium constant. These include the Born equation,⁷⁻⁹ Kirkwood-Buff theory,^{10,11} perturbation methods,¹²⁻²¹ quasichemical theory,²²⁻²⁴ scaled particle theory,²⁵ and correlation functions.^{26,27} Hydration entropy calculations are less commonly reported for ion hydration and may be evaluated from the the Born equation,⁸ the temperature derivative of the Gibbs energy change,²¹ subtracting off the enthalpy change from the Gibbs energy change,²⁸⁻³⁰ scaled-particle theory,^{25,31} correlation functions,^{26,27} or Langevin dipoles.^{32,33}

We have developed a method that is capable of calculating the entropy of a dilute aqueous solution from computer simulation which has so far been applied to noble gases³⁴ and small organic solutes.³⁵ It partitions the entropy into translational and rotational degrees of freedom for every molecule in the system. Configurational space in translation is discretised by the size of the water molecule into distinct minima whose entropy is derived from the force magnitude measured in a computer simulation. Rotational space for water molecules is discretised by hydrogen bonds between the solute and water molecules. Here we extend this approach to calculate the entropy of ion hydration as well as the Gibbs energy and enthalpy. Similar to most computational studies, we examine the alkali metal cations and halide anions because

of their simplicity and abundance of computational and experimental data. We examine how the entropy components vary for each ion and use this knowledge to understand commonly used concepts such as kosmotropes and chaotropes and give further insight into the Hofmeister series.^{4,5,36}

Method

Gibbs Energy, Entropy and Enthalpy of Hydration. Using the same theory derived elsewhere for the hydration of noble-gas solutes in water,³⁴ the standard Gibbs energy, entropy and enthalpy of hydration at temperature T for the transfer of an ion from the gas phase to a dilute aqueous solution at the same concentration are

$$\Delta G_{X+w}^{\circ} = E_{X,w}^{\text{sim}} - E_{w(1)}^{\text{sim}} - 3k_{\text{B}}T - k_{\text{B}}T \ln \left\{ \frac{1}{V_{\text{w}}} \left(\frac{2k_{\text{B}}T}{F_{X(\text{aq})}} \right)^3 \left(\frac{\Omega_{w(X)}^{\text{or}}}{\Omega_{w(1)}^{\text{or}}} \prod_{j=x,y,z} \frac{F_{w(1)}^j \tau_{w(1)}^j}{F_{w(X)}^j \tau_{w(X)}^j} \right)^{N_{w(X)}} \right\} \quad (1)$$

$$\Delta H_{X+w}^{\circ} = E_{X,w}^{\text{sim}} - E_{w(1)}^{\text{sim}} - k_{\text{B}}T (3/2 + \alpha_{\text{w}}T) \quad (2)$$

$$\Delta S_{X+w}^{\circ} = k_{\text{B}} \ln \left\{ \frac{1}{V_{\text{w}}} \left(\frac{2k_{\text{B}}T}{F_{X(\text{aq})}} \right)^3 \left(\frac{\Omega_{w(X)}^{\text{or}}}{\Omega_{w(1)}^{\text{or}}} \prod_{j=x,y,z} \frac{F_{w(1)}^j \tau_{w(1)}^j}{F_{w(X)}^j \tau_{w(X)}^j} \right)^{N_{w(X)}} \right\} + k_{\text{B}} (3/2 - \alpha_{\text{w}}T) \quad (3)$$

Eq. 2 is exact. Eqs. 1 and 3 assume that the environment of every molecule can be approximated by mean-field harmonic potentials and only include the contribution of water in the first hydration shell of the ions. $E_{X,w}^{\text{sim}}$ and $E_{w(1)}^{\text{sim}}$ are the energies of the solute in water and of bulk water, k_{B} is Boltzmann's constant, $V_{\text{w}} = 30.0 \text{ \AA}^3$ is the volume per water molecule at temperature 298 K and pressure 1 bar, $F_{X(\text{aq})}$ is half the force magnitude on the solute, $\Omega_{w(X)}^{\text{or}}$ and $\Omega_{w(1)}^{\text{or}}$ are the number of orientational minima of a water molecule in the ion's first shell and in bulk,

1
2
3
4
5
6
7 $F_{w(l)}^j$ and $\tau_{w(l)}^j$ are half the force and torque magnitudes on water for the $j = x, y, z$ axes, $N_{w(x)}$
8
9 is the number of water molecules perturbed by the solute, and α_w is water's thermal expansion
10
11 coefficient³⁷ which is calculated to be $2.8 \times 10^{-4} \text{ K}^{-1}$ for TIP4P/2005 at 298 K, close to the
12
13 experimental value³⁸ of $2.57 \times 10^{-4} \text{ K}^{-1}$. We note the presence of the $k_B T^2 \alpha_w$ term in Eq. 2,
14
15 amounting to 0.2 kJ mol^{-1} at 298 K, is usually ignored when enthalpy changes for solvation at
16
17 constant concentration are evaluated from simulation energies.

18
19
20
21 As in earlier work,^{34,39} $\Omega_{w(l)}^{\text{or}}$ for bulk water is evaluated using the equation

$$\Omega_{w(l)}^{\text{or}} = \frac{N_w^w (N_w^w - 1)}{2} \left(\frac{N_w^w - 2}{N_w^w} \right)^2 \quad (4)$$

22
23
24
25
26
27
28
29 where N_w^w is the number of first-shell water molecules, defined as being within the first minimum
30
31 of the oxygen-oxygen radial distribution function $g(r_{OO})$ for bulk water which lies at $r = 3.4 \text{ \AA}$.
32
33 This equation can be interpreted as the total number of uncorrelated ways that a water can
34
35 donate its two hydrogens, dividing by the symmetry number 2, and multiplying by the prob-
36
37 ability that the neighbouring water has the correct orientation to accept a hydrogen, squared
38
39 because the donating water has two hydrogen atoms. It represents the number of hydrogen-
40
41 bond arrangements per water molecule. In this work for ions, an extension of the theory to
42
43 calculate $\Omega_{w(x)}^{\text{or}}$ is necessary because ions disrupt the neighbouring hydrogen-bond structure
44
45 much more than noble gases. We again assume that only first-shell water is perturbed by the
46
47 ion. The number of first-shell water molecules $N_{w(x)}$ is defined as those within the first mini-
48
49 mum of the ion-oxygen radial distribution function $g(r_{XO})$. The first-shell assumption is based
50
51 on observations from experiment,^{36,40-42} other theoretical treatment,⁴³ and our own analysis of
52
53 second-shell water molecules whose entropies we found to be very similar to those in bulk. We
54
55 did observe more significant perturbations extending into the second-shell for divalent cations
56
57 but they are not the study of this work. For noble gases,³⁴ we had observed and consequently
58
59
60

1
2
3
4
5
6
7 assumed that first-shell water molecules have equal probabilities of forming hydrogen bonds to
8 any other water molecule, whether in the first shell or in bulk, and that no hydrogen bonds
9 are formed with the solute. There we defined a hydrogen bond as between a hydrogen and its
10 nearest non-bonded oxygen. For ions, the hydrogen-bond pattern of water would be expected
11 to be very different: first-shell water molecules would have different probabilities to form hy-
12 drogen bonds to the ion, to other first-shell water molecules or to bulk water. We denote these
13 probabilities as p_{HB}^i where i is the species involved in the hydrogen bond. These probabilities
14 are defined as

$$p_{\text{HB}}^i = \frac{N_{\text{HB}}^i}{N_{\text{w}(X)}^i} \quad (5)$$

15
16
17
18
19
20
21
22
23
24
25
26
27
28 where $N_{\text{w}(X)}^i$ is the number of water molecules in the first-shell of species i , defined as before,
29 and N_{HB}^i is the number that form a hydrogen bond. N_{HB}^i may be further decomposed into the
30 number of hydrogen bonds that the water accepts N_{a}^i and donates N_{d}^i from species i such that
31 $N_{\text{HB}}^i = N_{\text{a}}^i + N_{\text{d}}^i$. However, this decomposition is only carried out for interest and not needed
32 in the entropy calculation. Here we use a more general force-based hydrogen-bond definition
33 than those commonly used based on distance^{44,45} or energy⁴⁶ because we need to take into
34 account the differing abilities of water and ions to accept, and because these other definitions
35 contain arbitrary cut-off parameters that would give arbitrary entropies. We had employed this
36 force definition in earlier work on the entropies of small organic solutes in water.³⁵ We define a
37 hydrogen bond between a hydrogen and the atom with which it has the most attractive force.
38 Given that the hydrogen atoms of water do not have Lennard-Jones radii for the SPC/E force
39 field used here, only the Coulombic term contributes to the force. This amounts to saying that
40 a hydrogen forms a hydrogen bond to the acceptor atom A with the largest value of $q_{\text{A}}/r_{\text{AH}}^2$,
41 where r_{AH} is their distance of separation, and we neglect donation to cations because they have
42 repulsive interactions with the hydrogen atoms of water.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

We define an effective coordination number N_w^{eff} for a first-shell water whereby each value of $N_{w(X)}^i$ is weighted by its probability p_{HB}^i relative to the largest probability $p_{\text{HB}}^{i,\text{max}}$ summed over the three types of species i , namely X, first-shell water $w(X)$, and bulk water $w(l)$, giving

$$\begin{aligned} N_w^{\text{eff}} &= \sum_{i=X,w(X),w(l)} \left(\frac{p_{\text{HB}}^i}{p_{\text{HB}}^{i,\text{max}}} \times N_{w(X)}^i \right) \\ &= \sum_{i=X,w(X),w(l)} \frac{N_{\text{HB}}^i}{p_{\text{HB}}^{i,\text{max}}} \end{aligned} \quad (6)$$

It can be seen that $N_w^{\text{eff}} = N_w^w$ when the only species i is bulk water. N_w^{eff} is used in Eq. 4 in place of N_w^w to calculate the number of uncorrelated orientations. We retain the bulk probability that a neighbouring water molecule can accept a hydrogen bond because the effect of the ion on these probabilities has already been taken into account in defining N_w^{eff} . The only other change we make to Eq. 4 is in the probability that an ion can accept because the spherical anions considered here, unlike water, can always accept a hydrogen bond. Hence the probability that a neighbouring species can accept is not to the power 2 but rather to the power $2 - p_{\text{HB}}^X$. This gives the final equation

$$\Omega_{w(X)}^{\text{or}} = \frac{N_w^{\text{eff}}(N_w^{\text{eff}} - 1)}{2} \left(\frac{N_w^{\text{eff}} - 2}{N_w^{\text{eff}}} \right)^{2-p_{\text{HB}}^X} \quad (7)$$

For later analysis, we partition the entropy change in Eq. 3 into four terms

$$\text{ion :} \quad \Delta S_X^{\text{co}} = k_B \ln \left\{ \frac{1}{V_w} \left(\frac{2k_B T}{F_{X(\text{aq})}} \right)^3 \right\} + k_B (3/2 - \alpha_w T) \quad (8)$$

$$\text{water vibrational :} \quad \Delta S_w^{\text{vib}} = N_{w(X)} k_B \ln \left\{ \prod_{j=x,y,z} \frac{F_{w(l)}^j}{F_{w(X)}^j} \right\} \quad (9)$$

$$\text{water librational :} \quad \Delta S_w^{\text{lib}} = N_{w(X)} k_B \ln \left\{ \prod_{j=x,y,z} \frac{\tau_{w(l)}^j}{\tau_{w(X)}^j} \right\} \quad (10)$$

$$\text{water orientational :} \quad \Delta S_w^{\text{or}} = N_{w(X)} k_B \ln \left\{ \frac{\Omega_{w(X)}^{\text{or}}}{\Omega_{w(l)}^{\text{or}}} \right\}. \quad (11)$$

Molecular Dynamics Simulations. The ions examined in this study are the monovalent cations Li^+ , Na^+ , K^+ , Rb^+ , and Cs^+ , and anions F^- , Cl^- , Br^- , and I^- . The Lennard-Jones parameters for the ions, listed in Table 1, are the parameters used in the parm99 force field as implemented in AMBER 9.^{14,47-49} Simulations of bulk water in a cubic periodic box with 375 SPC/E water⁵⁰ and dilute solutions containing in addition one type of each ion are carried out using the AMBER9 software package.⁵¹ 500 steps of steepest-descent minimisation is followed by a molecular dynamics simulation in the NVT ensemble at 298 K using a Langevin thermostat. The systems are equilibrated for 2 ns in the NPT ensemble using a Berendsen barostat. The force and torque magnitudes on the ion and water, coordination numbers $N_{w(X)}$ and $N_{w(X)}^i$, and hydrogen-bond counts N_a^i and N_d^i are averaged over a further 1 ns. Longer simulations of 20 ns were used to calculate the energies $E_{X,w}^{\text{sim}}$ and $E_{w(l)}^{\text{sim}}$. In evaluating the energies, the size-dependent self-potential^{28,52,53} does not need to be accounted for separately⁵⁴ because it is already in the particle mesh Ewald implemented in AMBER 9. The thermodynamic finite-size correction^{30,52,55} is only a few kJ mol^{-1} and we omit it here. Simulations have SHAKE on all

bonds and a 2 fs timestep, periodic boundary conditions, 8 Å cutoff and particle mesh Ewald with default AMBER parameters. Equilibrated box sizes have sides of ~ 22.5 Å.

Results

Entropy Change of the Ion. Table 2 contains the ions' force magnitudes $F_{X(\text{aq})}$ and their contribution ΔS_X° to the entropies of hydration which are evaluated using Eq. 8. As expected, smaller ions with higher charge densities experience higher force magnitudes and lose more entropy. Anions lose more entropy than their isoelectronic cations because they can get closer to water's hydrogen than cations can to water's oxygen. For similar reasons, an anion's entropy loss is more sensitive to ionic radius (Table 1) than that of cations. To provide more insight into these entropies, we list in Table 2 the free volumes $v_{X(\text{aq})}^{\text{tr}} = 2k_B T / F_{X(\text{aq})}^3$ and the ion's contribution to the partial entropy in solution,³⁴ calculated using $s_{X(\text{aq})}^{\text{vib}} = k_B \ln(v_{X(\text{aq})}^{\text{tr}} / \Lambda_X^3) + 3k_B + \alpha_w k_B T$ where $\Lambda_X = h / (2\pi m_X k_B T)^{1/2}$ is the ion's thermal de Broglie wavelength, h is Planck's constant, and m_X is the ion's mass. The $v_{X(\text{aq})}^{\text{tr}}$ values for ions lie in the range 0.02–0.20 Å³ and have similar trends to ΔS_X° . Interestingly, the values for all ions except Cs⁺ are smaller than that of a water molecule in bulk water which is 0.16 Å³ and reflect the fact that aqueous ions experience up to about twice the force magnitude relative to water in water. These large forces and consequently larger frequencies of vibration hint that quantum effects may play a larger role for the smaller ions.⁵⁶ Table 2 also lists $s_{X(\text{aq})}^{\text{vib}}$ values using a quantum harmonic-oscillator formulation as in earlier work.^{57,58} Only for F⁻ and Li⁺ are the effects non-negligible with partial entropies 1.7 J K⁻¹ mol⁻¹ and 0.8 J K⁻¹ mol⁻¹ larger than the respective classical values. Thus this effect is comparable to that in the quantization of water's librational motion. Quantum effects are likely to be even stronger for ions of larger charge. As another point of comparison, the

1
2
3
4
5
6
7 $v_{X(aq)}^{tr}$ values for ions are much smaller than those of aqueous noble gas solutes³⁴ which range
8
9 from 1.8 Å³ xenon to 5 Å³ for neon to because their force magnitudes can be as low as a third
10 that of water's. The partial entropy of aqueous ions is seen to increase with size because of
11 both the ion's larger $v_{X(aq)}^{tr}$ and its larger mass but this dependence on mass cancels with the
12 gas-phase entropy for solvation.
13
14
15
16
17

18
19 **Entropy Change of Water.** The radial distribution functions $g(r_{OX})$ used to calculate $N_{w(X)}$
20 are shown in Figs. 1(a) and (b) and the cut-offs r_{XO}^{cut} are listed in Table 3. The resulting values
21 of $N_{w(X)}$ are also listed in Table 3 which are all consistent with experimental values.^{2,18} The
22 next set of figures in Table 3 are the x , y and z components of ΔS_w^{vib} and ΔS_w^{lib} calculated
23 using Eqs. 9 and 10. Many coordinate frames are possible for water. To be consistent with our
24 approach used for solutes,^{35,39} in contrast with the initial formulation,^{57,58} and also adopted by
25 Zielkiewicz,⁵⁹ the x , y and z axes are the principal axes ordered from lowest to highest moments
26 of inertia. Thus the x axis is parallel to the hydrogen-hydrogen vector, the y axis is along the
27 dipole moment, and the z axis is perpendicular to water's plane. Unlike the case of the noble-gas
28 solutes for which ΔS_w^{vib} and ΔS_w^{lib} increased marginally and essentially isotropically, we observe
29 a much richer behaviour for first-shell water around ions. First, anions' first-shell water loses
30 much more vibrational entropy than cations'. Second, anions' first-shell water loses a substantial
31 amount of librational entropy but cations' hardly lose any. Third, the size of the entropy loss
32 is greater for a higher charge density. The first two trends reflect the well-known asymmetry of
33 the water molecule and how cations and anions interact differentially with it.^{15,28,60-63} An anion
34 interacts with water's hydrogens, pinning the water down translationally and rotationally. A
35 cation interacts more weakly with water's less accessible and more central oxygen, such that the
36 translational and rotational restriction is less. If we consider the directional anisotropy in more
37 detail, the greatest translational loss for water around Li^+ is along the dipole axis, consistent
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 with a structure having water's dipole pointing away from Li^+ . This trend weakens for larger
8 ions and for the larger cations we observe a slight increase in all translational components,
9 similar to what was seen for noble gases.³⁴ This increase is largest in the z component of water,
10 suggesting that water molecules are partially lying down on the surface of the cations, again,
11 as they would for a non-polar solute. Little anisotropy is seen in the translational entropy loss
12 for anions which indicates that the interaction with the hydrogen can pin the whole molecule
13 down equally in every translational direction. Concerning libration for anions, the restriction is
14 greatest about the y and z axes but less about the x axis because rotation about the hydrogen-
15 hydrogen vector has less effect on the position of the hydrogen.
16
17
18
19
20
21
22
23
24
25

26 Next we consider the behaviour of the quantities needed to evaluate the orientational en-
27 tropy. In Table 4 are listed coordination numbers $N_{\text{w(X)}}^{\text{X}}$, $N_{\text{w(X)}}^{\text{w(X)}}$, and $N_{\text{w(X)}}^{\text{w(l)}}$ of the three types of
28 species X, w(X) and w(l) surrounding a first shell-water. Also included in Table 4 are the num-
29 ber of times these waters accept N_{a}^i and donate N_{d}^i , as well as the hydrogen-bond probabilities
30 p_{HB}^i calculated using Eq. 5. The ion is always in the coordination shell of first-shell water by
31 definition. We observe that $N_{\text{w(X)}}^{\text{w(X)}}$ decreases with ion size, as does $N_{\text{w(X)}}^{\text{w(l)}}$ to a smaller extent.
32 This is because of the flatter surface of larger ions that exclude more nearby water molecules
33 from a first-shell water. Concerning hydrogen-bond probabilities, water may donate to anions
34 and other water molecules but not to cations. We note that $N_{\text{d}}^{F^-}$ is actually calculated to be
35 1.03, indicating that on a small number of occasions, both of a water's hydrogens point to the
36 F^- . We neglect these instances and set $N_{\text{d}}^{F^-} = 1$. We next note for first-shell waters that the
37 number of donors equals the number of acceptors by definition. In addition, there is a strikingly
38 small number of hydrogen bonds between first-shell water molecules, a number that increases
39 with solute size. This is because triangles of hydrogen bonds and ion-water interactions are
40 much more strained compared to the more common pentagons, hexagons and heptagons. Non-
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 polar solutes, on the other hand, are engaged by and separate from water's hydrogen-bond
8 network, making a intra-first-shell hydrogen bond quite normal. The ions here lie somewhere
9 between these two extremes depending on their size. To the bulk, first-shell waters donate much
10 more around cations than they do around anions, and conversely for accepting hydrogen bonds.
11 This effect is consistent with the polarisation of water by ions and again decreases with ion
12 size. Yet even for the largest ions, the ratio between donors and acceptors is still 2:1. Thus the
13 orientational ordering extends beyond the first-shell but we condense this into the first-shell.
14 Interestingly, water around all anions almost always has one HB to the bulk, this being the
15 hydrogen not interacting with the ion or with first-shell water. In contrast, for cations, the
16 number of first-shell waters donating to bulk waters is nearer two, particularly for the smaller
17 ions. The hydrogen-bond probabilities represent the combination of these effects for donors and
18 acceptors normalised by the coordination numbers. They are close to the probability for bulk
19 water which is $p_{\text{HB}} = 0.85$, calculated using Eq. 5 and the values of $N_{\text{w}}^{\text{w}} = 4.73$ from Table 5
20 and $N_{\text{HB}}^{\text{w}} = 4$.

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36 To get the orientational entropy for water around ions, Table 5 contains the effective coor-
37 dination numbers $N_{\text{w}}^{\text{eff}}$ calculated using Eq. 6 and the numbers of orientations $\Omega_{\text{w}(X)}^{\text{or}}$ calculated
38 using Eq. 7. $\Omega_{\text{w}(X)}^{\text{or}}$ can be converted into the partial orientational entropy $s_{\text{w}(X)}^{\text{or}} = k_{\text{B}} \ln \Omega_{\text{w}(X)}^{\text{or}}$
39 per water molecule which is also in Table 5. The last two columns in Table 5 are the change in
40 orientational entropy per first-shell water molecule and the total for each ion, each calculated
41 using Eq. 11. $N_{\text{w}}^{\text{eff}}$ values around ions lie below that of water, primarily because it is difficult for
42 first-shell water molecules to form hydrogen bonds with other first-shell water molecules while
43 simultaneously interacting with the ion. This effect is strongest for the cations and smaller ions.
44 When converting these values to $\Omega_{\text{w}(X)}^{\text{or}}$, the first-shell waters of anions have much larger values
45 than those of the cations and even of bulk water because waters that donate to the anions are
46 always accepted. Waters around cations or in bulk water do not have this feature and are al-

ways dependent on neighbouring water molecules having the correct orientation to be accepted. Consequently, there is a loss of orientational entropy per water molecule around cations but a gain per molecule around anions. The total losses of orientational entropy for cations are similar because the smaller loss per water molecule is offset by the larger coordination number $N_{w(X)}$. The losses become less negative for larger cations and more positive for larger anions. This is because first-shell waters around larger ions form more hydrogen bonds with each other, an effect in turn made easier by the weaker interaction with the ion and the ion's flatter curvature which places other first-shell water molecules at an angle more compatible with water's tetrahedral angle of hydrogen-bonding. However, this gain in orientational entropy for anions may be exaggerated as shown by a comparison with other methods to calculate entropy and with experiment which we next examine.

Entropy, Enthalpy and Gibbs Energy of Hydration. A summary of the four entropy components calculated using Eqs. 8–11 are given in Table 6. These data sum up a key result that the loss of entropy of water around anions is dominated by vibrational and librational entropy while that of water around cations is dominated by orientational entropy. Also in Table 6 is the total ΔS_{X+w}° from Eq. 3 together with ΔS_{X+w}° from exponential averaging for the cations with the same Lennard-Jones parameters in SPC water,²¹ and from the Langevin-dipoles method.³² Exponential averaging measures the free energy difference between the hydrated and gas-phase ions. The Langevin-dipoles method is similar in spirit to ours in that they both evaluate the entropy directly from a single simulation. It is made up of a term dependent on the surface area and a term quantifying rotational restriction of the solvent dipoles. Incidentally, both sets of authors^{21,32} compare their entropy changes at constant concentration with those from experiment^{64,65} for the solvation process at 1 bar in the gas-phase to 1 M in aqueous solution for which the entropy change is $26.5 \text{ J K}^{-1} \text{ mol}^{-1}$ more negative. Agreement with values obtained from exponential averaging²¹ is good for the cations. Most values are within $6 \text{ J K}^{-1} \text{ mol}^{-1}$ and

1
2
3
4
5
6
7 the worst is K^+ which is too positive by $10 \text{ J K}^{-1} \text{ mol}^{-1}$. Similarly, there is good agreement with
8
9 the Langevin dipoles method for the cations, with all values within $9 \text{ J K}^{-1} \text{ mol}^{-1}$. Concerning
10
11 the anions, the agreement is worse with the Langevin dipoles. The overall trend is correct
12
13 but our values are spread over a wider range than those using the Langevin dipoles method,
14
15 particularly for the larger anions. As will be clearer when comparing with experiment in the
16
17 next section, it appears that our current model is too generous in the orientational entropy
18
19 of the anions which are able to accept hydrogen bonds. We cannot quantitatively compare
20
21 with the study of Horinek et al.³⁰ because they do not report any values representative of
22
23 comparable monovalent ions. Moreover, their inclusion of the surface-potential entropy term
24
25 of $96.5 \text{ J K}^{-1} \text{ mol}^{-1}$ appears to lead to unrealistically large entropy losses.
26
27

28
29 Finally in Table 7 we examine how ΔS_{X+w}° together with ΔH_{X+w}° from Eq. 2 and ΔG_{X+w}°
30
31 from Eq. 1 all compare with experiment as a test of the whole modelling procedure. Following
32
33 the rationale of Joung and Cheetham,⁶⁶ we use the experimental values of Schmid.⁶⁷ These
34
35 values do not include a contribution from the air-water surface potential which is consistent with
36
37 values calculated for infinitely periodic systems using Ewald with tin-foil boundary conditions
38
39 such as the one here. Only in the case of a spherical boundary potential are such terms
40
41 required.¹⁹ In addition, the values of other works are very close to Schmid et al.'s within an
42
43 additive offset.²⁰ They are listed for the same solvation process at constant concentration in
44
45 the gas and liquid phases and include enthalpy and entropy changes. The general trends versus
46
47 experiment are correct in that small ions have negative ΔS_{X+w}° , ΔH_{X+w}° and ΔG_{X+w}° which
48
49 become less negative for larger ions. However, there are some discrepancies. The agreement
50
51 with the entropies becomes worse for larger ions, not being negative enough for larger cations
52
53 and too negative for larger anions. Given the agreement for cations with exponential averaging²¹
54
55 described earlier, the difference for cations is likely to be a force field issue. The enthalpy changes
56
57 of cations match the trend of experiment but are consistently 25 kJ mol^{-1} too positive. This
58
59
60

probably reflects the experimental data^{64,65} that Aqvist used¹⁴ to derive the cations' Lennard-Jones parameters whose free energy values are about 40 kJ mol⁻¹ more negative than the ones being compared to here. If we also consider the Born correction of 11 kJ mol⁻¹ to account for ion-water interactions beyond the spherical boundary,^{14,21} then these terms together account for the discrepancy. For anions, the force field is probably partly responsible, particularly in light of the too-negative enthalpy changes. The disagreement in entropy with values from the Langevin dipoles method suggests that the model for the orientational entropy is also responsible for the difference, probably because an overestimation of the gain in $\Omega_{X(aq)}^{or}$ when a hydrogen donates to the ion. Clearly, an improved model for orientational entropy is required, an issue we discuss later.

Discussion

The increasing entropy loss with increasing charge density, either because of a larger charge or a smaller radius, is well known and is the basis of the Born equation.⁷ Our results are in accord with this trend. However, an important aspect of ionic hydration that the Born equation cannot explain is the nature of the entropy loss and how it might differ between cations, anions and non-polar solutes. One striking difference between non-polar solutes and ions concerns the heat capacity of solvation. It is generally negative for ions and positive for non-polar solutes.^{68,69} This is understood in terms of weaker, more distorted water-water hydrogen bonds in ionic solutions and stronger, more linear water-water hydrogen bonds in non-polar solutions. This is the reverse of the solute-water interactions which are stronger for ions than non-polar solutes. The influence on heat capacity of the water-water interactions, being more numerous than solute-water interactions, is evidently dominant. This different behaviour between ions and non-polar solutes would be a puzzle^{3,69} if there was only one kind

1
2
3
4
5
6
7 of entropy loss. The nature of the differing entropy losses is evident in the insightful study of
8 Lynden-Bell and Rasaiah.²⁸ The authors calculate the entropy of hydration as a function of ion
9 size and charge by letting the solute charge and radius be dynamic variables. They observed
10 a greater entropy loss for ions of either larger charge or smaller charge, and only a moderate
11 loss for ions of intermediate charges, whether a cation or anion. They also observe maxima in
12 the entropy loss as a function of solute radius for ions but not for non-polar solutes. Similar
13 trends have also been reported by others.^{25,26,70} These curves represent a balance between two
14 kinds of entropy loss, non-polar and ionic. Concerning the hydration of cations and anions,
15 there is a well-known difference^{15,28,60,62,63,71,72} arising from the asymmetric structure of the
16 water molecule. The two positively charged hydrogens are more accessible than the negatively
17 charged oxygen, giving it a positive electrostatic potential at its surface.^{15,62} Consequently,
18 anions are better hydrated than cations, meaning that the anion has a more favourable Gibbs
19 energy of solvation than a cation of the same size. Furthermore, there is a weaker dependence
20 of the entropy loss on cation size than anion size.^{28,61} Clearly, anions interact more strongly
21 with water's hydrogens than cations do with water's oxygen.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 Our results here (Table 6) and elsewhere³⁴ let us make the following interpretation of the
40 entropy losses: they are a consequence of stronger solute-water hydrogen bonds in the anionic
41 case and of fewer hydrogen-bond arrangements in the non-polar³⁴ and cationic cases. In other
42 words, the potential energy surface of aqueous anions have deeper vibrational and librational
43 energy wells while aqueous cations and non-polar solutes have fewer energy wells. This expla-
44 nation is consistent with the more favourable hydration of anions and the stronger dependence
45 on charge density. The small change in vibrational and librational entropy of cations is consis-
46 tent with their weaker interaction with water. The number of hydrogen-bond arrangements is
47 reduced for non-polar solutes and cations because of their inability to accept hydrogen bonds
48 from water. The number is lower for cations than non-polar solutes because first-shell water
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 molecules have difficulty hydrogen-bonding with each other and accepting from bulk water,
8
9 owing to the cation's strong attraction for the oxygens of first-shell waters. This effect for
10
11 anions is offset by anions' ability to always accept hydrogens which can actually increase the
12
13 number of hydrogen-bond arrangements, although perhaps excessively in the current model.
14
15 An explanation of the entropy loss for cations along similar lines was given by Singh and
16
17 Warshel³³ who used the Langevin dipoles method. They found that the entropy loss for Na⁺,
18
19 K⁺ and Ca²⁺ is in solvent configurational space rather than in solvent vibration modes because
20
21 water molecules around ions have a narrower variation in their dipole orientations but similar
22
23 coordinate fluctuations to those in bulk. They did not consider anions. The insensitivity of
24
25 ΔS_w^{or} for cations is also noteworthy. It occurs because the number of first-shell water molecules
26
27 increases more for cations than for anions. This in turn is because of the greater variation in
28
29 size in the cations than in the anions (Table 1). Thus, even though the entropy losses per water
30
31 molecule are reduced for larger ions of either type, the total entropy loss is not as diminished
32
33 for cations because more water molecules are affected (Table 5). This effect is even stronger
34
35 for non-polar solutes.³⁴ They reduce the entropy per water molecule even less but may have
36
37 up to 21 first-shell water molecules in the case of xenon. This helps explain the non-intuitively
38
39 greater entropy loss of two argon atoms compared to that of one K⁺ and one Cl⁻ ion.^{73,74}
40
41
42

43
44 In the absence of a full understanding of the entropy of solutions, key concepts used by
45
46 many to understand ionic hydration are structure-making kosmotropes and structure-breaking
47
48 chaotropes.^{5,6,72,75} Care must be taken when connecting them with entropy because the nega-
49
50 tive entropy of hydration for all ions would suggest that they are all kosmotropes. Rather, one
51
52 must look at the water's entropy component to make this connection, and there are different
53
54 ways of partitioning the entropy of solutions between water and the solute.^{34,39} Our approach
55
56 treats all molecules equivalently and assigns entropy to each molecule in the mean-field envi-
57
58 ronment of their neighbours. The solute and solvent entropy are still both negative for cations
59
60

1
2
3
4
5
6
7 and anions. However, if one focusses on water's vibrational and librational entropy (Table 6),
8
9 smaller ions, having negative entropy terms, match the term kosmotropes, and larger ions and
10 non-polar solutes,³⁴ having positive entropy terms, match the term chaotropes. The effect
11 is stronger for anions than for cations. The trend is consistent with the original naming of
12 structure-makers and structure-breakers by Frank and Evans⁵⁶ and the zero-points are consis-
13 tent with Na⁺ and Cl⁻ being intermediate.⁷⁶ An alternative approach is to assign the gas-phase
14 entropy to the solute and an excluded-volume entropy penalty to the solvent to compensate for
15 the solute's greater freedom.^{25,27,70,74} This gives quite a different interpretation of the entropy
16 change of solvation: it arises from an almost complete cancellation between large solute-water
17 and water-water hydration terms. For ions, the solute-water term is negative, indicating that
18 water molecules are very restricted relative to the solute. This is the term that corresponds
19 with kosmotropic and chaotropic behaviour because ions of higher charge density order the
20 surrounding water more relative to the ion. The water-water term, though, is positive because
21 the ion disrupts the water structure relative to how water is normally arranged relative to other
22 water molecules. For non-polar solutes, both these effects are strongly reduced and even have
23 the opposite sign such that the solute-water term is positive and the water-water term negative.
24 If one assesses water-structuring based on this term,^{27,77-80} contradictory results ensue: ions
25 of higher charge density, namely kosmotropes, break down the structure of water relative to
26 water rather than making it; chaotropes and non-polar solutes increase water-water structure.
27 Applying the concept of water-structuring beyond how it was originally intended has led many
28 to question the validity of the concepts of kosmotropes and chaotropes.^{75,79,80} Nonetheless,
29 the structure-making concept does not tell the full story in relation to the entropy of solu-
30 tions because it ignores, depending on which entropy decomposition one does, the number of
31 hydrogen-bond arrangements or the relative sizes of the water-water and water-solute terms.
32 Only a full analysis of the entropy can clarify these issues.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 Ions are known to have strong effects on solution properties. For example, kosmotropes in-
8
9 crease the solution's viscosity and surface tension.^{5,72} A particularly interesting and unresolved
10
11 effect of ions relates to the Hofmeister series^{4-6,36} which connects protein stability and solubility
12
13 in aqueous solution with an approximate sequence of anions and cations. Anionic kosmotropes
14
15 increase protein stability and promote aggregation, consistent with anions' strong attraction for
16
17 water. Anionic chaotropes interact more strongly with non-polar and chaotropic parts of the
18
19 protein such as ammonium and guanidinium groups and thereby solvate it.⁷² Cations, however,
20
21 have a mixed effect: strong chaotropes enhance protein unfolding and solubility, similar to an-
22
23 ionic chaotropes, but so do strong kosmotropes.⁵ A theory for this in terms of enthalpy is based
24
25 on the like-attracts-like rule in that a protein's anionic groups are kosmotropes but its cationic
26
27 groups are chaotropes;^{5,76} strong kosmotropic and strong chaotropic cations interact with the
28
29 kosmotropic and non-polar parts of the protein, respectively, while intermediate cations in-
30
31 teract less strongly. We suspect this difference between anions and cations also has entropic
32
33 origins in terms of the number of hydrogen-bond arrangements. Anions increase this number
34
35 while cations, similar to non-polar molecules, decrease it, meaning that cations may favour the
36
37 solvation of non-polar surfaces of the protein. In other words, the surface of the first hydration
38
39 shell of kosmotropic cations appears mostly hydrogen-like, giving it a more hydrophobic guise,
40
41 similar to strong chaotropes such as guanidinium and urea. Thus kosmotropic and chaotropic
42
43 cations would both have reasons to unfold or solvate proteins, thereby explaining the mixed
44
45 order of cations with respect to kosmotropic and chaotropic behaviour in the Hofmeister series
46
47 and the weaker influence of cations than anions. It is difficult to resolve unambiguously the
48
49 myriad of competing enthalpic and entropic effects involving ions, water and protein atoms
50
51 without studying proteins in aqueous electrolytes directly.
52

53
54
55 More accurate and extensive studies of these effects at the level of individual ions can be
56
57 made using better force fields and models for the entropy of aqueous solutions. A number of
58
59
60

1
2
3
4
5
6
7 force fields for ions have recently been developed, of both the fixed-charged^{18,66} and polarisable
8 nature.⁸¹ As we noted earlier, the inclusion of quantum effects would be necessary for ions of
9 higher charge density, particularly for transition-metal complexes with more covalent interac-
10 tions. Regarding methods to calculate the entropy of aqueous solutions, one of the deficiencies
11 in our theory is the dependence on arbitrary cut-offs in defining hydration shells. This is not
12 as serious a problem as fitted parameters which inevitably would give better agreement, but
13 are still unsatisfactory. These also make more problematic the study of aspherical solutes and
14 longer-range entropic contributions. Another deficiency is the restriction to one type of water
15 environment which had been based on the prevailing view that water structure is essentially
16 tetrahedral.^{4,82} This leads to a more awkward mean-field theory for the orientational entropy
17 that clearly averages too much over variability in water structure around ions. Recent work
18 by us has found a way past both these difficulties. We have proposed a new hydrogen-bond
19 definition that has no arbitrary parameters: a hydrogen donates to its nearest oxygen as long
20 as that oxygen-hydrogen distance is the shortest between the two water molecules involved.⁸³
21 This definition accurately resolves the hydrogen-bonding pattern in water and reveals multiple
22 water environments according to the number of hydrogen bonds formed rather than a single
23 distorted tetrahedral arrangement. It can be generalized in the case of different donors and
24 acceptors by using the electrostatic force as we have already done here. It points the way to a
25 much more flexible theory for solutions as a mixture of multiple water and solute environments
26 based on their hydrogen-bond patterns. The development of theory to do this is currently in
27 progress.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Conclusion

52
53
54
55 A method has been presented to calculate the entropy of hydration for alkali metal cations and
56 halide anions from molecular dynamics simulations of each ion in water and of bulk water. En-
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ropy changes are derived from the force magnitude on the ion, the force and torque magnitudes and coordination and hydrogen-bond numbers of its first hydration and bulk water molecules. The entropy results are in reasonable agreement with perturbation calculations for cations and partial agreement with the Langevin dipoles calculations for both types of ions. Agreement in the entropy change with experiment is better for smaller ions but is too negative for larger cations and not negative enough for larger anions. There is a much greater vibrational and librational entropy loss for the anions than cations which is consistent with stronger water-ion interactions and more favourable hydration for anions. The number of orientational minima, or equivalently, the number of hydrogen bond arrangements decreases substantially for cations but increases moderately for anions. This is because cations make it difficult for first-shell water molecules to accept hydrogen-bonds from other water molecules whereas anions affect this little and moreover can always accept a hydrogen. The terms of kosmotropes and chaotropes are shown to apply to the first-shell water's vibrational and librational entropy but not to the overall solution entropy. Finally, the number of hydrogen bond arrangements is proposed to make an important contribution to the Hofmeister ordering of ions in that anions increase this number but cations decrease it, and may help explain kosmotropic cations solvate and unfold proteins and why cations have a weaker influence than anions.

Acknowledgments

RHH and SJI are funded by EPSRC grant EP/E026222/1 and SJI is funded by an Overseas Research Scholarship and the School of Chemistry at the University of Manchester.

REFERENCES

1. Franks, F. *Water: a comprehensive treatise*, Vol. 3; Plenum Press: New York, 1973.
2. Ohtaki, H.; Radnai, T. *Chem. Rev.* **1993**, *93*, 1157–1204.
3. Schmid, R. *Monatsh. Chem.* **2001**, *132*, 1295–1326.
4. Ball, P. *Chem. Rev.* **2008**, *108*, 74–108.
5. Kunz, W.; Neueder, R. In *Specific Ion Effects*; Kunz, W., Ed.; World Scientific Publishing, 2009.
6. Zhang, Y. J.; Cremer, P. S. *Ann.. Rev.. Phys.. Chem.,. VOL. 61* **2010**, *61*, 63–83.
7. Born, M. *Z. Phys.* **1920**, *1*, 45.
8. Abraham, M. H.; Matteoli, E.; Liszi, J. *J. Chem. Soc. Farad. T. 1.* **1983**, *79*, 2781–2800.
9. Marcus, Y. *Biophys. Chem.* **1994**, *51*, 111–127.
10. Kirkwood, J. G.; Buff, F. P. *J. Chem. Phys.* **1951**, *19*, 774–777.
11. Ben-Naim, A. *Statistical Thermodynamics for Chemists and Biochemists*; Plenum: New York, 1992.
12. Straatsma, T. P.; Berendsen, H. J. C. *J. Chem. Phys.* **1988**, *89*, 5876–5886.
13. Migliore, M.; Corongiu, G.; Clementi, E.; Lie, G. C. *J. Chem. Phys.* **1988**, *88*, 7766–7771.
14. Aqvist, J. *J. Phys. Chem.* **1990**, *94*, 8021–8024.
15. Hummer, G.; Pratt, L. R.; Garcia, A. E. *J. Phys. Chem.* **1996**, *100*, 1206–1215.
16. Grossfield, A.; Ren, P. Y.; Ponder, J. W. *J. Am. Chem. Soc.* **2003**, *125*, 15671–15682.
17. Smith, E. J.; Bryk, T.; Haymet, A. D. J. *J. Chem. Phys.* **2005**, *123*, 034706.
18. Jensen, K. P.; Jorgensen, W. L. *J. Chem. Theory Comput.* **2006**, *2*, 1499–1509.
19. Lamoureux, G.; Roux, B. *J. Phys. Chem. B* **2006**, *110*, 3308–3322.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
20. Warren, G. L.; Patel, S. *J. Chem. Phys.* **2007**, *127*, 064509.
21. Carlsson, J.; Aqvist, J. *J. Phys. Chem. B* **2009**, *113*, 10255–10260.
22. Asthagiri, D.; Pratt, L. R.; Ashbaugh, H. S. *J. Chem. Phys.* **2003**, *119*, 2702–2708.
23. Rogers, D. M.; Beck, T. L. *J. Chem. Phys.* **2010**, *132*, 014505.
24. Roux, B.; Yu, H. B. *J. Chem. Phys.* **2010**, *132*, 234101.
25. Graziano, G. *Chem. Phys. Lett.* **2009**, *479*, 56–59.
26. Rashin, A. A.; Bukatin, M. A. *J. Phys. Chem.* **1994**, *98*, 386–389.
27. Kinoshita, M.; Yoshidome, T. *J. Chem. Phys.* **2009**, *130*, 144705.
28. Lynden-Bell, R. M.; Rasaiah, J. C. *J. Chem. Phys.* **1997**, *107*, 1981–1991.
29. Carlsson, J.; Aqvist, J. *Phys. Chem. Chem. Phys.* **2006**, *8*, 5385–5395.
30. Horinek, D.; Mamatkulov, S. I.; Netz, R. R. *J. Chem. Phys.* **2009**, *130*, 124507.
31. Sen, U. *J. Am. Chem. Soc.* **1980**, *102*, 2181–2188.
32. Florian, J.; Warshel, A. *J. Phys. Chem. B* **1999**, *103*(46), 10282–10288.
33. Singh, N.; Warshel, A. *J. Phys. Chem. B* **2009**, *113*, 7372–7382.
34. Irudayam, S. J.; Henchman, R. H. *J. Phys.: Cond. Matter* **2010**, *22*, 284108.
35. Irudayam, S. J.; Plumb, R. D.; Henchman, R. H. *Faraday Discuss.* **2010**, *45*, 467–485.
36. Collins, K. D.; Neilson, G. W.; Enderby, J. E. *Biophys. Chem.* **2007**, *128*, 95–104.
37. Abascal, J. L. F.; Vega, C. *J. Chem. Phys.* **2005**, *123*, 234505.
38. Kell, G. S. *J. Chem. Eng. Data* **1975**, *20*, 97–105.
39. Irudayam, S. J.; Henchman, R. H. *J. Phys. Chem. B* **2009**, *113*, 5871–5884.
40. Omta, A. W.; Kropman, M. F.; Woutersen, S.; Bakker, H. J. *Science* **2003**, *301*, 347–349.

- 1
- 2
- 3
- 4
- 5
- 6
- 7 41. Naslund, L. A.; Edwards, D. C.; Wernet, P.; Bergmann, U.; Ogasawara, H.; Pettersson, L.
8 G. M.; Myneni, S.; Nilsson, A. *J. Phys. Chem. A* **2005**, *109*, 5995–6002.
- 9
- 10
- 11 42. Cappa, C. D.; Smith, J. D.; Messer, B. M.; Cohen, R. C.; Saykally, R. J. *J. Phys. Chem.*
12 *B* **2006**, *110*, 5301–5309.
- 13
- 14
- 15 43. Abraham, M. H.; Liszi, J. *J. Chem. Soc. Farad. T. 1.* **1980**, *76*, 1219–1231.
- 16
- 17 44. Luzar, A.; Chandler, D. *Phys. Rev. Lett.* **1996**, *76*, 928–931.
- 18
- 19
- 20 45. Buch, V. *J. Chem. Phys.* **1992**, *96*, 3814–3823.
- 21
- 22 46. Rahman, A.; Stillinger, F. H. *J. Chem. Phys.* **1971**, *55*, 3336–3359.
- 23
- 24
- 25 47. Fox, T.; Kollman, P. A. *J. Phys. Chem. B* **1998**, *102*, 8070–8079.
- 26
- 27 48. Gough, C. A.; Debolt, S. E.; Kollman, P. A. *J. Comput. Chem.* **1992**, *13*, 963–970.
- 28
- 29 49. Weiner, S. J.; Kollman, P. A.; Nguyen, D. T.; Case, D. A. *J. Comput. Chem.* **1986**, *7*,
30 230–252.
- 31
- 32
- 33 50. Berendsen, H. J. C.; Grigera, J. R.; Straatsma, T. P. *J. Phys. Chem.* **1987**, *91*, 6269–6271.
- 34
- 35 51. Case, D. A.; Darden, T. A.; Cheatham III, T. E.; Simmerling, C. I.; Wang, J.; Duke, R. E.;
36 Luo, R.; Merz, K. M.; Pearlman, D. A.; Crowley, M.; Walker, R. C.; Zhang, W.; Wang, B.;
37 Hayik, S.; Roitberg, A.; Seabra, G.; Wong, K. F.; Paesani, F.; Wu, X.; Brozell, S.; Tsui,
38 V.; Gohlke, H.; Yang, L.; Tan, C.; Mongan, J.; Hornak, V.; Cui, G.; Beroza, P.; Mathews,
39 D, H.; Schafmeister, C.; Ross, W. S.; Kollman, P. A. *AMBER 9*; University of California,
40 San Francisco, 2006.
- 41
- 42
- 43
- 44
- 45
- 46 52. Hummer, G.; Pratt, L. R.; Garcia, A. E. *J. Chem. Phys.* **1997**, *107*, 9275–9277.
- 47
- 48 53. Figueirido, F.; DelBueno, G. S.; Levy, R. M. *J. Phys. Chem. B* **1997**, *101*, 5622–5623.
- 49
- 50 54. Darden, T.; Pearlman, D.; Pedersen, L. G. *J. Chem. Phys.* **1998**, *109*, 10921–10935.
- 51
- 52 55. Sakane, S.; Ashbaugh, H. S.; Wood, R. H. *J. Phys. Chem. B* **1998**, *102*, 5673–5682.
- 53
- 54
- 55
- 56 56. Frank, H. S.; Evans, M. W. *J. Chem. Phys.* **1945**, *13*, 507–532.
- 57
- 58
- 59
- 60

- 1
2
3
4
5
6
7 57. Henschman, R. H. *J. Chem. Phys.* **2007**, *126*, 064504.
8
9 58. Klefas-Stennett, M.; Henschman, R. H. *J. Phys. Chem. B* **2008**, *112*, 3769–3776.
10
11 59. Zielkiewicz, J. *J. Phys. Chem. B* **2008**, *112*, 7810–7815.
12
13
14 60. Latimer, W. M.; Pitzer, S. P.; Slansky, C. M. *J. Chem. Phys.* **1939**, *7*, 108–111.
15
16
17 61. Conway, B. E. *J. Solution Chem.* **1978**, *7*, 721–770.
18
19
20 62. Grossfield, A. *J. Chem. Phys.* **2005**, *122*, 024506.
21
22 63. Mobley, D. L.; Barber, A. E.; Fennell, C. J.; Dill, K. A. *J. Phys. Chem. B* **2008**, *112*,
23 2405–2414.
24
25
26 64. Noyes, R. M. *J. Am. Chem. Soc.* **1962**, *84*, 513–522.
27
28
29 65. Burgess, M. A. *Metal Ions in Solution*; Ellis Horwood: Chichester, 1978.
30
31 66. Joung, I. S.; Cheatham, T. E. *J. Phys. Chem. B* **2008**, *112*, 9020–9041.
32
33 67. Schmid, R.; Miah, A. M.; Sapunov, V. N. *Phys. Chem. Chem. Phys.* **2000**, *2*, 97–102.
34
35
36 68. Abraham, M. H.; Marcus, Y. *J. Chem. Soc. Farad. T. 1.* **1986**, *82*, 3255–3274.
37
38
39 69. Sharp, K. A.; Madan, B. *J. Phys. Chem. B* **1997**, *101*, 4343–4348.
40
41 70. Bergman, D. L.; Lyubartsev, A. P.; Laaksonen, A. *Phys. Rev. E* **1999**, *60*, 4482–4495.
42
43 71. Krestov, G. A. *Thermodynamics of Solvation: solution and dissolution, ions and solvents,*
44 *structure and energetics*; Horwood: New York, 1991.
45
46
47 72. Collins, K. D. *Biophys. J.* **1997**, *72*, 65–76.
48
49
50 73. Friedman, H. L.; Krishnam, C. V. In *Aqueous solutions of simple electrolytes*; Franks, F.,
51 Ed.; Plenum: London, 1973; pages 1–118.
52
53
54 74. Ben-Amotz, D.; Underwood, R. *Accounts Chem. Res.* **2008**, *41*, 957–967.
55
56
57 75. Tobias, D. J.; Hemminger, J. C. *Science* **2008**, *319*, 1197–1198.
58
59
60

- 1
2
3
4
5
6
7 76. Collins, K. D. *Methods* **2004**, *34*, 300–311.
8
9 77. Madan, B.; Sharp, K. *J. Phys. Chem.* **1996**, *100*, 7713–7721.
10
11 78. Hribar, B.; Southall, N. T.; Vlachy, V.; Dill, K. A. *J. Am. Chem. Soc.* **2002**, *124*, 12302–
12 12311.
13
14 79. Mancinelli, R.; Botti, A.; Bruni, F.; Ricci, M. A.; Soper, A. K. *J. Phys. Chem. B* **2007**,
15 *111*, 13570–13577.
16
17 80. Zangi, R. *J. Phys. Chem. B* **2010**, *114*, 643–650.
18
19 81. Yu, H. B.; Whitfield, T. W.; Harder, E.; Lamoureux, G.; Vorobyov, I.; Anisimov, V. M.;
20 MacKerell, A. D.; Roux, B. *J. Chem. Theory Comput.* **2010**, *6*, 774–786.
21
22 82. Finney, J. L. *Philos. T. Roy. Soc. B* **2004**, *359*, 1145–1163.
23
24 83. Henchman, R. H.; Irudayam, S. J. submitted for publication.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Captions

Figure 1. Ion-oxygen radial distribution functions $g(r_{\text{XO}})$ of (a) the five cations and (b) the four anions. Each vertical dashed line is the cut-off $r_{\text{XO}}^{\text{cut}}$ placed at the first minimum.

For Peer Review Only

Table 1. Lennard Jones Parameters of the Ions

cation	$r_{\min}/2$ (Å)	ϵ (kcal mol ⁻¹)	anion	$r_{\min}/2$ (Å)	ϵ (kcal mol ⁻¹)
Li ⁺	1.137	0.01830			
Na ⁺	1.868	0.00277	F ⁻	1.75	0.061
K ⁺	2.658	0.00033	Cl ⁻	1.95	0.265
Rb ⁺	2.956	0.00017	Br ⁻	2.22	0.320
Cs ⁺	3.395	0.00008	I ⁻	2.35	0.400

Table 2. Average Force Magnitude and Entropy Components of the Ions

ion	$F_{X(\text{aq})}$ (10 ⁻¹⁰ N)	$v_{X(\text{aq})}^{\text{tr}}$ (10 ⁻³⁰ Å ³)	$s_{X(\text{aq})}^{\text{vib, a}}$ (J K ⁻¹ mol ⁻¹)	ΔS_X° (J K ⁻¹ mol ⁻¹)
Li ⁺	2.80	0.03	17.6 (19.3)	-47.0
Na ⁺	2.18	0.05	38.8 (39.1)	-40.8
K ⁺	1.75	0.10	50.9 (51.0)	-35.3
Rb ⁺	1.61	0.13	62.7 (62.8)	-33.2
Cs ⁺	1.41	0.20	71.5 (71.5)	-30.0
F ⁻	3.09	0.02	27.7 (28.5)	-49.5
Cl ⁻	2.36	0.04	42.2 (42.5)	-42.7
Br ⁻	1.97	0.07	56.8 (56.9)	-38.3
I ⁻	1.83	0.09	64.4 (64.5)	-36.5

^a Values in parentheses are entropies for the quantum harmonic oscillator^{57,58}

Table 3. Cut-off Distance, Ion Coordination Number and Change in Vibrational and Librational Entropy of First-Shell Water

ion	$r_{\text{XO}}^{\text{cut}}$ (Å)	$N_{\text{w(X)}}$	$\Delta S_{\text{w}}^{\text{vib},x}$	$S_{\text{w}}^{\text{vib},y}$	$S_{\text{w}}^{\text{vib},z}$	$\Delta S_{\text{w}}^{\text{lib},x}$	$S_{\text{w}}^{\text{lib},y}$	$S_{\text{w}}^{\text{lib},z}$
			(J K ⁻¹ mol ⁻¹)			(J K ⁻¹ mol ⁻¹)		
Li ⁺	2.9	4.4	-2.7	-11.1	-2.0	-0.9	0.7	-1.7
Na ⁺	3.3	5.8	1.0	-1.8	1.0	0.0	0.8	0.5
K ⁺	3.6	6.7	2.0	2.1	3.7	0.1	0.2	0.4
Rb ⁺	3.8	7.4	1.7	2.7	4.2	0.3	0.1	0.5
Cs ⁺	4.0	8.3	1.7	3.3	5.1	-0.6	-0.8	-0.4
F ⁻	3.2	6.1	-10.8	-8.0	-9.0	-10.6	-15.3	-14.0
Cl ⁻	3.6	6.9	-2.4	-2.3	-2.7	-3.9	-5.6	-4.9
Br ⁻	3.9	7.5	1.7	0.4	0.5	-0.1	0.4	0.3
I ⁻	4.0	7.6	3.2	1.3	1.2	1.3	2.6	2.1

Table 4. Coordination and Hydrogen-Bond Numbers and Probabilities for First-Shell Water

ion	$N_{w(X)}^X$	$N_{w(X)}^{w(X)}$	$N_{w(X)}^{w(l)}$	N_a^X	$N_a^{w(X)}$	$N_a^{w(l)}$	N_d^X	$N_d^{w(X)}$	$N_d^{w(l)}$	p_{HB}^X	$p_{HB}^{w(X)}$	$p_{HB}^{w(l)}$
Li ⁺	1	2.38	3.04	0	0.02	0.36	0	0.02	1.98	0	0.02	0.77
Na ⁺	1	1.57	3.13	0	0.10	0.61	0	0.10	1.90	0	0.12	0.80
K ⁺	1	1.11	3.05	0	0.23	0.73	0	0.23	1.77	0	0.41	0.82
Rb ⁺	1	1.12	2.96	0	0.30	0.76	0	0.30	1.70	0	0.54	0.83
Cs ⁺	1	1.15	2.89	0	0.40	0.80	0	0.40	1.60	0	0.69	0.83
F ⁻	1	1.43	3.38	0	0.04	1.93	1.00	0.04	0.94	1.00	0.05	0.85
Cl ⁻	1	0.85	3.26	0	0.08	1.88	0.97	0.08	0.95	0.97	0.19	0.87
Br ⁻	1	0.65	3.18	0	0.13	1.83	0.92	0.13	0.95	0.92	0.40	0.88
I ⁻	1	0.56	3.20	0	0.14	1.84	0.89	0.14	0.97	0.89	0.51	0.88

Table 5. Effective Coordination Number, Number of Orientations, and Orientational Entropy for First-Shell Water



ion	N_w^{eff}	$\Omega_{X(\text{aq})}^{\text{or}}$	$s_{w(X)}^{\text{or}}$	$\Delta S_w^{\text{or}}/N_{w(X)}$ (J K ⁻¹ mol ⁻¹)	ΔS_w^{or}
Li ⁺	3.10	1.08	0.7	-8.3	-36.2
Na ⁺	3.37	1.33	2.3	-6.6	-38.0
K ⁺	3.60	1.56	3.7	-5.3	-35.4
Rb ⁺	3.69	1.65	4.2	-4.8	-35.3
Cs ⁺	3.84	1.82	5.0	-4.0	-33.0
F ⁻	3.86	3.24	9.8	0.8	5.0
Cl ⁻	4.08	3.56	10.6	1.6	11.0
Br ⁻	4.32	3.95	11.4	2.5	18.4
I ⁻	4.48	4.23	12.0	3.0	23.1
bulk	4.73	2.94	9.0	0	0

Table 6. Hydration Entropy Components and Comparison with Other Simulations ($\text{J K}^{-1} \text{mol}^{-1}$)

ion	ΔS_X°	ΔS_w^{vib}	ΔS_w^{lib}	ΔS_w^{or}	ΔS_{X+w}°		
					this work	ref 15	ref 16
Li^+	-47.0	-15.8	-1.9	-36.2	-101	-104	-93
Na^+	-40.8	0.2	1.3	-38.0	-77	-87	-82
K^+	-35.3	7.8	0.8	-35.4	-62	-56	-71
Rb^+	-33.2	8.6	0.9	-35.3	-59	-52	-67
Cs^+	-30.0	10.1	-1.8	-33.0	-55	-53	-60
F^-	-49.5	-27.9	-40.0	5.0	-112	-	-85
Cl^-	-42.7	-7.4	-14.4	11.0	-53	-	-70
Br^-	-38.3	2.6	0.6	18.4	-17	-	-64
I^-	-36.5	5.7	6.1	23.1	-2	-	-58

Table 7. Entropy, Enthalpy and Gibbs Energy of Hydration versus Experiment

ion	ΔS_{X+w}° ($\text{J K}^{-1} \text{mol}^{-1}$)		ΔH_{X+w}° (kJ mol^{-1})		ΔG_{X+w}° (kJ mol^{-1})	
	this work	ref 55	this work	ref 55	this work	ref 55
	Li^+	-100	-100	-481	-506	-451
Na^+	-77	-68	-364	-391	-340	-371
K^+	-61	-32	-280	-308	-261	-298
Rb^+	-58	-21	-258	-283	-241	-276
Cs^+	-54	-16	-224	-258	-207	-253
F^-	-112	-127	-639	-539	-605	-501
Cl^-	-53	-75	-499	-392	-483	-373
Br^-	-16	-49	-419	-361	-414	-346
I^-	-1	-31	-385	-321	-384	-311



Dr Richard H. Henchman
Manchester Interdisciplinary Biocentre
The University of Manchester
131 Princess Street
Manchester M1 7DN
United Kingdom

Tel: +44 (0)161 306 5194
Fax: +44 (0)161 306 5201
henchman@manchester.ac.uk
<http://www.people.man.ac.uk/~mcdssrh2>

8 October, 2010

Manuscript Title: Prediction and interpretation of the hydration entropies of monovalent cations and anions (Special Issue Paper for the Eight Liblice Conference on the Statistical Mechanics of Liquids).

Manuscript Authors: Sheeba Jem Irudayam and Richard H. Henchman

Dear Prof. Jackson,

Thank you for examining our manuscript and passing on the referee's comments to us. Our responses to the referee are the following:

"In this paper the hydration properties of several monovalent cations and anions is evaluated using an approximate theoretical expression. I think this paper can be published in Molecular Physics, although I must confess that I have certain doubts about the real practical use of the approach used by the authors."

In relation to the referee's doubts about the real practical use of our approach, we are not clear why the referee makes this judgment. The manuscript already demonstrates the practical use of the approach. Granted it is approximate (harmonic approximation, and only two water environments considered, namely first shell, each with a mean-field number of orientations) and it has only been implemented for a certain class of system (i.e. radially symmetric, dilute ions). However, a person wishing to study solvation has few alternative methods available to understand the entropy of solutions and even these have their own approximations. For example, the method of distribution functions requires approximations such as the omission of water-water correlations to keep dimensionalities at a practical. Despite these limitations,

1
2
3
4
5
6
7 our method gives new and quantitative insights into ionic solvation. Furthermore, it could be
8 directly extended to study the hydration of a solute of arbitrary complexity, again with these
9 same limitations. At the end of the manuscript we allude to a more general approach we are
10 developing using our new hydrogen bond definition that examines multiple water environments
11 and avoids the use of arbitrary cut-off parameters. I also made a brief mention in my seminar at
12 the Liblice Conference that we are not limited to the harmonic approximation. Work addressing
13 these two points is still being carried out and is not yet ready for publication.
14
15
16
17

18 “1. The authors should state that Eq.(1) is not an exact expression but rather an approximation.
19 In fact the approximate expression divides the free energy into a contribution of the ions and
20 a contribution of the water.”
21
22
23

24 We have made it clear in the manmsscript that Eq. 1 is an approximation. After these
25 equations, we write “Eq. 3 is exact. Eqs. 1 and 2 assume that the environment of every molecule
26 can be approximated by mean-field harmonic potentials and only include the contribution of
27 water in the first hydration shell of the ions.”
28
29
30

31 “2. The authors use the experimental value of α in Eq.(3) and not the value of α of the water
32 model used by them.”
33
34

35 Yes, to be consistent we now use the simulation value of α in Eq. (3) rather than the ex-
36 perimental value (0.000257 K^{-1}). This has been calculated by the developers of TIP4P/2005
37 to be 0.000028 K^{-1} (Abascal and Vega, JCP, 2005, 123, 234505) which is close to the exper-
38 imental value. This makes a negligible difference to the energy (0.02 kJ mol^{-1}) and entropy
39 ($0.06 \text{ J K}^{-1} \text{ mol}^{-1}$). Some entropy values in Table 2 (and the matching second column of Table
40 6) decreased by 0.1 J/K/mol . No change occurred in the final entropy or enthalpy values,
41 reported to lower precision. We also corrected an error in the size of the $\alpha k_{\text{B}} T^2$ term given in
42 the text to now read 0.2 kJ mol^{-1} instead of 0.6 kJ mol^{-1} which was the value of $\alpha k_{\text{B}} T$ but
43 with the units of energy. This was a typo and has no effect on the any of the other calculated
44 values.
45
46
47
48
49
50

51 “3. I do not understand why the mass is included in Table 2. Although the mass of a molecule
52 appears in the expression of the free energy, it is absolutely irrelevant when it comes to de-
53 scribing transfer properties (basically the mass contribution is the same in both phases). This
54 point should be clarified in the text.”
55
56
57
58
59
60

1
2
3
4
5
6
7 Yes, to calculate the solvation terms, we do not need the mass. The mass is only needed to
8 calculate the ion-entropy components in solution which is listed in the second-last column of
9 Table 2. However, we agree that the masses of monatomic ions are commonly available data
10 and do not need to be included here. Thus we have removed the masses from Table 2.
11
12

13
14 “4. The authors use a force field to perform simulations of the ions in water. After using this
15 force field the authors extract the required information to implement their theory (mean forces,
16 torques ...) . Is the force field used by the authors able to describe the hydration free energies
17 and entropies of the ions described in the paper? Could the authors comment on this?”
18
19

20
21 We compare our entropy results with those calculated elsewhere using thermodynamic integra-
22 tion with the same ion force field for cations in SPC water (Carlsson and Aqvist, ref 15) and
23 obtain good agreement. We have not found equivalent data for anions and nor have we done
24 the calculations using thermodynamic integration. However, we can assess the performance of
25 the ion force field compared to experiment by looking at the enthalpy: the solvation enthalpy
26 values are not negative enough for cations and too negative for anions. These discrepancies
27 would not affect entropies as much because entropy is much less sensitive to force field than
28 entropy e.g. going from Li^+ to Na^+ entails a ΔH of 117 kJ mol^{-1} but a $T\Delta S$ of only 7 kJ mol^{-1} .
29 Differences in force field parameters for a given ion are much subtler. Entropy is less sensitive
30 because it depends on the logarithm of configurational volume and the associated force field
31 parameters. Enthalpy, on the other hand, has an approximate linear dependence on force field
32 energy and charge parameters.
33
34

35
36
37
38
39
40 “5. The agreement between theory and experiment shown in Table 7 is rather impressive.
41 The authors used theory. I have the feeling that using SPC/E and the force field for the ions
42 described here the hydration properties would not match the experimental results. My feeling
43 is that if you use theory to describe experimental properties using information from a force
44 field that does not reproduce experimental properties, then you are basically fitting the theory
45 parameters to reproduce the experimental results. Could the authors comment on that?”
46
47
48
49

50
51 We do not fit any parameters to reproduce experimental results. We emphasise that all quanti-
52 ties in our entropy calculations are derived from the computer simulation which in turn is only
53 based on force-field parameters. This approach is consistent with our objective of predicting
54 free energy from a simulation without recourse to any fitted parameters. In our theory, there
55 are two arbitrarily assigned parameters which are the cut-off radii at the first minima in the
56
57
58
59
60



1
2
3
4
5
6
7 oxygen-oxygen and oxygen-ion radial distribution functions. These were chosen a priori and
8 not fitted.
9

10 Finally, we have rewritten the discussion to make it clearer and better organised and im-
11 proved the wording throughout the manuscript.
12

13
14 We hope these alterations satisfy you and the referee.
15

16 Sincerely,
17

18 Richard Henchman
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



82x125mm (600 x 600 DPI)