



HAL
open science

Asymptotics of a vanishing period: General existence theorem and basic properties of frescos

Daniel Barlet

► **To cite this version:**

Daniel Barlet. Asymptotics of a vanishing period: General existence theorem and basic properties of frescos. 2012. hal-00659253

HAL Id: hal-00659253

<https://hal.science/hal-00659253>

Preprint submitted on 12 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Asymptotics of a vanishing period : General existence theorem and basic properties of frescos.

Daniel Barlet.*

5/1/12.

Abstract

In this paper we introduce the word "fresco" to denote a $[\lambda]$ -primitive monogenic geometric (a,b)-module. The study of this "basic object" (generalized Brieskorn module with one generator) which corresponds to the minimal filtered (regular) differential equation satisfied by a relative de Rham cohomology class, began in [B.09] where the first structure theorems are proved. Then in [B.10] we introduced the notion of theme which corresponds in the $[\lambda]$ -primitive case to frescos having a unique Jordan-Hölder sequence. Themes correspond to asymptotic expansion of a given vanishing period, so to the image of a fresco in the module of asymptotic expansions. For a fixed relative de Rham cohomology class (for instance given by a smooth differential form d -closed and df -closed) each choice of a vanishing cycle in the spectral eigenspace of the monodromy for the eigenvalue $\exp(2i\pi.\lambda)$ produces a $[\lambda]$ -primitive theme, which is a quotient of the fresco associated to the given relative de Rham class itself.

The first part of this paper shows that, for any $[\lambda]$ -primitive fresco there exists an unique Jordan-Hölder sequence (called the principal J-H. sequence) with corresponding quotients giving the opposite of the roots of the Bernstein polynomial in a non decreasing order. Then we introduce and study the semi-simple part of a given fresco and we characterize the semi-simplicity of a fresco by the fact for any given order of the roots of its Bernstein polynomial we may find a J-H. sequence making them appear with this order. Then, using the parameter associated to a rank 2 $[\lambda]$ -primitive theme, we introduce inductively a numerical invariant, that we call the α -invariant, which depends polynomially on the isomorphism class of a fresco

*Barlet Daniel, Institut Elie Cartan UMR 7502
Université de Lorraine, CNRS, INRIA et Institut Universitaire de France,
BP 239 - F - 54506 Vandoeuvre-lès-Nancy Cedex.France.
e-mail : Daniel.Barlet@iecn.u-nancy.fr

(in a sens which has to be defined) and which allows to give an inductive way to produce a sub-quotient rank 2 theme of a given $[\lambda]$ -primitive fresco assuming non semi-simplicity.

In the last section we prove a general existence result which naturally associate a fresco to any relative de Rham cohomology class of a proper holomorphic function of a complex manifold onto a disc. This is, of course, the motivation for the study of frescos.

AMS CLASSIFICATION. 32 S 25, 32 S 40, 32 S 50.

KEY WORDS. Fresco, theme, (a,b)-module, asymptotic expansion, vanishing period, Gauss-Manin connection, filtered differential equation.

Contents

1 Preliminaries.	3
1.1 Definitions and characterization as $\tilde{\mathcal{A}}$ -modules.	3
1.2 The principal Jordan-Hölder sequence.	7
2 Semi-simplicity.	11
2.1 Semi-simple regular (a,b)-modules.	11
2.2 The semi-simple filtration.	14
2.3 Semi-simple frescos.	16
3 A criterion for semi-simplicity.	19
3.1 Polynomial dependance.	19
3.2 The α -invariant.	21
4 The existence theorem for frescos.	27
4.1 Preliminaries.	28
4.2 $\tilde{\mathcal{A}}$ -structures.	30
4.3 The finiteness theorem.	34

Introduction

Let $f : X \rightarrow D$ be an holomorphic function on a connected complex manifold. Assume that $\{df = 0\} \subset \{f = 0\} := X_0$. We consider X as a degenerating family of complex manifolds parametrized by $D^* := D \setminus \{0\}$ with a singular member X_0 at the origin of D . Let ω be a smooth $(p+1)$ -differential form on X satisfying $d\omega = 0 = df \wedge \omega$. Then in many interesting cases (see for instance [B.II] , [B.III] for the case of a function with 1-dimensional singular set and the section 4 for the general proper case) the relative family of de Rham cohomology classes induced on the fibers $(X_s)_{s \in D^*}$ of f by ω/df is solution of a minimal filtered differential equation

defined from the Gauss-Manin connection of f . This object, called a **fresco** is a monogenic regular (a,b)-module satisfying an extra condition, called "geometric", which encodes simultaneously the regularity at 0 of the Gauss-Manin connection, the monodromy theorem and B. Malgrange's positivity theorem.

We study the structure of such an object in order to determine the possible quotient themes of a given fresco. Such a theme corresponds to a possible asymptotic expansion of vanishing periods constructed from ω by choosing a vanishing cycle $\gamma \in H_p(X_{s_0}, \mathbb{C})$ and defining

$$F_\gamma(s) := \int_{\gamma_s} \omega/df$$

where γ_s is the (multivalued) horizontal family of cycles defined from γ in the fibers of f (see [M.74]).

Let me describe the content of this article.

After some easy preliminaries, we prove in section 1 that a $[\lambda]$ -primitive fresco E admits an *unique* Jordan-Hölder sequence, called the principal J-H. sequence, in which the opposite of the roots of the Bernstein polynomial of E appears in a non decreasing order. This uniqueness result is important because it implies, for instance, that the isomorphism classes of each quotient of terms of the principal J-H. sequence only depends on the isomorphism class of E .

In section 2 we define and study semi-simple regular (a,b)-modules and the corresponding semi-simple filtration. In the case of a $[\lambda]$ -primitive fresco we prove that semi-simplicity is characterized by the fact that we may find a J-H. sequence in which the opposite of the roots of the Bernstein polynomial appears in strictly decreasing order (but also in any given order).

In section 3 we answer to the following question : How to recognize from the principal J-H. sequence of a $[\lambda]$ -primitive fresco if it is semi-simple. Of course the answer is obvious if some rank 2 sub-quotient theme appears from this sequence. But when it is not the case, such a rank 2 sub-quotient theme may appear after commuting some terms in the J-H. sequence. The simplest example is when

$$E := \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b)(1 + \alpha.b^{p_1+p_2})^{-1}.(a - \lambda_2.b).(a - \lambda_3.b)$$

with $\lambda_{i+1} = \lambda_i + p_i - 1$ for $i = 1, 2$ with $p_i \in \mathbb{N}^*, i = 1, 2$ and $\alpha \in \mathbb{C}^*$. Using the identity in $\tilde{\mathcal{A}}$

$$(a - \lambda_2.b).(a - \lambda_3.b) = (a - (\lambda_3 + 1).b).(a - (\lambda_2 - 1).b)$$

it is easy to see that $(a - (\lambda_2 - 1).b).[1]$ generates a (normal) rank 2 theme in E , because we assume $\alpha \neq 0$.

We solve this question introducing for a $[\lambda]$ -primitive rank $k \geq 2$ fresco E such F_{k-1} and E/F_1 are semi-simple, where $(F_j)_{j \in [1, k]}$ is the principal J-H. sequence of E , the α -invariant $\alpha(E)$. This complex number only depends on the isomorphism class of E and is zero if and only if E is semi-simple. We also prove that when $\alpha(E)$ is not zero it gives the parameter of any normal rank 2 sub-theme of E .

In section 4 we give, using tools introduced in [B.II], a general existence theorem for the fresco associated to a relative de Rham cohomology class in the geometric situation described in the beginning of this introduction, assuming the function f proper.

1 Preliminaries.

1.1 Definitions and characterization as $\tilde{\mathcal{A}}$ -modules.

We are interested in "standard" formal asymptotic expansions of the following type

$$\sum_{q=1}^p \sum_{j=0}^N \mathbb{C}[[s]].s^{\lambda_q-1}.(\text{Log } s)^j$$

where $\lambda_1, \dots, \lambda_q$ are positive rational numbers, and in fact in vector valued such expansions. The two basic operations on such expansions are

- the multiplication by s that we shall denote a ,
- and the primitive in s without constant that we shall denote b .

This leads to consider on the set of such expansions a left module structure on the \mathbb{C} -algebra

$$\tilde{\mathcal{A}} := \left\{ \sum_{\nu=0}^{\infty} P_{\nu}(a).b^{\nu}, \quad \text{where } P_{\nu} \in \mathbb{C}[x] \right\}$$

defined by the following conditions

- The commutation relation $a.b - b.a = b^2$ which is the translation of the Leibnitz rule ;
- The continuity for the b -adic filtration of $\tilde{\mathcal{A}}$ of the left and right multiplications by a .

Define now for λ in $\mathbb{Q} \cap]0, 1]$ and $N \in \mathbb{N}$ the left $\tilde{\mathcal{A}}$ -module

$$\Xi_{\lambda}^{(N)} := \oplus_{j=0}^N \mathbb{C}[[s]].s^{\lambda-1}.(\text{Log } s)^j = \oplus_{j=0}^N \mathbb{C}[[a]].s^{\lambda-1}.(\text{Log } s)^j = \oplus_{j=0}^N \mathbb{C}[[b]].s^{\lambda-1}.(\text{Log } s)^j.$$

Of course we let a and b act on $\Xi_{\lambda}^{(N)}$ as explained above.

Define also, when Λ is a finite subset in $\mathbb{Q} \cap]0, 1]$, the $\tilde{\mathcal{A}}$ -module

$$\Xi_{\Lambda}^{(N)} := \oplus_{\lambda \in \Lambda} \Xi_{\lambda}^{(N)}.$$

More generally, if V is a finite dimensional complex vector space we shall put a structure of left $\tilde{\mathcal{A}}$ -module on $\Xi_{\Lambda}^{(N)} \otimes_{\mathbb{C}} V$ with the following rules :

$$a.(\varphi \otimes v) = (a.\varphi) \otimes v \quad \text{and} \quad b.(\varphi \otimes v) = (b.\varphi) \otimes v$$

for any $\varphi \in \Xi_{\Lambda}^{(N)}$ and any $v \in V$. It will be convenient to denote Ξ_{Λ} the $\tilde{\mathcal{A}}$ -module $\sum_{N \in \mathbb{N}} \Xi_{\Lambda}^{(N)}$.

Definition 1.1.1 A $\tilde{\mathcal{A}}$ -module is called a **fresco** when it is isomorphic to a submodule $\tilde{\mathcal{A}}.\varphi \subset \Xi_{\Lambda}^{(N)} \otimes V$ where φ is any element in $\Xi_{\Lambda}^{(N)} \otimes V$, for some choice of Λ, N and V as above.

A fresco is a **theme** when we may choose $V := \mathbb{C}$ in the preceding choice.

Now the characterization of frescos among all left $\tilde{\mathcal{A}}$ -modules is not so obvious. The following theorem is proved in [B.09].

Theorem 1.1.2 A left $\tilde{\mathcal{A}}$ -module E is a fresco if and only if it is a geometric (a, b) -module which is generated (as a $\tilde{\mathcal{A}}$ -module) by one element. Moreover the annihilator in $\tilde{\mathcal{A}}$ of any generator of E is a left ideal of the form $\tilde{\mathcal{A}}.P$ where P may be written as follows

$$P = (a - \lambda_1.b).S_1^{-1}.(a - \lambda_2.b).S_2^{-1} \dots (a - \lambda_k.b).S_k^{-1}, \quad k := \dim_{\mathbb{C}}(E/b.E)$$

where λ_j are rational numbers such that $\lambda_j + j > k$ for $j \in [1, k]$ and where S_1, \dots, S_k are invertible elements in the sub-algebra $\mathbb{C}[[b]]$ of $\tilde{\mathcal{A}}$.

Conversely, for such a $P \in \tilde{\mathcal{A}}$ the left $\tilde{\mathcal{A}}$ -module $E := \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.P$ is a fresco and it is a free rank k module on $\mathbb{C}[[b]]$.

Let me recall briefly for the convenience of the reader the definitions of the notions involved in the previous statement.

- A **(a,b)-module** E is a free finite rank $\mathbb{C}[[b]]$ module endowed with an \mathbb{C} -linear endomorphism a such that $a.S = S.a + b^2.S'$ for $S \in \mathbb{C}[[b]]$; or, in an equivalent way, a $\tilde{\mathcal{A}}$ -module which is free and finite type over the subalgebra $\mathbb{C}[[b]] \subset \tilde{\mathcal{A}}$.

It has a **simple pole** when a satisfies $a.E \subset b.E$. In this case the **Bernstein polynomial** B_E of E is defined as the minimal polynomial of $-b^{-1}.a$ acting on $E/b.E$.

- A (a, b) -module E is **regular** when it may be embedded in a simple pole (a, b) -module. In this case there is a minimal such embedding which is the inclusion of E in its saturation E^{\sharp} by $b^{-1}.a$. The **Bernstein polynomial** of a regular (a, b) -module is, by definition, the Bernstein polynomial of its saturation E^{\sharp} .
- A regular (a, b) -module E is called **geometric** when all roots of its Bernstein polynomial are rational and strictly negative.

Note that the formal completion in b of the Brieskorn module of a function with an isolated singularity is a geometric (a, b) -module. The last section of this article shows that this structure appears in a rather systematic way in the study of the Gauss-Manin connection of a proper holomorphic function on a complex manifold.

Definition 1.1.3 A submodule F in E is **normal** when $F \cap b.E = b.F$.

For any sub-module F of a (a,b) -module E there exists a minimal normal sub-module \tilde{F} of E containing F . We shall call it the **normalization** of F . It is easy to see that \tilde{F} is the pull-back by the quotient map $E \rightarrow E/F$ of the b -torsion of E/F .

When F is normal the quotient E/F is again a (a,b) -module. Note that a sub-module of a regular (resp. geometric) (a,b) -module is regular (resp. geometric) and when F is normal E/F is also regular (resp. geometric).

Lemma 1.1.4 *Let E be a fresco and F be a normal sub-module in E . Then F is a fresco and also the quotient E/F .*

PROOF. The only point to prove, as we already know that F and E/F are geometric (a,b) -modules thanks to [B.09], is the fact that F and E/F are generated as $\tilde{\mathcal{A}}$ -module by one element. This is obvious for E/F , but not for F . We shall use the theorem 1.1.2 for E/F to prove that F is generated by one element. Let e be a generator of E and let P as in the theorem 1.1.2 which generates the annihilator ideal of the image of e in E/F . Then $P.e$ is in F . We shall prove that $P.e$ generates F as a $\tilde{\mathcal{A}}$ -module. Let y be an element in F and write $y = u.e$ where u is in $\tilde{\mathcal{A}}$. As P is, up to an invertible element in $\mathbb{C}[[b]]$, a monic polynomial in a with coefficients in $\mathbb{C}[[b]]$, we may write $u = Q.P + R$ where Q and R are in $\tilde{\mathcal{A}}$ and R is a polynomial in a with coefficient in $\mathbb{C}[[b]]$ of degree $r < \deg(P) = \text{rank}(E/F)$. Now, as y is in F , the image in E/F of $u.e$ is 0, and this implies that R annihilates the image of e in E/F . So R lies in $\tilde{\mathcal{A}}.P$ and so $R = 0$. Then we have $u = Q.P$ and $y = Q.P.e$ proving our claim. ■

In the case of a fresco the Bernstein polynomial is more easy to describe, thanks to the following proposition proved in [B.09].

Proposition 1.1.5 *Soit $E = \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.P$ be a rank k fresco as described in the previous theorem. The Bernstein polynomial of E is the characteristic polynomial of $-b^{-1}.a$ acting on $E^\sharp/b.E^\sharp$. And the **Bernstein element** P_E of E , which is the element in $\tilde{\mathcal{A}}$ defined by the Bernstein polynomial B_E of E by the following formula*

$$P_E := (-b)^k . B_E(-b^{-1}.a)$$

is equal to $(a - \lambda_1.b) \dots (a - \lambda_k.b)$ for any such choice of presentation of E .

As an easy consequence, the k roots of the Bernstein polynomial of E are the opposite of the numbers $\lambda_1 + 1 - k, \dots, \lambda_k + k - k$. So the Bernstein polynomial is readable on the element P : the initial form of P in (a,b) is the Bernstein element P_E of E .

REMARK. If we have an exact sequence of (a,b) -modules

$$0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$$

where E is a fresco, then F and G are frescos and the Bernstein elements satisfy the equality $P_E = P_F.P_G$ in the algebra $\tilde{\mathcal{A}}$ (see [B.09] proposition 3.4.4). \square

EXAMPLE. The $\tilde{\mathcal{A}}$ -module $\Xi_\lambda^{(N)}$ is a simple pole (a,b) with rank $N + 1$. Its Bernstein polynomial is equal to $(x + \lambda)^{N+1}$.

The theme $\tilde{\mathcal{A}}.\varphi \subset \Xi_\lambda^{(N)}$ where $\varphi = s^{\lambda-1}.(Log s)^N$ has rank $N + 1$ and Bernstein element $(a - (\lambda + N).b)(a - (\lambda + N - 1).b) \dots (a - \lambda.b)$. Its saturation is $\Xi_\lambda^{(N)}$.

The dual of a regular (a,b)-module is regular, but duality does not preserve the property of being geometric because duality change the sign of the roots of the Bernstein polynomial. As duality preserves regularity (see [B.95]), to find again a geometric (a,b)-module it is sufficient to make the tensor product by a rank 1 (a,b)-module E_δ for δ a large enough rational number¹. The next lemma states that this "twist duality" preserves the notion of fresco.

Lemma 1.1.6 *Let E be a fresco and let δ be a rational number such that $E^* \otimes E_\delta$ is geometric. Then $E^* \otimes E_\delta$ is a fresco.*

The proof is obvious. ■

The following definition is useful when we want to consider only the part of asymptotic expansions corresponding to prescribe eigenvalues of the monodromy.

Definition 1.1.7 *Let Λ be a subset of $\mathbb{Q} \cap]0, 1]$. We say that a regular (a,b)-module E is $[\Lambda]$ -primitive when all roots of its Bernstein polynomial are in $-\Lambda + \mathbb{Z}$.*

The following easy proposition is proved in [B.09]

Proposition 1.1.8 *Let E be a regular (a,b)-module and Λ a subset of $\mathbb{Q} \cap]0, 1]$. Then there exists a maximal submodule $E_{[\Lambda]}$ in E which is $[\Lambda]$ -primitive. Moreover the quotient $E/E_{[\Lambda]}$ is a $[\Lambda^c]$ -primitive (a,b)-module, where we denote $\Lambda^c := \mathbb{Q} \cap]0, 1] \setminus \Lambda$.*

We shall mainly consider the case where Λ is a single element. Note that the $[\lambda]$ -primitive part of an (a,b)-module $E \subset \Xi_\lambda^{(N)} \otimes V$ corresponds to its intersection with $\Xi_\lambda^{(N)} \otimes V$.

1.2 The principal Jordan-Hölder sequence.

The classification of rank 1 regular (a,b)-module is very simple : each isomorphy class is given by a complex number and to $\lambda \in \mathbb{C}$ corresponds the isomorphy class of

¹to tensor by E_δ is the same that to replace a by $a + \delta.b$; see [B.I] for the general definition of the tensor product.

$E_\lambda := \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda.b)$. Then **Jordan-Hölder sequence** for a regular (a,b)-module E is a sequence

$$0 = F_0 \subset F_1 \subset \cdots \subset F_k = E$$

of normal sub-modules such that for each $j \in [1, k]$ the quotient F_j/F_{j-1} has rank 1. Then to each J-H. sequence we may associate an ordered sequence of complex numbers $\lambda_1, \dots, \lambda_k$ such that $F_j/F_{j-1} \simeq E_{\lambda_j}$.

EXAMPLE. A regular rank 1 (a,b)-module is a fresco if and only if it is isomorphic to E_λ for some $\lambda \in \mathbb{Q}^{+*}$. All rank 1 frescos are themes. The classification of rank 2 regular (a,b)-modules given in [B.93] gives the list of $[\lambda]$ -primitive rank 2 frescos which is the following, where $\lambda_1 > 1$ is a rational number :

$$E = E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(a - (\lambda_1 - 1).b) \quad (1)$$

$$E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(1 + \alpha.b^p)^{-1}.(a - (\lambda_1 + p - 1).b) \quad (2)$$

where $p \in \mathbb{N} \setminus \{0\}$ and $\alpha \in \mathbb{C}$.

The themes in this list are these in (1) and these in (2) with $\alpha \neq 0$. For a $[\lambda]$ -primitive fresco in case (2) the number α will be called the **parameter** of the rank 2 fresco. By convention we shall define $\alpha := 1$ in the case (1). \square

The following existence result is proved in [B.93]

Proposition 1.2.1 *For any regular (a,b)-module E of rank k there exists a J-H. sequence. The numbers $\exp(2i\pi.\lambda_j)$ are independant of the J-H. sequence, up to permutation. Moreover the number $\mu(E) := \sum_{j=1}^k \lambda_j$ is also independent of the choice of the J-H. sequence of E .*

EXERCICE. Let E be a regular (a,b)-module and $E' \subset E$ be a sub-(a,b)-module with the same rank than E . Show that E' has finite \mathbb{C} -codimension in E given by

$$\dim_{\mathbb{C}} E/E' = \mu(E') - \mu(E).$$

hint : make an induction on the rank of E . \square

For a $[\lambda]$ -primitive fresco a more precise result is proved in [B.09]

Proposition 1.2.2 *For any J-H. sequence of a rank k $[\lambda]$ -primitive fresco E the numbers $\lambda_j + j - k, j \in [1, k]$ are the opposite of the roots (with multiplicities) of the Bernstein polynomial of E . So, up to a permutation, they are independant of the choice of the J-H. sequence. Moreover, there always exists a J-H. sequence such that the associated sequence $\lambda_j + j$ is non decreasing.*

For a $[\lambda]$ -primitive theme the situation is extremely rigid (see [B.09]) :

Proposition 1.2.3 *Let E a rank k $[\lambda]$ -primitive theme. Then, for each j in $[0, k]$, there exists an unique normal rank j submodule F_j . The numbers associated to the unique J-H. sequence satisfy the condition that $\lambda_j + j, j \in [1, k]$ is an non decreasing sequence.*

Definition 1.2.4 *Let E be a $[\lambda]$ -primitive fresco of rank k and let*

$$0 = F_0 \subset F_1 \subset \cdots \subset F_k = E$$

*be a J-H. sequence of E . Then for each $j \in [1, k]$ we have $F_j/F_{j-1} \simeq E_{\lambda_j}$, where $\lambda_1, \dots, \lambda_k$ are in $\lambda + \mathbb{N}$. We shall say that such a J-H. sequence is **principal** when the sequence $[1, k] \ni j \mapsto \lambda_j + j$ is non decreasing.*

Proposition 1.2.5 *Let E be a $[\lambda]$ -primitive fresco. Then its principal J-H. sequence is unique.*

We shall prove the uniqueness by induction on the rank k of E .

We begin by the case of rank 2.

Lemma 1.2.6 *Let E be a rank 2 $[\lambda]$ -primitive fresco and let λ_1, λ_2 the numbers corresponding to a principal J-H. sequence of E (so $\lambda_1 + 1 \leq \lambda_2 + 2$). Then the normal rank 1 submodule of E isomorphic to E_{λ_1} is unique.*

PROOF. The case $\lambda_1 + 1 = \lambda_2 + 2$ is obvious because then E is a $[\lambda]$ -primitive theme (see [B.10] corollary 2.1.7). So we may assume that $\lambda_2 = \lambda_1 + p_1 - 1$ with $p_1 \geq 1$ and that E is the quotient $E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(a - \lambda_2.b)$ (see the classification of rank 2 frescos in 2.2), because the result is clear when E is a theme. We shall use the $\mathbb{C}[[b]]$ -basis e_1, e_2 of E where a is defined by the relations

$$(a - \lambda_2.b).e_2 = e_1 \quad (a - \lambda_1.b).e_1 = 0.$$

This basis comes from the isomorphism $E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(a - \lambda_2.b)$ deduced from the classification of rank 2 frescos with $e_2 = [1]$ and $e_1 = (a - \lambda_2.b).e_2$.

Let look for $x := U.e_2 + V.e_1$ such that $(a - \lambda_1.b).x = 0$. Then we obtain

$$b^2.U'.e_2 + U.(a - \lambda_2.b).e_2 + (\lambda_2 - \lambda_1).b.U.e_2 + b^2.V'.e_1 = 0$$

which is equivalent to the two equations :

$$b^2.U' + (p_1 - 1).b.U = 0 \quad \text{and} \quad U + b^2.V' = 0.$$

The first equation gives $U = 0$ for $p_1 \geq 2$ and $U \in \mathbb{C}$ for $p_1 = 1$. As the second equation implies $U(0) = 0$, in all cases $U = 0$ and $V \in \mathbb{C}$. So the solutions are in $\mathbb{C}.e_1$. ■

Remark that in the previous lemma, if we assume $p_1 \geq 1$ and E is not a theme, it may exist infinitely many different normal (rank 1) submodules isomorphic to E_{λ_2+1} . But then, $\lambda_2 + 2 > \lambda_1 + 1$. This happens in the following example.

EXAMPLE. Let $E := \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.P$ with

$$P := (a - \lambda_1.b).(a - \lambda_2.b)(1 + \alpha.b^{p_2})^{-1}.(a - \lambda_3.b)$$

where $\lambda_1 > 2$ is rational, p_1 and p_2 are in \mathbb{N}^* , $\lambda_2 = \lambda_1 + p_1 - 1$, $\lambda_3 = \lambda_2 + p_2 - 1$ and α is in \mathbb{C}^* .

Then using the identity (see the commuting lemma in [B.09])

$$(a - \lambda_1.b).(a - \lambda_2.b) = U^{-1}.(a - (\lambda_2 + 1).b).U^2.(a - (\lambda_1 - 1).b).U^{-1}$$

with $U := 1 + \rho.b^{p_1}$, $\rho \in \mathbb{C}^*$, it is easy to see that the rank 3 fresco E admits the rank 2 theme T with fundamental invariants $(\lambda_1 - 1, \lambda_3)$ and parameter $\rho.\alpha$. Remark that if we use the previous identity with $\rho = 0$ (so $U = 1$), then using the identity

$$(a - (\lambda_1 - 1).b).(1 + \alpha.b^{p_2})^{-1}.(a - \lambda_3.b) = \\ V^{-1}.(a - (\lambda_3 + 1).b).V^2.(1 + \alpha.b^{p_2})^{-1}.(a - (\lambda_1 - 2).b).V^{-1}$$

where $V = 1 + \beta.b^{p_2}$ and $\beta := (1 + p_2/p_1).\alpha$ we see that E contains a normal rank 2 sub-theme which has fundamental invariants equal to $(\lambda_2 + 1, \lambda_3 + 1)$ and parameter β . \square

PROOF OF PROPOSITION 1.2.5. As the result is obvious for $k = 1$, we may assume $k \geq 2$ and the result proved in rank $\leq k - 1$. Let $F_j, j \in [1, k]$ and $G_j, j \in [1, k]$ two J-H. principal sequences for E . As the sequences $\lambda_j + j$ and $\mu_j + j$ coincide up to the order and are both non decreasing, they coincide. Now let j_0 be the first integer in $[1, k]$ such that $F_{j_0} \neq G_{j_0}$. If $j_0 \geq 2$ applying the induction hypothesis to E/F_{j_0-1} gives $F_{j_0}/F_{j_0-1} = G_{j_0}/F_{j_0-1}$ and so $F_{j_0} = G_{j_0}$.

So we may assume that $j_0 = 1$. Let H be the normalization of $F_1 + G_1$. As F_1 and G_1 are normal rank 1 and distinct, then H is a rank 2 normal submodule. It is a $[\lambda]$ -primitive fresco of rank 2 with two normal rank 1 sub-modules which are isomorphic as $\lambda_1 = \mu_1$. Moreover the principal J-H. sequence of H begins by a normal submodule isomorphic to E_{λ_1} . So the previous lemma implies $F_1 = G_1$. So for any $j \in [1, k]$ we have $F_j = G_j$. \blacksquare

Definition 1.2.7 Let E be a $[\lambda]$ -primitive fresco and consider its principal J-H. sequence $F_j, j \in [1, k]$. Put $F_j/F_{j-1} \simeq E_{\lambda_j}$ for $j \in [1, k]$ (with $F_0 = \{0\}$). We shall call **fundamental invariants** of E the ordered k -tuple $(\lambda_1, \dots, \lambda_k)$.

Of course, if we have any J-H. sequence for a $[\lambda]$ -primitive fresco E with rank 1 quotients associated to the rational numbers μ_1, \dots, μ_k , it is easy to recover the fundamental invariants of E because the numbers $\mu_j + j, j \in [1, k]$ are the same than the numbers $\lambda_j + j$ up to a permutation. But as the sequence $\lambda_j + j$ is non decreasing with j , it is enough to put the $\mu_j + j$ in the non decreasing order with j to conclude.

Let $\lambda_1, \dots, \lambda_k$ be the fundamental invariants of a $[\lambda]$ -primitive fresco E_0 , and not $\mathcal{F}(\lambda_1, \dots, \lambda_k)$ the set of isomorphism classes of frescos with these fundamental invariants. The uniqueness of the principal J-H. sequence of a $[\lambda]$ -primitive fresco allows to define for each (i, j) $1 \leq i < j \leq k$ a map

$$q_{i,j} : \mathcal{F}(\lambda_1, \dots, \lambda_k) \rightarrow \mathcal{F}(\lambda_i, \dots, \lambda_j)$$

defined by $q_{i,j}([E]) = [F_j/F_{i-1}]$ where $(F_h)_{h \in [0,k]}$ is the principal J-H. sequence of E . This makes sens because any isomorphism $\varphi : E^1 \rightarrow E^2$ between two $[\lambda]$ -primitive frescos induces isomorphisms between each term of the corresponding principal J-H. sequences.

For instance classification of rank 2 $[\lambda]$ -primitive frescos gives (see example before proposition 1.2.1) for any rational number $\lambda_1 > 1$ and $p_1 \in \mathbb{N}$:

$$\begin{aligned} \text{for } p_1 = 0 \quad \mathcal{F}(\lambda_1, \lambda_1 - 1) &= \{pt\} \\ \text{for } p_1 \geq 1 \quad \alpha : \mathcal{F}(\lambda_1, \lambda_1 + p_1 - 1) &\xrightarrow{\simeq} \mathbb{C} \end{aligned}$$

and $\{pt\}$ is given by the isomorphism class of $\tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(a - (\lambda_1 - 1).b)$ and the isomorphism class associated to $\alpha^{-1}(x)$ for $x \in \mathbb{C}$ in the case $p_1 \geq 1$ is given by $\tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(1 + x.b^{p_1})^{-1}.(a - (\lambda_1 + p_1 - 1).b)$.

2 Semi-simplicity.

2.1 Semi-simple regular (a,b)-modules.

Definition 2.1.1 *Let E be a regular (a,b)-module. We say that E is **semi-simple** if it is a sub-module of a finite direct sum of rank 1 regular (a,b)-modules.*

Note that if E is a sub-module of a regular (a,b)-module it is necessary regular. As a direct sum of regular (a,b)-modules if regular, our assumption that E is regular is superfluous.

It is clear from this definition that a sub-module of a semi-simple (a,b)-module is semi-simple and that a (finite) direct sum of semi-simple (a,b)-modules is again semi-simple.

REMARK. A rather easy consequence of the classification of rank 2 (a,b)-modules given in [B.93] is that the rank 2 simple pole (a,b)-modules defined in the $\mathbb{C}[[b]]$ -basis x, y by the relations :

$$(a - (\lambda + p).b).x = b^{p+1}.y \quad \text{and} \quad (a - \lambda).y = 0$$

for any $\lambda \in \mathbb{C}$ and any $p \in \mathbb{N}$ are not semi-simple. We leave the verification of this point to the reader.

Let us begin by a characterization of the semi-simple (a,b)-modules which have a simple pole. First we shall prove that a quotient of a semi-simple (a,b)-module is semi-simple. This will be deduced from the following lemma.

Lemma 2.1.2 *Let E be a (a,b) -module which is direct sum of regular rank 1 (a,b) -modules, and let $F \subset E$ be a rank 1 normal sub-module. Then F is a direct factor of E .*

Corollary 2.1.3 *If E is a semi-simple regular (a,b) -module and F a normal sub-module of E , the quotient E/F is a (regular) semi-simple (a,b) -module.*

PROOF OF THE LEMMA. Let $E = \bigoplus_{j=1}^k E_{\lambda_j}$ and assume that $F \simeq E_{\mu}$. Let e_j be a standard generator of E_{λ_j} and e a standard generator of E_{μ} . Write

$$e = \sum_{j=1}^k S_j(b).e_j$$

and compute $(a - \mu.b).e = 0$ using the fact that $e_j, j \in [1, k]$ is a $\mathbb{C}[[b]]$ -basis of E and the relations $(a - \lambda_j.b).e_j = 0$ for each j . We obtain for each $j \in [1, k]$ the relation

$$b.S'_j - (\mu - \lambda_j).S_j = 0.$$

So, if $\mu - \lambda_j$ is not in \mathbb{N} , we have $S_j = 0$. When $\mu = \lambda_j + p_j$ with $p_j \in \mathbb{N}$ we obtain $S_j(b) = \rho_j.b^{p_j}$ for some $\rho_j \in \mathbb{C}$. As we assume that e is not in $b.E$, there exists at least one $j_0 \in [1, k]$ such that $p_{j_0} = 0$ and $\rho_{j_0} \neq 0$. Then it is clear that we have

$$E = F \oplus \left(\bigoplus_{j \neq j_0} E_{\lambda_j} \right)$$

concluding the proof. ■

PROOF OF THE COROLLARY. We argue by induction on the rank of F . In the rank 1 case, we have $F \subset E \subset \mathcal{E} := \bigoplus_{j=1}^k E_{\lambda_j}$. Let \tilde{F} the normalization of F in $\bigoplus_{j=1}^k E_{\lambda_j}$. Then the lemma shows that there exists a $j_0 \in [1, k]$ such that

$$\mathcal{E} = \tilde{F} \oplus \bigoplus_{j \neq j_0} E_{\lambda_j}.$$

Then, as $\tilde{F} \cap E = F$, the quotient map $\mathcal{E} \rightarrow \mathcal{E}/\tilde{F} \simeq \bigoplus_{j \neq j_0} E_{\lambda_j}$ induces an injection of E/F in a direct sum of regular rank 1 (a,b) -modules. So E/F is semi-simple. Assume now that the result is proved for F with rank $\leq d - 1$ and assume that F has rank d . Then using a rank 1 normal sub-module G in F , we obtain that F/G is a normal rank $d - 1$ sub-module of E/G . Using the rank 1 case we know that E/G is semi-simple, and the induction hypothesis gives that

$$E/F = (E/G) / (F/G)$$

is semi-simple. ■

Proposition 2.1.4 *Let E be a simple pole semi-simple (a,b) -module. Then E is a direct sum of regular rank 1 (a,b) -modules.*

PROOF. We shall prove the proposition by induction on the rank k of E . As the rank 1 case is obvious, assume the proposition proved for $k - 1$ and that E is a simple pole rank $k \geq 2$ semi-simple (a,b) -module. From the existence of J-H. sequence, we may find an exact sequence

$$0 \rightarrow F \rightarrow E \rightarrow E_\lambda \rightarrow 0$$

where F has rank $k - 1$. By definition F is semi-simple, but it also has a simple pole because $a.F \subset a.E \subset b.E$ implies that $a.F \subset F \cap b.E = b.F$ as F is normal in E . So by the induction hypothesis F is a direct sum of regular rank 1 (a,b) -modules.

Let $e \in E$ such that its image in E_λ is e_λ a standard generator of E_λ satisfying $a.e_\lambda = \lambda.b.e_\lambda$. Then we have $(a - \lambda.b).e \in F$. We shall first look at the case $k = 2$. So F is a rank 1 and we have $F \simeq E_\mu$ for some $\mu \in \mathbb{C}$. Let e_μ be a standard generator in F and put

$$(a - \lambda.b).e = S(b).e_\mu.$$

Our simple pole assumption on E implies $S(0) = 0$ and we may write $S(b) = b.\tilde{S}(b)$. We look for $T \in \mathbb{C}[[b]]$ such that $\varepsilon := e + T(b).e_\mu$ satisfies

$$(a - \lambda.b).\varepsilon = 0.$$

If such a T exists, then we would have $E = E_\mu \oplus E_\lambda$ where E_λ is the submodule generated by ε , because it is clear that we have $E = F \oplus \mathbb{C}[[b]].e$ as a $\mathbb{C}[[b]]$ -module. To find T we have to solve in $\mathbb{C}[[b]]$ the differential equation

$$b.\tilde{S}(b) + (\mu - \lambda).b.T(b) + b^2.T'(b) = 0.$$

If $\lambda - \mu$ is not a non negative integer, such a T exists and is unique. But when $\lambda = \mu + p$ with $p \in \mathbb{N}$, the solution exists if and only if the coefficient of b^p in \tilde{S} is zero. If it is not the case, define \tilde{T} as the solution of the differential equation

$$\tilde{S}(b) - \alpha.b^p + b.\tilde{T}'(b) - p.\tilde{T}(b) = 0$$

where α is the coefficient of b^p in \tilde{S} and where we choose \tilde{T} by asking that it has no b^p term. Then $\varepsilon_1 := e + \tilde{T}(b).e_{\lambda-p}$ satisfies

$$(a - (\lambda - p).b).\varepsilon_1 = \alpha.b^{p+1}.e_{\lambda-p}.$$

Then, after changing $\lambda - p$ in λ and $e_{\lambda-p}$ in $\alpha.e_{\lambda-p}$, we recognize one of the rank 2 modules which appears in the previous remark and which is not semi-simple. So we have a contradiction. This concludes the rank 2 case.

Now consider the case $k \geq 3$ and using the induction hypothesis write

$$F \simeq \bigoplus_{j=1}^{k-1} E_{\mu_j}$$

and denote e_j a standard generator for E_{μ_j} . Write

$$(a - \lambda.b).e = \sum_{j=1}^{k-1} S_j(b).e_j.$$

The simple pole assumption again gives $S_j(0) = 0$ for each j ; now we look for T_1, \dots, T_{k-1} in $\mathbb{C}[[b]]$ such that, defining $\varepsilon := e + \sum_{j=1}^{k-1} T_j(b).e_j$, we have

$$(a - \lambda.b).\varepsilon = 0.$$

For each j this leads to the differential equation

$$b.\tilde{S}_j(b) + (\mu_j - \lambda).b.T_j(b) + b^2.T_j'(b) = 0$$

where we put $S_j(b) = b.\tilde{S}_j(b)$. We are back, for each given $j \in [1, k-1]$, to the previous problem in the rank 2 case. So if for each $j \in [1, k-1]$ we have a solution $T_j \in \mathbb{C}[[b]]$ it is easy to conclude that

$$E \subset F \oplus \mathbb{C}[[b]].\varepsilon \simeq F \oplus E_\lambda.$$

If for some j_0 there is no solution, the coefficient of b^{j_0} in \tilde{S}_{j_0} does not vanish and then the image of E in $E / \bigoplus_{j \neq j_0} E_{\mu_j}$ is a rank 2 not semi-simple (a,b)-module. This contradicts the corollary 2.1.3 and concludes the proof. \blacksquare

REMARK. Let E be a semi-simple rank k (a,b)-module and let E^\sharp its saturation by $b^{-1}.a$, then E^\sharp is semi-simple because there exists $N \in \mathbb{N}$ with an inclusion $E^\sharp \subset b^{-N}.E$. Also E^\flat its maximal simple pole sub-module is semi-simple. Then we have

$$E^\flat \simeq \bigoplus_{j=1}^k E_{\lambda_j + d_j} \subset E \subset E^\sharp \simeq \bigoplus_{j=1}^k E_{\lambda_j}$$

where d_1, \dots, d_k are non negative integers.

Corollary 2.1.5 *Let E be a semi-simple (a,b)-module. Then its dual E^* is semi-simple.*

PROOF. For a regular (a,b)-module E we have $(E^\flat)^* = (E^*)^\sharp$. But the dual of E_λ is $E_{-\lambda}$ and the duality commutes with direct sums. So we conclude that $E^* \subset (E^\flat)^*$ and so E^* is a semi-simple. \blacksquare

REMARK. The tensor product of two semi-simple (a,b)-modules is semi-simple as a consequence of the fact that $E_\lambda \otimes E_\mu = E_{\lambda+\mu}$.

2.2 The semi-simple filtration.

Definition 2.2.1 Let E be a regular (a,b) -module and x an element in E . We shall say that x is **semi-simple** if $\tilde{\mathcal{A}}.x$ is a semi-simple (a,b) -module.

It is clear that any element in a semi-simple (a,b) -module is semi-simple. The next lemma shows that the converse is true.

Lemma 2.2.2 Let E be a regular (a,b) -module such that any $x \in E$ is semi-simple. Then E is semi-simple.

PROOF. Let e_1, \dots, e_k be a $\mathbb{C}[[b]]$ -basis of E . Then each $\tilde{\mathcal{A}}.e_j$ is semi-simple, and the map $\bigoplus_{j=1}^k \tilde{\mathcal{A}}.e_j \rightarrow E$ is surjective. So E is semi-simple thanks to the corollary 2.1.3 and the comment following the definition 2.1.1. ■

Lemma 2.2.3 Let E be a regular (a,b) -module. The subset $S_1(E)$ of semi-simple elements in E is a normal submodule in E .

PROOF. As a direct sum and quotient of semi-simple (a,b) -modules are semi-simple, it is clear that for x and y semi-simple the sum $\tilde{\mathcal{A}}.x + \tilde{\mathcal{A}}.y$ is semi-simple. So $x + y$ is semi-simple. This implies that $S_1(E)$ is a submodule of E . If $b.x$ is in $S_1(E)$, then $\tilde{\mathcal{A}}.b.x$ is semi-simple. Then $\tilde{\mathcal{A}}.x \simeq \tilde{\mathcal{A}}.b.x \otimes E_{-1}$ is also semi-simple, and so $S_1(E)$ is normal in E . ■

Definition 2.2.4 Let E be a regular (a,b) -module. Then the submodule $S_1(E)$ of semi-simple elements in E will be called the **semi-simple part** of E . Defining $S_j(E)$ as the pull-back on E of the semi-simple part of $E/S_{j-1}(E)$ for $j \geq 1$ with the initial condition $S_0(E) = \{0\}$, we define a sequence of normal submodules in E such that $S_j(E)/S_{j-1}(E) = S_1(E/S_{j-1}(E))$. We shall call it the **semi-simple filtration** of E . The smallest integer d such we have $S_d(E) = E$ will be called the **semi-simple depth** of E and we shall denote it $d(E)$.

REMARKS.

- i) As $S_1(E)$ is the maximal semi-simple sub-module of E it contains any rank 1 sub-module of E . So $S_1(E) = \{0\}$ happens if and only if $E = \{0\}$.
- ii) Then the ss-filtration of E is strictly increasing for $0 \leq j \leq d(E)$.
- iii) It is easy to see that for any submodule F in E we have $S_j(F) = S_j(E) \cap F$. So $S_j(E)$ is the subset of $x \in E$ such that $d(x) := d(\tilde{\mathcal{A}}.x) \leq j$ and $d(E)$ is the supremum of $d(x)$ for $x \in E$. □

Lemma 2.2.5 *Let $0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$ be an exact sequence of regular (a,b) -modules such that F and G are semi-simple and such that for any roots λ, μ of the Bernstein polynomial of F and G respectively we have $\lambda - \mu \notin \mathbb{Z}$. Then E is semi-simple.*

PROOF. By an easy induction we may assume that F is rank 1 and also that G is rank 1. Then the conclusion follows from the classification of the rank 2 regular (a,b) -module. ■

APPLICATION. A regular (a,b) -module E is semi-simple if and only if for each $\lambda \in \mathbb{C}$ its primitive part $E[\lambda]$ is semi-simple.

The following easy facts are left as exercises for the reader.

1. Let $0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$ be a short exact sequence of regular (a,b) -modules. Then we have the inequalities

$$\sup\{d(F), d(G)\} \leq d(E) \leq d(F) + d(G).$$

2. Let $E \subset \Xi_\lambda^{(N)} \otimes V$ be a sub-module where V is a finite dimensional complex space. Then for each $j \in [1, N + 1]$ we have

$$S_j(E) = E \cap \left(\Xi_\lambda^{(j-1)} \otimes V \right).$$

3. In the same situation as above, the minimal N for such an embedding of a geometric $[\lambda]$ -primitive E is equal to $d(E) - 1$.
4. In the same situation as above, the equality $S_1(E) = E \cap \Xi_\lambda^{(0)} \otimes V$ implies $\text{rank}(S_1(E)) \leq \dim_{\mathbb{C}} V$. It is easy to see that any $[\lambda]$ -primitive semi-simple geometric (a,b) -module F may be embedded in $\Xi_\lambda^{(0)} \otimes V$ for some V of dimension $\text{rank}(F)$. But it is also easy to prove that when $F = S_1(E)$ with E as above, such an embedding may be extended to an embedding of E into $\Xi_\lambda^{(d(E)-1)} \otimes V$. □

2.3 Semi-simple frescos.

We begin by a simple remark : A $[\lambda]$ -primitive theme is semi-simple if and only if it has rank ≤ 1 . This is an easy consequence of the fact that the saturation of a rank 2 $[\lambda]$ -primitive theme is one of the rank 2 (a,b) -modules considered in the remark following the definition 2.1.1 which are not semi-simple

Lemma 2.3.1 *A geometric (a,b) -module E is **semi-simple** if and only if any quotient of E which is a $[\lambda]$ -primitive theme for some $[\lambda] \in \mathbb{Q}/\mathbb{Z}$ is of rank ≤ 1 .*

PROOF. The condition is clearly necessary as a quotient of a semi-simple (a,b)-module is semi-simple (see corollary 2.1.3), thanks to the remark above. Using the application of the lemma 2.2.5, it is enough to consider the case of a $[\lambda]$ -primitive fresco to prove that the condition is sufficient. Let $\varphi : E \rightarrow \Xi_\lambda^{(N)} \otimes V$ be an embedding of E which exists thanks to the embedding theorem 4.2.1. of [B.09], we obtain that each component of this map in a basis v_1, \dots, v_p of V has rank at most 1 as its image is a $[\lambda]$ -primitive theme. Then each of these images is isomorphic to some $E_{\lambda+q}$ for some integer q . So we have in fact an embedding of E in a direct sum of $E_{\lambda+q_i}$ and E is semi-simple. ■

REMARK. The preceding proof shows that a fresco is a non zero $[\lambda]$ -primitive theme for some $[\lambda] \in \mathbb{Q}/\mathbb{Z}$ if and only if $S_1(E)$ has rank 1. □

Lemma 2.3.2 *Let E be a semi-simple fresco with rank k and let $\lambda_1, \dots, \lambda_k$ be the numbers associated to a J-H. sequence of E . Let μ_1, \dots, μ_k be a twisted permutation² of $\lambda_1, \dots, \lambda_k$. Then there exists a J-H. sequence for E with quotients corresponding to μ_1, \dots, μ_k .*

PROOF. As the symmetric group \mathfrak{S}_k is generated by the transpositions $t_{j,j+1}$ for $j \in [1, k-1]$, it is enough to show that, if E has a J-H. sequence with quotients given by the numbers $\lambda_1, \dots, \lambda_k$, then there exists a J-H. sequence for E with quotients $\lambda_1, \dots, \lambda_{j-1}, \lambda_{j+1} + 1, \lambda_j - 1, \lambda_{j+2}, \dots, \lambda_k$ for $j \in [1, k-1]$. But $G := F_{j+1}/F_{j-1}$ is a rank 2 sub-quotient of E with an exact sequence

$$0 \rightarrow E_{\lambda_j} \rightarrow G \rightarrow E_{\lambda_{j+1}} \rightarrow 0.$$

As G is a rank 2 semi-simple fresco, it admits also an exact sequence

$$0 \rightarrow G_1 \rightarrow G \rightarrow G/G_1 \rightarrow 0$$

with $G_1 \simeq E_{\lambda_{j+1}+1}$ and $G/G_1 \simeq E_{\lambda_j-1}$. Let $q : F_{j+1} \rightarrow G$ be the quotient map. Now the J-H. sequence for E given by

$$F_1, \dots, F_{j-1}, q^{-1}(G_1), F_{j+1}, \dots, F_k = E$$

satisfies our requirement. ■

Proposition 2.3.3 *Let E be a $[\lambda]$ -primitive fresco. A necessary and sufficient condition in order that E is semi-simple is that it admits a J-H. sequence with quotient corresponding to μ_1, \dots, μ_k such that the sequence $\mu_j + j$ is strictly decreasing.*

²This means that the sequence $\mu_j + j, j \in [1, k]$ is a permutation (in the usual sens) of $\lambda_j + j, j \in [1, k]$.

REMARKS.

1. As a fresco is semi-simple if and only if for each $[\lambda]$ its $[\lambda]$ -primitive part is semi-simple, this proposition gives also a criterium to semi-simplicity for any fresco.
2. This criterium is a very efficient tool to produce easily examples of semi-simple frescos.

PROOF. Remark first that if we have, for a $[\lambda]$ -primitive fresco E , a J-H. sequence $F_j, j \in [1, k]$ such that $\lambda_j + j = \lambda_{j+1} + j + 1$ for some $j \in [1, k - 1]$, then F_{j+1}/F_{j-1} is a sub-quotient of E which is a $[\lambda]$ -primitive theme of rank 2. So E is not semi-simple. As a consequence, when a $[\lambda]$ -primitive fresco E is semi-simple the principal J-H. sequence corresponds to a strictly increasing sequence $\lambda_j + j$. Now, thanks to the previous lemma we may find a J-H. sequence for E corresponding to the strictly decreasing order for the sequence $\lambda_j + j$.

No let us prove the converse. We shall use the following lemma.

Lemma 2.3.4 *Let F be a rank k semi-simple $[\lambda]$ -primitive fresco and let $\lambda_j + j$ the strictly increasing sequence corresponding to its principal J-H. sequence. Let $\mu \in [\lambda]$ such that $0 < \mu + k < \lambda_1 + 1$. Then any fresco E in an exact sequence*

$$0 \rightarrow F \rightarrow E \rightarrow E_\mu \rightarrow 0$$

is semi-simple (and $[\lambda]$ -primitive).

PROOF. The case $k = 1$ is obvious, so assume that $k \geq 2$ and that we have a rank 2 quotient $\varphi : E \rightarrow T$ where T is a $[\lambda]$ -primitive theme. Then $\text{Ker } \varphi \cap F$ is a normal submodule of F of rank $k - 2$ or $k - 3$ (for $k \geq 3$). If $\text{Ker } \varphi \cap F$ is of rank $k - 3$, the rank of $F/(\text{Ker } \varphi \cap F)$ is 2 and it injects in T via φ . So $F/(\text{Ker } \varphi \cap F)$ is a rank 2 $[\lambda]$ -primitive theme. As it is semi-simple, because F is semi-simple, we get a contradiction.

So the rank of $F/(\text{Ker } \varphi \cap F)$ is 1 and we have an exact sequence

$$0 \rightarrow F/(\text{Ker } \varphi \cap F) \rightarrow T \rightarrow E/F \rightarrow 0.$$

Put $F/(\text{Ker } \varphi \cap F) \simeq F_1(T) \simeq E_\nu$. Because T is a $[\lambda]$ -primitive theme, we have the inequality $\nu + 1 \leq \mu + 2$. Looking at a J-H. sequence of E ending by

$$\cdots \subset \text{Ker } \varphi \cap F \subset \varphi^{-1}(F_1(T)) \subset E$$

we see that $\nu + k - 1$ is in the set $\{\lambda_j + j, j \in [1, k]\}$ and as $\lambda_1 + 1$ is the infimum of this set we obtain $\lambda_1 + 1 \leq \mu + k$ contradicting our assumption. ■

END OF PROOF OF THE PROPOSITION 2.3.3. Now we shall prove by induction on the rank of a $[\lambda]$ -primitive fresco E that if it admits a J-H. sequence corresponding to a strictly decreasing sequence $\mu_j + j$, it is semi-simple. As the result is obvious in rank 1, we may assume $k \geq 1$ and the result proved for k . So let E be a fresco of rank $k + 1$ and let $F_j, j \in [1, k + 1]$ a J-H. sequence for E corresponding to the strictly decreasing sequence $\mu_j + j, j \in [1, k + 1]$. Put $F_j/F_{j-1} \simeq E_{\mu_j}$ for all $j \in [1, k + 1]$, define $F := F_k$ and $\mu := \mu_{k+1}$; then the induction hypothesis gives that F is semi-simple and we apply the previous lemma to conclude. ■

The following interesting corollary is an obvious consequence of the previous proposition.

Corollary 2.3.5 *Let E be a fresco and let $\lambda_1, \dots, \lambda_k$ be the numbers associated to any J-H. sequence of E . Let μ_1, \dots, μ_d be the numbers associated to any J-H. sequence of $S_1(E)$. Then, for $j \in [1, k]$, there exists a rank 1 normal submodule of E isomorphic to $E_{\lambda_j + j - 1}$ if and only if there exists $i \in [1, d]$ such that we have $\lambda_j + j - 1 = \mu_i + i - 1$.*

Of course, this gives the list of all isomorphy classes of rank 1 normal submodules contained in E . So, using shifted duality, we get also the list of all isomorphy classes of rank 1 quotients of E . It is interesting to note that this is also the list of the possible initial exponents for maximal logarithmic terms which appear in the asymptotic expansion of a given relative de Rham cohomology class after integration on any vanishing cycle in the spectral subspace of the monodromy associated to the eigenvalue $\exp(2i\pi.\lambda)$.

3 A criterion for semi-simplicity.

3.1 Polynomial dependance.

All \mathbb{C} -algebras have a unit.

When we consider a sequence of algebraically independent variables $\rho := (\rho_i)_{i \in \mathbb{N}}$ we shall denote by $\mathbb{C}[\rho]$ the \mathbb{C} -algebra generated by these variables. Then $\mathbb{C}[\rho][[b]]$ will be the commutative \mathbb{C} -algebra of formal power series

$$\sum_{\nu=0}^{\infty} P_{\nu}(\rho).b^{\nu}$$

where $P_{\nu}(\rho)$ is an element in $\mathbb{C}[\rho]$ so a polynomial in $\rho_0, \dots, \rho_{N(\nu)}$ where $N(\nu)$ is an integer depending on ν . So each coefficient in the formal power serie in b depends only on a finite number of the variables ρ_i .

Definition 3.1.1 Let E be a (a,b) -module and let $e(\rho)$ be a family of elements in E depending on a family of variables $(\rho_i)_{i \in \mathbb{N}}$. We say that $e(\rho)$ **depends polynomially on** ρ if there exists a fixed $\mathbb{C}[[b]]$ -basis e_1, \dots, e_k of E such that

$$e(\rho) = \sum_{j=1}^k S_j(\rho).e_j$$

where S_j is for each $j \in [1, k]$ an element in the algebra $\mathbb{C}[\rho][[b]]$.

REMARKS.

1. It is important to remark that when $e(\rho)$ depends polynomially of ρ , then $a.e(\rho)$ also. Then for any $u \in \tilde{\mathcal{A}}$, again $u.e(\rho)$ depends polynomially on ρ .
2. It is easy to see that we obtain an equivalent condition on the family $e(\rho)$ by asking that the coefficient of $e(\rho)$ are in $\mathbb{C}[\rho][[b]]$ in a $\mathbb{C}[[b]]$ -basis of E whose elements depend polynomially of ρ .
3. The invertible elements in the algebra $\mathbb{C}[\rho][[b]]$ are exactly those elements with a constant term in b invertible in the algebra $\mathbb{C}[\rho]$. As we assume that the variables $(\rho)_{i \in \mathbb{N}}$ are algebraically independent, the invertible elements are those with a constant term in b in \mathbb{C}^* . \square

Proposition 3.1.2 Let E be a rank k $[\lambda]$ -primitive fresco and let $\lambda_1, \dots, \lambda_k$ its fundamental invariants. Let $e(\rho)$ be a family of generators of E depending polynomially on a family of algebraically independant variables $(\rho_i)_{i \in \mathbb{N}}$. Then there exists S_1, \dots, S_k in $\mathbb{C}[\rho][[b]]$ such that $S_j(\rho)[0] \equiv 1$ for each $j \in [1, k]$ and such that the annihilator of $e(\rho)$ in E is generated by the element of $\tilde{\mathcal{A}}$

$$P(\rho) := (a - \lambda_1.b).S_1(\rho)^{-1} \dots S_{k-1}(\rho)^{-1}.(a - \lambda_k.b).S_k(\rho)^{-1}.$$

PROOF. The key result to prove this proposition is the rank 1 case. In this case we may consider a standard generator e_1 of E which is a $\mathbb{C}[[b]]$ -basis of E and satisfies

$$(a - \lambda_1.b).e_1 = 0.$$

Then, by definition, we may write

$$e(\rho) = S_1(\rho).e_1$$

where S_1 is in $\mathbb{C}[\rho][[b]]$ is invertible in this algebra, so has a constant term in \mathbb{C}^* . Up to normalizing e_1 , we may assume that $S_1(\rho)[0] \equiv 1$ and then define

$$P(\rho) := (a - \lambda_1.b).S_1(\rho)^{-1}.$$

It clearly generates the annihilator of $e(\rho)$ for each ρ .

Assume now that the result is already proved for the rank $k - 1 \geq 1$. Then consider the family $[e(\rho)]$ in the quotient E/F_{k-1} where F_{k-1} is the rank $k - 1$ sub-module of E in its principal J-H. sequence. Remark first that $[e(\rho)]$ is a family of generators of E/F_{k-1} which depends polynomially on ρ . This is a trivial consequence of the fact that we may choose a $\mathbb{C}[[b]]$ -basis e_1, \dots, e_k in E such that e_1, \dots, e_{k-1} is a $\mathbb{C}[[b]]$ basis of F_{k-1} and e_k maps to a standard generator of $E/F_{k-1} \simeq E_{\lambda_k}$. Then the rank 1 case gives $S_k \in \mathbb{C}[\rho][[b]]$ with $S_k(\rho)[0] = 1$ and such that $(a - \lambda_k \cdot b) \cdot S_k(\rho)^{-1} \cdot e(\rho)$ is in F_{k-1} for each ρ . But then it is a family of generators of F_{k-1} which depends polynomially on ρ and the inductive assumption allows to conclude. \blacksquare

Fix now the fundamental invariants $\lambda_1, \dots, \lambda_k$ for a $[\lambda]$ -primitive fresco.

Definition 3.1.3 Consider now a complex valued function f defined on a subset \mathcal{F}_0 of the isomorphism classes $\mathcal{F}(\lambda_1, \dots, \lambda_k)$ of $[\lambda]$ -primitive frescos with fundamental invariants $\lambda_1, \dots, \lambda_k$. We shall say that f **depends polynomially on the isomorphism class** $[E] \in \mathcal{F}_0$ **of the fresco** E if the following condition is satisfied :

Let s be the collection of algebraically independant variables corresponding to the non constant coefficients of k elements S_1, \dots, S_k in $\mathbb{C}[[b]]$ satisfying $S_j(0) = 1 \quad \forall j \in [1, k]$ and consider for each value of s the rank k $[\lambda]$ -primitive fresco $E(s) := \tilde{A}/\tilde{A} \cdot P(s)$ where

$$P(s) := (a - \lambda_1 \cdot b) S_1(s)^{-1} \dots (a - \lambda_k \cdot b) \cdot S_k(s)^{-1}$$

where $S_1(s), \dots, S_k(s)$ correspond to the given values for s .

Then there exists a polynomial $F \in \mathbb{C}[s]$ such that for each value of s such that $[E(s)]$ is in \mathcal{F}_0 , the value of $F(s)$ is equal to $f([E(s)])$.

EXAMPLE. Let $(s_i)_{i \in \mathbb{N}^*}$ a family of algebraically independant variables and let $S(s) := 1 + \sum_{i=1}^{\infty} s_i \cdot b^i \in \mathbb{C}[s][[b]]$. Define $E(s) := \tilde{A}/\tilde{A} \cdot (a - \lambda_1 \cdot b) \cdot S(s)[b]^{-1} \cdot (a - \lambda_2 \cdot b)$ where $\lambda_1 > 1$ is rational and $\lambda_2 := \lambda_1 + p_1 - 1$ with $p_1 \in \mathbb{N}^*$. Define $\alpha(s) := s_{p_1}$. Then the number $\alpha(s)$ depends only of the isomorphism class of the fresco $E(s)$ and defines a function on $\mathcal{F}(\lambda_1, \lambda_2)$ which depends polynomially on $[E] \in \mathcal{F}(\lambda_1, \lambda_2)$. \square

3.2 The α -invariant.

The first proposition will be the induction step in the construction of the α -invariant.

Proposition 3.2.1 Fix $k \geq 2$. Denote $\mathcal{F}_0(\lambda_1, \dots, \lambda_k)$ the subset of $\mathcal{F}(\lambda_1, \dots, \lambda_k)$ of isomorphism class of rank k $[\lambda]$ -primitive fresco E with invariants $\lambda_1, \dots, \lambda_k$ such that F_{k-1} and E/F_1 are semi-simple where $(F_j)_{j \in [1, k]}$ is the principal J-H. sequence of E .

We assume that, for $k \geq 3$ we have :

- for rank $\leq k - 1$ frescos the α invariant is defined on the corresponding subset \mathcal{F}_0 ;
- for $[E] \in \mathcal{F}_0$ the number $\alpha(E)$ is zero if and only if E is semi-simple
- and that $[E] \mapsto \alpha(E)$ depends polynomially of $[E] \in \mathcal{F}_0$.

Fix E with $[E]$ in $\mathcal{F}_0(\lambda_1, \dots, \lambda_k)$. Let e be a generator of E such that $(a - \lambda_{k-1}.b).(a - \lambda_k.b).e$ lies in F_{k-2} . Then the sub-module G_{k-1} of E generated by $(a - (\lambda_{k-1} - 1).b).e$ is a normal rank $k - 1$ sub-module of E which is in $\mathcal{F}_0(\lambda_1, \dots, \lambda_{k-2}, \lambda_k + 1)$ and $\alpha(G_{k-1})$ is independant of the choice of such a generator e . Moreover, it defines a polynomial function on $\mathcal{F}_0(\lambda_1, \dots, \lambda_k)$.

PROOF. We shall prove first that if e is a generator of E such that $(a - \lambda_{k-1}.b).(a - \lambda_k.b).e$ lies in F_{k-2} then $G_{k-1} := \tilde{\mathcal{A}}.(a - (\lambda_{k-1} - 1).b).e$ is a normal rank $k - 1$ sub-module of E such that $[G_{k-1}]$ belongs to $\mathcal{F}_0(\lambda_1, \dots, \lambda_{k-2}, \lambda_k + 1)$. Using the identity in $\tilde{\mathcal{A}}$:

$$(a - \lambda_{k-1}.b).(a - \lambda_k.b) = (a - (\lambda_k + 1).b).(a - (\lambda_{k-1} - 1).b)$$

we see that G_{k-1} contains F_{k-2} , that $G_{k-1}/F_{k-2} \simeq E_{\lambda_k+1}$ and G_{k-1} admits

$$F_1 \subset \dots \subset F_{k-2} \subset G_{k-1}$$

as principal J-H. sequence. Then the fundamental invariants for G_{k-1} are equal to $\lambda_1, \dots, \lambda_{k-2}, \lambda_k + 1$, and the corank 1 term F_{k-2} is semi-simple. As G_{k-1}/F_1 is a sub-module of E/F_1 it is semi-simple and we have proved that

$$[G_{k-1}] \in \mathcal{F}_0(\lambda_1, \dots, \lambda_{k-2}, \lambda_k + 1).$$

CLAIM. If ε is another generator of E such that $(a - \lambda_{k-1}.b).(a - \lambda_k.b).\varepsilon$ is in F_{k-2} we have

$$\varepsilon = \rho.e + \sigma.b^{p_{k-1}-1}.(a - \lambda_k.b).e \quad \text{modulo } F_{k-2}$$

where (ρ, σ) is in $\mathbb{C}^* \times \mathbb{C}$.

Write $\varepsilon = U.e_k + V.e_{k-1}$ modulo F_{k-2} where $e_k := e$ and $e_{k-1} := (a - \lambda_k.b).e$ and where U, V are in $\mathbb{C}[[b]]$. Now

$$(a - \lambda_k.b).\varepsilon = b^2.U'.e_k + U.e_{k-1} + (\lambda_{k-1} - \lambda_k).b.V.e_{k-1} + b^2.V'.e_{k-1} \quad \text{modulo } F_{k-2}$$

and, as $(a - \lambda_{k-1}.b).b^2.U'.e_k \in F_{k-1}$ implies $U' = 0$, we have $U = \rho \in \mathbb{C}^*$, and we obtain

$$(a - \lambda_k.b).\varepsilon = [\rho + b^2.V' - (p_{k-1} - 1).b.V].e_{k-1} \quad \text{modulo } F_{k-2}.$$

If $T := \rho + b^2.V' - (p_{k-1} - 1).b.V$ we have

$$(a - \lambda_{k-1}.b).(a - \lambda_k.b).\varepsilon = b^2.T'.e_{k-1} \quad \text{modulo } F_{k-2}$$

and so T is a constant and equal to ρ . Then $V = \sigma.b^{p^{k-1}-1}$ for some complex number σ and our claim is proved.

For $\tau \in \mathbb{C}$ define

$$\varepsilon(\tau) := e_k + \tau.b^{p^{k-1}-1}.e_{k-1}$$

and let

$$G_{k-1}^\tau := \tilde{\mathcal{A}}.(a - (\lambda_{k-1} - 1).b).(e_k + \tau.b^{p^{k-1}-1}.e_{k-1}) = \tilde{\mathcal{A}}.(a - (\lambda_{k-1} - 1).b).\varepsilon(\tau).$$

Remark also that for any choice of ε such that $(a - \lambda_{k-1}.b).(a - \lambda_k.b).\varepsilon$ is in F_{k-2} the sub-module generated by ε is equal to G_{k-1}^τ for some $\tau \in \mathbb{C}$.

The generator $(a - (\lambda_k + 1).b).(a - (\lambda_{k-1} - 1).b).\varepsilon(\tau)$ of F_{k-2} depends polynomially on τ . Then the proposition 3.1.2 gives $Q(\tau)$ depending polynomially on τ which annihilates for each τ this generator. Then $Q(\tau).(a - (\lambda_k + 1).b)$ annihilates the generator $(a - (\lambda_{k-1} - 1).b).\varepsilon(\tau)$ of G_{k-1}^τ . Remark now that for any $\tau \in \mathbb{C}$ the fresco G_{k-1}^τ has its principal J-H. sequence given by $F_1 \subset \dots \subset F_{k-2} \subset G_{k-1}^\tau$. So G_{k-1}^τ/F_1 and F_{k-2} are semi-simple, and by our induction hypothesis, the number $\alpha(G_{k-1}^\tau)$ is well defined and $\tau \mapsto \alpha(G_{k-1}^\tau)$ is a polynomial in τ .

The key point will be now to show that either $\alpha(G_{k-1}^\tau)$ is identically zero or it never vanishes. This will prove that this number is independant of τ concluding the proof.

But if for some $\tau \in \mathbb{C}$ we have $\alpha(G_{k-1}^\tau) = 0$, then G_{k-1}^τ is semi-simple. Now the sub-module $F_{k-1} + G_{k-1}^\tau$ has rank k and is semi-simple. So its normalization is also semi-simple. But this normalization is E . So any sub-module of E is semi-simple and we have $\alpha(G_{k-1}^{\tau'}) = 0$ for any τ' .

This also shows that when $\alpha(G_{k-1}^\tau) \neq 0$ for some τ then it is not zero for any choice of τ . Now a polynomial never vanishing is a constant, and this implies that we may define $\alpha(E) := \alpha(G_{k-1}^\tau)$ for any value of τ .

To conclude the proof we have to show that, with this definition, the number $\alpha(E)$ depends polynomially on E . For that purpose it is enough, thanks to the previous computation, to produce a polynomial function $F \in \mathbb{C}[s]$ such that for each s with $E(s) \in \mathcal{F}_0(\lambda_1, \dots, \lambda_k)$ we have $F(s) = \alpha([E(s)])$.

The first remark is that we may assume that $S_k = 1$. Then, thanks to the previous computation and the induction on the rank, it is enough to show that we may also reduce to the case where $S_{k-1} = 1$ because in this case we have

$$\alpha(E[s]) = \alpha([G_{k-1}(s)])$$

where

$$G_{k-1}(s) = \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).S_1(s)^{-1} \dots S_{k-2}(s)^{-1}.(a - (\lambda_k + 1).b).$$

Denote $s_{k-1}^i, i \in \mathbb{N}^*$ the variables associated to the non constant coefficients of S_{k-1} . Looking to a generator of the form

$$\tilde{e} := e_k + X.e_{k-1}$$

with $X \in \mathbb{C}[s_{k-1}][[b]]$ such that $(a - \lambda_{k-1}.b).(a - \lambda_k.b).\tilde{e}$ is in F_{k-2} this leads to the equation

$$b^2.X' - (p_{k-1} - 1).b.X = 1 - S_{k-1}(s)$$

which has an unique solution $X \in \mathbb{C}[s_{k-1}][[b]]$ without term in $b^{p_{k-1}-1}$ as we assume $E(s)/F_1(s)$ semi-simple, hypothesis which implies that S_{k-1} has no $b^{p_{k-1}}$ term. With this generator \tilde{e} we have $(a - \lambda_{k-1}.b).(a - \lambda_k.b).\tilde{e}$ which is a generator of F_{k-2} which depends polynomially on s_{k-1}^i . Now we may consider F_{k-2} as a fix fresco (so we fix the coefficients in S_1, \dots, S_{k-2} and apply the proposition 3.1.2 to obtain a $Q(s) := (a - \lambda_1.b).S_1(s)^{-1} \dots (a - \lambda_{k-2}.b).S_{k-2}(s)^{-1}$ which annihilates $(a - \lambda_{k-1}.b).(a - \lambda_k.b).\tilde{e}$ in $E(s)$ and depends polynomially on s . We may conclude by the induction assumption because we know that we have

$$\alpha([E(s)]) = \alpha([G_{k-1}(s)])$$

where $G_{k-1}(s) = \tilde{\mathcal{A}}.(a - \lambda_{k-1} - 1).b.\tilde{e}(s)$ and because we know that the generator $(a - \lambda_{k-1} - 1).b.\tilde{e}(s)$ is annihilated by $Q(s).(a - (\lambda_k + 1).b)$ which depends polynomially on s . \blacksquare

Theorem 3.2.2 *Let $\lambda_1, \dots, \lambda_k$ be the fundamental invariants of a rank k $[\lambda]$ -primitive fresco such that $p_j \geq 1$ for each $j \in [1, k-1]$, and let $\mathcal{F}_0(\lambda_1, \dots, \lambda_k)$ the subset of $\mathcal{F}(\lambda_1, \dots, \lambda_k)$ of isomorphism classes of E satisfying the condition that F_{k-1} and E/F_1 are semi-simple, where $(F_j)_{j \in [1, k]}$ is the principal J-H. sequence of E . Then there exists a unique polynomial function*

$$\alpha : \mathcal{F}_0(\lambda_1, \dots, \lambda_k) \rightarrow \mathbb{C}$$

with the following properties :

- i) $\alpha([E]) = 0$ if and only if E is semi-simple.
- ii) When E has a generator e such that $(a - \lambda_{k-1}.b).(a - \lambda_k.b).e$ is in F_{k-2} , we have $\alpha(E) = \alpha(G_{k-1})$ where $G_{k-1} := \tilde{\mathcal{A}}.(a - \lambda_{k-1} - 1).b.e$.
- iii) When $E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(1 + x.b^{p_1})^{-1}.(a - \lambda_2.b)$ then $\alpha([E]) = x$.

PROOF. We define α for the rank 2 case as the coefficient of b^{p_1} in S_1 in any standard presentation $E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).S_1^{-1}.(a - \lambda_2.b)$. This is given by the easy computation which gives the rank 2 classification for $[\lambda]$ -primitive frescos, and fulfills condition iii). Then thanks to the previous proposition we may define $\alpha(E)$ for any rank $k \geq 3$ $[\lambda]$ -primitive fresco E such that F_{k-1} and E/F_1 are semi-simple via condition ii). The condition i) and the fact that α is a polynomial function are proved in the induction step given by the proposition. \blacksquare

Let me give in rank 3 a polynomial $F \in \mathbb{C}[s]$ such that we have $F(s) = \alpha([E(s)])$ for those s for which $E(s)$ is in $\mathcal{F}_0(\lambda_1, \lambda_2, \lambda_3)$.

Assume $p_1 \geq 1$ and $p_2 \geq 1$. Then the necessary and sufficient condition to be in $\mathcal{F}_0(\lambda_1, \lambda_2, \lambda_3)$ is $s_{p_1}^1 = s_{p_2}^2 = 0$.

Lemma 3.2.3 *Let $E := \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).S_1^{-1}.(a - \lambda_2.b).S_2^{-1}.(a - \lambda_3.b)$ such that $s_{p_1}^1 = s_{p_2}^2 = 0$; the complex number $\alpha(E)$ is the coefficient of $b^{p_1+p_2}$ in $p_2.V.S_1$ for $j = 1, 2$ where $V \in \mathbb{C}[[b]]$ is a solution of the differential equation*

$$b.V' = p_2.(V - S_2).$$

The proof is left to the reader as an exercise.

This gives the following formula for F :

$$F(s) = p_2 \cdot \sum_{j \neq p_2, j=0}^{p_1+p_2} s_{p_1+p_2-j}^1 \cdot \frac{s_j^2}{p_2 - j}.$$

Remark that for $S_2 = 1$ we find that $F(s)$ reduces to $s_{p_1+p_2}^1$. Using the commutation relation $(a - \lambda_2.b).(a - \lambda_3.b) = (a - (\lambda_3 + 1).b).(a - (\lambda_2 - 1).b)$ we find in this case that $E(s)$ has a normal rank 2 sub-module with fundamental invariants $\lambda_1, \lambda_3 + 1$ and parameter $s_{p_1+p_2}^1$ which is precisely $\alpha(E(s))$.

CONSEQUENCE. If we have rank 4 $[\lambda]$ -primitive fresco E with its principal J-H. sequence such that F_{j+1}/F_{j-1} is semi-simple for $j \in [1, 3]$, we may look inductively to $\alpha(F_3)$. If it is zero then we may look at $\alpha(E/F_1)$, and if it is zero we may look at $\alpha(E)$. If this is zero then E is semi-simple, and so, the inductive vanishing of all α -invariants of sub-quotients of the principal J-H. sequence give a necessary and sufficient condition for semi-simplicity of E .

This easily extends to any rank k $[\lambda]$ -primitive fresco. □

Now we shall show that in the previous situation when the α -invariant of E is not zero it is the parameter of any normal rank 2 sub-theme of E . Although such a rank 2 normal sub-theme is not unique (in general), its isomorphism class is uniquely determined from the fundamental invariants of E and from the α -invariant of E .

Proposition 3.2.4 *Let E be a rank $k \geq 2$ $[\lambda]$ -primitive fresco such that F_{k-1} and E/F_1 are semi-simple and with $\alpha(E) \neq 0$. Note $\lambda_1, \dots, \lambda_k$ the fundamental invariants of E and $p(E) := \sum_{j=1}^{p-1} p_j$. Then there exists at least one rank 2 normal sub-theme in E and each rank 2 normal sub-theme of E is isomorphic to*

$$\tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(1 + \alpha(E).b^{p(E)})^{-1}.(a - (\lambda_k + k - 2).b). \quad (\textcircled{a})$$

PROOF. The case $k = 2$ is clear, so we may assume that $k \geq 3$ and we shall prove the proposition by induction on k . So assume that the proposition is known for the rank $k - 1 \geq 2$ and let E be a rank k $[\lambda]$ -primitive fresco satisfying our assumptions. The fact that there exists a rank 2 normal sub-theme is consequence of the induction hypothesis as G^τ for any $\tau \in \mathbb{C}$ is normal rank $k - 1$ in E and satisfies again our assumptions.

Recall that the fundamental invariants of G^τ are μ_1, \dots, μ_{k-1} with $\mu_j = \lambda_j$ for $j \in [1, k - 2]$ and $\mu_{k-1} = \lambda_k + 1$. We have also $\alpha(G^\tau) = \alpha(E)$ for each τ , thanks to the proof of the theorem 3.2.2. As we have $p(G^\tau) = p(E)$ for each τ , and $\mu_{k-1} = \mu_1 + p(E) - 1$ the inductive hypothesis implies that any rank 2 normal sub-theme of any G^τ is isomorphic to $(@)$. Then, to complete the proof, it is enough to show that any rank 2 normal sub-theme of E is contained in some G^τ .

Let T be a rank 2 normal sub-theme of E . We shall first prove that its fundamental invariants are equal to $\lambda_1, \lambda_k + k - 2$. As E/F_1 is semi-simple, the image of T by the quotient map $E \rightarrow E/F_1$ has rank 1 and this implies that $F_1 \cap T$ is a rank 1 normal sub-module. Then this implies that F_1 is the unique normal rank 1 submodule of T , proving that $\lambda_1(T) = \lambda_1$. We shall prove now that $\lambda_2(T) = \lambda_k + k - 2$.

First remark that the uniqueness of the principal J-H. sequence of E implies the uniqueness of the quotient map $E \rightarrow E/F_{k-1} \simeq E_{\lambda_k}$ because any surjective map $q : E \rightarrow E_{\lambda_k}$ will produce principal a J-H. sequence for E by adjoining a principal J-H. sequence for $\text{Ker } q$. So a quotient E/H admits a surjective map on E_{λ_k} if (and only if) $H \subset F_{k-1}$. So H has to be semi-simple. Then the quotient E/H has no surjective map on E_{λ_k} .

The exact sequence of frescos

$$0 \rightarrow T/F_1 \rightarrow E/F_1 \rightarrow E/T \rightarrow 0$$

gives the equality in $\tilde{\mathcal{A}}$:

$$P_{T/F_1} \cdot P_{E/T} = P_{E/F_1} = (a - \lambda_2 \cdot b) \dots (a - \lambda_k \cdot b).$$

But as E/T is semi-simple and does not has a quotient isomorphic to E_{λ_k} this implies $T/F_1 \simeq E_{\lambda_k + k - 2}$, proving our claim.

So T is isomorphic to $\tilde{\mathcal{A}}/\tilde{\mathcal{A}} \cdot (a - \lambda_1 \cdot b) \cdot (1 + \beta \cdot b^{p(E)})^{-1} \cdot (a - (\lambda_k + k - 2) \cdot b)$ for some $\beta \in \mathbb{C}^*$. Let $x \in T$ be a generator of T which is annihilated by

$$(a - \lambda_1 \cdot b) \cdot (1 + \beta \cdot b^{p(E)})^{-1} \cdot (a - (\lambda_k + k - 2) \cdot b).$$

It satisfies

$$(a - (\lambda_k + k - 2) \cdot b) \cdot x \in F_1 \subset F_{k-2}. \quad (*)$$

We shall determine all $x \in E$ which satisfies the condition $(*)$ modulo F_{k-2} . Fix a generator $e := e_k$ of E such that $e_{k-1} := (a - \lambda_k \cdot b) \cdot e_k$ is in F_{k-1} and satisfies

$(a - \lambda_{k-1}.b).e_{k-1} \in F_{k-2}$. Then we look for $U, V \in \mathbb{C}[[b]]$ such that

$$x := U.e_k + V.e_{k-1} \quad \text{satisfies} \quad (a - (\lambda_k + k - 2).b).x \in F_{k-2}.$$

This leads to the equations

$$\begin{aligned} b.U' - (k - 2).b.U &= 0 \\ U + b^2.V' - (p_{k-1} + k - 3).V &= 0 \end{aligned}$$

and so we get

$$U = \rho.b^{k-2} \quad \text{and} \quad V = \frac{\rho}{p_{k-1}}.b^{k-3} + \sigma.b^{p_{k-1}+k-3}$$

Note that for $\rho = 0$ we would have $x \in F_{k-1}$ and this is not possible for the generator of a rank 2 theme as F_{k-1} is semi-simple. Then assuming $\rho \neq 0$ we may write

$$x = \frac{\rho.b^{k-3}}{p_{k-1}} \left[p_{k-1}.b.e_k + e_{k-1} + \frac{p_{k-1}.\sigma}{\rho}.b^{p_{k-1}}.e_{k-1} \right] \quad \text{modulo } F_{k-2}.$$

Now recall that the generator of G_τ is given by

$$(a - (\lambda_{k-1} - 1).b).\varepsilon(\tau)$$

where $\varepsilon(\tau) = e_k + \tau.b^{p_{k-1}-1}.e_{k-1}$. A simple computation gives

$$(a - (\lambda_{k-1} - 1).b).\varepsilon(\tau) = p_{k-1}.b.e_k + e_{k-1} + p_{k-1}.\tau.b^{p_{k-1}}.e_{k-1} \quad \text{modulo } F_{k-2}.$$

As we know that each G_τ contains F_{k-2} , we conclude that any such x is in G_τ for $\tau = \sigma/\rho$, concluding the proof. \blacksquare

Using duality we may deduce from this result the following corollary.

Corollary 3.2.5 *In the situation of the previous proposition E as a rank 2 quotient theme and any such rank 2 quotient theme is isomorphic to*

$$\tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - (\lambda_1 - k + 2).b).(1 + \beta(E).b^{p(E)})^{-1}.(a - \lambda_k.b)$$

where

$$\beta(E) := (-1)^k \frac{p_1.(p_1 + p_2) \dots (p_1 + \dots + p_{k-2})}{p_{k-1}.(p_{k-2} + p_{k-1}) \dots (p_2 + \dots + p_{k-1})} \alpha(E)$$

The proof is left as an exercise for the reader who may use the following remark.

REMARK. Let E be a rank 2 $[\lambda]$ -primitive theme with fundamental invariants λ_1, λ_2 and parameter $\alpha(E)$. For $\delta \in \mathbb{N}, \delta \gg 1$ $E^* \otimes E_\delta$ is a 2 $[1 - \lambda]$ -primitive theme with fundamental invariants $(\delta - \lambda_2, \delta - \lambda_1)$ and parameter $-\alpha(E)$.

Then in the situation of the proposition 3.2.4 $E^* \otimes E_\delta, \delta \gg 1$ satisfies again our assumptions and we have $\alpha(E^* \otimes E_\delta) = \beta(E)$.

4 The existence theorem for frescos.

The aim of this section is to prove the the following existence theorem for the fresco associated to a relative de Rham cohomology class :

Theorem 4.0.6 *Let X be a connected complex manifold of dimension $n+1$ where n is a natural integer, and let $f : X \rightarrow D$ be an non constant proper holomorphic function on an open disc D in \mathbb{C} with center 0 . Let us assume that df is nowhere vanishing outside of $X_0 := f^{-1}(0)$.*

Let ω be a $\mathcal{C}^\infty - (p+1)$ -differential form on X such that $d\omega = 0 = df \wedge \omega$. Denote by E the geometric $\tilde{\mathcal{A}}$ -module $\mathbb{H}^{p+1}(X, (\hat{K}^\bullet, d^\bullet))$ and $[\omega]$ the image of ω in $E/B(E)$. Then $\tilde{\mathcal{A}}.[\omega] \subset E/B(E)$ is a fresco.

Note that this result is an obvious consequence of the finiteness theorem 4.3.4 that we shall prove below. It gives the fact that E is naturally an $\tilde{\mathcal{A}}$ -module which is of finite type over the subalgebra $\mathbb{C}[[b]]$ of $\tilde{\mathcal{A}}$, and so its b -torsion $B(E)$ is a finite dimensional \mathbb{C} -vector space. Moreover, the finiteness theorem asserts that $E/B(E)$ is a geometric (a,b)-module.

4.1 Preliminaries.

Here we shall complete and precise the results of the section 2 of [B.II]. The situation we shall consider is the following : let X be a connected complex manifold of dimension $n+1$ and $f : X \rightarrow \mathbb{C}$ a non constant holomorphic function such that $\{x \in X / df = 0\} \subset f^{-1}(0)$. We introduce the following complexes of sheaves supported by $X_0 := f^{-1}(0)$

1. The formal completion "in f " $(\hat{\Omega}^\bullet, d^\bullet)$ of the usual holomorphic de Rham complex of X .
2. The sub-complexes $(\hat{K}^\bullet, d^\bullet)$ and $(\hat{I}^\bullet, d^\bullet)$ of $(\hat{\Omega}^\bullet, d^\bullet)$ where the subsheaves \hat{K}^p and \hat{I}^{p+1} are defined for each $p \in \mathbb{N}$ respectively as the kernel and the image of the map

$$\wedge df : \hat{\Omega}^p \rightarrow \hat{\Omega}^{p+1}$$

given par exterior multiplication by df . We have the exact sequence

$$0 \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow (\hat{\Omega}^\bullet, d^\bullet) \rightarrow (\hat{I}^\bullet, d^\bullet)[+1] \rightarrow 0. \quad (1)$$

Note that \hat{K}^0 and \hat{I}^0 are zero by definition.

3. The natural inclusions $\hat{I}^p \subset \hat{K}^p$ for all $p \geq 0$ are compatible with the différential d . This leads to an exact sequence of complexes

$$0 \rightarrow (\hat{I}^\bullet, d^\bullet) \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow ([\hat{K}/\hat{I}]^\bullet, d^\bullet) \rightarrow 0. \quad (2)$$

4. We have a natural inclusion $f^*(\hat{\Omega}_{\mathbb{C}}^1) \subset \hat{K}^1 \cap Ker d$, and this gives a sub-complex (with zero differential) of $(\hat{K}^\bullet, d^\bullet)$. As in [B.07], we shall consider also the complex $(\tilde{K}^\bullet, d^\bullet)$ quotient. So we have the exact sequence

$$0 \rightarrow f^*(\hat{\Omega}_{\mathbb{C}}^1) \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow (\tilde{K}^\bullet, d^\bullet) \rightarrow 0. \quad (3)$$

We do not make the assumption here that $f = 0$ is a reduced equation of X_0 , and we do not assume that $n \geq 2$, so the cohomology sheaf in degree 1 of the complex $(\hat{K}^\bullet, d^\bullet)$, which is equal to $\hat{K}^1 \cap Ker d$ does not coincide, in general with $f^*(\hat{\Omega}_{\mathbb{C}}^1)$. So the complex $(\tilde{K}^\bullet, d^\bullet)$ may have a non zero cohomology sheaf in degree 1.

Recall now that we have on the cohomology sheaves of the following complexes $(\hat{K}^\bullet, d^\bullet)$, $(\hat{I}^\bullet, d^\bullet)$, $([\hat{K}/\hat{I}]^\bullet, d^\bullet)$ and $f^*(\hat{\Omega}_{\mathbb{C}}^1)$, $(\tilde{K}^\bullet, d^\bullet)$ natural operations a and b with the relation $a.b - b.a = b^2$. They are defined in a naïve way by

$$a := \times f \quad \text{and} \quad b := \wedge df \circ d^{-1}.$$

The definition of a makes sens obviously. Let me precise the definition of b first in the case of $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ with $p \geq 2$: if $x \in \hat{K}^p \cap Ker d$ write $x = d\xi$ with $\xi \in \hat{\Omega}^{p-1}$ and let $b[x] := [df \wedge \xi]$. The reader will check easily that this makes sens. For $p = 1$ we shall choose $\xi \in \hat{\Omega}^0$ with the extra condition that $\xi = 0$ on the smooth part of X_0 (set theoretically). This is possible because the condition $df \wedge d\xi = 0$ allows such a choice: near a smooth point of X_0 we can choose coordinates such $f = x_0^k$ and the condition on ξ means independance of x_1, \dots, x_n . Then ξ has to be (set theoretically) locally constant on X_0 which is locally connected. So we may kill the value of such a ξ along X_0 .

The case of the complex $(\hat{I}^\bullet, d^\bullet)$ will be reduced to the previous one using the next lemma.

Lemma 4.1.1 *For each $p \geq 0$ there is a natural injective map*

$$\tilde{b} : \mathcal{H}^p(\hat{K}^\bullet, d^\bullet) \rightarrow \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$$

which satisfies the relation $a.\tilde{b} = \tilde{b}.(b + a)$. For $p \neq 1$ this map is bijective.

PROOF. Let $x \in \hat{K}^p \cap Ker d$ and write $x = d\xi$ where $\xi \in \hat{\Omega}^{p-1}$ (with $\xi = 0$ on X_0 if $p = 1$), and set $\tilde{b}([x]) := [df \wedge \xi] \in \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$. This is independant on the choice of ξ because, for $p \geq 2$, adding $d\eta$ to ξ does not modify the result as $[df \wedge d\eta] = 0$. For $p = 1$ remark that our choice of ξ is unique.

This is also independant of the the choice of x in $[x] \in \mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ because adding $\theta \in \hat{K}^{p-1}$ to ξ does not change $[df \wedge \xi]$.

Assume $\tilde{b}([x]) = 0$ in $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$; this means that we may find $\alpha \in \hat{\Omega}^{p-2}$ such $df \wedge \xi = df \wedge d\alpha$. But then, $\xi - d\alpha$ lies in \hat{K}^{p-1} and $x = d(\xi - d\alpha)$ shows that $[x] = 0$. So \tilde{b} is injective.

Assume now $p \geq 2$. If $df \wedge \eta$ is in $\hat{I}^p \cap Ker d$, then $df \wedge d\eta = 0$ and $y := d\eta$ lies

in $\hat{K}^p \cap \text{Ker } d$ and defines a class $[y] \in \mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ whose image by \tilde{b} is $[df \wedge \eta]$. This shows the surjectivity of \tilde{b} for $p \geq 2$.

For $p = 1$ the map \tilde{b} is not surjective (see the remark below).

To finish the proof let us to compute $\tilde{b}(a[x] + b[x])$. Writing again $x = d\xi$, we get

$$a[x] + b[x] = [f.d\xi + df \wedge \xi] = [d(f.\xi)]$$

and so

$$\tilde{b}(a[x] + b[x]) = [df \wedge f.\xi] = a.\tilde{b}([x])$$

which concludes the proof. ■

Denote by $i : (\hat{I}^\bullet, d^\bullet) \rightarrow (\hat{K}^\bullet, d^\bullet)$ the natural inclusion and define the action of b on $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ by $b := \tilde{b} \circ \mathcal{H}^p(i)$. As i is a -linear, we deduce the relation $a.b - b.a = b^2$ on $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ from the relation of the previous lemma.

The action of a on the complex $([\hat{K}/\hat{I}]^\bullet, d^\bullet)$ is obvious and the action of b is zero.

The action of a and b on $f^*(\hat{\Omega}_{\mathbb{C}}^1) \simeq E_1 \otimes \mathbb{C}_{X_0}$ are the obvious one, where E_1 is the rank 1 (a, b) -module with generator e_1 satisfying $a.e_1 = b.e_1$ (or, equivalently, $E_1 := \mathbb{C}[[z]]$ with $a := \times z$, $b := \int_0^z$ and $e_1 := 1$).

Remark that the natural inclusion $f^*(\hat{\Omega}_{\mathbb{C}}^1) \hookrightarrow (\hat{K}^\bullet, d^\bullet)$ is compatible with the actions of a and b . The actions of a and b on $\mathcal{H}^1(\hat{K}^\bullet, d^\bullet)$ are simply induced by the corresponding actions on $\mathcal{H}^1(\hat{K}^\bullet, d^\bullet)$.

REMARK. The exact sequence of complexes (1) induces for any $p \geq 2$ a bijection

$$\partial^p : \mathcal{H}^p(\hat{I}^\bullet, d^\bullet) \rightarrow \mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$$

and a short exact sequence

$$0 \rightarrow \mathbb{C}_{X_0} \rightarrow \mathcal{H}^1(\hat{I}^\bullet, d^\bullet) \xrightarrow{\partial^1} \mathcal{H}^1(\hat{K}^\bullet, d^\bullet) \rightarrow 0 \quad (\textcircled{a})$$

because of the de Rham lemma. Let us check that for $p \geq 2$ we have $\partial^p = (\tilde{b})^{-1}$ and that for $p = 1$ we have $\partial^1 \circ \tilde{b} = \text{Id}$. If $x = d\xi \in \hat{K}^p \cap \text{Ker } d$ then $\tilde{b}([x]) = [df \wedge \xi]$ and $\partial^p[df \wedge \xi] = [d\xi]$. So $\partial^p \circ \tilde{b} = \text{Id} \quad \forall p \geq 0$. For $p \geq 2$ and $df \wedge \alpha \in \hat{I}^p \cap \text{Ker } d$ we have $\partial^p[df \wedge \alpha] = [d\alpha]$ and $\tilde{b}[d\alpha] = [df \wedge \alpha]$, so $\tilde{b} \circ \partial^p = \text{Id}$. For $p = 1$ we have $\tilde{b}[d\alpha] = [df \wedge (\alpha - \alpha_0)]$ where $\alpha_0 \in \mathbb{C}$ is such that $\alpha|_{X_0} = \alpha_0$. This shows that in degree 1 \tilde{b} gives a canonical splitting of the exact sequence (\textcircled{a}) .

4.2 $\tilde{\mathcal{A}}$ -structures.

Let us consider now the \mathbb{C} -algebra

$$\tilde{\mathcal{A}} := \left\{ \sum_{\nu \geq 0} P_\nu(a).b^\nu \right\}$$

where $P_\nu \in \mathbb{C}[z]$, and the commutation relation $a.b - b.a = b^2$, assuming that left and right multiplications by a are continuous for the b -adic topology of $\tilde{\mathcal{A}}$. Define the following complexes of sheaves of left $\tilde{\mathcal{A}}$ -modules on X :

$$\begin{aligned}
& (\Omega'^\bullet[[b]], D^\bullet) \quad \text{and} \quad (\Omega''^\bullet[[b]], D^\bullet) \quad \text{where} \tag{4} \\
\Omega'^\bullet[[b]] & := \sum_{j=0}^{+\infty} b^j . \omega_j \quad \text{with} \quad \omega_0 \in \hat{K}^p \\
\Omega''^\bullet[[b]] & := \sum_{j=0}^{+\infty} b^j . \omega_j \quad \text{with} \quad \omega_0 \in \hat{I}^p \\
D\left(\sum_{j=0}^{+\infty} b^j . \omega_j\right) & = \sum_{j=0}^{+\infty} b^j . (d\omega_j - df \wedge \omega_{j+1}) \\
a . \sum_{j=0}^{+\infty} b^j . \omega_j & = \sum_{j=0}^{+\infty} b^j . (f . \omega_j + (j-1) . \omega_{j-1}) \quad \text{with the convention} \quad \omega_{-1} = 0 \\
b . \sum_{j=0}^{+\infty} b^j . \omega_j & = \sum_{j=1}^{+\infty} b^j . \omega_{j-1}
\end{aligned}$$

It is easy to check that D is $\tilde{\mathcal{A}}$ -linear and that $D^2 = 0$. We have a natural inclusion of complexes of left $\tilde{\mathcal{A}}$ -modules

$$\tilde{i} : (\Omega''^\bullet[[b]], D^\bullet) \rightarrow (\Omega'^\bullet[[b]], D^\bullet).$$

Remark that we have natural morphisms of complexes

$$\begin{aligned}
u & : (\hat{I}^\bullet, d^\bullet) \rightarrow (\Omega''^\bullet[[b]], D^\bullet) \\
v & : (\hat{K}^\bullet, d^\bullet) \rightarrow (\Omega'^\bullet[[b]], D^\bullet)
\end{aligned}$$

and that these morphisms are compatible with i . More precisely, this means that we have the commutative diagram of complexes

$$\begin{array}{ccc}
(\hat{I}^\bullet, d^\bullet) & \xrightarrow{u} & (\Omega''^\bullet[[b]], D^\bullet) \\
\downarrow i & & \downarrow \tilde{i} \\
(\hat{K}^\bullet, d^\bullet) & \xrightarrow{v} & (\Omega'^\bullet[[b]], D^\bullet)
\end{array}$$

The following theorem is a variant of theorem 2.2.1. of [B.II].

Theorem 4.2.1 *Let X be a connected complex manifold of dimension $n+1$ and $f : X \rightarrow \mathbb{C}$ a non constant holomorphic function such that*

$$\{x \in X / df = 0\} \subset f^{-1}(0).$$

Then the morphisms of complexes u and v introduced above are quasi-isomorphisms. Moreover, the isomorphisms that they induce on the cohomology sheaves of these complexes are compatible with the actions of a and b .

This theorem builds a natural structure of left $\tilde{\mathcal{A}}$ -modules on each of the complex $(\hat{K}^\bullet, d^\bullet), (\hat{I}^\bullet, d^\bullet), ([\hat{K}/\hat{I}]^\bullet, d^\bullet)$ and $f^*(\hat{\Omega}_{\mathbb{C}}^1), (\tilde{K}^\bullet, d^\bullet)$ in the derived category of bounded complexes of sheaves of \mathbb{C} -vector spaces on X .

Moreover the short exact sequences

$$\begin{aligned} 0 \rightarrow (\hat{I}^\bullet, d^\bullet) \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow ([\hat{K}/\hat{I}]^\bullet, d^\bullet) \rightarrow 0 \\ 0 \rightarrow f^*(\hat{\Omega}_{\mathbb{C}}^1) \rightarrow (\hat{K}^\bullet, d^\bullet), (\hat{I}^\bullet, d^\bullet) \rightarrow (\tilde{K}^\bullet, d^\bullet) \rightarrow 0 \end{aligned}$$

are equivalent to short exact sequences of complexes of left $\tilde{\mathcal{A}}$ -modules in the derived category.

PROOF. We have to prove that for any $p \geq 0$ the maps $\mathcal{H}^p(u)$ and $\mathcal{H}^p(v)$ are bijective and compatible with the actions of a and b . The case of $\mathcal{H}^p(v)$ is handled (at least for $n \geq 2$ and f reduced) in prop. 2.3.1. of [B.II]. To seek for completeness and for the convenience of the reader we shall treat here the case of $\mathcal{H}^p(u)$.

First we shall prove the injectivity of $\mathcal{H}^p(u)$. Let $\alpha = df \wedge \beta \in \hat{I}^p \cap Ker d$ and assume that we can find $U = \sum_{j=0}^{+\infty} b^j . u_j \in \Omega''^{p-1}[[b]]$ with $\alpha = DU$. Then we have the following relations

$$u_0 = df \wedge \zeta, \quad \alpha = du_0 - df \wedge u_1 \quad \text{and} \quad du_j = df \wedge u_{j+1} \quad \forall j \geq 1.$$

For $j \geq 1$ we have $[du_j] = b[du_{j+1}]$ in $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$; using corollary 2.2. of [B.II] which gives the b -separation of $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$, this implies $[du_j] = 0, \forall j \geq 1$ in $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$. For instance we can find $\beta_1 \in \hat{K}^{p-1}$ such that $du_1 = d\beta_1$. Now, by de Rham, we can write $u_1 = \beta_1 + d\xi_1$ for $p \geq 2$, where $\xi_1 \in \hat{\Omega}^{p-2}$. Then we conclude that $\alpha = -df \wedge d(\xi_1 + \zeta)$ and $[\alpha] = 0$ in $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$.

For $p = 1$ we have $u_1 = 0$ and $[\alpha] = [-df \wedge d\xi_1] = 0$ in $\mathcal{H}^1(\hat{I}^\bullet, d^\bullet)$.

We shall show now that the image of $\mathcal{H}^p(u)$ is dense in $\mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$ for its b -adic topology. Let $\Omega := \sum_{j=0}^{+\infty} b^j . \omega_j \in \Omega''^p[[b]]$ such that $D\Omega = 0$. The following relations holds $d\omega_j = df \wedge \omega_{j+1} \quad \forall j \geq 0$ and $\omega_0 \in \hat{I}^p$. The corollary 2.2. of [B.II] again allows to find $\beta_j \in \hat{K}^{p-1}$ for any $j \geq 0$ such that $d\omega_j = d\beta_j$. Fix $N \in \mathbb{N}^*$. We have

$$D\left(\sum_{j=0}^N b^j . \omega_j\right) = b^N . d\omega_N = D(b^N . \beta_N)$$

and $\Omega_N := \sum_{j=0}^N b^j . \omega_j - b^N . \beta_N$ is D -closed and in $\Omega''^p[[b]]$. And we have $\Omega - \Omega_N \in b^N . \mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$, so the sequence $(\Omega_N)_{N \geq 1}$ converges to Ω in $\mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$ for its b -adic topology. Let us show that each Ω_N is in the image of $\mathcal{H}^p(u)$.

Write $\Omega_N := \sum_{j=0}^N b^j . \omega_j$. The condition $D\Omega_N = 0$ implies $d\omega_N = 0$ and $d\omega_{N-1} = df \wedge \omega_N = 0$. If we write $w_N = dv_N$ we obtain $d(w_{N-1} + df \wedge v_N) = 0$ and $\Omega_N - D(b^N . v_N)$ is of degree $N - 1$ in b . For $N = 1$ we are left with $w_0 + b . w_1 - (-df \wedge v_1 + b . dv_1) = w_0 + df \wedge v_1$ which is in $\hat{I}^p \cap Ker d$ because

$$dw_0 = df \wedge dv_1.$$

To conclude it is enough to know the following two facts

- i) The fact that $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ is complete for its b -adic topology.
- ii) The fact that $Im(\mathcal{H}^p(u)) \cap b^N \cdot \mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet) \subset Im(\mathcal{H}^p(u) \circ b^N) \quad \forall N \geq 1.$

Let us first conclude the proof of the surjectivity of $\mathcal{H}^p(u)$ assuming i) and ii). For any $[\Omega] \in \mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$ we know that there exists a sequence $(\alpha_N)_{N \geq 1}$ in $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ with $\Omega - \mathcal{H}^p(u)(\alpha_N) \in b^N \cdot \mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$. Now the property ii) implies that we may choose the sequence $(\alpha_N)_{N \geq 1}$ such that $[\alpha_{N+1}] - [\alpha_N]$ lies in $b^N \cdot \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$. So the property i) implies that the Cauchy sequence $([\alpha_N])_{N \geq 1}$ converges to $[\alpha] \in \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$. Then the continuity of $\mathcal{H}^p(u)$ for the b -adic topologies coming from its b -linearity, implies $\mathcal{H}^p(u)([\alpha]) = [\Omega]$.

The compatibility with a and b of the maps $\mathcal{H}^p(u)$ and $\mathcal{H}^p(v)$ is an easy exercise.

Let us now prove properties i) and ii).

The property i) is a direct consequence of the completion of $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ for its b -adic topology given by the corollary 2.2. of [B.II] and the b -linear isomorphism \tilde{b} between $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ and $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ constructed in the lemma 2.1.1. above.

To prove ii) let $\alpha \in \hat{I}^p \cap Ker d$ and $N \geq 1$ such that

$$\alpha = b^N \cdot \Omega + DU$$

where $\Omega \in \Omega''^p[[b]]$ satisfies $D\Omega = 0$ and where $U \in \Omega''^{p-1}[[b]]$. With obvious notations we have

$$\begin{aligned} \alpha &= du_0 - df \wedge u_1 \\ &\dots \\ 0 &= du_j - df \wedge u_{j+1} \quad \forall j \in [1, N-1] \\ &\dots \\ 0 &= \omega_0 + du_N - df \wedge u_{N+1} \end{aligned}$$

which implies $D(u_0 + b \cdot u_1 + \dots + b^N \cdot u_N) = \alpha + b^N \cdot du_N$ and the fact that du_N lies in $\hat{I}^p \cap Ker d$. So we conclude that $[\alpha] + b^N \cdot [du_N]$ is in the kernel of $\mathcal{H}^p(u)$ which is 0. Then $[\alpha] \in b^N \cdot \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$. ■

REMARK. The map

$$\beta : (\Omega'[[b]]^\bullet, D^\bullet) \rightarrow (\Omega''[[b]]^\bullet, D^\bullet)$$

defined by $\beta(\Omega) = b \cdot \Omega$ commutes to the differentials and with the action of b . It induces the isomorphism \tilde{b} of the lemma 4.1.1 on the cohomology sheaves. So it is a quasi-isomorphism of complexes of $\mathbb{C}[[b]]$ -modules.

To prove this fact, it is enough to verify that the diagram

$$\begin{array}{ccc} (\hat{K}^\bullet, d^\bullet) & \xrightarrow{v} & (\Omega'[[b]]^\bullet, D^\bullet) \\ \downarrow \tilde{b} & & \downarrow \beta \\ (\hat{I}^\bullet, d^\bullet) & \xrightarrow{u} & (\Omega''[[b]]^\bullet, D^\bullet) \end{array}$$

induces commutative diagrams on the cohomology sheaves.

But this is clear because if $\alpha = d\xi$ lies in $\hat{K}^p \cap \text{Ker } d$ we have $D(b.\xi) = b.d\xi - df \wedge \xi$ so $\mathcal{H}^p(\beta) \circ \mathcal{H}^p(v)([\alpha]) = \mathcal{H}^p(u) \circ \mathcal{H}^p(\tilde{b})([\alpha])$ in $\mathcal{H}^p(\Omega''[[b]]^\bullet, D^\bullet)$. \blacksquare

4.3 The finiteness theorem.

Let us recall some basic definitions on the left modules over the algebra $\tilde{\mathcal{A}}$.

Now let E be any left $\tilde{\mathcal{A}}$ -module, and define $B(E)$ as the b -torsion of E . that is to say

$$B(E) := \{x \in E / \exists N \quad b^N .x = 0\}.$$

Define $A(E)$ as the a -torsion of E and

$$\hat{A}(E) := \{x \in E / \mathbb{C}[[b]].x \subset A(E)\}.$$

Remark that $B(E)$ and $\hat{A}(E)$ are sub- $\tilde{\mathcal{A}}$ -modules of E but that $A(E)$ is not stable by b .

Definition 4.3.1 A left $\tilde{\mathcal{A}}$ -module E is called **small** when the following conditions hold

1. E is a finite type $\mathbb{C}[[b]]$ -module ;
2. $B(E) \subset \hat{A}(E)$;
3. $\exists N / a^N .\hat{A}(E) = 0$;

Recall that for E small we have always the equality $B(E) = \hat{A}(E)$ (see [B.I] lemme 2.1.2) and that this complex vector space is finite dimensional. The quotient $E/B(E)$ is an (a,b) -module called **the associate (a,b) -module** to E .

Conversely, any left $\tilde{\mathcal{A}}$ -module E such that $B(E)$ is a finite dimensional \mathbb{C} -vector space and such that $E/B(E)$ is an (a,b) -module is small.

The following easy criterium to be small will be used later :

Lemma 4.3.2 A left $\tilde{\mathcal{A}}$ -module E is small if and only if the following conditions hold :

1. $\exists N / a^N .\hat{A}(E) = 0$;
2. $B(E) \subset \hat{A}(E)$;

3. $\cap_{m \geq 0} b^m \cdot E \subset \hat{A}(E)$;

4. $\text{Ker } b$ and $\text{Coker } b$ are finite dimensional complex vector spaces.

As the condition 3 in the previous lemma has been omitted in [B.II] (but this does not affect this article because this lemma was used only in a case where this condition 3 was satisfied, thanks to proposition 2.2.1. of *loc. cit.*), we shall give the (easy) proof.

PROOF. First the conditions 1 to 4 are obviously necessary. Conversely, assume that E satisfies these four conditions. Then condition 2 implies that the action of b on $\hat{A}(E)/B(E)$ is injective. But the condition 1 implies that $b^{2N} = 0$ on $\hat{A}(E)$ (see [B.I]). So we conclude that $\hat{A}(E) = B(E) \subset \text{Ker } b^{2N}$ which is a finite dimensional complex vector space using condition 4 and an easy induction. Now $E/B(E)$ is a $\mathbb{C}[[b]]$ -module which is separated for its b -adic topology. The finiteness of $\text{Coker } b$ now shows that it is a free finite type $\mathbb{C}[[b]]$ -module concluding the proof. \blacksquare

Definition 4.3.3 We shall say that a left $\tilde{\mathcal{A}}$ -module E is **geometric** when E is small and when its associated (a,b) -module $E/B(E)$ is geometric.

The main result of this section is the following theorem, which shows that the Gauss-Manin connection of a proper holomorphic function produces geometric $\tilde{\mathcal{A}}$ -modules associated to vanishing cycles and nearby cycles.

Theorem 4.3.4 Let X be a connected complex manifold of dimension $n+1$ where n is a natural integer, and let $f : X \rightarrow D$ be a non constant proper holomorphic function on an open disc D in \mathbb{C} with center 0. Let us assume that df is nowhere vanishing outside of $X_0 := f^{-1}(0)$.

Then the $\tilde{\mathcal{A}}$ -modules

$$\mathbb{H}^j(X, (\hat{K}^\bullet, d^\bullet)) \quad \text{and} \quad \mathbb{H}^j(X, (\hat{I}^\bullet, d^\bullet))$$

are geometric for any $j \geq 0$.

In the proof we shall use the \mathcal{C}^∞ version of the complex $(\hat{K}^\bullet, d^\bullet)$. We define K_∞^p as the kernel of $\wedge df : \mathcal{C}^{\infty,p} \rightarrow \mathcal{C}^{\infty,p+1}$ where $\mathcal{C}^{\infty,j}$ denote the sheaf of \mathcal{C}^∞ -forms on X of degree p , let \hat{K}_∞^p be the f -completion and $(\hat{K}_\infty^\bullet, d^\bullet)$ the corresponding de Rham complex.

The next lemma is proved in [B.II] (lemma 6.1.1.)

Lemma 4.3.5 The natural inclusion

$$(\hat{K}^\bullet, d^\bullet) \hookrightarrow (\hat{K}_\infty^\bullet, d^\bullet)$$

induce a quasi-isomorphism.

REMARK. As the sheaves \hat{K}_∞^\bullet are fine, we have a natural isomorphism

$$\mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet)) \simeq H^p(\Gamma(X, \hat{K}_\infty^\bullet), d^\bullet).$$

Let us denote by X_1 the generic fiber of f . Then X_1 is a smooth compact complex manifold of dimension n and the restriction of f to $f^{-1}(D^*)$ is a locally trivial \mathcal{C}^∞ bundle with typical fiber X_1 on $D^* = D \setminus \{0\}$, if the disc D is small enough around 0. Fix now $\gamma \in H_p(X_1, \mathbb{C})$ and let $(\gamma_s)_{s \in D^*}$ the corresponding multivalued horizontal family of p -cycles $\gamma_s \in H_p(X_s, \mathbb{C})$. Then, for $\omega \in \Gamma(X, \hat{K}_\infty^p \cap Ker d)$, define the multivalued holomorphic function

$$F_\omega(s) := \int_{\gamma_s} \frac{\omega}{df}.$$

Let now

$$\Xi := \bigoplus_{\alpha \in \mathbb{Q} \cap]-1, 0], j \in [0, n]} \mathbb{C}[[s]] \cdot s^\alpha \cdot \frac{(Log s)^j}{j!}.$$

This is an $\tilde{\mathcal{A}}$ -modules with a acting as multiplication by s and b as the primitive in s without constant. Now if \hat{F}_ω is the asymptotic expansion at 0 of F_ω , it is an element in Ξ , and we obtain in this way an $\tilde{\mathcal{A}}$ -linear map

$$Int : \mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet)) \rightarrow H^p(X_1, \mathbb{C}) \otimes_{\mathbb{C}} \Xi.$$

To simplify notations, let $E := \mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet))$. Now using Grothendieck theorem [G.65], there exists $N \in \mathbb{N}$ such that $Int(\omega) \equiv 0$, implies $a^N \cdot [\omega] = 0$ in E . As the converse is clear we conclude that $\hat{A}(E) = Ker(Int)$. It is also clear that $B(E) \subset Ker(Int)$ because Ξ has no b -torsion. So we conclude that E satisfies properties 1 and 2 of the lemma 4.3.2. The property 3 is also true because of the regularity of the Gauss-Manin connection of f .

END OF THE PROOF OF THEOREM 4.3.4. To show that $E := \mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet))$ is small, it is enough to prove that E satisfies the condition 4 of the lemma 4.3.2. Consider now the long exact sequence of hypercohomology of the exact sequence of complexes

$$0 \rightarrow (\hat{I}^\bullet, d^\bullet) \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow ([\hat{K}/\hat{I}]^\bullet, d^\bullet) \rightarrow 0.$$

It contains the exact sequence

$$\mathbb{H}^{p-1}(X, ([\hat{K}/\hat{I}]^\bullet, d^\bullet)) \rightarrow \mathbb{H}^p(X, (\hat{I}^\bullet, d^\bullet)) \xrightarrow{\mathbb{H}^p(i)} \mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet)) \rightarrow \mathbb{H}^p(X, ([\hat{K}/\hat{I}]^\bullet, d^\bullet))$$

and we know that b is induced on the complex of $\tilde{\mathcal{A}}$ -modules quasi-isomorphic to $(\hat{K}^\bullet, d^\bullet)$ by the composition $i \circ \tilde{b}$ where \tilde{b} is a quasi-isomorphism of complexes of $\mathbb{C}[[b]]$ -modules. This implies that the kernel and the cokernel of $\mathbb{H}^p(i)$ are isomorphic (as \mathbb{C} -vector spaces) to $Ker b$ and $Coker b$ respectively. Now to prove that E satisfies condition 4 of the lemma 4.3.2 it is enough to prove finite dimensionality for the vector spaces $\mathbb{H}^j(X, ([\hat{K}/\hat{I}]^\bullet, d^\bullet))$ for all $j \geq 0$.

But the sheaves $[\hat{K}/\hat{I}]^j \simeq [Ker\,df/Im\,df]^j$ are coherent on X and supported in X_0 . The spectral sequence

$$E_2^{p,q} := H^q(H^p(X, [\hat{K}/\hat{I}]^\bullet), d^\bullet)$$

which converges to $\mathbb{H}^j(X, ([\hat{K}/\hat{I}]^\bullet, d^\bullet))$, is a bounded complex of finite dimensional vector spaces by Cartan-Serre. This gives the desired finite dimensionality.

To conclude the proof, we want to show that $E/B(E)$ is geometric. But this is an easy consequence of the regularity of the Gauss-Manin connexion of f and of the Monodromy theorem, which are already incoded in the definition of Ξ : the injectivity on $E/B(E)$ of the $\tilde{\mathcal{A}}$ -linear map Int implies that $E/B(E)$ is geometric.

Remark now that the piece of exact sequence above gives also the fact that $\mathbb{H}^p(X, (\hat{I}^\bullet, d^\bullet))$ is geometric, because it is an exact sequence of $\tilde{\mathcal{A}}$ -modules. \blacksquare

REMARK. It is easy to see that the properness assumption on f is only used for two purposes :

– To have a (global) \mathcal{C}^∞ Milnor fibration on a small punctured disc around 0, with a finite dimensional cohomology for the Milnor fiber.

– To have compactness of the singular set $\{df = 0\}$, which contains the supports of the coherent sheaves $(Ker\,df/Im\,df)^i$.

This allows to give with the same proof an analogous finiteness result in many other situations.

Bibliography

- [Br.70] Brieskorn, E. *Die Monodromie der Isolierten Singularitäten von Hyperflächen*, Manuscripta Math. 2 (1970), p. 103-161.
- [B.84] Barlet, D. *Contribution du cup-produit de la fibre de Milnor aux pôles de $|f|^{2\lambda}$* , Ann. Inst. Fourier (Grenoble) t. 34, fasc. 4 (1984), p. 75-107.
- [B.93] Barlet, D. *Théorie des (a,b)-modules I*, in Complex Analysis and Geometry, Plenum Press, (1993), p. 1-43.
- [B.95] Barlet, D. *Théorie des (a,b)-modules II. Extensions*, in Complex Analysis and Geometry, Pitman Research Notes in Mathematics Series 366 Longman (1997), p. 19-59.
- [B.05] Barlet, D. *Module de Brieskorn et forme hermitiennes pour une singularité isolée d'hypersurface*, revue de l'Inst. E. Cartan (Nancy) 18 (2005), p. 19-46.
- [B.I] Barlet, D. *Sur certaines singularités non isolées d'hypersurfaces I*, Bull. Soc. math. France 134 (2), (2006), p.173-200.

- [B.II] Barlet, D. *Sur certaines singularités d'hypersurfaces II*, J. Alg. Geom. 17 (2008), p. 199-254.
- [B.08] Barlet, D. *Two finiteness theorem for regular (a,b) -modules*, preprint Institut E. Cartan (Nancy) (2008) n^o5, p. 1-38, arXiv:0801.4320 (math. AG and math. CV)
- [B.09] Barlet, D. *Périodes évanescentes et (a,b) -modules monogènes*, Bollettino U.M.I. (9) II (2009), p. 651-697.
- [B.III] Barlet, D. *Sur les fonctions à lieu singulier de dimension 1*, Bull. Soc. math. France 137 (4), (2009), p. 587-612.
- [B.10] Barlet, D. *Le thème d'une période évanescence*, preprint Institut E. Cartan (Nancy) (2009) n^o33, p. 1-57.
- [B.-S. 04] Barlet, D. et Saito, M. *Brieskorn modules and Gauss-Manin systems for non isolated hypersurface singularities*, J. Lond. Math. Soc. (2) 76 (2007) n^o1 p. 211-224.
- [Bj.93] Björk, J-E, *Analytic D-modules and applications*, Kluwer Academic publishers (1993).
- [G.65] Grothendieck, A. *On the de Rham cohomology of algebraic varieties* Publ. Math. IHES 29 (1966), p. 93-101.
- [K.76] Kashiwara, M. *b-function and holonomic systems*, Inv. Math. 38 (1976) p. 33-53.
- [M.74] Malgrange, B. *Intégrale asymptotique et monodromie*, Ann. Sc. Ec. Norm. Sup. 7 (1974), p. 405-430.
- [M.75] Malgrange, B. *Le polynôme de Bernstein d'une singularité isolée*, in Lect. Notes in Math. 459, Springer (1975), p. 98-119.
- [S.89] Saito, M. *On the structure of Brieskorn lattices*, Ann. Inst. Fourier 39 (1989), p. 27-72.