



**HAL**  
open science

## Continuous mappings between spaces of arcs

Goulwen Fichou, Masahiro Shiota

► **To cite this version:**

Goulwen Fichou, Masahiro Shiota. Continuous mappings between spaces of arcs. 2012. hal-00659415v2

**HAL Id: hal-00659415**

**<https://hal.science/hal-00659415v2>**

Preprint submitted on 13 Dec 2012 (v2), last revised 29 May 2013 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# CONTINUOUS MAPPINGS BETWEEN SPACES OF ARCS

GOULWEN FICHOU AND MASAHIRO SHIOTA

ABSTRACT. A blow-analytic homeomorphism is an arc-analytic subanalytic homeomorphism, and therefore it induces a bijective mapping between spaces of analytic arcs. We tackle the question of the continuity of this induced mapping between the spaces of arcs, giving a positive and a negative answer depending of the topology involved. We generalise the result to spaces of definable arcs in the context of o-minimal structures, obtaining notably a uniform continuity property.

## INTRODUCTION

The blow-analytic equivalence between real analytic function germs [12] is an interesting counterpart in the real setting to the topological equivalence between complex analytic function germs. For  $f, g : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}, 0)$  analytic function germs, we say that  $f$  and  $g$  are blow-analytically equivalent if there exists a blow-analytic homeomorphism germ  $\phi : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^n, 0)$  such that  $f = g \circ \phi$ . A homeomorphism  $\phi : U \rightarrow V$  between open subsets  $U$  and  $V$  of  $\mathbb{R}^n$  is called a blow-analytic homeomorphism if there exist two finite sequences of blowings-up along smooth analytic centres  $\pi : M \rightarrow U$  and  $\sigma : N \rightarrow V$  and an analytic diffeomorphism  $\Phi : M \rightarrow N$  such that  $\phi \circ \pi = \sigma \circ \Phi$ .

If the definition of blow-analytic equivalence via sequences of blowings-up makes it difficult to study, it has also very nice properties (cf. [9] for a survey). In particular a blow-analytic homeomorphism  $\phi : U \rightarrow V$  is arc-analytic [13], namely if  $\gamma \in \mathbb{R}\{t\}^n$  is a n-uplet of convergent power series on a neighbourhood of  $0 \in \mathbb{R}$  with  $\gamma(0) \in U$ , then  $\phi \circ \gamma$  is analytic. In particular a blow-analytic homeomorphism  $\phi$  induces a bijective mapping  $\phi_*$  between the spaces of analytic arcs at the origin of  $\mathbb{R}^n$ . The nice behaviour of the blow-analytic equivalence with respect to arcs and more generally spaces of arcs have already produced very interesting invariants (such as the Fukui invariants [7], zeta functions [10, 4]), a complete classification in dimension two [11], or explained some relations with respect to bi-Lipschitz property [8].

The first question we address in this paper is very natural in this context: given a blow-analytic homeomorphism  $\phi$ , is the induced mapping  $\phi_*$  between the spaces of analytic arcs continuous? It is natural to hope that such a homeomorphism induces a homeomorphism between arcs, even if the definition of a blow-analytic homeomorphism via sequences of blowings-up makes it difficult to handle directly. We offer in this paper two answers to this question.

A first answer is that the induced mapping is not continuous, even at the level of truncated arcs, when we considered  $\mathbb{R}\{t\}^n$  endowed with the product topology. We

---

*Date:* December 13, 2012.

*2010 Mathematics Subject Classification.* 14P15.

*Key words and phrases.* blow-analytic homeomorphism, arc space, real closed field, o-minimal structure.

The first author has been supported by the ANR project ANR-08-JCJC-0118-01.

show the existence of a counter-example in dimension two in section 1.3, where the sequence of blowings-up consists simply of the blowing-up  $\pi : M \rightarrow \mathbb{R}^2$  at the origin of  $\mathbb{R}^2$ . We produce the analytic diffeomorphism of  $M$  using classical real analytic sheaf theory combined with a cardinality argument.

A second answer is that the induced mapping is continuous... if  $\mathbb{R}\{t\}^n$  is endowed with the  $t$ -adic topology (cf. Theorem 1.3)! The result is actually a simple consequence of the Hölder property of subanalytic maps. However this question has a natural generalisation in the context of o-minimal structures over a real closed field where such a Hölder property is no longer available. Nevertheless, the tameness of an o-minimal structure should guaranty that the continuity of a mapping over a real closed field continues to hold when we naturally extend the field to another real closed field, and the mapping to a mapping over the extended field. And actually, if the Hölder property suffices to obtain the continuity in the case of subanalytic mappings over real numbers, a generalisation of Lojasiewicz Inequality to locally closed definable sets (Proposition 3.5), and an adapted version for general definable sets (Proposition 3.7) enable to control the behaviour of arcs in the o-minimal setting in order to keep the continuity at the level of spaces of definable arcs.

We propose moreover another approach, more natural in the non necessary locally closed case (e.g. a bijection coming from a resolution of the singularities as in Example 3.15), and following the model-theoretic point of view developed in [2]. In particular, we study more in details in section 3.3 the transport of properties between the initial o-minimal structure over a given real closed field and the new o-minimal structure on the real closed field of germs of definable arcs at the origin. We obtain moreover in a very simple way a uniform continuity property in Proposition 3.20.

**Acknowledgements.** The first author wish to thank K. Kurdyka for valuable remarks.

## 1. BLOW-ANALYTIC HOMEOMORPHISMS AND CONTINUITY

If a blow-analytic homeomorphism induces a continuous mapping at the level of spaces of arcs considered with the  $t$ -adic topology (cf. Theorem 1.3), we prove the existence of a blow-analytic homeomorphism which does not induce a continuous mapping at the level of spaces of arcs when we considered it with the product topology. The counter-example is produced in section 1.3 using a cardinality result concerning analytic vector fields detailed in section 1.4.

### 1.1. Blow-analytic homeomorphisms.

**Definition 1.1.** Let  $U$  and  $V$  be open subsets of  $\mathbb{R}^n$ . A homeomorphism  $\phi : U \rightarrow V$  is called a *blow-analytic homeomorphism* if there exist two finite sequences of blowings-up along smooth analytic centres  $\pi : M \rightarrow U$  and  $\sigma : N \rightarrow V$  and an analytic diffeomorphism  $\Phi : M \rightarrow N$  such that  $\phi \circ \pi = \sigma \circ \Phi$ .

Denote by  $\mathbb{R}\{t\}$  the one-variable convergent power series ring and  $\mathfrak{m}$  its maximal ideal. We consider in this section  $\mathbb{R}\{t\}^n$  equipped either with the  $t$ -adic topology or with the product topology. We regard  $\mathbb{R}\{t\}^n$  as the family of analytic curve germs  $c : [0; \epsilon) \rightarrow \mathbb{R}^n$ , with  $\epsilon > 0 \in \mathbb{R}$ , at 0 and let  $\mathcal{A}_0(\mathbb{R}^n)$  denote those curve germs  $c$  with  $c(0) = 0$ . Set

$$\mathcal{A}_U(\mathbb{R}^n) = \{c \in \mathbb{R}\{t\}^n : c(0) \in U\}$$

for an open subset  $U$  of  $\mathbb{R}^n$ . We identify  $\mathcal{A}_U(\mathbb{R}^n)$  with  $U \times \mathcal{A}_0(\mathbb{R}^n)$  by the correspondence

$$\mathcal{A}_U(\mathbb{R}^n) \ni c \rightarrow (c(0), c - c(0)) \in U \times \mathcal{A}_0(\mathbb{R}^n).$$

**1.2.  $t$ -adic topology.** Let  $h : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^n, 0)$  be an arc-analytic mapping germ [13], namely any analytic arc  $\gamma : (-\epsilon; \epsilon) \rightarrow \mathbb{R}^n$ , where  $\epsilon > 0$ , with  $\gamma(0) = 0$  is sent by  $h$  into an analytic arc  $h \circ \gamma : (-\epsilon'; \epsilon') \rightarrow \mathbb{R}^n$  for some  $\epsilon' > 0$ . Denote by  $\mathcal{A}_0(\mathbb{R}^n)$  those arcs  $\gamma$  in  $\mathbb{R}\{t\}^n$  with  $\gamma(0) = 0$ . Then  $h$  defines a mapping from  $\mathcal{A}_0(\mathbb{R}^n)$  to  $\mathcal{A}_0(\mathbb{R}^n)$ , denoted by  $h_*$  in the sequel. We considered  $\mathcal{A}_0(\mathbb{R}^n)$  with its  $t$ -adic topology.

**Remark 1.2.** A blow-analytic homeomorphism is arc-analytic since any analytic arc  $c : [0; \epsilon) \rightarrow U$  may be lifted via a sequence of blowings-up along smooth analytic centres  $\pi : M \rightarrow U$  to an analytic arc  $d : [0; \epsilon) \rightarrow M$  such that  $c = \pi \circ d$ . In particular a blow-analytic homeomorphism  $\phi : U \rightarrow V$  induces a mapping

$$\phi_* : \mathcal{A}_U(\mathbb{R}^n) \ni c \rightarrow \phi \circ c \in \mathcal{A}_V(\mathbb{R}^n)$$

Note that this mapping is moreover bijective.

**Theorem 1.3.** *Let  $h : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^n, 0)$  be a subanalytic homeomorphism. Assume  $h$  and  $h^{-1}$  are arc analytic. The induced mapping  $h_* : \mathcal{A}_0(\mathbb{R}^n) \rightarrow \mathcal{A}_0(\mathbb{R}^n)$  is a uniformly continuous homeomorphism with respect to the  $t$ -adic topology.*

*Proof.* As a subanalytic homeomorphism  $h$  is Hölder, so there exist  $\alpha \in \mathbb{Q}$ , with  $\alpha \in ]0, 1]$ , such that

$$|h(x) - h(y)| \leq c|x - y|^\alpha$$

for  $x, y$  close to 0, where  $c > 0$ . In particular, if  $\gamma, \delta : [0, \epsilon) \rightarrow (\mathbb{R}^n, 0)$  are analytic arcs, then

$$\text{ord}(h \circ \gamma(t) - h \circ \delta(t)) \geq \alpha \text{ord}(\gamma(t) - \delta(t)),$$

and this gives the continuity with respect to the  $t$ -adic topology.  $\square$

Theorem 1.3 extends to a global version on compact analytic manifolds. Let  $M \subset \mathbb{R}^n$  be a compact analytic manifold, and denote by  $\mathcal{A}(M)$  those arcs in  $\mathbb{R}\{t\}^n$  with origin in  $M$ . We consider  $\mathcal{A}(M)$  endowed with the topology induced by that of  $\mathbb{R}\{t\}^n$ .

**Theorem 1.4.** *Let  $M \subset \mathbb{R}^m$  and  $N \subset \mathbb{R}^n$  be compact analytic manifolds and  $h : M \rightarrow N$  be a subanalytic homeomorphism. Assume that  $h$  and  $h^{-1}$  are arc analytic. The induced mapping  $h_* : \mathcal{A}(M) \rightarrow \mathcal{A}(N)$  is a uniformly continuous homeomorphism.*

If the ambient spaces are no longer compact, we keep the continuity of the induced mapping between analytic arcs since continuity is a local property:

**Theorem 1.5.** *Let  $h : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a subanalytic homeomorphism. Assume  $h$  and  $h^{-1}$  are arc analytic. Then  $h_* : \mathbb{R}\{t\}^n \rightarrow \mathbb{R}\{t\}^n$  is a homeomorphism.*

**Remark 1.6.** The exponential function gives a counter-example to uniform continuity. Actually, let  $\gamma_s \in \widetilde{\mathbb{R}}$  be defined by  $\gamma_s(t) = \frac{1}{s} + st$ , for  $s > 0$ . Then  $\exp \circ \gamma_s$  is a family of analytic arcs with origin  $\exp(\gamma_s(0)) = \exp(\frac{1}{s})$  going to  $\infty$  as  $s$  goes to zero. However

$$\left(\frac{\partial \gamma_s}{\partial t}\right)_{t=0} = s \exp \frac{1}{s}$$

goes to  $\infty$  as  $s$  goes to zero, so the family is not uniformly continuous.




FIGURE 1. Blowing-up and curves.

**1.3. Product topology.** For a blow-analytic homeomorphism  $\phi$  as in Remark 1.2, it is natural to hope that the induced mapping  $\phi_*$  is a homeomorphism when we considered the product topology. However, it is not difficult to find a counter-example!

In the sequel, we focus therefore on some particular blow-analytic homeomorphisms, namely those blow-analytic homeomorphisms which induce a blow-analytic equivalence between some analytic functions (we refer to [9] for similar distinctions between different kind of blow-analytic homeomorphisms).

**Definition 1.7.** Let  $f, g : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}, 0)$  be analytic function germs. We call  $f$  and  $g$  *blow-analytically equivalent* if there exist open subsets  $U$  and  $V$  of  $\mathbb{R}^n$  containing 0 and a blow-analytic homeomorphism  $\phi : U \rightarrow V$  such that  $\phi(0) = 0$  and  $f = g \circ \phi_0$ , where  $\phi_0$  denotes the germ of  $\phi$  at 0. We call  $\phi_0 : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^n, 0)$  a *blow-analytic homeomorphism germ*.

The problem we address is then as follows.

**Question 1.8.** Let  $\phi$  be a blow-analytic homeomorphism germ which realises the blow-analytic equivalence of two non-zero analytic function germs. Is the induced mapping  $\phi_* : \mathcal{A}_0(\mathbb{R}^n) \ni c \rightarrow \phi \circ c \in \mathcal{A}_0(\mathbb{R}^n)$  continuous in the product topology?

In the following we give a negative answer to Question 1.8. We prove the existence of a counter-example via a cardinality argument rather than by an explicit construction. For the counter-example, we fix  $n = 2$  and define  $f, g : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$  by  $f(x, y) = g(x, y) = y$ . We define a family of curves  $c_\epsilon \in \mathcal{A}_0(\mathbb{R}^2)$  by

$$c_\epsilon(t) = (\epsilon t, t^2)$$

for  $\epsilon \in \mathbb{R}$ . Then  $c_\epsilon$  converges to  $c_0$  in  $\mathcal{A}_0(\mathbb{R}^2)$  for the product topology as  $\epsilon$  goes to 0.

Let  $\pi : M \rightarrow \mathbb{R}^2$  denote the blowing-up of  $\mathbb{R}^2$  along center  $\{0\}$ . We will prove the existence of an analytic diffeomorphism  $\Phi$  of  $M$  such that:

- (i)  $\Phi$  induces a blow-analytic homeomorphism  $\phi$  of  $\mathbb{R}^2$ ,
- (ii)  $f \circ \phi = f$  and
- (iii)  $(\phi_0)_*(c_\epsilon) \not\rightarrow (\phi_0)_*(c_0)$  as  $\epsilon \rightarrow 0$ .

Let  $M_1 = \pi^{-1}(0)$  denote the exceptional divisor of  $\pi$  and  $M_2$  denote the closure of  $\pi^{-1}(\mathbb{R} \times \{0\} - \{0\})$  in  $M$ .

**Lemma 1.9.** *Let  $\Phi$  be an analytic diffeomorphism of  $M$  such that (i) and (ii) hold. Then  $\Phi(M_1) = M_1$  and  $\Phi(M_2) = M_2$ .*

*Proof.* Condition (i) is equivalent to the condition  $\Phi(M_1) = M_1$  while condition (ii) is equivalent to the condition

$$(ii') \quad f \circ \pi \circ \Phi = f \circ \pi$$

Then  $(f \circ \pi)^{-1}(0) = M_1 \cup M_2$  and hence condition (ii') implies  $\Phi(M_2) = M_2$ .  $\square$

Assume that there exists an analytic diffeomorphism  $\Phi$  with the conditions (i), (ii) and (iii) satisfied. We describe when (iii) holds by a coordinate system. Denote by  $p$  the intersection point  $M_1 \cap M_2 = \{p\}$  and consider  $\pi$  around  $p$ . Choose a chart in  $M$  such that in the chart  $(M, M_1, M_2, p)$  is equal to  $(\mathbb{R}^2, \{0\} \times \mathbb{R}, \mathbb{R} \times \{0\}, (0, 0))$  and the restriction of  $\pi$  to the chart coincides with the mapping

$$\mathbb{R}^2 \ni (u, v) \rightarrow (u, uv) = (x, y) \in \mathbb{R}^2$$

that we still denote by  $\pi$ . Note that in this chart  $\text{Im } c_\epsilon$ , for  $\epsilon \neq 0$ , is included in  $\text{Im } \pi$  whereas  $\text{Im } c_0$  is not.

The restriction of  $\Phi$  to the above chart is of the form  $(u\Phi_1(u, v), v\Phi_2(u, v))$  for some analytic functions  $\Phi_1$  and  $\Phi_2$  on  $\mathbb{R}^2$  which vanish nowhere by Lemma 1.9.

**Lemma 1.10.** *In this chart, the image by  $\phi$  of the family of curves  $c_\epsilon$ , for  $\epsilon > 0$ , is given by*

$$\phi_*(c_\epsilon)(t) = (\epsilon t \Phi_1(\epsilon t, \frac{t}{\epsilon}), t^2)$$

*Proof.* In this chart we obtain for  $\epsilon > 0$ :

$$\begin{aligned} \pi^*(c_\epsilon)(t) &= (\epsilon t, \frac{t}{\epsilon}), \quad \Phi_*\pi^*(c_\epsilon)(t) = (\epsilon t \Phi_1(\epsilon t, \frac{t}{\epsilon}), \frac{t}{\epsilon} \Phi_2(\epsilon t, \frac{t}{\epsilon})), \\ \phi_*(c_\epsilon)(t) &= \pi_*\Phi_*\pi^*(c_\epsilon)(t) = (\epsilon t \Phi_1(\epsilon t, \frac{t}{\epsilon}), t^2 \Phi_1(\epsilon t, \frac{t}{\epsilon}) \Phi_2(\epsilon t, \frac{t}{\epsilon})), \\ f \circ \pi(u, v) &= uv. \end{aligned}$$

The last equality together with condition (ii') implies

$$uv\Phi_1(u, v)\Phi_2(u, v) = f \circ \pi \circ \Phi(u, v) = f \circ \pi(u, v) = uv$$

hence  $\Phi_1\Phi_2 \equiv 1$ . In particular  $\phi_*(c_\epsilon)(t) = (\epsilon t \Phi_1(\epsilon t, \frac{t}{\epsilon}), t^2)$ .  $\square$

As a consequence, to obtain a negative answer to Question 1.8, it suffices to find a global analytic diffeomorphism  $\Phi$  such that  $\epsilon t \Phi_1(\epsilon t, \frac{t}{\epsilon})$  does not converge to any element of  $\mathfrak{m}$  as  $\epsilon \rightarrow 0$  in  $\mathbb{R}$ .

**Remark 1.11.**

- (1) Note that  $\phi_*(c_0)$  does exist, even if we do not know how to describe it.
- (2) Note that the non-convergence of  $\pi^*(c_\epsilon)$  does not imply necessarily the non-convergence of  $\phi_*(c_\epsilon)$  whereas the non-convergence of  $\epsilon t \Phi_1(\epsilon t, \frac{t}{\epsilon})$  does.

Describe  $\Phi_1$  around 0 as a convergent power series  $\sum_{m,n \in \mathbf{N}} a_{m,n} u^m v^n$  with  $a_{m,n} \in \mathbb{R}$ . Then the first component of  $\phi_*(c_\epsilon)(t)$  for  $\epsilon > 0$  is given by

$$\epsilon t \Phi_1(\epsilon t, \frac{t}{\epsilon}) = \sum_{m,n \in \mathbf{N}} a_{m,n} t^{m+n+1} \epsilon^{m-n+1}.$$

**Lemma 1.12.** *If there exist  $m$  and  $n$  in  $\mathbf{N}$  such that  $n > m + 1$  and  $a_{m,n} \neq 0$ , then  $\epsilon t \Phi_1(\epsilon t, \frac{t}{\epsilon})$  does not converge to any element of  $\mathfrak{m}$  as  $\epsilon \rightarrow 0$ .*

*Proof.* Assume that there exist such pairs  $(m, n)$  with  $n > m + 1$  and  $a_{m,n} \neq 0$ . Let  $L$  denote the family of all such pairs. Denote by  $L_1$  the subfamily of  $L$  consisting of  $(m, n)$  such that the sum  $m + n$  is the smallest possible in  $L$  and by  $L_2$  the subfamily of  $L_1$  of  $(m, n)$  such that  $n - m$  is the largest possible in  $L_1$ . Then  $L_1$  is non-empty and finite, and  $L_2$  is non-empty and consists of a unique element, say  $(m_0, n_0)$ . Then  $\epsilon t \Phi_2(\epsilon t, \frac{t}{\epsilon})$  is of the form

$$\epsilon t \Phi_2(\epsilon t, \frac{t}{\epsilon}) = a_{m_0, n_0} t^{m_0 + n_0 + 1} (\epsilon^{m_0 - n_0 + 1} + \delta_{m_0 + n_0 + 1}(\epsilon)) + \sum_{k=m_0 + n_0 + 2}^{\infty} \delta_k(\epsilon) t^k$$

where the functions  $\delta_k$  are functions in the variable  $\epsilon$  such that

$$|\delta_{m_0 + n_0 + 1}(\epsilon)| \leq c |\epsilon| |\epsilon^{m_0 - n_0 + 1}|$$

for some  $c > 0 \in \mathbb{R}$  and  $\epsilon \neq 0$  near 0. Therefore  $\epsilon t \Phi_2(\epsilon t, \frac{t}{\epsilon})$  does not converge to any element of  $\mathfrak{m}$  as  $\epsilon \rightarrow 0$ .  $\square$

For simplicity, we strengthen the condition

$$a_{m,n} \neq 0 \text{ for some } n > m + 1$$

to the condition

$$a_{0,n} \neq 0 \text{ for some } n > 1$$

In particular in that case

$$\Phi_1(0, v) = a_{0,0} + a_{0,1}v + \sum_{n \in \mathbb{N}} a_{0,n}v^n.$$

**Lemma 1.13.** *There exists  $\Phi_1$  satisfying  $a_{0,n} \neq 0$  for some  $n > 1$ .*

*Proof.* We find  $\Phi_1$  with this condition by reduction to absurdity. Assume that no  $\Phi_1$  satisfies this condition. Then the family of all such  $\Phi_1(0, v)$  is a subfamily of the family of  $v$ -polynomials of degree less or equal to 1 and hence has at most the cardinal number of the continuum. Moreover, since  $\Phi_1 \Phi_2 \equiv 1$  by the proof of Lemma 1.10, the family of all  $\Phi_2(0, v)$  also has at most the cardinal number of the continuum.

However the restriction of  $\Phi$  to  $M_1$  is given by  $\Phi(0, v) = (0, v \Phi_2(0, v))$ , and we obtain a contradiction with Proposition 1.14 below.  $\square$

**1.4. Cardinality of sets of vector fields.** The aim of this section is to prove the cardinality argument to finish the proof of the negative answer to Question 1.8.

**Proposition 1.14.** *Let  $T$  denote the set of all  $C^\omega$  diffeomorphism  $\Phi$  of  $M$  such that  $\Phi(M_1) = M_1$ ,  $\Phi(M_2) = M_2$  and  $f \circ \pi = f \circ \pi \circ \Phi$ , and  $T_1$  the set of restrictions of such  $\Phi$  to  $M_1$ . Then the cardinal number of  $T_1$  is larger than  $\mathbb{R}$ .*

If it is difficult to construct directly  $C^\omega$  diffeomorphisms of  $M$ , it is easier to construct  $C^\omega$  vector fields via analytic sheaf theory. In particular, we will use Cartan Theorem A and B, Oka Theorem for real analytic manifold (cf. [6] for example), together with Whitney Approximation Theorem (any  $C^0$  function on an analytic manifold can be approximated in the Whitney topology by a  $C^\omega$  function [15]).

Recall that any  $C^\omega$  vector field  $w_1$  on (a compact)  $M_1$  generates a one-parameter group of transformations  $\psi_t$ ,  $t \in \mathbb{R}$  of  $M_1$ , and the correspondence  $w_1 \rightarrow \psi_1$  is a countable-to-one correspondence from the family of  $C^\omega$  vector fields on  $M_1$  to the family of  $C^\omega$  diffeomorphisms on  $M_1$ .



Using this fact, Proposition 1.14 is a consequence of Propositions 1.15 and 1.17 below.

An integrable vector field on  $M$  means that the vector field generates a one-parameter group of transformations.

**Proposition 1.15.** *Any  $C^\omega$  vector field  $w_1$  on  $M_1$  singular at  $p$  can be extended to an integrable  $C^\omega$  vector field  $w$  on  $M$  so that  $f \circ \pi$  takes a constant value on each integral curve of  $w$ .*

In order to prove Proposition 1.15, we introduce a weaker lemma.

**Lemma 1.16.** *Any  $C^\omega$  vector field  $w_1$  on  $M_1$  singular at  $p$  can be extended to a  $C^\omega$  vector field  $w$  on  $M$  so that  $f \circ \pi$  takes a constant value on each local integral curve of  $w$ .*

*Proof of Proposition 1.15.* Assume that there exists a  $C^\omega$  vector field  $w$  on  $M$  satisfying the conclusion of Lemma 1.16. To prove Proposition 1.15 it is sufficient to construct a bounded vector field satisfying that  $f \circ \pi$  takes a constant value on each local integral curve of  $w$ .

Let  $U$  be a sufficiently small neighbourhood of  $M_1$  in  $M$  and  $\delta$  a sufficiently small positive  $C^0$  function on  $M$  such that  $\delta w$  is bounded, where  $M$  is endowed with the Whitney  $C^0$  topology. Assume there exists a positive  $C^\omega$  function  $h$  on  $M$  such that:

- (1)  $h \leq 1$  on  $M$ ,
- (2)  $h = 1$  on  $M_1$  and
- (3)  $h \leq \delta$  on  $M - U$ .

Then  $hw$  is integrable and hence satisfies the requirements on  $w$ .

Hence we have to construct such a positive  $C^\omega$  function  $h$  satisfying (1), (2) and (3). Actually it will be convenient to construct  $g = 1 - h$  in place of  $h$  using the fact that  $g$  need to be an element of the ideal of the analytic function ring on  $M$  vanishing on  $M_1$ .

Let  $\mathcal{O}$  denote the sheaf of  $C^\omega$  function germs on  $M$  and  $\mathcal{J}$  the sheaf of  $\mathcal{O}$ -ideals consisting of germs vanishing on  $M_1$ . Then by Cartan Theorem A each stalk  $\mathcal{J}_q$ , for  $q \in M$ , is generated by the global cross-sections  $H^0(M, \mathcal{J})$  of  $\mathcal{J}$ . Note that  $H^0(M, \mathcal{J})$  is finitely generated since  $M_1$  is compact. Moreover the common zero set of the germs in  $\mathcal{J}_q$  is equal to the germ of  $M_1$  at  $q$ . Therefore there exist finitely many elements of  $H^0(M, \mathcal{J})$  whose common zero set is equal to  $M_1$ . Define  $g_1$  to be the square sum of these elements. Then  $g_1$  is a non-negative  $C^\omega$  function on  $M$  with zero set equal to  $M_1$ , so that (2) is satisfied by  $1 - g_1$ .

Next we shrink  $g_1$  in order to satisfy (1). Let  $\alpha$  be a  $C^0$  function on  $M$  such that  $\alpha g_1 < 1$  on  $M$ . By Whitney Approximation Theorem,  $\alpha$  can be approximated by a  $C^\omega$  function  $\alpha_1$  in the Whitney  $C^0$  topology such that  $\alpha_1 g_1 \leq 1$  on  $M$ . Set  $g_2 = \alpha_2 g_1$ .

Finally, we modify  $g_2$  so that  $1 - g_2$  satisfies (3). Let  $V$  be a neighbourhood of  $M_1$  in  $M$  such that  $\bar{V} \subset \text{Int } U$ . Let  $\alpha_3$  be a positive  $C^0$  function on  $M$  such that

- $\alpha_3 < 1$  on  $V$ ,
- $\alpha_3 < \frac{1}{g_2}$  on  $M - V$  and
- $\alpha_3 > \frac{1}{g_2} - \frac{\delta}{2g_2}$  on  $M - U$ .

By Whitney Approximation Theorem, approximate  $\alpha_3$  by a positive  $C^\omega$  function  $\alpha_4$  on  $M$  so that



- $\alpha_4 < 1$  on  $V$ ,
- $\alpha_4 < \frac{1}{g_2}$  on  $M - V$  and
- $\alpha_4 \geq \frac{1}{g_2} - \frac{\delta}{g_2}$  on  $M - U$ .

Then  $1 - g_2\alpha_4 \leq \delta$  on  $M - U$  and  $h = 1 - g_2\alpha_4$  satisfies (1), (2) and (3) as expected.  $\square$

*Proof of Lemma 1.16.* Let  $\mathcal{I}$  denote the sheaf of  $\mathcal{O}$ -modules of  $C^\omega$  vector field germs  $\xi$  on  $M$  such that  $\xi(f \circ \pi) \equiv 0$ . In particular  $H^0(M, \mathcal{I})$  is the family of  $C^\omega$  vector fields on  $M$  such that  $f \circ \pi$  takes a constant value on each local integral curve. Set

$$\mathcal{I}_0 = \{\xi \in \mathcal{I} : \xi = 0 \text{ on } M_1\}$$

and  $\mathcal{I}_1 = \mathcal{I}/\mathcal{I}_0$ . We can regard each element of  $\mathcal{I}_1$  as a  $C^\omega$  vector field germ on  $M_1$  and each element of  $H^0(M, \mathcal{I}_1)$  as a  $C^\omega$  vector field on  $M_1$ .

By Oka Theorem  $\mathcal{I}$ ,  $\mathcal{I}_0$  and  $\mathcal{I}_1$  are coherent. Therefore by Cartan Theorem B the mapping  $H^0(M, \mathcal{I}) \rightarrow H^0(M, \mathcal{I}_1)$  is surjective. Therefore it is enough to prove that  $w_1$  belongs to  $H^0(M, \mathcal{I}_1)$  in order to prove Lemma 1.16.

To that purpose, it suffices to prove that each  $C^\omega$  vector field germ  $\xi_1$  on  $M_1$  at each point  $q$  of  $M_1$  can be extended to a  $C^\omega$  vector field germ  $\xi$  on  $M$  at  $q$  so that  $\xi(f \circ \pi) \equiv 0$ , under the condition that if  $p = q$  then  $\xi_1$  is singular at  $q$ .

In the case  $q \neq p$ , we have that  $f \circ \pi \equiv 0$  on  $M_1$  and  $f \circ \pi$  is non-singular at  $q$ , therefore the family  $\{(f \circ \pi)^{-1}(a) : a \in \mathbb{R}\}$  around  $q$  is a  $C^\omega$  foliation that contains  $M_1$ .

Assume now that  $q = p$  and  $\xi_1$  is singular at  $p$ . We use the local coordinate system  $(u, v)$  at  $p = 0$  introduced in Lemma 1.10. Set  $\xi_1 = \beta(0, v) \frac{\partial}{\partial v}$  on  $\{0\} \times \mathbb{R}$ , where  $\beta$  is a  $C^\omega$  function on  $\{0\} \times \mathbb{R}$  such that  $\beta(0, 0) = 0$ . Define  $\xi$  to be

$$-\frac{u\beta(0, v)}{v} \frac{\partial}{\partial u} + \beta(0, v) \frac{\partial}{\partial v}$$

for  $(u, v) \in \mathbb{R}^2$  with  $v \neq 0$  and to be

$$-u \frac{\partial \beta}{\partial v}(0, 0) \frac{\partial}{\partial u}$$

for  $(u, v) \in \mathbb{R}^2$  with  $v = 0$ . Then  $\xi$  gives the required extension.  $\square$

**Proposition 1.17.** *The set of all  $C^\omega$  vector fields on  $M_1$  singular at  $p$  has a cardinal number larger than  $\mathbb{R}$ .*

*Proof.* First, there exists at least one non-zero  $C^\omega$  vector field  $w$  on  $M_1$  singular at  $p$  by Cartan Theorem A. Now, for distinct  $C^\omega$  functions  $h_1$  and  $h_2$  on  $M_1$ , the vector fields  $h_1w$  and  $h_2w$  are distinct and singular at  $p$ . We conclude using the fact that  $M_1$  is analytically diffeomorphic to  $S^1$  and that the family of  $C^\omega$  functions on  $S^1$  has a cardinal number larger than  $\mathbb{R}$ .  $\square$

**Remark 1.18.**

- (1) One can prove that the family of  $C^\omega$  functions on  $M_1$  has a cardinal number larger than  $\mathbb{R}$  as follows. It suffices to consider the case of  $\mathbb{R}$  because the correspondence  $C^\omega(\mathbb{R}) \ni h(v) \rightarrow h(\sin v) \in C^\omega(\mathbb{R})$  is injective, so that we can regard the image of the correspondence as the family of  $C^\omega$  functions on  $S^1$ . In the case of  $\mathbb{R}$ , the family of convergent power series with infinite radius of convergence is of cardinal number larger than  $\mathbb{R}$ .

- (2) We can modify above  $\Phi$  such that it becomes a Nash diffeomorphism by [5]. However, the above proof does no longer work in the Nash category since the family of all Nash functions on a Nash manifold has the cardinal number of the continuum.

## 2. SET OF GERMS OF DEFINABLE FUNCTIONS

In order to deal with a generalisation of Theorem 1.3 in the context of o-minimal structures in section 3, we introduce the real closed field of definable germs of arcs over a given real closed field in section 2.3. We begin with some basic facts about real closed fields and recall the unique Euclidean topology on a real closed field, following [1].

**2.1. Real closed fields.** An ordered field is a field  $F$  equipped with an ordering, namely there exists a total order relation  $\leq$  on  $F$  satisfying  $x + z \leq y + z$  if  $x \leq y$  and the product of positive elements is positive. Fields such as  $\mathbb{Q}$  and  $\mathbb{R}$  are ordered fields.

**Example 2.1.** There exist several orderings on the field  $\mathbb{R}(t)$  of real rational fractions. For example, we may choose that  $a_n t^n + a_{n+1} t^{n+1} + \dots + a_m t^m$ , with  $a_n \neq 0$  is positive if and only if  $a_n > 0$ . In particular  $t$  is positive and smaller than any positive real number. The situation is similar with the field of formal power series  $\mathbb{R}((t))$ .

A real field is a field that can be ordered. A real closed field is a real field that does not admit any non trivial real extension. A real closed field  $\mathbf{R}$  is characterised by the fact that  $\mathbf{R}$  admits a unique ordering such that the positive elements coincide with the squares and every polynomial in  $\mathbf{R}[X]$  of odd degree has a root in  $\mathbf{R}$ . Every ordered field admits a real closure, unique up to a unique isomorphism. Of course  $\mathbb{R}$  is real closed. The real closure of  $\mathbb{Q}$  consists of the field of real algebraic numbers  $\mathbb{R}_{alg}$ .

**Example 2.2.** The field of (formal) real Puiseux series  $\mathbb{R}((t^{1/\infty})) = \bigcup_{q \in \mathbb{N}^*} \mathbb{R}((t^{1/q}))$ , namely the set of formal expressions

$$\sum_{p=n}^{+\infty} a_p t^{p/q} \text{ with } n \in \mathbb{Z}, q \in \mathbb{N}^*, a_p \in \mathbb{R},$$

is real closed. A positive Puiseux series is a series of the form  $\sum_{p=n}^{+\infty} a_p t^{p/q}$  with  $a_n > 0$ .

The real closure of the field of rational functions  $\mathbb{R}(t)$  is the field of algebraic Puiseux series  $\mathbb{R}((t^{1/\infty}))_{alg}$ , i.e. Puiseux series algebraic over  $\mathbb{R}(t)$ . The field of convergent Puiseux series  $\mathbb{R}\{t^{1/\infty}\}$  is also a real closed field. The following inclusions hold:

$$\mathbb{R}_{alg} \subset \mathbb{R} \subset \mathbb{R}((t^{1/\infty}))_{alg} \subset \mathbb{R}\{t^{1/\infty}\} \subset \mathbb{R}((t^{1/\infty}))$$

**2.2. Topology.** A real closed field  $\mathbf{R}$  induces an Euclidean topology on  $\mathbf{R}^n$  as follows: for  $x \in \mathbf{R}^n$ , defined  $|x| = \sqrt{x_1^2 + \dots + x_n^2}$ . Then the open balls

$$B(x, r) = \{y \in \mathbf{R}^n : |y - x| < r\}$$

for  $x \in \mathbf{R}^n$  and  $r \in \mathbf{R}$  with  $r > 0$ , form a basis of open subsets. The Euclidean topology is the unique topology compatible with the real structure, in the sense that

a real closed field admits a unique ordering. It is called the Euclidean topology of the real closed field.

**Example 2.3.**

- (1) The subset of  $\mathbb{R}((t^{1/\infty}))_{alg}$  defined by

$$\left\{ \sum_{k \in \mathbb{N}} a_k t^{k/p} : a_k \in \mathbb{R}, p \in \mathbb{N}^* \right\}$$

is open. Actually it is an infinite union of open intervals:

$$\bigcup_{a \in (0; +\infty) \cap \mathbb{R}} (-a; a).$$

- (2) As a real closed field,  $\mathbb{R}((t^{1/\infty}))_{alg}$  inherits to an Euclidean topology. An element  $\gamma$  in  $\mathbb{R}((t^{1/\infty}))_{alg}$  of the form  $\gamma(t) = \sum_{p \geq m} a_p t^{p/q}$ , with  $a_p \in \mathbb{R}$ ,  $q \in \mathbb{N}^*$ , and  $a_m \neq 0$  is small in this topology if and only if  $m$  is large.

Note that the Euclidean topology of a real closed field may behave very differently from that of real numbers: a real closed field is not necessarily connected, and  $[0, 1]$  is not compact in  $\mathbb{R}_{alg}$  nor in  $\mathbb{R}((t^{1/\infty}))$ .

**Remark 2.4.** On the field of (convergent) power series  $\mathbb{R}\{t\}$  one can consider several topologies such as the product topology or  $t$ -adic topology studied in section 1. Note that the  $t$ -adic topology on  $\mathbb{R}\{t\}$  coincides with the restriction to  $\mathbb{R}\{t\}$  of the Euclidean topology of the field of convergent Puiseux series  $\mathbb{R}\{t^{1/\infty}\}$ .

**2.3. o-minimal structures expanding  $\mathbf{R}$ .** We introduce an o-minimal structure expanding a real closed field following [2].

Let  $\mathbf{R}$  be a real closed field. An o-minimal structure expanding  $\mathbf{R}$  is a collection  $\mathcal{S} = (\mathcal{S}^n)_{n \in \mathbb{N}}$ , where each  $\mathcal{S}^n$  is a set of subsets of the affine space  $\mathbf{R}^n$ , satisfying the following axioms:

- (1) All algebraic subsets of  $\mathbf{R}^n$  are in  $\mathcal{S}^n$ .
- (2) For every  $n$ ,  $\mathcal{S}^n$  is a Boolean subalgebra of subsets of  $\mathbf{R}^n$ .
- (3) If  $A \in \mathcal{S}^m$  and  $B \in \mathcal{S}^n$ , then  $A \times B \in \mathcal{S}^{m+n}$ .
- (4) If  $p : \mathbf{R}^{n+1} \rightarrow \mathbf{R}^n$  is the projection on the first  $n$  coordinates and  $A \in \mathcal{S}^{n+1}$ , then  $p(A) \in \mathcal{S}^n$ .
- (5) An element of  $\mathcal{S}^1$  is a finite union of points and open intervals  $(a; b) = \{x \in \mathbf{R} : a < x < b\}$ , with  $a, b \in \mathbf{R} \cup \{\pm\infty\}$ .

**Example 2.5.**

- (1) Let  $\mathbf{R}$  be a real closed field. The most simplest o-minimal structure expanding  $\mathbf{R}$  is the structure whose definable sets are the semialgebraic sets.
- (2) A globally subanalytic subset  $A$  of  $\mathbb{R}^n$  is a subanalytic subset  $A$  of  $\mathbb{R}^n$  which is subanalytic at infinity. Namely, if we embed  $\mathbb{R}^n$  in  $\mathbb{S}^n$  (via some rational regular embedding) and consider  $\mathbb{S}^n$  in  $\mathbb{R}^{n+1}$ , then we ask that  $A \subset \mathbb{R}^n \subset \mathbb{S}^n \subset \mathbb{R}^{n+1}$  is a subanalytic subset of  $\mathbb{R}^{n+1}$ . The collection of globally subanalytic sets forms an o-minimal structure expanding  $\mathbb{R}$ .

A definable function is a function defined on a definable set whose graph is definable. We will use several times the curve selection lemma in o-minimal structures in the sequel.

**Theorem 2.6** (Curve selection lemma). *Let  $\mathbf{R}$  be a real closed field and let be given an o-minimal structure expanding  $\mathbf{R}$ . Let  $A$  be a definable subset of  $\mathbf{R}^n$  and  $x \in \overline{A}$ . There exists a continuous definable mapping  $\gamma : [0; 1) \rightarrow \mathbf{R}^n$  such that  $\gamma(0) = x$  and  $\gamma((0; 1)) \subset A$ .*

We will also need to reparametrize definable arcs ([3], Exercise (1.9) p 49).

**Lemma 2.7.** *Let  $\gamma : (0; \epsilon) \rightarrow \mathbf{R}$  be a non-constant continuous definable function. There exist numbers  $\epsilon_1$  and  $\epsilon_2$  in  $\mathbf{R}$  with  $\epsilon_1 < \epsilon_2$  and a continuous definable bijection  $\delta : (\epsilon_1; \epsilon_2) \rightarrow (0; \epsilon_3)$ , with  $0 < \epsilon_3 < \epsilon$  such that  $\gamma \circ \delta(t) = t$  for any  $t \in (\epsilon_1; \epsilon_2)$ .*

**2.4. Germs of definable functions.** Given a real closed field and an o-minimal structure expanding it, we introduce its field of germs of continuous definable curves at the origin in section 2.4. We derive an o-minimal structure expanding it and study in section 3.3 the transport of some properties from the initial o-minimal structure to the new one.

Let  $\mathbf{R}$  be a real closed field. We fix an o-minimal structure over  $\mathbf{R}$  expanding  $\mathbf{R}$ . Let  $\tilde{\mathbf{R}}$  be the set of germs at  $0 \in \mathbf{R}$  of continuous definable functions from  $(0; \infty)$  to  $\mathbf{R}$ .

**Lemma 2.8.** *The set  $\tilde{\mathbf{R}}$  is a real closed field.*

*Proof.* The set  $\tilde{\mathbf{R}}$  is a field. Actually a non zero definable function on a neighbourhood of 0 in  $(0; \infty)$  nowhere vanishes on a sufficiently small neighbourhood of 0, so its inverse is well defined and definable.

To prove that  $\tilde{\mathbf{R}}$  is a real closed field, it is sufficient to prove that  $\tilde{\mathbf{R}}[i] = \frac{\tilde{\mathbf{R}}[Y]}{(Y^2+1)}$  is an algebraically closed field. First we prove that the ring  $\frac{\tilde{\mathbf{R}}[Y]}{(Y^2+1)}$  is a field. For that it suffices to show that  $(Y^2 + 1)$  is a prime ideal. Otherwise,  $Y^2 + 1$  is the product of two polynomial functions because  $\tilde{\mathbf{R}}[Y]$  is a unique factorization ring and hence  $Y^2 + 1$  is of the form  $(Y - f_1) \times (Y - f_2)$  and  $Y^2 - f_1^2$  for some  $f_1$  and  $f_2$  in  $\tilde{\mathbf{R}}$ , which is impossible. For  $P(t, X) \in \tilde{\mathbf{R}}[i][X]$ , write

$$P(t, X) = \sum_{j=0}^d f_j(t) X^j$$

with  $f_j \in \tilde{\mathbf{R}}[i]$ . Choose  $\epsilon > 0$  such that every  $f_j$ , for  $j = 0, \dots, d$ , is defined on  $(0; \epsilon)$ . For  $t \in (0; \epsilon)$  fixed, the polynomial  $P(t, X) \in R[i][X]$  admits roots in  $R[i]$  and the map

$$\{(t, x_1, \dots, x_d) \in (0; \epsilon) \times R[i] : P(t, x) = 0\} \rightarrow (0; \epsilon)$$

is finite-to-one. By the o-minimal Hardt triviality Theorem (which is proved in the same way as the semialgebraic case, cf. [1], Theorem 9.3.2), this mapping admits a definable continuous section on a neighbourhood of 0 in  $(0; \epsilon)$ . It furnishes a root of  $P$  in  $\tilde{\mathbf{R}}[i]$ .  $\square$

**Remark 2.9.**

- (1) Note that  $\tilde{\mathbf{R}}$  is even a field extension of  $\mathbf{R}$  by assigning to a number  $x \in \mathbf{R}$  the constant function  $\tilde{x}$  with image  $x$ .
- (2) A positive element in  $\tilde{\mathbf{R}}$  is a square; in particular a definable continuous function germ  $\gamma : (0; \infty) \rightarrow \mathbf{R}$  at 0 is positive in  $\tilde{\mathbf{R}}$  if and only if there exists  $\epsilon > 0$  in  $\mathbf{R}$  such that  $\gamma(t) \geq 0$  for  $t \in (0; \epsilon)$ .

**Example 2.10.**

- (1) Note that [1] the field of germs of continuous semialgebraic functions defined on intervals of the form  $(0; \epsilon) \subset \mathbb{R}$  is isomorphic to the field of algebraic Puiseux series  $\mathbb{R}((t^{1/\infty}))_{alg}$ . The subring of algebraic formal power series  $\mathbb{R}[[t]]_{alg}$  corresponds to the germs at the origin of analytic semialgebraic functions defined on intervals  $[0; \epsilon)$ .

In particular, if  $\mathbf{R} = \mathbb{R}$  and the o-minimal structure is defined by the semialgebraic sets, then  $\widetilde{\mathbf{R}}$  is the field of algebraic Puiseux series  $\mathbb{R}((t^{1/\infty}))_{alg}$ .

Note that the same holds true for any real closed field  $\mathbf{R}$  in place of  $\mathbb{R}$ .

- (2) If  $\mathbf{R} = \mathbb{R}$  and the o-minimal structure is given by the globally subanalytic sets, then  $\widetilde{\mathbf{R}}$  is the field of convergent Puiseux series  $\mathbb{R}\{t^{1/\infty}\}$ .

For a definable set  $X \subset \mathbf{R}^n$ , for  $n \in \mathbb{N}$ , denote by  $\widetilde{X}$  the set of germs at  $0 \in \mathbf{R}$  of continuous definable functions from  $(0; \infty)$  to  $X$ . For  $x \in X$ , we denote again  $\widetilde{x} \in \widetilde{X}$  the germ of the constant function equal to  $x$ .

**Lemma 2.11.** *Let  $f : \mathbf{R} \rightarrow \mathbf{R}$  be definable. Let  $\widetilde{f} : \widetilde{\mathbf{R}} \rightarrow \widetilde{\mathbf{R}}$  be defined by  $\widetilde{f}(\gamma(t)) = f \circ \gamma(t)$  for  $\gamma \in \widetilde{\mathbf{R}}$ . Then*

$$\text{graph } \widetilde{f} = \widetilde{\text{graph } f}.$$

*Proof.* It suffices to remark that  $\widetilde{\mathbf{R}^2} = \widetilde{\mathbf{R}}^2$  since

$$\begin{aligned} \widetilde{\text{graph } f} &= \{\gamma : (0; \epsilon) \rightarrow \mathbf{R}^2 : \gamma \text{ continuous definable and } \text{Im } \gamma \subset \text{graph } f\} \subset \widetilde{\mathbf{R}^2} \\ \text{graph } \widetilde{f} &= \{(\gamma_1, \gamma_2) \in \widetilde{\mathbf{R}}^2 : \gamma_2 = f \circ \gamma_1\} \subset \widetilde{\mathbf{R}}^2 \end{aligned}$$

□

**Proposition 2.12.** *The collection of subsets of  $\widetilde{\mathbf{R}}^n$  given by  $\widetilde{X}$ , for  $X \subset \mathbf{R}^n$  definable and  $n \in \mathbb{N}$ , defines an o-minimal structure expanding  $\widetilde{\mathbf{R}}$ .*

*Proof.* Let  $A$  and  $B$  be definable sets in  $\mathbf{R}^n$ . What we need to prove is the following:

- (i)  $\widetilde{A \cup B} = \widetilde{A} \cup \widetilde{B}$ ,  $\widetilde{A \cap B} = \widetilde{A} \cap \widetilde{B}$ ,  $\widetilde{A \setminus B} = \widetilde{A} \setminus \widetilde{B}$ ,  $\widetilde{A \times B} = \widetilde{A} \times \widetilde{B}$ ,
- (ii) if  $p$  denotes the projection  $p : \mathbf{R}^{n+1} \rightarrow \mathbf{R}^n$  onto the first  $n$  coordinates, and  $C \subset \mathbf{R}^{n+1}$  is definable, then  $\widetilde{p(C)} = \widetilde{p}(\widetilde{C})$ ,

(iii)

$$\{(x, y, z) \in \mathbf{R}^3 : x = yz\} = \{(\alpha, \beta, \gamma) \in \widetilde{\mathbf{R}}^3 : \alpha = \beta\gamma\},$$

(iv)

$$\{(x, y, z) \in \mathbf{R}^3 : x = y + z\} = \{(\alpha, \beta, \gamma) \in \widetilde{\mathbf{R}}^3 : \alpha = \beta + \gamma\},$$

(v) if  $I \subset \mathbf{R}$  is an interval, then  $\widetilde{I} \subset \widetilde{\mathbf{R}}$  is an interval,

(vi) if  $S \subset \mathbf{R}$  is a singleton, then  $\widetilde{S} \subset \widetilde{\mathbf{R}}$  is a singleton.

*Proof of (i).* The inclusion  $\widetilde{A} \cup \widetilde{B} \subset \widetilde{A \cup B}$  is obvious. Conversely, take  $\gamma \in \widetilde{A \cup B}$  with  $\gamma$  defined and continuous on  $(0; \epsilon)$ . Assume  $\gamma \notin \widetilde{A}$ . Then  $\gamma^{-1}(A)$  is a finite union of points and intervals, therefore there exists  $0 < \epsilon_1 < \epsilon$  such that  $\gamma((0; \epsilon_1)) \cap A = \emptyset$ . Then  $\gamma((0; \epsilon_1)) \subset B$  and  $\gamma \in \widetilde{B}$ .

The equality  $\widetilde{A \cap B} = \widetilde{A} \cap \widetilde{B}$  is obvious.

The inclusion  $\widetilde{A \setminus B} \subset \widetilde{A} \setminus \widetilde{B}$  is obvious. Take  $\gamma \in \widetilde{A} \setminus \widetilde{B}$ , and assume  $\gamma$  defined and continuous on  $(0; \epsilon)$ . Then  $\gamma^{-1}(B) \subset (0; \epsilon)$  is a finite union of points and

intervals, that does not contain any interval of the form  $(0; s)$  since  $\gamma \notin \widetilde{B}$ . Therefore  $\gamma((0; s)) \subset A \setminus B$ . and  $\gamma \in \widetilde{A \setminus B}$ .

The equality  $\widetilde{A \times B} = \widetilde{A} \times \widetilde{B}$  is obvious.

Proof of (ii). Take  $\gamma \in \widetilde{C}$  and assume that  $\gamma$  is defined and continuous on  $(0; \epsilon)$ , with  $\gamma((0; \epsilon)) \subset C$ . Then  $p \circ \gamma : (0; \epsilon) \rightarrow p(C)$  is a definable continuous function so that  $\widetilde{p(\gamma)} \in \widetilde{p(C)}$ . Therefore  $\widetilde{p(\widetilde{C})} \subset \widetilde{p(C)}$ .

Conversely, take  $\gamma \in \widetilde{p(C)}$ , and choose  $\epsilon > 0$  so that  $\gamma$  is a definable continuous function on  $(0; \epsilon)$  with value in  $p(C)$ . Note that we can assume that  $C$  is bounded, embedding if necessary  $\mathbf{R}^{n+1}$  in  $\mathbb{S}^{n+1} \subset \mathbf{R}^{n+2}$ . Then

$$D = \{(t, c) \in (0; \epsilon) \times C : p(c) = \gamma(t)\}$$

is a definable set which boundary has a non-empty intersection with  $\{0\} \times C$  since  $C$  is bounded. By the curve selection lemma, there exists  $\delta = (\delta_1, \delta_2) \in \widetilde{D}$ , i.e. there exists  $0 < \epsilon_1 < \epsilon$  such that  $p(\delta_2(t)) = \gamma(\delta_1(t))$  for  $t \in (0; \epsilon_1)$ . By Lemma 2.7 there exists  $\alpha \in \widetilde{\mathbf{R}}$  defined on  $(0; \epsilon_2)$ , with  $0 < \epsilon_2 < \epsilon_1$ , such that  $\delta_1 \circ \alpha(t) = t$  for  $t \in (0; \epsilon_2)$ . As a consequence  $\delta_2 \circ \alpha \in \widetilde{C}$  satisfies  $\widetilde{p}(\delta_2 \circ \alpha) = \gamma$ , therefore  $\gamma \in \widetilde{p(\widetilde{C})}$ .

The proofs of (iii) and (iv) are very similar, so let us prove (iii). Take

$$\gamma = (\gamma_1, \gamma_2, \gamma_3) \in \widetilde{\{(x, y, z) \in \mathbf{R}^3 : x = yz\}}.$$

There exists  $\epsilon > 0$  such that  $\gamma$  is defined on  $(0; \epsilon)$  and for  $t \in (0; \epsilon)$  the equality  $\gamma_1(t) = \gamma_2(t)\gamma_3(t)$  holds. Therefore  $\gamma_1 = \gamma_2\gamma_3$  and

$$\widetilde{\{(x, y, z) \in \mathbf{R}^3 : x = yz\}} \subset \widetilde{\{(\alpha, \beta, \gamma) \in \widetilde{\mathbf{R}}^3 : \alpha = \beta\gamma\}}.$$

If conversely  $\alpha, \beta, \gamma \in \widetilde{\mathbf{R}}$  satisfies  $\alpha = \beta\gamma$ , then

$$(\alpha, \beta, \gamma) \in \widetilde{\{(x, y, z) \in \mathbf{R}^3 : x = yz\}}.$$

Proof of (v). Assume  $I = (a; b)$  with  $a < b \in \mathbf{R}$ . Then  $\widetilde{I} = (\widetilde{a}; \widetilde{b})$  where  $\widetilde{a}, \widetilde{b} \in \widetilde{\mathbf{R}}$  denotes the constant germs equal to  $a$  and  $b$ . Indeed, if  $\gamma \in \widetilde{I}$ , there exists  $\epsilon > 0$  such that  $\gamma$  is defined on  $(0; \epsilon)$  and  $\gamma(t) \in (a; b)$  for  $t \in (0; \epsilon)$ . In particular  $\widetilde{a}(t) = a < \gamma(t) < b = \widetilde{b}(t)$  for  $t \in (0; \epsilon)$  so  $\gamma \in (\widetilde{a}; \widetilde{b})$ . The proof of the converse inclusion is similar.

Proof of (vi). An element of  $\widetilde{S}$  is a germ of continuous definable curve with image in  $S = \{s\}$ , so is equal to the function germ  $\widetilde{s}$  constant equal to  $s$ . □

**Remark 2.13.** The convex set given in example 2.3 is not a definable subset of  $\widetilde{\mathbb{R}} = \mathbb{R}((t^{1/\infty}))$  since it can not be described by a finite union of intervals.

### 3. CONTINUITY

We prove in this section that a continuous definable map between definable sets induces a continuous map between the spaces of definable arcs. Using Lojasiewicz inequality, we give a proof of this fact generalising the Hölder argument of Theorem 1.3. We propose another approach, studying more in details the transport of properties between the initial o-minimal structure and the new one constructed in section 2.4. These results are a geometric interpretation of the model-theoretical aspects developed in [2], Chapter 5.

**3.1. Łojasiewicz inequality in the closed and bounded case.** We recall Łojasiewicz inequality in the context of o-minimal structures, not necessarily polynomially bounded. We refer to [14] for the real case, and note that the real closed field case follows similarly.

Let  $\mathbf{R}$  be a real closed field and fix an o-minimal structure expanding it.

**Theorem 3.1.** *Let  $X \subset \mathbf{R}^n$  be a definable set, closed and bounded. Let  $\phi_1, \phi_2 : X \rightarrow \mathbf{R}$  be non negative continuous definable functions satisfying  $\phi_2^{-1}(0) \subset \phi_1^{-1}(0)$ . There exists a strictly increasing continuous definable function  $\rho : [0, +\infty) \rightarrow [0, +\infty)$  such that*

$$\forall x \in X, \quad \phi_2(x) \geq \rho \circ \phi_1(x).$$

**Remark 3.2.** In the polynomially bounded case, we may choose  $\rho$  of the form  $\rho(s) = c|s|^r$ , with  $c, r > 0$ .

We apply Łojasiewicz inequality to prove the continuity between the spaces of arcs in the closed and bounded case.

**Proposition 3.3.** *Let  $X$  and  $Y$  be definable sets, with  $X$  closed and bounded. Let  $f : X \rightarrow Y$  be a continuous definable map. Then  $\tilde{f} : \tilde{X} \rightarrow \tilde{Y}$  is continuous.*

**Remark 3.4.** In the polynomially bounded case, the map  $f$  is Hölder by Łojasiewicz inequality. In particular, in the case of the field of Puiseux series equipped with the  $t$ -adic topology, this result becomes clear.

*Proof of Proposition 3.3.* For  $\gamma \in \tilde{X}$ , there exists  $\epsilon$  such that  $\gamma$  admits a representative  $\gamma : [0, \epsilon) \rightarrow X$ . Let  $B \subset X$  be a closed neighbourhood of  $\gamma(0)$  in  $X$ . Define two functions  $\phi_1, \phi_2$  on  $B \times B$  by

$$\phi_1(x_1, x_2) = |f(x_1) - f(x_2)|$$

and

$$\phi_2(x_1, x_2) = |x_1 - x_2|.$$

Then  $\phi_1$  and  $\phi_2$  are continuous definable,  $\phi_2^{-1}(0) \subset \phi_1^{-1}(0)$  and therefore by Łojasiewicz inequality there exists a strictly increasing continuous definable function  $\rho : [0, +\infty) \rightarrow [0, +\infty)$  such that

$$\forall (x_1, x_2) \in B \times B, \quad \phi_2(x_1, x_2) \geq \rho \circ \phi_1(x_1, x_2),$$

which means

$$\forall (x_1, x_2) \in B \times B, \quad |x_1 - x_2| \geq \rho(|f(x_1) - f(x_2)|).$$

As a consequence,

$$\forall (x_1, x_2) \in B \times B, \quad |f(x_1) - f(x_2)| \leq \rho^{-1}(|x_1 - x_2|),$$

and  $\rho^{-1}$  is a definable continuous function with  $\rho^{-1}(0) = 0$ . In particular, for any arc  $\delta$  closed to  $\gamma$ , there exists a representative  $\delta : [0, \epsilon) \rightarrow X$  of  $\delta$  with value in  $B$  and therefore

$$\forall t \in [0, \epsilon), \quad |f \circ \gamma(t) - f \circ \delta(t)| \leq \rho^{-1}(|\gamma(t) - \delta(t)|),$$

which proves the continuity of  $\tilde{f}$ . □



**3.2. A generalised Łojasiewicz inequality.** As a first step, we slightly generalise Theorem 3.1 in order to get rid of the boundedness assumption of  $X$  in Proposition 3.3.

**Proposition 3.5.** *Let  $X \subset \mathbf{R}^n$  be a locally closed definable set. Let  $\phi_1, \phi_2 : X \rightarrow \mathbf{R}$  be non negative continuous definable functions satisfying  $\phi_2^{-1}(0) \subset \phi_1^{-1}(0)$ . There exist a strictly increasing continuous definable function  $\rho : [0, +\infty) \rightarrow [0, +\infty)$  and a positive continuous definable function  $c : X \rightarrow \mathbf{R}$  such that  $c|_{\phi_2^{-1}(0)} = 1$  and*

$$\forall x \in X, \quad \phi_2(x) \geq c(x)\rho \circ \phi_1(x).$$

*Proof.* Note first that we can find a definable homeomorphism mapping  $X$  onto a closed definable set, so that we can assume  $X$  closed. Let  $U$  denote a definable neighbourhood of  $\phi_1^{-1}(0)$  in  $X$  such that  $\phi_1|_U$  is bounded and  $\phi_1|_{U - \phi_1^{-1}(0)} : U - \phi_1^{-1}(0) \rightarrow \phi_1(U - \phi_1^{-1}(0))$  is proper. Such a neighbourhood  $U$  always exists; for example, let  $\phi_3$  denote the function on  $X$  measuring the distance from a fixed point of  $\phi_1^{-1}(0)$ . Then the set  $U = \{x \in X : \phi_1(x)\phi_3(x) \leq 1\}$  is convenient.

Let  $Y$  denote the image of  $U$  under the map  $(\phi_1, \phi_2) : X \rightarrow \mathbf{R}^2$ . Then  $Y$  is a closed definable subset of a sufficiently small closed neighbourhood of the origin in  $\mathbf{R}^2$  by the properness condition. Moreover  $Y \cap (\mathbf{R} \times \{0\}) = \{(0, 0)\}$  and therefore there exists a continuous definable function  $\tau$  on  $\text{Im } \phi_1|_U$  such that  $\tau^{-1}(0) = \{0\}$  and  $x_2 \geq \tau(x_1)$  for  $(x_1, x_2) \in Y$ , so that  $\phi_2 \geq \tau \circ \phi_1$  on  $U$ . One can find a strictly increasing continuous definable function  $\rho : [0, +\infty) \rightarrow [0, +\infty)$  such that  $\rho \leq \tau$  on  $\phi_1|_U(U)$ . Therefore  $\phi_2 \geq \rho \circ \phi_1$  on  $U$ .

To conclude, it suffices to choose a positive continuous definable function  $c$  so that  $c|_{\phi_2^{-1}(0)} = 1$  and  $c \leq \frac{\phi_2}{\rho \circ \phi_1}$  on  $X \setminus \{\phi_2^{-1}(0)\}$ .  $\square$

Let  $M$  be a definable set in some o-minimal structure. We define  $\widehat{M}$  by

$$\widehat{M} = \{\gamma \in \widetilde{M} : \gamma(0) \in M\}.$$

If  $h : M \rightarrow N$  is a definable map between definable sets  $M$  and  $N$ , define  $\widehat{h} : \widehat{M} \rightarrow \widehat{N}$  as the restriction of  $\widetilde{h}$  to  $\widehat{M}$ . Note that neither  $\widehat{M}$  nor  $\widehat{h}$  are definable.

**Corollary 3.6.** *Let  $X$  and  $Y$  be definable sets, with  $X \subset \mathbf{R}^n$  locally closed. Let  $f : X \rightarrow Y$  be a continuous definable map. Then  $\widehat{f} : \widehat{X} \rightarrow \widehat{Y}$  is continuous.*

*Proof.* Using proposition 3.5 and proceeding similarly to the proof of Proposition 3.3, we obtain that for  $\gamma, \delta \in \widehat{X}$ , there exist  $\epsilon > 0$  such that

$$\forall t \in [0, \epsilon), \quad |f \circ \gamma(t) - f \circ \delta(t)| \leq \rho^{-1}\left(\frac{|\gamma(t) - \delta(t)|}{c(\gamma(t), \delta(t))}\right).$$

Let  $U$  be defined by

$$U = \{(x_1, x_2) \in X^2 : c(x_1, x_2) > \frac{1}{2}\}.$$

Fixing  $\gamma$ , there exists  $l \in \mathbb{N}$  such that if  $|\gamma(t) - \delta(t)| < t^l$  for  $t \in [0, \epsilon)$ , then  $(\gamma(t), \delta(t)) \in U$  for  $t \in [0, \epsilon)$ . Therefore for  $\delta \in B(\gamma, t^l)$  we obtain

$$\forall t \in [0, \epsilon), \quad |f \circ \gamma(t) - f \circ \delta(t)| \leq \rho^{-1}(2|\gamma(t) - \delta(t)|)$$

which proves the continuity of  $\widehat{f}$ .  $\square$

We can refine the argument in order to deal with the non necessarily locally closed case as follows.

**Proposition 3.7.** *Let  $X \subset \mathbf{R}^n$  be a definable set. Let  $\phi_1, \phi_2 : X \rightarrow \mathbf{R}$  be non negative continuous definable functions satisfying  $\phi_2^{-1}(0) \subset \phi_1^{-1}(0)$ . Then there exist an increasing sequence of closed and bounded definable sets  $K_i$  in  $X$ , a strictly increasing continuous definable function  $\rho : [0, +\infty) \rightarrow [0, +\infty)$  and positive continuous definable function  $c_i : K_i \rightarrow \mathbf{R}$  such that  $\cup_i K_i = X$  and  $\phi_2 \geq c_i \rho \circ \phi_1$  on each  $K_i$ .*

**Remark 3.8.** Note in particular that the function  $\rho$  does not depend on the sets  $K_i$ .

*Proof of Proposition 3.7.* Set  $A = X \setminus (\overline{X} \setminus X)$ . By triangulating  $X$ , we obtain a closed definable subset  $X'$  of some Euclidean space and a surjective definable continuous map  $\tau : X' \rightarrow X$  such that  $\tau|_{\tau^{-1}(A)}$  is a homeomorphism onto  $A$ . It remains to apply Proposition 3.5 on  $X'$ .  $\square$

As a direct consequence, Corollary 3.6 can be generalised as follows.

**Corollary 3.9.** *Let  $X$  and  $Y$  be definable sets. Let  $f : X \rightarrow Y$  be a continuous definable map. Then  $\hat{f} : \hat{X} \rightarrow \hat{Y}$  is continuous.*

**3.3. Transport of properties.** We develop now the other approach based on the extension of fields from a real closed field  $\mathbf{R}$  to its field of germs of definable arcs  $\tilde{\mathbf{R}}$ . First, we note that the closedness and boundedness of a definable set are preserved under the extension.

**Lemma 3.10.** *Let  $A$  be a definable subset of a definable set  $B$ . Then  $A$  is closed in  $B$  if and only if  $\tilde{A}$  is closed in  $\tilde{B}$ .*

*Proof.* It is equivalent to prove that  $A$  is open in  $B$  if and only if  $\tilde{A}$  is open in  $\tilde{B}$ .

Assume  $A$  is open in  $B$ . Take  $\gamma \in \tilde{A}$ . There exists  $\epsilon > 0$  such that  $\gamma$  is well defined on  $(0; \epsilon)$  and  $\gamma((0; \epsilon)) \subset A$ . The openness of  $A$  enables to construct a strictly positive definable function germ by

$$r(t) = \sup\{s \in (0; 1] : |x - \gamma(t)| < s \Rightarrow x \in A\}$$

for  $t \in (0; \epsilon)$ . Reducing  $\epsilon$  if necessary, we can suppose  $r$  continuous. Then the open ball  $B(\gamma, r)$  is included in  $\tilde{A}$  and therefore  $\tilde{A}$  is open.

Conversely, to prove that  $A$  is open in  $B$  if  $\tilde{A}$  is open in  $\tilde{B}$ , let us assume that  $A$  is not open at  $a \in A$ . Then  $a$  belongs to the closure of  $B \setminus A$ , therefore by the curve selection lemma there exists  $\gamma : [0; \epsilon) \rightarrow B$  continuous definable with  $\gamma((0; \epsilon)) \subset B \setminus A$  and  $\gamma(0) = a$ . If  $\tilde{A}$  is open, let  $B(\tilde{a}; r) = \{\delta \in B : |\delta - \tilde{a}| < r\} \subset \tilde{A}$  be an open neighbourhood of the constant germ  $\tilde{a}$  equal to  $a$ , with  $r : (0; \epsilon_1) \rightarrow \mathbf{R}$  a strictly positive definable germ, and  $0 < \epsilon_1 < \epsilon$ . We get a contradiction if  $\gamma \in B(\tilde{a}; r)$ . Otherwise, we reparametrize  $\alpha$  as follows. Define  $\alpha : (0; \epsilon_1) \rightarrow \mathbf{R}$  by

$$\alpha(t) = \sup\{s \in (0; t) : |\gamma(s) - a| < \frac{r(t)}{2}\}$$

Then  $\alpha$  is definable and strictly positive since  $\gamma(0) = a$  and  $\gamma$  is continuous. We can moreover assume that  $\alpha$  is continuous, decreasing  $\epsilon_1$  if necessary. Then  $\gamma \circ \alpha((0; \epsilon_1)) \subset B \setminus A$  and  $\gamma \circ \alpha \in B(\tilde{a}; r)$ .  $\square$

**Lemma 3.11.** *Let  $A$  be a definable set in  $\mathbf{R}^n$ . Then  $A$  is bounded if and only if  $\tilde{A}$  is bounded in  $\tilde{\mathbf{R}}^n$ .*

*Proof.* Assume  $A$  is bounded in  $\mathbf{R}^n$ . Then there exist  $c_1, c_2 \in \mathbf{R}$  such that  $A \subset [c_1; c_2]^n$ . For  $\gamma \in \tilde{A}$ , there exists  $\epsilon$  such that  $\gamma(t) \in [c_1; c_2]^n$  for  $t \in (0; \epsilon)$ . In particular  $\gamma \in [\tilde{c}_1; \tilde{c}_2]^n$  and  $\tilde{A}$  is bounded.

Assume conversely that  $\tilde{A}$  is bounded, namely there exists  $b \in \tilde{\mathbf{R}}$  such that  $|\gamma| < b$  for any  $\gamma \in \tilde{A}$ . Then considering all constant functions in  $\tilde{A}$ , we see that  $A$  is bounded.  $\square$

The injectivity and surjectivity of a definable map are also preserved under the extension of fields.

**Lemma 3.12.** *Let  $f : X \rightarrow Y$  be a definable map. Then  $f$  is injective if and only if  $\tilde{f}$  is injective. Similarly,  $f$  is surjective if and only if  $\tilde{f}$  is surjective.*

*Proof.* Assume  $f$  is injective. Let  $\gamma_1, \gamma_2 \in \tilde{X}$  satisfies  $\tilde{f}(\gamma_1) = \tilde{f}(\gamma_2)$ . There exists  $\epsilon > 0$  such that  $\gamma_1$  and  $\gamma_2$  are defined on  $(0; \epsilon)$  and then for  $t \in (0; \epsilon)$  we have  $f(\gamma_1(t)) = f(\gamma_2(t))$ . Therefore  $\gamma_1(t) = \gamma_2(t)$  for  $t \in (0; \epsilon)$  by injectivity of  $f$  and  $\tilde{f}$  is injective.

Assume  $\tilde{f}$  injective. Let  $x_1, x_2 \in X$  satisfy  $f(x_1) = f(x_2)$ . If  $\tilde{x}_i$  denotes the constant curve equal to  $x_i$ , for  $i \in \{1, 2\}$ , then  $\tilde{f}(\tilde{x}_1) = \tilde{f}(\tilde{x}_2)$ . Therefore  $\tilde{x}_1 = \tilde{x}_2$  and thus  $x_1 = x_2$  and  $f$  is injective.

Assume now that  $f$  is surjective. Let  $\gamma \in \tilde{Y}$  and take  $\epsilon > 0$  such that  $\gamma$  is defined and continuous on  $(0; \epsilon)$ . Consider the set

$$A = \{(x, t) \in f^{-1}(\gamma((0; \epsilon))) \times (0; \epsilon) : f(x) = \gamma(t)\}.$$

Then  $A$  is a definable set, and we may assume  $A$  is bounded (embedding the ambient space  $\mathbf{R}^n$  of  $X$  in  $\mathbb{S}^n$  if necessary), so that  $(\overline{A} \setminus A) \cap (\mathbf{R}^n \times \{0\})$  is not empty. By the curve selection lemma, there exists  $\delta = (\delta_1, \delta_2) \in \tilde{A}$ , defined on  $(0; \epsilon')$  where  $\epsilon' < \epsilon$ , such that  $\delta((0; \epsilon')) \subset A$  and  $\delta$  admits a limit at 0, denoted by  $\delta(0)$ , and  $\delta(0) \in (\overline{A} \setminus A) \cap (\mathbf{R}^n \times \{0\})$ . Then  $f \circ \delta_1(t) = \gamma(\delta_2(t))$  by definition of  $A$ . Denote by  $\alpha \in \tilde{\mathbf{R}}$  a continuous definable function germ such that  $\delta_2 \circ \alpha(t) = t$ , obtained by Lemma 2.7. Then  $\delta_1 \circ \alpha \in \tilde{X}$  is an inverse of  $\gamma$  by  $\tilde{f}$  and  $\tilde{f}$  is surjective.

Assume finally that  $\tilde{f}$  is surjective. Take  $y \in Y$ . Then the constant curve  $\tilde{y}$  equal to  $y$  has a pre-image  $\gamma$  by  $\tilde{f}$ . There exists  $\epsilon > 0$  such that  $\gamma$  is well defined on  $(0; \epsilon)$ ,  $\gamma((0; \epsilon)) \subset X$  and  $f(\gamma(t)) = y$ . Therefore  $f$  is surjective.  $\square$

**3.4. Continuity.** Let  $\mathbf{R}$  be a real closed field and consider an o-minimal structure expanding  $\mathbf{R}$ . A definable mapping  $f : X \rightarrow Y$  between definable sets  $X, Y$  is said continuous if it is continuous for the Euclidean topology inherited to the real closed field.

**Proposition 3.13.** *Let  $f : X \rightarrow Y$  be definable and  $Y$  be bounded. Then  $f$  is continuous if and only if  $\text{graph } f$  is closed in  $X \times \overline{Y}$ .*

*Proof.* Assume  $f$  is continuous. Let  $(a, b)$  be in the closure of the graph of  $f$  in  $X \times \overline{Y}$ . By the curve selection lemma there exists a continuous curve  $\gamma = (\gamma_1, \gamma_2) : [0; 1] \rightarrow X \times \overline{Y}$  with  $\gamma(0) = (a, b)$  and  $\gamma((0; 1]) \subset \text{graph } f$ . Then for  $t \in (0; 1]$  we

have  $f \circ \gamma_1(t) = \gamma_2(t)$ . By continuity we obtain, passing to the limit as  $t$  goes to zero, that  $f(a) = b$ . In particular the graph of  $f$  is closed in  $X \times \overline{Y}$ .

Conversely, assume  $f$  is not continuous at  $a \in X$ . For  $\epsilon > 0$ , define  $\delta_\epsilon$  to be the supremum on  $x \in X$  with  $|x - a| \leq \epsilon$  of  $|f(x) - f(a)|$ . Then  $(0, a)$  is in the closure of the definable set

$$\cup_{\epsilon > 0} \{\epsilon\} \times \{x \in X : |x - a| \leq \epsilon \text{ and } |f(x) - f(a)| \geq \frac{\delta_\epsilon}{2}\}.$$

By the o-minimal curve selection lemma, there exists a continuous definable curve  $\gamma : [0; 1] \rightarrow X$  such that  $\gamma(0) = a$  and  $f \circ \gamma$  is not continuous at 0.

Denote by  $\Gamma$  the image  $\gamma((0; 1])$  of  $\gamma$  and by  $A$  the set

$$\{(x, f(x)) : x \in \Gamma\} \subset X \times Y.$$

Then  $\overline{A} \setminus A$  is not empty because  $Y$  is bounded. It is then equal to a point of the form  $(a, b)$  with  $b \neq f(a)$ , therefore the graph of  $f$  is not closed in  $X \times \overline{Y}$ .  $\square$

**Corollary 3.14.** *Let  $f : X \rightarrow Y$  be a continuous definable map. Then  $\tilde{f} : \tilde{X} \rightarrow \tilde{Y}$  is continuous.*

*Proof.* Without loss of generality we can assume that  $Y$  is bounded (as  $Y \subset \mathbf{R}^n \subset \mathbb{S}^n \subset \mathbf{R}^{n+1}$ ). Now the set  $\tilde{Y}$  is bounded by Lemma 3.11, and moreover  $\overline{\tilde{Y}} = \tilde{\overline{Y}}$ . Finally the graph of  $\tilde{f}$  is closed in  $\tilde{X} \times \overline{\tilde{Y}}$  by Lemma 3.10 since the graph of  $f$  is closed in  $X \times \overline{Y}$  by Proposition 3.13.  $\square$

**Example 3.15.** As a illustrative example, consider the Whitney umbrella  $W$  in  $\mathbb{R}^3$  defined by  $x^2z = y^2$ . Its two-dimensional part is the image of  $\mathbb{R}^2$  by the map  $h : (u, v) \mapsto (u, uv, v^2)$ , giving a resolution of the singularities of  $W$ . Any positive element of the  $z$ -axis has two pre-images by  $h$ , and so  $h$  defines a semi-algebraic bijection between the semi-algebraic sets  $(\mathbb{R}^2 \setminus \{u = 0\}) \cup \{(0, 0)\}$  and  $(W \setminus \{x = y = 0\}) \cup \{(0, 0, 0)\}$ , which are not even locally closed. Corollary 3.14 enables to assert that nevertheless,  $h$  induces a homeomorphism between the associated spaces of arcs.

We recover the results in section 1.2 as a consequence of Corollary 3.14 by considering the field of real numbers with the o-minimal structure given by globally subanalytic sets. For instance for Theorem 1.5:

**Corollary 3.16.** *Let  $h : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a subanalytic homeomorphism. Then the map  $\tilde{h} : \mathbb{R}\{t^{1/\infty}\}^n \rightarrow \mathbb{R}\{t^{1/\infty}\}^n$  is a homeomorphism with respect to the  $t$ -adic topology. If moreover we assume that  $h$  and  $h^{-1}$  are arc-analytic, then  $h_* : \mathcal{A}_0(\mathbb{R}^n) \rightarrow \mathcal{A}_0(\mathbb{R}^n)$  is a homeomorphism.*

*Proof.* The continuity of  $\tilde{h}$  follows from Corollary 3.14. Moreover  $h_*$  is well-defined by arc-analyticity of  $h$  and  $h^{-1}$ , and so  $h_*$  is continuous by restriction.  $\square$

**Lemma 3.17.** *Let  $C \subset \mathbf{R}^n$  be a closed and bounded definable subset of  $\mathbf{R}^n$  and  $g : C \rightarrow \mathbf{R}$  be a strictly positive definable function. There exists  $r \in \mathbf{R}$  with  $r > 0$  such that  $g \geq r$ .*

*Proof.* Assume the conclusion does not hold. Then

$$A = \{(t, c) \in (0; 1] \times C : g(c) < t\}$$

is a non empty definable subset of  $[0; 1] \times C$ , with a non empty boundary. Therefore there exists a continuous definable function germ  $\delta = (\delta_1, \delta_2) : [0; \epsilon) \rightarrow [0; 1] \times C$  with  $\delta((0; \epsilon)) \subset A$  and  $\delta_1(0) = 0$  by the curve selection lemma. Note that  $\delta_2(t)$  admits a limit  $x$  as  $t$  goes to zero since  $C$  is bounded, and the limit belongs to  $C$  since  $C$  is closed.

As  $\delta((0; \epsilon)) \subset A$  the following inequalities hold:

$$0 < g \circ \delta_2(t) < \delta_1(t)$$

for  $t \in (0; \epsilon)$ , and passing to the limit as  $t$  goes to 0 we obtain  $0 \leq g(x) \leq 0$ , in contradiction with the assumption on  $g$ .  $\square$

**Proposition 3.18.** *Let  $X$  and  $Y$  be bounded and closed definable sets and  $f : X \rightarrow Y$  be a continuous definable function. Then  $f$  is uniformly continuous.*

*Proof.* Let  $\epsilon \in \mathbf{R}$  such that  $\epsilon > 0$ . As  $f$  is continuous, we may define for  $x \in X$  a definable function  $g$  on  $X$  by

$$g(x) = \sup\{s \in (0; 1] : a \in X, |x - a| < s \Rightarrow |f(x) - f(a)| < \epsilon\}.$$

We conclude applying Lemma 3.17 to  $g$ .  $\square$

We recover by this way Theorem 1.4.

**Corollary 3.19.** *Let  $M \subset \mathbb{R}^m$  and  $N \subset \mathbb{R}^n$  be compact analytic manifolds and  $h : M \rightarrow N$  be a subanalytic homeomorphism. Then  $\tilde{h} : \tilde{M} \rightarrow \tilde{N}$  is a uniformly continuous homeomorphism. If moreover  $h$  and  $h^{-1}$  are arc-analytic, then  $h_* : \mathcal{A}(M) \rightarrow \mathcal{A}(N)$  is a uniformly continuous homeomorphism.*

*Proof.* The uniform continuity of  $\tilde{h}$  follows from Corollary 3.14 and Proposition 3.18. Then  $h_*$  is well-defined by arc-analyticity of  $h$  and  $h^{-1}$ , and finally  $h_*$  is uniformly continuous by restriction.  $\square$

Actually, we can even drop the compactness condition in Corollary 3.19 as follows.

**Proposition 3.20.** *Let  $M \subset \mathbb{R}^m$  and  $N \subset \mathbb{R}^n$  be globally subanalytic manifolds and  $h : M \rightarrow N$  be a globally subanalytic homeomorphism. Then  $\hat{h} : \hat{M} \rightarrow \hat{N}$  is a uniformly continuous homeomorphism. If moreover  $h$  and  $h^{-1}$  are arc-analytic, then  $h_* : \mathcal{A}(M) \rightarrow \mathcal{A}(N)$  is a uniformly continuous homeomorphism.*

The proof is a direct consequence of the following lemma.

**Lemma 3.21.** *Let  $f : X \rightarrow Y$  be a continuous definable map between closed definable subsets of  $\mathbf{R}^n$ . Then  $\hat{f} : \hat{X} \rightarrow \hat{Y}$  is uniformly continuous.*

*Proof.* Let  $a \in \tilde{R}$  be an element larger than any element of  $\mathbf{R}$ . since  $\tilde{X} \cap [-a, a]^n$  is closed and bounded, the result follows from Proposition 3.18.  $\square$

## REFERENCES

- [1] J. Bochnak, M. Coste, M.F. Roy, Real algebraic geometry, Springer, 1998
- [2] M. Coste, *An introduction to o-minimal geometry*, Dip. Mat. Univ. Pisa, Dottorato di Ricerca in Matematica, Istituti Editoriali e Poligrafici Internazionali, Pisa 2000
- [3] L. van den Dries, *Tame Topology and O-minimal Structures*, London Math. Soc. Lecture Note 248, Cambridge Univ. Press 1998
- [4] G. Fichou, *Motivic invariants of Arc-Symmetric sets and Blow-Nash Equivalence*, Compositio Math. 141 (2005) 655-688

- [5] G. Fichou, M. Shiota, *On almost Blow-analytic equivalence*, Proceedings of London Math. Soc. 103 (3) (2011) 676-709
- [6] G. Fichou, M. Shiota, *Analytic equivalence of normal crossing functions on a real analytic manifold*, Proceedings of London Math. Soc. 104 (6) (2012) 1121-1170
- [7] T. Fukui, *Seeking invariants for blow-analytic equivalence*, Compositio Math. 105 (1997), no. 1, 95-108
- [8] T. Fukui, K. Kurdyka, A. Parusiński, *Inverse Function Theorems for Arc-analytic Homeomorphisms*, arXiv:1003.0826, march 2010
- [9] T. Fukui, L. Paunescu, *On blow-analytic equivalence*, Arc spaces and additive invariants in real algebraic geometry, Panoramas et Synthèses, SMF 26 (2008) 87-125
- [10] S. Koike, A. Parusiński, *Motivic-type Invariants of Blow-analytic Equivalence*, Ann. Institut Fourier 53 , 7(2003), 2061-2104
- [11] S. Koike, A. Parusiński, *Blow-analytic equivalence of two variable real analytic function germs*, J. Algebraic Geom. 19 (2010), no. 3, 439-472
- [12] T.-C. Kuo, *On classification of real singularities*, Invent. Math. 82 (1985), no. 2, 257-262
- [13] K. Kurdyka, *Ensembles semi-algébriques symétriques par arcs*, Math. Ann. 282, 445-462 (1988)
- [14] K. Kurdyka, *On gradients of functions definable in o-minimal structures*, Ann. Inst. Fourier 48 (1998), no. 3, 769-783
- [15] W. Whitney, *Analytic extensions of differentiable functions defined in closed sets*, Trans. Amer. Math. Soc., 36 (1934), 63-89

IRMAR (UMR 6625), UNIVERSITÉ DE RENNES 1, CAMPUS DE BEAULIEU, 35042 RENNES CEDEX, FRANCE

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, CHIKUSA, NAGOYA, 464-8602, JAPAN