



HAL
open science

In vitro activity of -chlorotaurine (NCT) in combination with NHCl against

Ursula Fürnkranz, Markus Nagl, Waldemar Gottardi, Michael Duchêne, Horst Aspöck, Julia Walochnik

► **To cite this version:**

Ursula Fürnkranz, Markus Nagl, Waldemar Gottardi, Michael Duchêne, Horst Aspöck, et al.. In vitro activity of -chlorotaurine (NCT) in combination with NHCl against. *International Journal of Antimicrobial Agents*, 2011, 37 (2), pp.171. 10.1016/j.ijantimicag.2010.09.006 . hal-00659902

HAL Id: hal-00659902

<https://hal.science/hal-00659902>

Submitted on 14 Jan 2012


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: In vitro activity of *N*-chlorotaurine (NCT) in combination with NH₄Cl against *Trichomonas vaginalis*

Authors: Ursula Fűrnkranz, Markus Nagl, Waldemar Gottardi, Michael Duchêne, Horst Aspöck, Julia Walochnik



PII: S0924-8579(10)00416-4
DOI: doi:10.1016/j.ijantimicag.2010.09.006
Reference: ANTAGE 3432

To appear in: *International Journal of Antimicrobial Agents*

Received date: 28-6-2010
Revised date: 26-8-2010
Accepted date: 18-9-2010

Please cite this article as: Fűrnkranz U, Nagl M, Gottardi W, Duchêne M, Aspöck H, Walochnik J, In vitro activity of *N*-chlorotaurine (NCT) in combination with NH₄Cl against *Trichomonas vaginalis*, *International Journal of Antimicrobial Agents* (2010), doi:10.1016/j.ijantimicag.2010.09.006

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

In vitro activity of *N*-chlorotaurine (NCT) in combination with NH₄Cl against *Trichomonas vaginalis*

Ursula Fűrnkranz ^a, Markus Nagl ^b, Waldemar Gottardi ^b, Michael Duchêne ^a, Horst Aspöck ^a, Julia Walochnik ^{a,*}

^a *Institute of Specific Prophylaxis and Tropical Medicine, Center for Pathophysiology, Infectiology and Immunology, Medical University of Vienna, Kinderspitalgasse 15, 1090 Vienna, Austria*

^b *Department of Hygiene, Microbiology and Social Medicine, Division of Hygiene and Medical Microbiology, Innsbruck Medical University, Fritz-Pregl-Strasse 3, 6020 Innsbruck, Austria*

ARTICLE INFO

Article history:

Received 28 June 2010

Accepted 18 September 2010

Keywords:

Trichomonas vaginalis

Susceptible

N-Chlorotaurine

Oxidant

In vitro

* Corresponding author. Tel.: +43 1 40490 79446; fax: +43 1 40490 79435.

E-mail address: julia.walochnik@meduniwien.ac.at (J. Walochnik).

Accepted Manuscript

ABSTRACT

Trichomoniasis, caused by the protozoan *Trichomonas vaginalis*, is usually treated with metronidazole, however resistance is on the rise. In this study, *N*-chlorotaurine (NCT), a new endogenous mild active chlorine compound for topical use, killed *T. vaginalis* in vitro within 15 min of treatment at a concentration of 55 mM (1%), which is well tolerated by human tissue. The activity of NCT was further enhanced by addition of ammonium chloride (NH₄Cl). A combination of 5.5 mM (0.1%) NCT plus 19 mM (0.1%) NH₄Cl killed 100% of trichomonads within 5 min.

1. Introduction

Trichomonas vaginalis is the causative agent of trichomoniasis, the most common non-viral sexually transmitted disease worldwide. Typical symptoms are vaginitis and cervicitis in women; in men the infection can lead to chronic prostatitis syndrome [1]. Metronidazole is the drug of choice for trichomoniasis, however treatment during pregnancy is problematic [2] and resistance is increasing [3,4].

In a previous study, we demonstrated the amoebicidal effect of *N*-chlorotaurine (NCT) (Cl-HN-CH₂-CH₂-SO₃Na), thereby establishing the activity of this compound against protozoa [5]. NCT, the *N*-chloro derivative of the amino acid taurine produced by granulocytes and monocytes during the oxidative burst, is a long-lived oxidant that can be synthesised chemically [6] and applied topically for treatment of infections [7]. It has broad-spectrum microbicidal activity against bacteria, viruses, fungi and protozoa [5,7–9]. Phase I and II clinical studies have revealed high tolerability and therapeutic effects of regular applications of 55 mM (1%) NCT in the eye, sinuses, oral cavity, outer ear canal, skin ulcerations and urinary bladder [7,10].

The antimicrobial activity of NCT against bacteria [8], fungi [8,9] and free-living amoebae [5] can be enhanced markedly by addition of ammonium chloride (NH₄Cl) owing to the formation of monochloramine (NH₂Cl) in equilibrium by transfer of the active chlorine atom [6–9]. NH₂Cl is more lipophilic than NCT and penetrates microorganisms more rapidly. For special indications, e.g. treatment of viral conjunctivitis, addition of NH₄Cl to NCT might be an advantage [7].

The aim of the current study was to investigate the activity of NCT and its combination with NH_4Cl against *T. vaginalis* in vitro.

2. Material and methods

A metronidazole-susceptible (ATCC 30001) and a metronidazole-resistant (ATCC 50138) strain of *T. vaginalis* were investigated for their susceptibilities to NCT. Both strains were grown under microaerophilic conditions in trypticase–yeast extract–maltose (TYM) medium [11] at 37 °C and were subcultured every 72 h.

NCT was produced synthetically at Innsbruck Medical University (Innsbruck, Austria) and was used alone as well as in combination with NH_4Cl as previous studies had revealed a high synergistic potential for this substance [6,8]. NCT (molecular weight 181.57 g/mol) [6] was used at final concentrations of between 5.5 mM (0.1%) and 55 mM (1%) dissolved in either TYM or phosphate-buffered saline (PBS). Trichomonads were treated for 5, 15 and 60 min at 37 °C. Short intervals were chosen because stability testing revealed that the oxidative capacity of 55 mM NCT is reduced by 15% after 1 min, by 90% after 1 h and by >95% after 2 h in TYM. NH_4Cl was purchased from Sigma-Aldrich (Vienna, Austria) and was used in the molar ratio 1:3.4 and 1:29 (NCT: NH_4Cl) as in previous studies [5,8]. Control experiments revealed no antiparasitic effects of NH_4Cl alone. Additional experiments with 5.5 mM (0.1%) NCT plus 19 mM (0.1%) NH_4Cl (1:3.4) in PBS instead of TYM medium were performed to investigate the efficacy in the absence of chlorine consumption caused by unspecific reactions of NCT with the culture medium.

Viability of trichomonads was followed by phase contrast microscopy and trypan blue (0.4%) staining, and 50% and 90% effective concentrations (EC_{50} and EC_{90} values) were calculated using non-linear regression models. Cytotoxicity of NCT against vaginal epithelial cells (ATCC; CRL 2616) was investigated using the EZ4U test (Biomedica GmbH, Vienna, Austria), which is a tetrazolium reduction assay. All experiments were carried out in triplicate and were repeated in two independent set-ups. For each experiment, a control series without NCT and additives was included.

Single-factor analysis of variance (ANOVA) and Tukey's HSD test (SPSS 14.0; SPSS Inc., Chicago, IL) were used for statistical analyses. P -values of <0.05 were considered significant.

3. Results and discussion

NCT showed cytotoxic effects against both strains of *T. vaginalis* tested. Treatment with 55 mM NCT (1%), a concentration well tolerated by all human tissues tested so far [7], led to 100% killing of both strains tested within 15 min in TYM medium (Fig. 1). At lower concentrations in this medium, strain ATCC 30001 was more susceptible than strain ATCC 50138. The EC_{50} of NCT was 9.0 mM for strain ATCC 30001, whilst it was 28.7 mM for strain ATCC 50138 after 1 h of treatment. EC_{90} values were 9.8 mM and 36.7 mM, respectively.

The activity of NCT was markedly enhanced by the addition of NH_4Cl , explained by formation of the small and more lipophilic monochloramine (NH_2Cl), which penetrates pathogens better than NCT and leads to faster oxidation of intracellular proteins [6–8]. In strain ATCC 30001, only one-third of the concentration of NCT was needed to

achieve 100% killing when NH_4Cl was added, i.e. 17 mM (0.3%) NCT plus 57 mM (0.3%) NH_4Cl (Fig. 1); moreover, the killing time was reduced to 5 min. This effect was even greater when experiments were carried out in PBS instead of culture medium. In PBS, co-treatment with 5.5 mM (0.1%) NCT plus 19 mM (0.1%) NH_4Cl also resulted in complete killing of trichomonads within 5 min (data not shown). Inactivation of NCT and the formed NH_2Cl by reaction with constituents of the TYM medium must be assumed to contribute significantly to delayed killing in this medium, particularly at concentrations below 10 mM. Such chlorine consumption in the presence of organic material, in particular reducing thio compounds, is well known [6,7]. It may also explain the relatively sharp threshold between highly active and inactive concentrations of NCT (Fig. 1). In strain ATCC 50138, higher concentrations of NH_4Cl (molar ratio 1:29) were necessary to achieve enhancement of activity in TYM medium.

The metronidazole-sensitive strain (ATCC 30001) was more susceptible to NCT than the metronidazole-resistant strain (ATCC 50138), and this difference was statistically significant ($P < 0.001$). An approximately three- to four-fold concentration of NCT required to kill 50% of the metronidazole-sensitive trichomonads was required to achieve the same killing rate in the metronidazole-resistant strain. Although the modes of action are different, since both metronidazole and NCT produce oxidative stress one could hypothesise that the more resistant strain has a better anti-oxidative protection system. This may be the case, but at the applied millimolar concentrations of NCT such mechanisms would not be sufficient. Since penetration of the pathogen rather than oxidation of the surface is the decisive step for killing by NCT [7], we

assume that several factors, particularly differences in the membrane composition, may play a major role as has already been observed in other microorganisms [12].

In vivo studies have shown that 55 mM NCT (1%) is well tolerated by human mucosal tissue [7]. The in vitro tolerability of human cells in cell culture to NCT is known to be significantly (ca. 50–100-fold) lower [13–15]. However, vaginal epithelial cells were observed to be relatively resistant to NCT in the current study. The EC_{50} of NCT for these cells in vitro was 5.5 mM and the survival rate was 25% after treatment with 55 mM NCT for 1 h (Fig. 2). Tolerability to NCT was altogether significantly higher than to metronidazole, the standard drug for treatment of trichomoniasis. Increased tolerability of human cells to the agent in vivo compared with that in vitro can be explained by the fact that within tissues the cells are coherently organised in a structure affording better protection to oxidative agents. This has already been proven with human skin in vivo as well as human epidermis carcinoma cells in vitro [10].

The stability of the aqueous solution of NCT depends on temperature. At 37 °C its oxidation capacity decreases to zero within 3 weeks, whereas it remains at 75% after storage at room temperature for 2 months [6]. At sites of inflammation, oxidation capacity decreases more quickly owing to halogen-consuming proteins. After 15 min the oxidation capacity is reduced by 30–50% and after 2 h by up to 90% [8].

However, as trichomonads are killed within 15 min, and even faster with the addition of NH_4Cl , these decomposition rates are not likely to affect a possible therapeutic approach against trichomoniasis. Moreover, since human exudates demonstrated enhanced rather than decreased microbicidal activity of 0.5–1% NCT after topical

application [8,9], it may be expected that NCT is also microbicidal against *T. vaginalis* in vivo after topical application.

In conclusion, we have shown that NCT has strong trichomonacidal effects that are further enhanced by the addition of NH₄Cl. Further investigations of its utility in *T. vaginalis* infections are required.

Acknowledgments

The authors thank Susanne Glöckl, Iveta Häfeli and Jacek Pietrzak (Department of Medical Parasitology, Medical University of Vienna, Vienna, Austria) for excellent technical assistance.

Funding

This study was supported partly by the International Relations Office of the Medical University of Vienna (Austria). The contributions of MN and WG were supported by the Austrian Science Fund (FWF) (grant no. L313-B13).

Competing interests

None declared.

Ethical approval

Not required.


References

- [1] Skerk V, Krhen I, Schonwald S, Cajic V, Markovinovic L, Roglic S, et al. The role of unusual pathogens in prostatitis syndrome. *Int J Antimicrob Agents* 2004;24(Suppl 1):S53–6.
- [2] Kazy Z, Puho E, Czeizel AE. Teratogenic potential of vaginal metronidazole treatment during pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2005;123:174–8.
- [3] Dunne RL, Dunn LA, Upcroft P, O'Donoghue PJ, Upcroft JA. Drug resistance in the sexually transmitted protozoan *Trichomonas vaginalis*. *Cell Res* 2003;13:239–49.
- [4] Schwebke JR, Barrientes FJ. Prevalence of *Trichomonas vaginalis* isolates with resistance to metronidazole and tinidazole. *Antimicrob Agents Chemother* 2006;50:4209–10.
- [5] Fürnkranz U, Nagl M, Gottardi W, Köhler M, Aspöck H, Walochnik J. Cytotoxic activity of *N*-chlorotaurine on *Acanthamoeba* spp. *Antimicrob Agents Chemother* 2008;52:470–6.
- [6] Gottardi W, Nagl M. Chemical properties of *N*-chlorotaurine sodium, a key compound in the human defense system. *Arch Pharm (Weinheim)* 2002;335:411–21.
- [7] Gottardi W, Nagl M. *N*-Chlorotaurine, a natural antiseptic with outstanding tolerability. *J Antimicrob Chemother* 2010;65:399–409.
- [8] Nagl M, Gottardi W. Enhancement of the bactericidal efficacy of *N*-chlorotaurine of inflammation samples and selected N–H compounds. *Hyg Med* 1996;21:597–605.
- [9] Nagl M, Lass-Flörl C, Neher A, Gunkel A, Gottardi W. Enhanced fungicidal activity of *N*-chlorotaurine in nasal secretion. *J Antimicrob Chemother* 2001;47:871–4.

- [10] Nagl M, Nguyen VA, Gottardi W, Ulmer H, Höpfl R. Tolerability and efficacy of *N*-chlorotaurine in comparison with chloramine T for the treatment of chronic leg ulcers with a purulent coating: a randomized phase II study. *Br J Dermatol* 2003;149:590–7.
- [11] Clark CG, Diamond LS. Methods for cultivation of luminal parasitic protists of clinical importance. *Clin Microbiol Rev* 2002;15:329–41.
- [12] Beach DH, Holz GG Jr, Anekwe GE. Lipids of *Leishmania* promastigotes. *J Parasitol* 1979;65:201–16.
- [13] Kontny E, Grabowska A, Kowalczewski J, Kurowska M, Janicka I, Marcinkiewicz J, et al. Taurine chloramine inhibition of cell proliferation and cytokine production by rheumatoid arthritis fibroblast-like synoviocytes. *Arthritis Rheum* 1999;42:2552–60.
- [14] Marcinkiewicz J, Nowak B, Grabowska A, Bobek M, Petrovska L, Chain B. Regulation of murine dendritic cell functions in vitro by taurine chloramine, a major product of the neutrophil myeloperoxidase–halide system. *Immunology* 1999;98:371–8.
- [15] Park E, Levis WR, Quinn MR, Park SY, Schuller-Levis GB. Regulation of nitric oxide induced by mycobacterial lipoarabinomannan in murine macrophages: effects of interferon- β and taurine-chloramine. *Int J Lepr Other Mycobact Dis* 2000;68:444–51.

Fig. 1. Treatment of *Trichomonas vaginalis* ATCC 30001 with *N*-chlorotaurine (NCT) and co-treatment with NH₄Cl in trypticase–yeast extract–maltose (TYM) medium: ▼, 55 mM (1%) NCT; ▲, 10 mM (0.18%) NCT; ○, 5.5 mM (0.1%) NCT; ◆, 17 mM (0.3%) NCT + 57 mM (0.3%) NH₄Cl; - - * - -, control. Mean value of three independent experiments; the standard deviation ranged between 0.14 and 0.21 log₁₀ for values exceeding 4 log₁₀ (not shown). *P* < 0.01 versus control for all zero survival values.

Fig. 2. Survival of vaginal epithelial cells treated for 1 h with *N*-chlorotaurine (NCT) or metronidazole.



Script

