



**A ^1H , ^{13}C , ^{31}P and ^{15}N NMR study of
(pyrrolidine-2,2-diyl)bisphosphonic acid,
tetraalkyl(pyrrolidine-2,2-diyl)bisphosphonates and
acyclic tetraethyl bisphosphonates**

Gilles Olive, Marcel H.P. van Genderen

► **To cite this version:**

Gilles Olive, Marcel H.P. van Genderen. A ^1H , ^{13}C , ^{31}P and ^{15}N NMR study of (pyrrolidine-2,2-diyl)bisphosphonic acid, tetraalkyl(pyrrolidine-2,2-diyl)bisphosphonates and acyclic tetraethyl bisphosphonates. *Magnetic Resonance in Chemistry*, 2000, 38 (5), pp.379-381. hal-00660649

HAL Id: hal-00660649

<https://hal.science/hal-00660649>

Submitted on 17 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A ^1H , ^{13}C , ^{31}P and ^{15}N NMR study of (pyrrolidine-2,2-diyl)bisphosphonic acid, tetraalkyl(pyrrolidine-2,2-diyl)bisphosphonates and acyclic tetraethyl bisphosphonates

Gilles Olive[†], Marcel H.P. van Genderen^{‡*}

[†] Laboratory of Polymer Chemistry and Coatings Technology

[‡] Laboratory of Macromolecular and Organic Chemistry

Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands.

Tel: +31-40-2473139. Fax: +31-40-2451036.

* To whom correspondence should be addressed.

ABSTRACT:

A multinuclear NMR study (^1H , ^{13}C , ^{31}P , ^{15}N) was performed on a series of new cyclic pyrrolidine bisphosphonates and acyclic bisphosphonates. Values are reported and discussed for the chemical shifts and coupling constants of the various nuclei.

KEYWORDS: NMR; ^1H NMR; ^{13}C NMR; ^{31}P NMR; ^{15}N NMR; tetraalkyl bisphosphonates; bisphosphonic acid

INTRODUCTION

Gem-bisphosphonates or *gem*-bisphosphonic acids can complex well with calcium and magnesium.¹ As a result their use in various areas, for instance in the medical field², as plant growth regulators³ and in the nuclear industry², has been known for some time. We report here a multinuclear NMR study, in particular nitrogen-15, of new tetraalkyl bisphosphonates in a cyclic series and acyclic forms. It is interesting that these compounds can be labeled with ¹⁵N for a direct study in biological material, which has been a considerable problem for several decades.⁴

EXPERIMENTAL

The synthesis of compounds **1** to **6** (Figure 1) has been described^{5,6} and the synthesis and full characterization of **7** and **8** will be published in a forthcoming paper.

¹H, ¹³C and ³¹P NMR spectra were measured at 25 °C on Bruker AC-100, AC-200 and AMX-400 spectrometers. ¹H spectra were recorded at 100, 200 and 400 MHz. ¹³C spectra were recorded at 25.18, 50.32 and 100.65 MHz. ³¹P spectra were recorded at 40.54 MHz. ¹⁵N NMR spectra were measured at 25 °C on a Varian Unity Inova 500 spectrometer at 50.65 MHz. In all cases, a field-frequency lock on the solvent ²H signal was used. Chemical shifts are reported in ppm downfield from the standards, which were: TMS ($\delta = 0$) or residual HDO in aqueous NaOD ($\delta = 4.81$) for ¹H spectra, TMS ($\delta = 0$) for ¹³C spectra, external 85% H₃PO₄ ($\delta = 0$) for ³¹P spectra, and external CD₃NO₂ ($\delta = 0$) for ¹⁵N spectra. Samples were prepared in C₆D₆, CDCl₃ or aqueous NaOD (50 mg Na/ml D₂O), in concentrations of 50-100 mg/ml for ¹H, ¹³C and ³¹P spectra in 5-mm tubes, and in a 80:20 (v/v) ratio of compound and solvent for natural abundance ¹⁵N measurements in 10-mm tubes. Acquisition parameters for ¹H spectra were: pulse width 2-6 μ s (45° pulse), 16 transients, 4 s acquisition time, 2 s relaxation delay, spectral width 10 ppm (1-4 kHz), 8-32 K data points, 0.24 Hz resolution. ¹H spectra were obtained with line broadening (lb = 0.1). Acquisition parameters for ¹³C spectra were: pulse width 3 μ s (45° pulse), 100 transients, 0.7-1.4 s acquisition time, 5 s relaxation delay, spectral width 6-24 kHz, 16-32 K data points, 0.74-1.45 Hz resolution. ¹³C spectra

were obtained with line broadening ($lb = 1$) and 1H broadband decoupling. Acquisition parameters for ^{31}P spectra were: pulse width 5 μs (90° pulse), 80 transients, 5 s acquisition time, 3 s relaxation delay, spectral width 16 kHz, 16 K data points, 2 Hz resolution. ^{31}P spectra were obtained with 1H broadband decoupling. Acquisition parameters for ^{15}N spectra were: pulse width 10 μs (90° pulse: 20.5 μs), 400 transients, 1.9 s acquisition time, 10 s relaxation delay, spectral width 8 kHz, 30000 data points, 0.5 Hz resolution. ^{15}N spectra were obtained with line broadening ($lb = 1$) and inverse-gated 1H broadband decoupling.



RESULTS AND DISCUSSION

For all bisphosphonates, except **1**, the ^{15}N spectrum (Table 1) shows a triplet, indicating that the two phosphorus nuclei are equivalent. This means that the ring pseudorotation is very fast for **2** to **5**, as previously reported for the nitroxide⁷ (confirmed by the fact that we obtained triplets in ^{13}C NMR). The two-bond coupling constants between phosphorus and nitrogen are in the range 2-9 Hz, which agrees with the few instances of $^2J_{\text{NP}}$ reported in the literature.^{8,9} It is evident that the linear compounds **6**, **7** and **8** have smaller coupling constants (6.3, 6.1 and 5.2 Hz respectively) than the cyclic structures.


Except for **1** (due to a different solvent) and for the picrate of **2**, the ^{15}N chemical shifts of the cyclic forms **2-4** are all similar (max. change 0.5 ppm) and no correlation with the structure was found. Cyclic **5** has a different chemical shift because it is a tertiary amine (the donating methyl group induces an upfield shift). All ^{15}N chemical shifts (Table 1) for amines **2-8** are very different from the starting materials (respectively -264.3 ppm for the pyrrolidine-2-one, -271.9 ppm for *N*-methyl pyrrolidone, -262.1 ppm for the precursor of **6**, -242.0 ppm for the precursor of **7**, and -244.5 ppm for the precursor of **8**).

It is also not possible to make a correlation between the structure and the chemical shift in the ^{31}P spectra (Table 1). In all cases we see only one line in the ^{31}P spectra. Hence the two phosphorus nuclei are equivalent and the ring pseudorotation is very fast for **1-5**. For ^1H (Table 2) and ^{13}C NMR (Table 3) there are some cases (for example for **3**) where the methyl groups are not equivalent,


presumably free rotation around the P-C bond is prevented.¹⁰ The analysis of the ¹³C spectra (Table 3) for the α and β-carbons of the phosphonate ester groups, that is an AA'X system, has been described in reference 5.



6



7



8

Figure 1. Structures and numbering of the compounds studied.
Letters refer to protons and numbers to carbons

Table 1. ^{31}P and ^{15}N NMR spectral data for bisphosphonates **1-8**: $\delta(^{31}\text{P})$ and $\delta(^{15}\text{N})$ (ppm)

(Parentheses indicate coupling constants in Hz)

	^{31}P			^{15}N	
	δ	Solvent	Ref	δ	Solvent
1	15.9	Na/D ₂ O ^d	5	-312.8	Na/D ₂ O ^d
2	22.5	CDCl ₃	5	-340.7 (8.7) ^a	C ₆ D ₆
Picrate of 2	16.2	CDCl ₃	5	-12.2 ^b	CDCl ₃
				-15.4 ^c -322.1 (2.2)	
3	21.2	CDCl ₃	5	-340.3 (9.3)	C ₆ D ₆
4	22.7	CDCl ₃	6	-340.5 (8.5)	C ₆ D ₆
5	21.8	CDCl ₃	6	-344.8 (8.3)	C ₆ D ₆
6	21.5	CDCl ₃	6	-340.4 (6.3)	C ₆ D ₆
7	22.5	CDCl ₃		-324.3 (6.1)	C ₆ D ₆
8	23.3	CDCl ₃		-330.8 (5.2)	C ₆ D ₆

^a -341.6 (8.6) in CDCl₃, ^b nitro ortho, ^c nitro para, ^d 50 mg Na/ml D₂O

Table 2. ¹H NMR spectral data for bisphosphonates **1-8**: δ(¹H) (ppm)
(Parentheses indicate the pattern of coupling and the coupling constants in Hz)

	a	b	c	d	e	f	g	Solvent Frequency	Ref
1	3.21 (t, 7.0)	1.85 (quint., 7.0)	2.17 (sept., 7.1 ^e ; 14.5 ^f)					Na/D ₂ O ^c 200	5
2	2.88 (t, 6.5)	1.69 (quint., 6.8)	2.42 (tt, 7.2 ^e ; 17.7 ^f)	4.17 (m)	1.11 (t, 7.1 ^e) 1.10 (t, 7.1 ^e)			C ₆ D ₆ 400	5
Picrate of 2	3.64 (t, 6.9)	2.15-2.67 (m)		4.24 (m)	1.36 (t, 7.1 ^e) 1.33 (t, 7.1 ^e)	8.86 ^a (s)	7.99 ^b (s)	CDCl ₃ 100	5
3	2.95 (t, 6.5)	1.75 (quint., 6.8)	2.37 (tt, 7.3 ^e ; 17.7 ^f)	5.00 (m) 4.87 (m)	1.31 (d, 6.2 ^e) 1.28 (d, 6.3 ^e) 1.27 (d, 6.4 ^e) 1.22 (d, 6.1 ^e)			C ₆ D ₆ 400	5
4	2.97 (t, 6.5)	1.76 (quint., 6.9)	2.49 (tt, 7.3 ^e ; 17.8 ^f)	4.27 (m) 4.22 (m)	1.56 (m)	1.33 (sext., 7.5 ^e) 1.32 (sext., 7.3 ^e)	0.83 (t, 7.4 ^e) 0.82 (t, 7.4 ^e)	C ₆ D ₆ 400	6
5	2.84 (t, 6.6)	1.84 (quint., 6.6)	2.43 (m)	4.21 (m)	1.34 (t, 7.1 ^e)	2.79 (t, 1.7)		CDCl ₃ 200	6
6	4.11 (m)	1.12 (t, 7.5 ^e)	2.13 (tq, 7.5 ^e ; 15.0 ^f)	4.21 (m)	1.32 (t, 7.1 ^e) 1.31 (t, 7.1 ^e)	7.24 (m)		CDCl ₃ 400	6
7		1.24 (s)	1.74 (t, 18.4 ^f)	4.1 - 4.3 (m)	1.33 (q, 7.0)			CDCl ₃ 400	
8	2.9 (m)	1.3 ^d (m) 1.1 ^d (m)	1.41 (t, 17.4 ^f)	4.02 (m)	1.15 (t, 7.2 ^e)	0.87 (d, 6.4 ^e)	0.68 (t, 7.4 ^e)	CDCl ₃ 400	

^a aromatics, ^b -NH₂⁺, ^c 50 mg Na/ml D₂O, ^d the two protons are not equivalent, ^e H-H coupling, ^f P-H coupling

Table 3. ^{13}C NMR spectral data for bisphosphonates **1-8**: $\delta(^{13}\text{C})$ (ppm)
(Parentheses indicate coupling constants with ^{31}P in Hz)

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	Solvent Frequency	Ref
1	48.6	25.4	31.3	65.8 (t, 118.0)					Na/D ₂ O ⁱ 50	5
2	47.7 (t, 4.0)	26.5 (t, 3.1)	31.2 (t, 3.0)	62.8 (t, 151.8)	63.4 (t, 5.3) ^f 62.7 (t, 5.8) ^f	16.6 (t, 5.5) ^g 16.5 (t, 7.2) ^g			C ₆ D ₆ 100	5
Picrate of 2	49.3 (t ^e)	24.2 (t ^e)	31.2 (t ^e)	63.3 (t, 145.4)	65.0 (d ^e)	16.4 (t ^e)	160.6 ^a 141.0 ^c 128.7 ^d 126.2 ^b		CDCl ₃ 100	
3	47.7 (t, 4.6)	26.4 (t, 3.3)	31.0 (t, 3.4)	63.1 (t, 151.1)	71.7 (t, 6.2) ^f 70.8 (t, 6.7) ^f	24.7 24.4 24.0 (t, 6.2) ^g 23.8 (t, 6.7) ^g			C ₆ D ₆ 100	5
4	48.1 (t, 4.3)	26.8 (t, 3.2)	31.5 (t, 3.0)	63.2 (t, 152.0)	67.8 (t, 3.2) ^f 67.1 (t, 3.5) ^f	33.5 (t, 2.7) ^g 33.4 (t, 2.7) ^g	19.4	14.1	C ₆ D ₆ 100	6
5	55.3 (t, 4.7)	23.6 (t ^e)	32.2 (4.6)	64.2 (153.0)	62.2 (m)	16.1 (t ^e)	37.3 (t ^e)		CDCl ₃ 50	6
6	47.9 (t, 6.5)	8.5 (t, 6.5)	23.5 (t ^e)	62.8 (t, 140.1)	63.1 (t, 3.2) ^f 62.8 (t, 3.1) ^f	16.5 (t ^e)	140.5 ^a 128.3 ^b 128.2 ^c 127.0 ^d		CDCl ₃ 100	6
7	52.6 (t, 5.7)	32.4	15.8 (t, 5.5)	58.9 (t, 145.0)	64.0 (t, 3.7) ^f 62.5 (t, 4.1) ^f	16.3 (m)			CDCl ₃ 100	
8	48.4 (t, 6.1)	32.1	15.6 (t, 4.2)	57.5 (t, 143.0)	63.2 (t, 3.8) ^f 62.3 (t, 3.8) ^f	16.2 (t, 3.0) ^g	10.1	22.3	CDCl ₃ 100	

^a quaternary aromatic, ^b meta aromatics, ^c ortho aromatics, ^d para aromatic, ^e coupling can not be resolved because they are too small, ^f $^2J_{\text{CP}}(+^4J_{\text{CP}})$ ^h, ^g $^3J_{\text{CP}}$ ^h, ^h see ref 5, ⁱ 50 mg Na/ml D₂O

REFERENCES

1. Ebrahimipour, A.; Ebetino, F. H.; Sethuraman, G.; Nancollas, G. H. , *Determination of solubility and calcium ion stability constants of a phosphonoalkylphosphinate (PAP) and bisphosphonates (BPs) such as EHDP, Risedronate, Alendronate, 3-pic AMBP, and 3-pic AMPMP*; Plenum Press, New York (1995).
2. Ohanessian, J.; Avenel, D.; El Manouni, D.; Benramdane, M., *Phosphorus, Sulfur Silicon Relat. Elem.* **129**, 99 (1997).
3. Kantoci, D.; Denike, J. K.; Wechter, W. J., *Synth. Commun.* **26**, 2037 (1996).
4. Sparidans, R. W.; den Hartigh, J., *Pharm. World Sci.* **21**, 1 (1999).
5. Olive, G.; Le Moigne, F.; Mercier, A.; Rockenbauer, A.; Tordo, P., *J. Org. Chem.* **63**, 9095 (1998).
6. Olive, G.; Le Moigne, F.; Mercier, A.; Tordo, P., *Synth. Commun.* **In Press**, (1999).
7. Rockenbauer, A.; Mercier, A.; Le Moigne, F.; Olive, G.; Tordo, P., *J. Phys. Chem. Part A* **101**, 7965 (1997).
8. *Ann. Rep. NMR Spectr.* Webb, G. A., Ed.; Academic Press: London, Vol. 18, pp 713 (1986).
9. Berger, S.; Braun, S.; Kalinowski, H.-O., *NMR Spektroskopie von Nichtmetallen - Band 2 : ¹⁵N-NMR Spektroskopie*; Georg Thieme Verlag, Stuttgart; pp 111 and 112 (1992).

10. Olive, G. R. F. M., Synthèse de Nouvelles Nitrones du Type Pyrroline-*N*-Oxyde et leur Utilisation en Spin-Trapping (98AIX30024), *Thesis of Organic Chemistry*, Aix-Marseille III (1998).