



HAL
open science

In situ and ex situ studies of the gas phase in PAMPRE, a dusty plasma experiment for the study of Titan's atmosphere

Ella Sciamma-O'Brien, Nathalie Carrasco, A. Buch, Cyril Szopa, Guy
Cernogora

► To cite this version:

Ella Sciamma-O'Brien, Nathalie Carrasco, A. Buch, Cyril Szopa, Guy Cernogora. In situ and ex situ studies of the gas phase in PAMPRE, a dusty plasma experiment for the study of Titan's atmosphere. European Planetary Science Congress 2009, Sep 2009, Potsdam, Germany. hal-00669171

HAL Id: hal-00669171

<https://hal.science/hal-00669171>

Submitted on 9 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

In situ and ex situ studies of the gas phase in PAMPRE, a dusty plasma experiment for the study of Titan's atmosphere

E. Sciamma-O'Brien (1), N. Carrasco (1), A. Buch (2), C. Szopa (1) and G. Cernogora (1)

(1) Université Versailles Saint Quentin, Université Pierre et Marie Curie, CNRS, LATMOS, Réduit de Verrières BP 3, 91371 Verrières le Buisson cedex, France,

(2) Ecole Centrale de Paris, Grande voie des Vignes, F-92295 Châtenay-Malabry Cedex, France.
(ella.obrien@latmos.ipsl.fr/ Fax: +33-169-202-999)

Abstract

The complex organic chemistry that takes place in Titan's dense atmosphere, composed mainly of N_2 and CH_4 , and that leads to the production of aerosols, called tholins, has been puzzling scientists for many years. Until recently, tholins were thought to be formed in the lower stratosphere, at altitudes about a few hundred kilometres, where the atmosphere is dense and neutral. The recent observations of the INMS and the CAPS instruments on Cassini showed that these organic aerosols are actually formed at higher altitudes, in the ionosphere, where an active chemistry coupling ion and neutral species occurs [1].

To try and understand this chemistry, we use a dusty plasma experiment that reproduces in a laboratory setting, the atmospheric reactivity on Titan [2]. This experiment, called PAMPRE, uses a Capacitively Coupled Plasma (CCP) discharge, produced in a continuous gas flow, to induce the chemistry between N_2 and CH_4 . More complex molecules are then formed in the plasma and lead to the production of solid particles, analogues of Titan's aerosols.

In the study presented here, different nitrogen-methane mixtures (from 1% to 10% CH_4) have been used to study the influence of methane on the chemistry and on the subsequent production of aerosols. Using a quadrupole mass spectrometer, in situ measurements of the evolution of the gas phase composition have been achieved. An initial methane consumption phase, lasting up to a hundred seconds, has been observed. This CH_4 consumption phase is longer than the time of appearance of tholins in the discharge [3]. This phase is followed by a stationary regime where the

methane concentration is lower than in the initial gas mixture and no longer changes. This regime corresponds to the continuous working conditions (CWC) where the tholins are produced. For each initial N_2 - CH_4 gas mixture (from 1% to 10% CH_4), we have used in situ mass spectrometry measurements to determine the real CH_4 concentration in the gas mixture at CWC. In situ measurements of the gas phase in PAMPRE are fundamental since they have made it possible for us to determine which initial CH_4 concentration needs to be used to produce tholins in experimental conditions close to Titan's conditions. Using mass spectrometry, we have also been able to detect some of the products of the complex chemistry like the production of a molecule of mass 27, attributed to the production of HCN in the plasma.

Since mass spectrometry cannot detect the minor compounds present in the gas phase, a cold trap which condenses and accumulates these minor compounds at liquid nitrogen temperature has been installed, as a complementary ex situ diagnostic. This trap is similar to the one used by Coll et al. in their DC plasma discharge experiment [4]. Once accumulated, the condensed molecules are then evaporated into a gas chromatograph coupled to a mass spectrometer (GC-MS). Hydrocarbons and nitrides have been detected ex situ using this method. HCN has been detected both in situ by mass spectrometry and ex situ by GC-MS.

References

- [1] Waite, J.H. Jr., Young, D. T., Cravens, T.E., Coates, A.J., Crary, F.J., Magee, B., Westlake, J. (2007) *Science*, 316, 870-875.
- [2] Szopa, C., Cernogora, G., Boufendi, L., Correia, J.-J. and Coll, P. (2006) *Planet. Space Sci.*, 54, 394-404.
- [3] Alcouffe, G., Cavarroc, M., Cernogora, G., Ouni, F., Jolly, A., Boufendi, L., Szopa, C., submitted to *Plasma Sources Sci. Technol.*
- [4] Coll, P., Coscia, D., Smith, N., Gazeau, M.-C., Ramirez, S.I., Cernogora, G., Israel, G., Raulin, F. (1999) *Planet. Space Sci.*, 47, 1331-1340.

Acknowledgment

This work is supported by grants from the OPV program of the French National Center for Scientific Research (CNRS), the French Planetology National Program (PNP) and the University of Versailles Saint Quentin (PRES UniverSud).