



HAL
open science

Amyloid beta impairs mitochondrial anterograde transport and degenerates synapses in Alzheimer's disease neurons

Marcus J. Calkins, P. Hemachandra Reddy

► **To cite this version:**

Marcus J. Calkins, P. Hemachandra Reddy. Amyloid beta impairs mitochondrial anterograde transport and degenerates synapses in Alzheimer's disease neurons. *Biochimica et Biophysica Acta - Molecular Basis of Disease*, 2011, 1812 (4), pp.507. 10.1016/j.bbadis.2011.01.007 . hal-00670752

HAL Id: hal-00670752

<https://hal.science/hal-00670752>

Submitted on 16 Feb 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Amyloid beta impairs mitochondrial anterograde transport and degenerates synapses in Alzheimer's disease neurons

Marcus J. Calkins, P. Hemachandra Reddy

PII: S0925-4439(11)00020-2
DOI: doi: [10.1016/j.bbadis.2011.01.007](https://doi.org/10.1016/j.bbadis.2011.01.007)
Reference: BBADIS 63233

To appear in: *BBA - Molecular Basis of Disease*

Received date: 15 December 2010
Revised date: 6 January 2011
Accepted date: 10 January 2011



Please cite this article as: Marcus J. Calkins, P. Hemachandra Reddy, Amyloid beta impairs mitochondrial anterograde transport and degenerates synapses in Alzheimer's disease neurons, *BBA - Molecular Basis of Disease* (2011), doi: [10.1016/j.bbadis.2011.01.007](https://doi.org/10.1016/j.bbadis.2011.01.007)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Amyloid Beta Impairs Mitochondrial Anterograde Transport and Degenerates Synapses in Alzheimer's Disease Neurons

¹Marcus J. Calkins and ^{1,2}P. Hemachandra Reddy

¹Neurogenetics Laboratory, Division of Neuroscience, Oregon National Primate Research Center, Oregon Health & Science University, 505 NW 185th Avenue, Beaverton, OR 97006

²Department of Physiology and Pharmacology, Oregon Health & Science University, 3181 SW Sam Jackson Park Road, Portland, OR 97239

Address for correspondence and reprint requests:

P. Hemachandra Reddy, PhD
Neurogenetics Laboratory
Neuroscience Division
Oregon National Primate Research Center
West Campus, Oregon Health & Science University
505 NW 185th Avenue
Beaverton, OR 97006
Tel: 503 418 2625
Fax: 503 418 2701
E-mail: reddyh@ohsu.edu

Abbreviations

A β PP, A β precursor protein; AD, Alzheimer's disease; ATP, adenosine triphosphate; Drp1, dynamin related protein 1; Fis1, mitochondrial fission 1.

Abstract

Loss of synapses and synaptic damage are the best correlates of cognitive decline identified in patients with Alzheimer's disease (AD), and mitochondrial oxidative damage and synaptic pathology have been identified as early events in the progression of AD. The progressive accumulation of amyloid beta ($A\beta$) in synapses and synaptic mitochondria are hypothesized to cause synaptic degeneration and cognitive decline in patients with AD. However, the precise mechanistic link between $A\beta$ and mitochondria is not well understood. The purpose of this study was to better understand the effects of $A\beta$ on mitochondrial axonal transport and synaptic alterations in AD. Using mouse hippocampal neurons and $A\beta_{25-35}$ peptide, we studied axonal transport of mitochondria, including mitochondrial motility, mitochondrial length and size, mitochondrial index per neurite, and synaptic alterations of the hippocampal neurons. In the PBS-treated neurons, $36.4 \pm 4.7\%$ of the observed mitochondria were motile, with $21.0 \pm 1.3\%$ moving anterograde and $15.4 \pm 3.4\%$ moving retrograde and the average speed of movement was $12.1 \pm 1.8 \mu\text{m}/\text{min}$. In contrast, in the $A\beta$ -treated neurons, the number of motile mitochondria were significantly less, at $20.4 \pm 2.6\%$ ($P < 0.032$), as were those moving anterograde ($10.1 \pm 2.6\%$, $P < 0.016$) relative to PBS-treated neurons, suggesting that the $A\beta_{25-35}$ peptide impairs axonal transport of mitochondria in AD neurons. In the $A\beta$ -treated neurons, the average speed of motile mitochondria was also less, at $10.9 \pm 1.9 \mu\text{m}/\text{min}$, and mitochondrial length was significantly decreased. Further, synaptic immunoreactivity was also significantly less in the $A\beta$ -treated neurons relative to the PBS-treated neurons, indicating that $A\beta$ affects synaptic viability. These findings suggest that, in neurons affected by AD, $A\beta$ is toxic, impairs mitochondrial movements, reduces mitochondrial length, and causes synaptic degeneration.



Introduction

Alzheimer's disease (AD) is the most common neurodegenerative disorder in the aged population. It is characterized by the progressive decline of memory, decline in cognitive functions such as memory, and changes in behavior and personality [1]. It is a major health concern in society mainly, especially because half of individuals 85 years of age and older are expected to develop AD [2]. AD is associated with multiple cellular changes in the brain, including the loss of synapses/synaptic pathology, mitochondrial structural/functional abnormalities, inflammatory responses, extracellular amyloid beta (A β) deposits, and intracellular neurofibrillary tangles [1,3-10]. Among these changes, mitochondrial oxidative damage and synaptic pathology are reported as early events in AD progression [11-15].

In AD, A β is a major component of neuritic plaques found in brain regions known to be responsible for learning and memory. A β , the 39-43 amino acid residue protein, is generated by the proteolysis of A β precursor protein (A β PP) by the sequential enzymatic actions β -secretase, and γ secretase. In AD, levels of A β are steady-state and are controlled by the production, the clearance, and the degradation of A β . Decreased clearance of A β or the overproduction of A β may lead to an accumulation of A β in subcellular compartments, including synapses, mitochondria, and may impair functions of subcellular organelles and damage neurons [4,5,16]. Thus, agents or drugs capable of increasing the clearance or decreasing the production of A β may be important therapeutic strategies in preventing AD development and progression.

Several researchers have reported that A β is responsible for damaging mitochondria and synapses in neurons affected by AD [17-24]. Monomeric and oligomeric forms of A β have been found in mitochondrial membranes and to interact with mitochondrial proteins, induce free radical production, alter mitochondrial enzymes, disrupt the electron transport chain, inhibit adenosine triphosphate (ATP) production, and damage mitochondria [17-19,24-26]. Although these results clearly associate A β with subcellular components such as mitochondria, the precise link between A β and mitochondria is not clear and how A β damages mitochondria is also not well understood.

Mitochondrial shape and structure are maintained by mitochondrial fission and fusion [27-29]. Fission and fusion mechanisms are equally balanced in healthy neurons. Mitochondria alter their shape and size to move, through mitochondrial trafficking, from the cell body to the axons, dendrites and synapses via anterograde fashion, and back to the cell body via retrograde manner [30]. Synapses are the sites of high-energy demand. Mitochondria supply energy to synapses, which is required for neural communications and several synaptic functions, including the release of neurotransmitters, synaptic vesicles between pre- and post-synaptic neurons. In a healthy neuron, abundant numbers of healthy, functionally active mitochondria are present at synapses, whereas in diseased neurons, such as in AD-affected neurons, small numbers of

defective mitochondria are present [31]. In the AD neuron, A β accumulates in the mitochondria, induces free radical production, and activates the two mitochondrial fission proteins dynamin-related protein 1 (Drp1) and mitochondrial fission 1 (Fis1) [29]. Activated Drp1 and Fis1 have been associated with excessive fragmentation of mitochondria that may not move to synapses and do not supply the necessary ATP at nerve terminals. Further, these defective mitochondria may not be able to participate in mitochondrial fusion, may not be able to synthesize healthy mitochondria, and may, ultimately, prematurely die. The continual production of excessive numbers of defective mitochondria in neurons may ultimately damage synapses and cause synaptic neurodegeneration [29,31].

Using electron and confocal microscopy, gene expression analysis, and biochemical methods, we treated mouse neuroblastoma (N2a) cells with the A β 25-35 peptide, and studied mitochondrial structure and function; expressions of the fission genes Drp1 and Fis1 and the 3 fusion genes Mfn1, Mfn2, and Opa1; and neurite outgrowth in [32]. In the neurons treated with only A β , we found increased expressions of fission genes and decreased expressions of fusion genes, indicating abnormal mitochondrial dynamics in neurons treated with A β . Our immunocytochemistry of N2a cells treated with A β revealed increased immunoreactivity of Drp1 and Fis1, suggesting that A β elevates fission genes and fragments mitochondria. Electron microscopy of the N2a cells incubated with A β revealed a significantly increased number of defective mitochondria, indicating that A β fragments mitochondria. Biochemical analysis revealed A β in association with defective mitochondria. Neurite outgrowth was significantly decreased in those N2a cells that we incubated with A β , indicating that A β affects neurite outgrowth.

In the research reported here, we sought to determine the effects of A β on axonal transport of mitochondria in mouse hippocampal neurons. We measured the total number of mitochondria, the length and size of mitochondria, the mitochondrial index (length of the mitochondria per neurite length), and the synaptic immunoreactivity of AD neurons treated with the A β 25-35 peptide compared to control neurons (A β 35-25) and those treated with the vehicle (PBS).

Materials and Methods

Neuronal Culture

Hippocampal neurons were cultured, as previously described [32]. Briefly, hippocampi were dissected from C57BL/6 day 1 pups in room temperature HABG (Hibernate E medium [Brain Bits, LLC] supplemented with 1X B-27 [Invitrogen] and 0.5 mM glutamine [Invitrogen, CA]). The tissues were dissociated with 2 mg/ml papain at 30°C for 30min in a dissociation medium (Hibernate E medium without Ca⁺⁺, supplemented with 0.5 mM glutamine). Digested tissue was triturated, using a fire-polished, silicon-coated Pasteur pipette in 2 ml HABG. Non-dissociated tissue was allowed to settle for 5 min, after which

time the supernatant was passed through a 70-mm nylon mesh, into a 50-mL conical tube and centrifuged for 2 min at 200g. The pellet was gently resuspended in a maintenance medium (Neurobasal A medium [Invitrogen, CA] supplemented with 1X B-27 minus antioxidants, and 0.5 mM L-glutamine) and plated onto a poly-D-lysine (Sigma-Aldrich) –coated, chambered coverglass (Nunc). The medium was completely replaced after 1 hr, and then half of the medium was replaced every 3days.

Time Lapse Photography of Mitochondrial Motility and Data Analysis

Mitochondria were labeled by transfecting pDsRed2-mito (Clontech) into the hippocampal neurons at day 2 (DIV) with lipofectamine 2000 (Invitrogen), according to the manufacturer's protocol. DsRed-labeled neuronal mitochondria were observed in cultures up to 21 days after transfection (data not shown). A β ₂₅₋₃₅, A β ₃₅₋₂₅ (control) peptides were dissolved in 1XPBS. To increase the toxicity of A β peptide, we incubated A β ₂₅₋₃₅, and A β ₃₅₋₂₅ (control) peptides at 37°C for overnight, and this overnight incubation is expected to convert monomeric A β to oligomeric A β . We treated neurons with 20 μ M A β ₂₅₋₃₅ (oligomeric) A β ₃₅₋₂₅ (control), and a vehicle (PBS) at 10 DIV and then imaged 24 hours later. Axonal processes were determined by morphological characteristics. Axons were identified as processes stemming from the soma that are two to three times longer than other processes [32,33]. Recordings were made on axonal segments about 20-100 μ m from the soma. Series of time-lapse images were captured every 5 sec, using a Leica SP5 AOBS confocal microscope with a heated 37°C, 5% CO₂ controlled stage for a total of 5min. Z-stacks for each time point were collapsed to maximum projections, and the time series were archived as avi files. ImageJ software, with a Multiple Kymograph plug-in, was used to analyze the avi files. Mitochondrial movements (direction and speed) were determined from the kymographic images. Mitochondria were considered stationary if they did not move more than 2 μ m during the entire recording period. Each series of images was recorded for at least three randomly selected Ds-Red-mito labeled cells per culture and four independent cultures per condition.

Mitochondrial Content of Neurites

After imaging the mitochondria in the neuronal cultures, the cultures were fixed with 4% paraformaldehyde for 5 min at room temperature and then washed with PBS. The plastic chambers were carefully removed, and the coverslips were mounted on slides with a ProLong Gold mounting medium. Images of cell bodies with neurites extending at least 100 μ m were collected, using a Leica SP5 AOBS confocal microscope with a 63x objective. GFP and Ds-Red were analyzed using measurement tools in ImageJ to determine the neurite mitochondrial index, the average mitochondrial length, and the number of

mitochondria per neurite length). Data were collected from at least six cells per culture and four independent cultures per condition

Immunofluorescence analysis of synaptic proteins

To determine the toxicity of A β in hippocampal neurons, we performed immunocytochemical analysis of Drp1 with mitochondria-encoded protein Cytochrome b (Cyt. B), and synaptophysin, which is marker for AD, with microtubule associated protein 2 (MAP2). Briefly, we plated cells on 13-mm round coverslips coated with poly-D-lysine contained within wells of a 24-well plate. After treatment, the medium was removed, and cells were fixed with 4% paraformaldehyde in PBS for 10-15 min at room temperature. Coverslips were washed with PBS, and cell membranes were permeabilized with 0.1% Triton X-100 in PBS for 5 min, after which time a blocking solution was applied (2% normal goat serum, 1% BSA in PBS). All subsequent incubations were carried out in a humidified environment. Samples were blocked for 2 hr at room temperature and then incubated with a primary antibody diluted in a blocking solution overnight at 4°C. Drp1 (1:200, rabbit polyclonal, Novus Biologicals, Inc.), and synaptophysin (1:200, mouse monoclonal, Millipore/Chemicon, Temecula, CA) were probed. After cells were incubated with the primary antibody, cells were washed three times with PBS and then incubated with either goat-anti-rabbit-Alexa488 or goat-anti-mouse-Alexa568 (both 1:500, Invitrogen/Molecular Probes) for 2 hr at room temperature. For co-labeling with Cyt. B (1:50, mouse monoclonal, Invitrogen) or MAP2 (1:1000, rabbit polyclonal, Milipore), cells were then incubated with the second primary antibody for 3 hr at room temperature. After washing three times with PBS, coverslips were incubated with the corresponding secondary antibody for 2 hr at room temperature. Antibody was removed, and DAPI (600nM in PBS) was added to the cells for 5 min. Cells were washed three times in PBS, and then coverslips were mounted on slides using a ProLong Gold antifade mounting reagent (Invitrogen). Cells were imaged, using a Zeiss Axioskop 40 FL microscope.

Statistical Analysis

Data from independent cultures were compared, using a two-tailed, unpaired student's t-test with significance level set at $p \leq 0.05$.

Results

Mitochondrial motility is less after A β treatment. To determine the effect of A β peptide on mitochondrial transport, we incubated hippocampal neurons with 20 μ M A β 25-35, A β 35-25 (control), or PBS for 24 hours, and then quantified mitochondrial motility in axons. The average speed of mitochondrial movement was 12.1 ± 1.8 mm/min. In the A β 25-35-treated cultures, we observed significantly decreased total mitochondrial

motility ($20.4 \pm 2.6\%$ – mean \pm SE, $P < 0.032$) and mitochondria moving anterograde ($10.1 \pm 2.6\%$ – mean \pm SE, $P < 0.016$) relative to the PBS vehicle-treated neurons. A decrease in retrograde-moving mitochondria was also observed ($10.3 \pm 1.5\%$ – mean \pm SE) in the A β 25-35-treated neurons relative to the PBS vehicle-treated neurons. However, this decrease did not reach statistical significance ($P = 0.234$). The average speed of mitochondria undergoing the A β 25-35 treatment was only slightly decreased ($10.9 \pm 1.9 \mu\text{m}/\text{min}$ – mean \pm SE, $P = 0.66$) compared to the mitochondria from the vehicle-treated neurons. In the PBS vehicle-treated neurons, $36.4 \pm 4.7\%$ (mean \pm SE) of the observed mitochondria were motile, with $21.0 \pm 1.3\%$ moving anterograde and $15.4 \pm 3.4\%$ moving retrograde.

As shown in Fig. 1, the A β 35-25-treated neurons (the controls) did not show any change in total mitochondrial motility – neither anterograde nor retrograde– relative to the PBS vehicle-treated neurons, indicating that mitochondrial alterations are specific to A β 25-35 specific.

Mitochondrial distribution is altered after A β treatment. Following our observations of the decrease in overall anterograde mitochondrial motility, we examined whether A β treatment also induced changes in mitochondrial distribution within neurites. Cultures treated for 24 hours with the PBS vehicle or the A β 25-35 were harvested, and the neuritic mitochondrial content was analyzed. We found that A β treatment significantly decreased the mitochondrial index in neurons treated with A β ($40.1 \pm 3.1\%$) relative to the PBS-treated (PBS, $40.1 \pm 3.1\%$ – mean \pm SE; A β , $18.4 \pm 2.5\%$ – mean \pm SE, $P < 0.001$) (Figs.2 and 3). We found the mitochondrial index did not change in neurons treated with A β 35-25 peptide, indicating that the decrease in the mitochondrial index that we found in neurons treated A β 25-35 peptide is, indeed, associated with A β .

The mitochondrial index reflects the total mass of mitochondria per unit length of a neurite and can be affected by changes in either the number or the size of mitochondria. We therefore analyzed mitochondrial length and density (the number of mitochondria per length of). As shown in Fig. 2, the average mitochondrial length was significantly decreased in neurons with the A β peptide relative to the PBS-treated neurons (PBS, $2.47 \pm 0.21 \mu\text{m}$ – mean \pm SE; A β , $1.37 \pm 0.10 \mu\text{m}$ – mean \pm SE, $P < 0.001$) (Fig.2). Mitochondrial density decreased from 17.3 ± 0.9 mitochondria/100 mm in the PBS-treated neurons to 13.6 ± 1.5 mitochondria/100mm relative to the A β -treated neurons ($P = 0.054$). Overall, we found that mitochondrial mass was greatly reduced after receiving A β treatment relative PBS vehicle treated control neurons, and that this effect is largely due to decreased mitochondrial size (Figs. 2 and 3). As shown in Fig. 3, we found fragmented mitochondria in Ds-red labeled hippocampal neurons treated with A β relative to PBS treated control neurons.

Immunoreactivity of synaptic protein was less in A β 25-35- treated neurons

To determine the effect of A β 25-35 treatment on the immunoreactivity of synaptophysin and Drp1 in hippocampal neurons, we treated neurons with A β 25-35 and with PBS for 24 hrs, and then performed immunostaining. As shown in Fig. 4, we found the immunoreactivity of synaptophysin significantly decreased, particularly at synapses in those neurons treated with A β 25-35 ($54.2\pm 5.9\%$, mean \pm SE- $P<0.013$, Fig. 4a), relative to the synapses in neurons treated with the PBS vehicle ($100\pm 5.9\%$, mean \pm SE).

To determine, the effect of A β 25-35 on synapses, we performed double-labeling immunostaining analysis using synaptic marker, MAP2 and Drp1 in neurons treated with A β 25-35 and with PBS for 24 hrs. Similar to synaptophysin, immunoreactivity of MAP2 is decreased in neurons treated with A β 25-35 (Fig. 4a image E) relative to PBS treated neurons (image B), indicating that A β 25-35 peptide may be involved in synaptic degeneration.

To determine the effect of A β 25-35 on mitochondria, particularly mitochondria localized at synapses, we performed double-labeling analysis of Drp1 (enriched in synapses) and Cyt.B (mitochondria-encoded protein) antibodies. As shown in Fig. 5, we found fragmented and punctuated immunoreactivity of Drp1 in neurons treated with A β 25-35 relative to the Drp1 immunoreactivity in neurons treated with A β 35-25 or the PBS vehicle (Fig. 5). Drp1 expression was higher in synapses, spines, and branches that were developing in reverse A β peptide or PBS vehicle treated neurons, whereas in A β 25-35 peptide treated neurons, we found decreased and fragmented immunoreactivity, particularly at synapses indicating mitochondrial fragmentation, and degenerating synapses. Further, Drp1 is colocalized with Cyt. B further confirming that fragmented immunoreactivity represents mitochondrial fragmentation caused by A β 25-35 peptide.

Discussion

In this study, we analyzed the effect of A β treatment on mitochondrial motility and distribution in hippocampal neurons from mice. We found mitochondrial motility greatly reduced by the exposure of the hippocampal neurons to A β , and the reduction in anterograde transport greater than the reduction in retrograde movement. Overall, a reduction in mitochondrial transport in the hippocampal neurons was observed and mitochondrial transport was impaired in the distal regions of the neurites. We also found decreased synaptic branching and growth in neurons treated with A β . This would lead to decreased ATP production, followed by synaptic dysfunction and degeneration. These findings lead us to conclude that A β impairs axonal transport of mitochondria and may cause synaptic degeneration.

Amyloid beta and impaired axonal transport of mitochondria. A β is generated in neurons, wherever APP and β - and γ - secretases are present, which is in several intracellular sites, including Golgi apparatus, endoplasmic reticulum, endosomal-lysosomal systems, and multivesicular bodies. Recent studies of AD patients and transgenic AD mice found intracellular A β present in AD-affected brain regions and that A β 1-42 participates in fibrillogenesis and the formation of A β plaques. Increasing evidence suggests that A β accumulates in cellular compartments, including mitochondria, and that A β interfere with normal functions of mitochondria and synapses. Findings from the present study help elucidate a mechanistic relationship among A β , synaptic damage, and mitochondrial damage in neurons affected by AD. We found a dramatic reduction (55%) in the total mitochondria in hippocampal neurons exposed to A β , and we found significantly reduced anterograde mitochondrial movement (cell body to synapses) in the A β -treated hippocampal neurons relative to the PBS-treated neurons. We also found the length of mitochondria significantly decreased in the A β -treated hippocampal neurons compared to the PBS-treated or the A β 35-25-treated hippocampal neurons. Finally, we found the immunoreactivity of synaptophysin significantly decreased in the A β -treated neurons, compared to the PBS-treated hippocampal neurons.

Reductions in the total number of mitochondria and of anterograde-moving mitochondria are likely responsible for the synaptic degeneration observed in the AD neurons since a large number of healthy and functionally active mitochondria are necessary for ATP trafficking to synapses where energy-demand is high. Reduction in anterograde-moving mitochondria appear to reduce the supply of ATP at synapses, which may cause synaptic degeneration that marks AD neurons. Our finding pointing to A β -induced impaired mitochondrial transport is consistent with other studies showing decreased mitochondrial transport [23,35-36] and organelle transport [37-39] in A β -treated neurons.

Wang et al. [23] reported that A β oligomer (ADDLs)-treated hippocampal neurons from mouse showed significantly reduced mitochondrial anterograde ($0.80 \pm .67$) and retrograde axonal transport ($0.44 \pm .21$), compared to hippocampal neurons not so treated ($5.34 \pm .47$ for anterograde and $3.31 \pm .02$ for retrograde transport). Neurons treated with A β 42-1 exhibited a fast axonal transport of mitochondria that was similar to the axonal transport found in the control neurons.

Du and colleagues [36] found a 30% increase in the percentage of stationary mitochondria in A β -treated neurons relative to mitochondria in untreated neurons. Further, the total number of mitochondria moving in both directions (anterograde and retrograde) was significantly reduced in the A β -treated neurons compared to those treated with the PBS vehicle or the control (A β 42-1 treated). After A β treatment, the anterograde moving mitochondria decreased by <23% compared to those in the PBS-treated neurons and in the control neurons ($P < 0.05$). In contrast, the percentage of retrograde mobile mitochondria increased those neurons treated with A β 1-42.

The velocity of mitochondrial motility was also less in neurons treated with A β relative to that of the PBS-treated neurons, indicating that A β affects mitochondrial velocity. Our findings are consistent with those from Du et al. [34], who also reported significantly reduced mitochondrial anterograde velocity in neurons treated with A β 1-42 relative to control and A β 42-1-treated neurons.

Overall, findings from our study, together with those from Wang et al. [23] and Du et al. [36], indicate that the A β peptide is toxic and impairs axonal transport of mitochondria. Further, our findings are consistent with findings from other mitochondrial trafficking studies by Chang et al. [40] and Rui et al. [35], reporting that mitochondrial transport and distribution were affected by hippocampal neurons exposed to the A β peptide, glutamate and zinc.

By shifting the balance of anterograde/retrograde transport toward retrograde, a decrease in axonal and neuritic mitochondria would be expected. Indeed, we observed this effect, finding a large reduction in mitochondrial mass in the A β -treated cells (Fig. 2); more specifically, we found the average length of mitochondria length was decreased, but the total number of mitochondria per each length was less affected. These findings suggest that factors dictating mitochondrial distribution along an axon are unaffected by A β treatment. However, mitochondrial dynamics are likely shifted toward fission.

Amyloid beta and abnormal mitochondrial dynamics. We [32] and others [21,22] reported increased mitochondrial fission and decreased fusion in neurons affected by AD. In other words, abnormal mitochondrial dynamics occurs in neurons affected by AD. This may be due to the association of A β with mitochondria. Mitochondrial A β is known to induce free radical production and to activate the fission proteins Drp1 and Fis1, and to fragment mitochondria [32]. When mitochondrial fission is activated, fusion proteins decrease.

Our immunostaining analysis of Drp1 showed fragmented and punctuated immunoreactivity in A β -treated neurons, indicating increased mitochondrial fission and decreased fusion. As described elsewhere [32], in A β -treated neurons, mitochondrial fusion is reduced following the induction of apoptosis. Overexpression of mitofusins can reduce apoptosis. Further, it has been previously suggested that mitochondrial fusion and transport are closely linked, such that disruptions in either process affect the other. Recently, Miskow et al. [41] showed that Mfn2 interacts with the Miro-Milton complex, and this interaction is required for axonal transport of mitochondria in dorsal root ganglia neurons. Conversely, mitochondrial fragmentation induced by Pink1 knockdown can be rescued by the overexpression of Miro or Milton [42]. In this light, it is not surprising that we observed both altered mitochondrial transport and altered mitochondrial dynamics.

Decreased synaptic growth in A β -treated neurons. Our quantitative analysis of synaptophysin immunoreactivity revealed that synaptophysin was significantly decreased in A β -treated neurons, indicating that A β affects synaptic growth, an observation supported by other studies [43-46]. Further, MAP2 immunoreactivity was decreased in A β -treated neurons, further supporting that synaptic degeneration is present in A β -treated neurons. This reduced synaptic growth was primarily due to the decrease mitochondrial trafficking, particularly the anterograde movement of mitochondria, which is known to ultimately lead to the production of ATP at synapses and to the degeneration of synapses in AD neurons.

In summary, we found reduced mitochondrial mass, reduced mitochondrial motility, and reduced mitochondrial anterograde transport in neurons exposed to the A β ₂₅₋₃₅ peptide. We also found a reduction in the number of mitochondria in distal regions of neurons and decreased synaptic branching and growth in the A β -treated neurons. These findings lead us to conclude that A β impairs axonal transport of mitochondria and may cause synaptic degeneration.

Acknowledgements

This research presented was supported by NIH grants AG028072, AG026051, and RR00163, Alzheimer Association grant IIRG-09-92429, and Medivation, Inc. We also thank Dr. Anda Cornea for her assistance with confocal imaging and mitochondrial transport assessment.

ACCE

Figure legends

Figure 1. Amyloid beta treatment reduces mitochondrial movements. Axons from mature hippocampal neurons transfected with DsRed-mito and GFP, then treated for 24 hours with vehicle, A β , or reverse peptide were imaged to evaluate mitochondrial movements. Total proportion of moving mitochondria, proportion of mitochondria moving anterograde, and proportion of mitochondria moving retrograde were calculated (A). The speed of motion was also calculated for all moving mitochondria (B). Calculations were based on analysis of kymographs. Representative kymographs are shown for the three experimental groups (C). N= 4 independent cultures. * $p < 0.05$ compared to vehicle treated, and statistical variation is shown as mean \pm SE.

Figure 2. Amyloid beta treatment causes a reduction in mitochondrial mass within neurites.

Mitochondria from DsRed-mito transfected hippocampal neurons were analyzed after vehicle, A-beta, or reverse peptide treatment. Mitochondrial index was calculated as the percent of neuritic length occupied by mitochondria. A β treated cultures showed pronounced reduction in mitochondrial index (A). The distribution of mitochondria was evaluated as number of mitochondria per neuritic length (B). Mitochondrial length was reduced by A β treatment (C). N = 4 independent cultures. ** $p < 0.01$. Statistical variation is shown as mean \pm SE.

Figure 3. Mitochondria are more fragmented after Amyloid beta treatment. DsRed-mito transfected hippocampal neurons were imaged after vehicle, A β , or reverse peptide treatment. Representative images are shown in upper panels. Enlargements of neurites are shown in the lower panels.

Figure 4. Double-labeling analysis of Synaptophysin and MAP2 in neurons treated with amyloid beta.

Hippocampal neurons treated with PBS vehicle or A β were immunostained for synaptophysin and MAP2 (a). Vehicle treated cells (upper panel) showed typical strong synaptophysin (A), MAP2 (B) immunoreactivities and colocalization of synaptophysin and MAP2 (C). A β -treated cells (lower panel) showed reduced density of synaptophysin puncta (D), MAP2 (E) and merged (C). Image **b** shows quantification of synaptophysin immunoreactivity. Significantly decreased synaptophysin was found in A β -treated neurons ($P < 0.005$). Image **c** shows enlarged portion of a neurite from PBS vehicle treated neuron (A) and A β -treated neuron (B).

Figure 5. Drp1 distribution relocates from areas of active neurite outgrowth to neuritic processes after amyloid beta treatment. Hippocampal neurons treated with PBS vehicle or A β were immunostained for

Drp1, mitochondrial-encoded protein, Cyt. B and nuclear marker, DAPI. Vehicle treated cells (upper panel) showed intense immunoreactivities of Drp1 (A), Cyt. B (B), DAPI (C) and merged (D) in neurite growth cones (white arrows). Drp1 is colocalized with Cyt. B (D) (white arrows). After A β treatment (lower panel), Drp1 staining is reduced (E), and Drp1 is colocalized with mitochondria in merged image (H).

ACCEPTED MANUSCRIPT

References

- [1]. D.J. Selkoe, Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev.* 81(2001) 741-66.
- [2]. Alzheimer Association Report, Alzheimer's Disease Facts and Figures (2010).
- [3]. M.P. Mattson, Pathways towards and away from Alzheimer's disease. *Nature.* 430 (2004) 631-639.
- [4]. F.M. LaFerla, K.N. Green, S. Oddo, Intracellular amyloid-beta in Alzheimer's disease. *Nat Rev Neurosci.* (2007) 499-509.
- [5]. P.H. Reddy, M.F. Beal, Amyloid beta, mitochondrial dysfunction and synaptic damage: implications for cognitive decline in aging and Alzheimer's disease. *Trends Mol Med.* 14 (2008) 45-53.
- [6]. J.N. Keller, Z. Pang, J.W. Geddes, J.G. Begley, A. Germeyer, G. Waeg, M.P. Mattson, Impairment of glucose and glutamate transport and induction of mitochondrial oxidative stress and dysfunction in synaptosomes by amyloid beta-peptide: role of the lipid peroxidation product 4-hydroxynonenal. *J Neurochem.* 69(1997), 273-84.
- [7]. R.H. Swerdlow, J.K. Parks, D.S. Cassarino, D.J. Maguire, R.S. Maguire, J.P. Jr., Bennett, R.E. Davis, W.D. Jr. Parker, Cybrids in Alzheimer's disease: a cellular model of the disease? *Neurology.* 49(1997), 918-25.
- [8]. R.H. Swerdlow, S.M. Khan, A "mitochondrial cascade hypothesis" for sporadic Alzheimer's disease. *Med Hypotheses.* 63(2004), 8-20.
- [9]. R.H. Swerdlow, S.M. Khan, The Alzheimer's disease mitochondrial cascade hypothesis: an update. *Exp Neurol.* 218(2009), 308-315.
- [10]. M.J. Barsoum, H. Yuan, A.A. Gerencser, G. Liot, Y. Kushnareva, S. Gräber, I. Kovacs, W.D. Lee, J. Waggoner, J. Cui, A.D. White, B. Bossy, J.C. Martinou, R.J. Youle, S.A. Lipton, M.H. Ellisman, G.A. Perkins, E. Bossy-Wetzel. Nitric oxide-induced mitochondrial fission is regulated by dynamin-related GTPases in neurons. *EMBO J.* 25 (2006) 3900-3911.
- [12]. C. Bertoni-Freddari, P. Fattoretti, T. Casoli, W. Meier-Ruge, J. Ulrich, Morphological adaptive response of the synaptic junctional zones in the human dentate gyrus during aging and Alzheimer's disease. *Brain Res.* 517 (1990) 69-75.
- [11]. R.D. Terry, E. Masliah, D.P. Salmon, N. Butters, R. DeTeresa, R. Hill, L.A. Hansen, R. Katzman, Physical basis of cognitive alterations in Alzheimer's disease: synapse loss is the major correlate of cognitive impairment. *Ann Neurol.* 30(1991) 572-580.
- [13]. S.T. DeKosky, S.W. Scheff, Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity. *Ann Neurol.* 27(1990) 457-464.
- [14]. S.W. Scheff, S.T. DeKosky, D.A. Price, Quantitative assessment of cortical synaptic density in Alzheimer's disease. *Neurobiol Aging.* (1990) 29-37.
- [15]. A. Nunomura, G. Perry, G. Aliev, K. Hirai, A. Takeda, E.K. Balraj, P.K. Jones, H. Ghanbari, T. Wataya, S. Shimohama, S. Chiba, C.S. Atwood, R.B. Petersen, M.A. Smith, Oxidative damage is the earliest event in Alzheimer disease. *J Neuropathol Exp Neurol.* 60 (2001) 759-767.

- [16]. G.K. Gouras, C.G. Almeida, R.H. Takahashi, Intraneuronal Abeta accumulation and origin of plaques in Alzheimer's disease. *Neurobiol Aging*. 26 (2005) 1235-1244.
- [17]. C. Caspersen, N. Wang, J. Yao, A. Sosunov, X. Chen, J.W. Lustbader, H.W. Xu, D. Stern, G. McKhann, S.D. Yan, Mitochondrial Abeta: a potential focal point for neuronal metabolic dysfunction in Alzheimer's disease. *FASEB J*. 19(2005):2040-1.
- [18]. M. Manczak, T.S. Anekonda, E. Henson, B.S. Park, J. Quinn, P.H. Reddy, Mitochondria are a direct site of A beta accumulation in Alzheimer's disease neurons: implications for free radical generation and oxidative damage in disease progression. *Hum Mol Genet*. 15 (2006) 1437-49.
- [19]. L. Devi, B.M. Prabhu, D.F. Galati, A.G. Avadhani, H.K. Anandatheerthavarada, Accumulation of amyloid precursor protein in the mitochondrial import channels of human Alzheimer's disease brain is associated with mitochondrial dysfunction. *J Neurosci*. 26 (2006) 9057-9068.
- [20]. X. Wang, B. Su, H. Fujioka, X. Zhu, Dynamin-like protein 1 reduction underlies mitochondrial morphology and distribution abnormalities in fibroblasts from sporadic Alzheimer's disease patients. *Am J Pathol*. 173 (2008) 470-482.
- [21]. X. Wang, B. Su, S.L. Siedlak, P.I. Moreira, H. Fujioka, Y. Wang, G. Casadesus, X. Zhu, Amyloid-beta overproduction causes abnormal mitochondrial dynamics via differential modulation of mitochondrial fission/fusion proteins. *Proc Natl Acad Sci U S A*. 105(2008)19318-19323.
- [22]. X. Wang, B. Su, H.G. Lee, X. Li, G. Perry, M.A. Smith, X. Zhu, Impaired balance of mitochondrial fission and fusion in Alzheimer's disease. *J Neurosci*. 29 (2009) 9090-9103.
- [23]. X. Wang, G. Perry, M.A. Smith, X. Zhu, Amyloid-beta-derived diffusible ligands cause impaired axonal transport of mitochondria in neurons. *Neurodegenerative Dis* 7 (2010) 56-59.
- [24]. H. Du, L. Guo, F. Fang, D. Chen, A.A. Sosunov, G.M. McKhann, Y. Yan, C. Wang, H. Zhang, J.D. Molkentin, F.J. Gunn-Moore, J.P. Vonsattel, O. Arancio, J.X. Chen, S.D. Yan. Cyclophilin D deficiency attenuates mitochondrial and neuronal perturbation and ameliorates learning and memory in Alzheimer's disease. *Nat Med*. 14 (2008) 1097-105.
- [25]. P.H. Reddy, S. McWeeney, B.S. Park, M. Manczak, R.V. Gutala, D. Partovi, Y. Jung, V. Yau, R. Searles, M. Mori, J. Quinn, Gene expression profiles of transcripts in amyloid precursor protein transgenic mice: up-regulation of mitochondrial metabolism and apoptotic genes is an early cellular change in Alzheimer's disease. *Hum Mol Genet*. 13 (2004) 1225-1240.
- [26]. J.W. Lustbader, M. Cirilli, C. Lin, H.W. Xu, K. Takuma, N. Wang, C. Caspersen, X. Chen, S. Pollak, M. Chaney, F. Trinchese, S. Liu, F. Gunn-Moore, L.F. Lue, D.G. Walker, P. Kuppasamy, Z.L. Zewier, O. Arancio, D. Stern, S.S. Yan, H. Wu, ABAD directly links Abeta to mitochondrial toxicity in Alzheimer's disease. *Science*. 304 (2004) 448-452.
- [27]. P.H. Reddy, Mitochondrial dysfunction in aging and Alzheimer's disease: strategies to protect neurons. *Antioxid Redox Signal*. 9 (2007), 1647-1658.
- [28]. M.P. Mattson, M. Gleichmann, A. Cheng, Mitochondria in neuroplasticity and neurological disorders. *Neuron*. 60 (2008) 748-766.

- [29]. P.H. Reddy, T.P. Reddy, M. Manczak, M.J. Calkins, U. Shirendeb, P.Mao, Dynamin-Related Protein 1 and Mitochondrial Fragmentation in Neurodegenerative Diseases. *Brain Res Rev*, in press.
- [30]. P.J. Hollenbeck, W.M. Saxton, The axonal transport of mitochondria. *J Cell Sci.* 118 (2005) 5411-5419.
- [31]. P.H. Reddy, Amyloid beta, mitochondrial structural and functional dynamics in Alzheimer's disease. *Exp Neurol.* 218 (2009) 286-292.
- [32]. M. Manczak, P. Mao, M.J. Calkins, A. Cornea, A.P. Reddy, M.P. Murphy, H.H. Szeto, B. Park, P.H. Reddy, Mitochondria-targeted antioxidants protect against amyloid-beta toxicity in Alzheimer's disease neurons. *J Alz Dis* 20 (2010) S609-S631.
- [33]. A. Martínez, A. J. Lübke, J.A. Del Río, E. Soriano, M. Frotscher, Regional variability and postsynaptic targets of chandelier cells in the hippocampal formation of the rat. *J Comp Neurol.* (1996) 376, 28-44.
- [34]. M. R. Brown, P.G. Sullivan, J.W. Geddes, Synaptic mitochondria are more susceptible to Ca²⁺ overload than nonsynaptic mitochondria. *J Biol Chem* (2006) 281, 11658–11668.
- [35]. Y. Rui, P. Tiwari, Z. Xie, J.Q. Zheng, Acute impairment of mitochondrial trafficking by beta-amyloid peptides in hippocampal neurons. *J Neurosci* 26 (2006) 10480-10487.
- [36]. H. Du, L. Guo, S. Yan, A.A. Sosunov, G.M. McKhann, S.S. Yan, Early deficits in synaptic mitochondria in an Alzheimer's disease mouse model. *Proc Nat Acad Sci USA* 107(2010) 18670-18675.
- [37]. G. Pigino, G. Morfini, Y. Atagi, A. Deshpande, C. Yu, L. Jungbauer, M. SaDu, J. Busciglio, S. Brady, Disruption of fast axonal transport is a pathogenic mechanism for intraneuronal amyloid beta. *Proc Nat Acad Sci USA* 106 (2009) 5907-5912.
- [38]. H. Decker, K.Y. Lo, S.M. Unger, S.T. Ferreira, M.A. Silverman, Amyloid-beta peptide oligomers disrupt axonal transport through an NMDA receptor-dependent mechanism that is mediated by glycogen synthase kinase 3beta in primary cultured hippocampal neurons. *J Neurosci* 30 (2010) 9166-9171
- [39]. K.A. Vossel, K. Zhang, J. Brodbeck, A.C. Daub, P. Sharma, S. Finkbeiner, B. Cui, L. Mucke, Tau reduction prevents Abeta-induced defects in axonal transport. *Science* 330 (2010) 198.
- [40]. D.T. Chang, A.S. Honick, I.J. Reynolds, Mitochondrial trafficking to synapses in cultured primary cortical neurons. *J Neurosci.* 26 (2006) 7035-7045.
- [41]. A. Misko, S. Jiang, I. Wegorzewska, J. Milbrandt, R.H. Baloh, Mitofusion 2 is necessary for transport of axonal mitochondrial and interacts with the Miro/Milton complex. *J Neurosci* 30 (2010) 4232-4240.
- [42]. A. Weihofen, K.J. Thomas, B.L. Ostaszewski, M.R. Cookson, D.J. Selkoe, Pink1 forms a multiprotein complex with Miro and Milton, linking Pink1 function to mitochondrial trafficking. *Biochemistry* 48 (2009) 2045-2052
- [43]. R.V. Gutala, P.H. Reddy, The use of real-time PCR analysis in a gene expression study of Alzheimer's disease post-mortem brains. *J Neurosci Methods.* 132 (2004) 101-107.

[44]. P.H. Reddy, G. Mani, B.S. Park, J. Jacques, G. Murdoch, W. Jr. Whetsell, J. Kaye, M. Manczak, Differential loss of synaptic proteins in Alzheimer's disease: implications for synaptic dysfunction. *J Alzheimers Dis.* (2005) 103-117; discussion 173-80.

[45]. P.D. Coleman, P.J. Yao, Synaptic slaughter in Alzheimer's disease. *Neurobiol Aging.* 24 (2003) 1023-1027.

[46]. C.G. Almeida, D. Tampellini, R.H. Takahashi, P. Greengard, M.T. Lin, E.M. Snyder, G.K. Gouras, Beta-amyloid accumulation in APP mutant neurons reduces PSD-95 and GluR1 in synapses. *Neurobiol Dis.* 20(2005) 187-198.

ACCEPTED MANUSCRIPT




Figure 1




Figure 2




Figure 3

ACCEPTED MANUSCRIPT




Figure 4




Figure 5

Research highlights

1. We investigated toxic effects of amyloid beta ($A\beta$) on mitochondrial axonal transport in $A\beta$ -treated mouse hippocampal neurons.
2. Mitochondrial mass, mitochondrial motility and mitochondrial anterograde transport were reduced in $A\beta$ -treated neurons.
3. A reduction in the number of mitochondria in distal regions of neurons and synaptic branching were observed.
4. We conclude that $A\beta$ impairs axonal transport of mitochondria and may cause synaptic degeneration.

ACCEPTED MANUSCRIPT