

A moment inequality of the Marcinkiewicz-Zygmund type for some weakly dependent random fields

Lionel Truquet

▶ To cite this version:

Lionel Truquet. A moment inequality of the Marcinkiewicz-Zygmund type for some weakly dependent random fields. Statistics and Probability Letters, 2010, 80 (21-22), pp.1673. 10.1016/j.spl.2010.07.011 . hal-00671842

HAL Id: hal-00671842 https://hal.science/hal-00671842

Submitted on 19 Feb2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

A moment inequality of the Marcinkiewicz-Zygmund type for some weakly dependent random fields

Lionel Truquet

PII:	S0167-7152(10)00205-1
DOI:	10.1016/j.spl.2010.07.011
Reference:	STAPRO 5749

To appear in: Statistics and Probability Letters

Received date:4 June 2010Revised date:4 July 2010Accepted date:9 July 2010



Please cite this article as: Truquet, L., A moment inequality of the Marcinkiewicz-Zygmund type for some weakly dependent random fields. *Statistics and Probability Letters* (2010), doi:10.1016/j.spl.2010.07.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A moment inequality of the Marcinkiewicz-Zygmund type for some weakly dependent random fields

Lionel Truquet

July 4, 2010

Abstract

The goal of this note is to give a new moment inequality for sums of some weakly dependent random fields. Our result extends moment bounds given by Wu (2007) [14] or Liu & Lin (2008) [11] for causal autoregressive processes and follows using recursive applications of the Burkhölder inequality for martingales.

AMS subject classification: Primary: 60G60, secondary: 60F99. **Keywords:** random fields, weak dependence, moment inequalities.

1 Introduction

In this paper, we consider random fields X indexed by \mathbb{Z}^d , $d \ge 1$ and which can be written as a Bernoulli shift, e.g

$$X_t = H\left((\xi_{t-j})_{j \in \mathbb{Z}^d} \right), \quad t \in \mathbb{Z}^d, \tag{1}$$

where ξ is an independent and identically distribued E-valued random field and H is a real-valued measurable function. We consider here $E = \mathbb{R}^k$ with $k \in \mathbb{N}^* \cup \{\infty\}$ but a more general measurable space can be considered. A random field of the form (1) is strictly stationary. The most well-known examples of such random fields are linear random fields, i.e

$$X_t = \sum_{j \in \mathbb{Z}^d} a_j \xi_{t-j}, \quad t \in \mathbb{Z}^d.$$
⁽²⁾

Spatial AR processes which write as linear random fields were extensively studied in spatial statistics (see [9] for a nice presentation).

Others (nonlinear) examples of such random fields are given in [8] as solutions of autoregressive équations of the form:

$$X_t = F\left((X_{t-j})_{j \in \mathbb{Z}^d}; \xi_t \right), \quad t \in \mathbb{Z}^d,$$

where F is a Lipschitzian function.

In this note, we prove moment inequalities of the Marcinkiewicz-Zygmund type for the partial sums of a random field X of the form (1). More precisely we prove that for a real number q > 1,

$$||S(B)||_q \le C_q^d \sum_{k \in \mathbb{Z}^d} ||p_{d,k_d}^{(\xi)} \circ \dots \circ p_{1,k_1}^{(\xi)}(X_0)||_q |B|^{1/q'},$$
(3)

where B is a finite subset of \mathbb{Z}^d , $S(B) = \sum_{j \in B} X_j$, $q' = \min(q, 2)$ and

for $(s, \ell) \in \{1, \ldots, d\} \times \mathbb{Z}, p_{s,\ell}^{(\xi)}$ denotes the projection operator defined for an integrable random variable Z by

$$p_{s,\ell}^{(\xi)}(Z) = Z - \mathbb{E}\left(Z/\sigma(\xi_j/j_s \neq \ell)\right).$$
(4)

The constant C_q is the universal constant of the Burkhölder inequality (see [10]) and then it does not depend on the distribution of the random field. $\|\cdot\|_q$ denotes the usual norm of \mathbb{L}^q . Denoting

$$A_{q} = \sum_{k \in \mathbb{Z}^{d}} \| p_{d,k_{d}}^{(\xi)} \circ \dots \circ p_{1,k_{1}}^{(\xi)}(X_{0}) \|_{q},$$
(5)

 A_q is a postive constant depending only on q and the distribution of the random field ξ . Obviously inequality (3) is interesting only when $A_q < \infty$. The latter condition basically indicates short-range dependence. Then we study in details the finiteness of the constant A_q when the random field X writes as a functional of linear fields.

An inequality of the form (3) is a useful tool to study the behaviour of sums of dependent random variables. Strong laws of large numbers, strong invariance principle and other variants of limit theorems are often based on this kind of inequality. Some existing methods to obtain such inequalities use mixing coefficients. ϕ -mixing random fields are considered in Dedecker [3] and Delyon [5], an inequality for α -mixing random fields is given in Doukhan [6]. Unfortunately, for random fields defined by (1), mixing conditions lead to restrictive assumptions on the random field distribution (see [9] for the case of linear random fields) or are impossible to check (see the counterexample of Andrews [1] when d = 1). On the other hand, general weak dependence conditions formulated in terms of covariances of Lipschitzian functionals can be applied to random fields defined by (1) and corresponding moment inequalities are available. We refer to [4] and [7] for weak dependence conditions that generalize strong mixing and to [2] for a generalization of associated random fields. Nevertheless, despite their generality, such approaches do not exploit the particular form of the random field when it writes as a functional of independent random variables and lead to complicated and restrictive conditions for the representation (1). When d = 1, martingales decompositions and Burkhölder inequality for martingales have been used to prove simple moment bounds for partial sums of causal processes of the form (2). We refer to [14] and [11] for details. In this paper, we generalize this approach to random fields and we prove inequality (3) using a simple recursive application of the Burkhölder inequality. We next study in details the application of such inequalities to partial sums of transforms of linear random fields.

2 Moment bounds

We first state basic properties of specific conditional expectations and operators $p_{s,\ell}$ defined in (4).

Lemma 1 Let Z be an integrable random variable, ξ an i.i.d random field and q a real number with $q \ge 1$. Then:

1. If $A, B \subset \mathbb{Z}^d$, $\mathbb{E}\left(\mathbb{E}\left(Z/\xi_A\right)/\xi_B\right) = \mathbb{E}\left(Z/\xi_{A\cap B}\right).$ 2. If $A \subset \mathbb{Z}^d$ and $B \subset \mathbb{Z}^d \setminus A$:

$$\|\mathbb{E}\left(Z/\xi_{A\cup B}\right) - \mathbb{E}\left(Z/\xi_{A}\right)\|_{q} \le \|Z - \mathbb{E}\left(Z/\xi_{\mathbb{Z}^{d}\setminus B}\right)\|_{q}.$$

3. If
$$(s, s', \ell, \ell') \in \{1, \ldots, d\}^2 \times \mathbb{Z} \times \mathbb{Z}$$

$$\|p_{s,\ell}^{(\xi)} \circ p_{s',\ell'}^{(\xi)}(Z)\|_q \le 2\min\left(\|p_{s,\ell}^{(\xi)}(Z)\|_q, \|p_{s',\ell'}^{(\xi)}(Z)\|_q\right).$$

Proof of Lemma 1

- 1. The result follows from the independence properties of the random field ξ .
- 2. From the point 1., we have:

$$\mathbb{E}\left(Z/\xi_{A\cup B}\right) - \mathbb{E}\left(Z/\xi_A\right) = \mathbb{E}\left(Z - \mathbb{E}\left(Z/\xi_{\mathbb{Z}^d/B}\right)/\xi_{A\cup B}\right),$$

and the result is a consequence of the Jensen inequality.

3. We have from the point 1.:

$$p_{s,\ell}^{(\xi)} \circ p_{s',\ell'}^{(\xi)}(Z) = Z - \mathbb{E}\left(Z/\xi_A\right) + \mathbb{E}\left(Z - \mathbb{E}\left(Z/\xi_A\right)/\xi_B\right)$$

for $A = \{j \in \mathbb{Z}^d / j_s \neq \ell\}$ and $B = \{j \in \mathbb{Z}^d / j_{s'} \neq \ell'\}$ and from the triangular inequality and Jensen inequality, we obtain:

$$\|p_{s,\ell}^{(\xi)} \circ p_{s',\ell'}^{(\xi)}(Z)\|_q \le 2\|p_{s,\ell}^{(\xi)}(Z)\|_q.$$

Exchanging A and B, the result follows. \Box

The main result of the paper can now be stated.

Theorem 1 Let X be a real centered random field indexed by \mathbb{Z}^d , $d \ge 1$ and such that $\mathbb{E}|X_t| < \infty$ for all $t \in \mathbb{Z}^d$. Assume furthermore that there exists an i.i.d random field ξ such that $\sigma(X) \subset \sigma(\xi)$. Then for a real number q > 1, we have with $q' = \min(q, 2)$:

$$\|S(B)\|_{q} \le C_{q}^{d} \sum_{k \in \mathbb{Z}^{d}} \left(\sum_{t \in B} \|p_{d, t_{d} + k_{d}}^{(\xi)} \circ \dots \circ p_{1, t_{1} + k_{1}}^{(\xi)}(X_{t})\|_{q}^{q'} \right)^{1/q'}, \tag{6}$$

where C_q denotes the universal constant of the Burkhölder inequality for martingales.

Remarks

1. The proof of inequality (6) do not use the representation (1) and the stationarity assumption. Note that for a stationary random field, inequality (6) coincides with (3).

2. Let X be a random field of the form (1) and such that $A_q < \infty$ for a real number q > 1. Then the Móricz theorem [12] can be used to deduce from the bounds (3) a moment inequality for partial maxima when B is a block of \mathbb{Z}^d , i.e

$$B = ((a_1, b_1] \times \ldots \times (a_d, b_d]) \cap \mathbb{Z}^d.$$

More precisely, if q > 2, we have:

$$\|\max_{W \triangleleft B} |S(W)|\|_q \le (5/2)^{d/q} \left(1 - 2^{\frac{2-q}{2q}}\right)^{-d} C_q^d A_q |B|^{1/2},$$

where the notation $W \triangleleft B$ means $W \subset B$ and W is a block of \mathbb{Z}^d with the same minimal vertex than B, i.e

$$W = ((a_1, c_1] \times \ldots \times (a_d, c_d]) \cap \mathbb{Z}^d.$$

In fact the result of Móricz is analytic, in the sense that it does not involve any dependence properties. When $q \in (1, 2)$, it gives the bound

$$\|\max_{W \triangleleft B} |S(W)|\|_q \le 5^{d/q} 2^{d(q-1)/q} C_q^d A_q |B|^{1/q} \prod_{i=1}^d \log(b_i - a_i).$$

Proof of Theorem 1

1. We first prove the result for d = 1. For $j \in \mathbb{Z}$ and an integrable random variable Z, we define:

$$\mathcal{P}_i(Z) = \mathbb{E}\left(Z/\mathcal{F}_i\right) - \mathbb{E}\left(Z/\mathcal{F}_{i-1}\right),$$

where $\mathcal{F}_i = \sigma(\xi_j / j \leq i)$. We use the following decomposition

$$X_{i} = X_{i} - \mathbb{E} (X_{i}/\mathcal{F}_{i}) + \mathbb{E} (X_{i}/\mathcal{F}_{i})$$
$$= \sum_{j \ge 1} \mathcal{P}_{i+j}(X_{i}) + \sum_{j \ge 0} \mathcal{P}_{i-j}(X_{i})$$
$$= \sum_{j \in \mathbb{Z}} \mathcal{P}_{i+j}(X_{i})$$

Then we have $S(B) = \sum_{j \in \mathbb{Z}} \sum_{i \in B} \mathcal{P}_{i+j}(X_i)$, and $(\mathcal{P}_{i+j}(X_i))_{i \in \mathbb{Z}}$ is a martingale difference for the filtration $(\mathcal{F}_{i+j})_{i \in \mathbb{Z}}$. The Burkhölder inequality leads to:

$$\|S(B)\|_q \le \sum_{j \in \mathbb{Z}} \|\sum_{i \in B} \mathcal{P}_{i+j}(X_i)\|_q \le C_q \sum_{j \in \mathbb{Z}} \left\{ \mathbb{E} \left| \sum_{i \in B} \mathcal{P}_{i+j}(X_i)^2 \right|^{q/2} \right\}^{1/q}$$

If $q \ge 2$, the triangular inequality for the $\mathbb{L}^{q/2}$ norm leads to

$$||S(B)||_q \le C_q \sum_{j \in \mathbb{Z}} \left\{ \sum_{i \in B} ||\mathcal{P}_{i+j}(X_i)||_q^2 \right\}^{1/2}.$$

Setting $A = \{\ell \in \mathbb{Z}/\ell \le i+j-1\}$ and $B = \{i+j\}$, an application of point 2 in Lemma 1 yields to

$$|\mathcal{P}_{i+j}(X_i)||_q \le ||p_{1,i+j}^{(\xi)}(X_i)||_q$$

Inequality (6) follows.

If now $q \in (1,2]$, inequality (6) is a consequence of the previous remark and the bound

$$\left(\sum_{i\in B} \mathcal{P}_{i+j}(X_i)^2\right)^{q/2} \le \sum_{i\in B} |\mathcal{P}_{i+j}(X_i)|^q.$$

2. We show the result for $d \ge 2$ using induction on d. Suppose that the bound (6) holds for any random field indexed by \mathbb{Z}^s , $s \le d-1$ and satisfying the assumptions of Theorem 1. We have $X = (X_{i,j})_{i \in \mathbb{Z}, j \in \mathbb{Z}^{d-1}}$ and

$$S(B) = \sum_{i \in A} Y_i, \qquad Y_i = \sum_{j \in B_i} X_{i,j},$$

where $B = \bigcup_{i \in A} \{i\} \times B_i$. It is obvious that $(Y_i)_{i \in \mathbb{Z}}$ satisfies the assumptions of Theorem 1 with the random field $\xi'_i = \xi_{i,\mathbb{Z}^{d-1}}, i \in \mathbb{Z}$. Then, using inequality (6) for d = 1, we obtain

$$\|S(B)\|_{q} \le C_{q} \sum_{k_{1} \in \mathbb{Z}} \left(\sum_{i \in A} \|p_{1,i+k_{1}}^{(\xi')}(Y_{i})\|_{q}^{q'} \right)^{1/q'}.$$
(7)

Note also that for $\ell \in \mathbb{Z}$, we have $p_{1,\ell}^{\xi'} = p_{1,\ell}^{\xi}$. Then for $k_1 \in \mathbb{Z}$ and $i \in A$, we have $p_{1,i+k_1}^{(\xi')}(Y_i) = \sum_{j \in B_i} p_{1,k_1}^{(\xi)}(X_{i,j})$. Since the random field $\left(p_{1,k_1}^{(\xi)}(X_{i,j})\right)_{j \in B_i}$ is indexed by \mathbb{Z}^{d-1} and satisfies the assumptions of Theorem 1 with the i.i.d random field $\xi'' = (\xi_{\mathbb{Z},j})_{j \in \mathbb{Z}^{d-1}}$, we can apply the induction hypothesis. We observe that for $s \in \{1, \ldots, d-1\}, p_{s,k}^{(\xi'')} = p_{s+1,k}^{(\xi)}$. Then we get the following bound:

$$\|p_{1,i+k_{1}}^{(\xi)}(Y_{i})\|_{q}$$

$$\leq C_{q}^{d-1} \sum_{k_{2},\dots,k_{d} \in \mathbb{Z}} \left(\sum_{j \in B_{i}} \|p_{d,j_{d-1}+k_{d}}^{(\xi)} \circ \cdots \circ p_{2,j_{1}+k_{2}}^{(\xi)} \left(p_{1,i+k_{1}}^{(\xi)}(X_{i,j})\right)\|_{q}^{q'} \right)^{1/q'}$$

$$\leq C_{q}^{d-1} \sum_{k_{2},\dots,k_{d} \in \mathbb{Z}} \left(\sum_{j \in B_{i}} \|p_{d,j_{d-1}+k_{d}}^{(\xi)} \circ \cdots \circ p_{2,j_{1}+k_{2}}^{(\xi)} \circ p_{1,i+k_{1}}^{(\xi)}(X_{i,j})\|_{q}^{q'} \right)^{1/q'}.$$

Then, using the triangular inequality for the norm $||x|| = \left(\sum_i |x_i|^{q'}\right)^{1/q'}$, we obtain

$$\left(\sum_{i\in A} \|p_{1,i+k_1}^{(\xi)}(Y_i)\|_q^{q'}\right)^{1/q'} \le C_q^{d-1} \sum_{k_2,\dots,k_d\in\mathbb{Z}} \left(\sum_{i\in A} \sum_{j\in B_i} \|p_{d,j_{d-1}+k_d}^{(\xi)} \circ \cdots \circ p_{1,i+k_1}^{(\xi)}(X_{i,j})\|_q^{q'}\right)^{1/q'}$$

and the bound (6) follows from (7) using the equality $\sum_{i \in A} \sum_{j \in B_i} = \sum_{(i,j) \in B} \square$

The iterations involving the operators $p_{s,\ell}^{(\xi)}$ are not always easy to evaluate. The following corollary will be useful to verify the condition $A_q < \infty$ in most of examples.

Corollary 1 Let X be a real random field satisfying the assumptions of Theorem 1. Then

$$A_q \le 2^{d-1} \sum_{s=1}^d \sum_{n \in \mathbb{N}} (2n+1)^{d-1} \sum_{|k_s|=n} c_{s,k_s},\tag{8}$$

where for $s \in \{1, ..., d\}$,

$$c_{s,k_s} = \left(\sum_{t \in B} \|p_{s,t_s+k_s}^{(\xi)}(X_t)\|_q^{q'}\right)^{1/q}$$

Remarks

1. When the random field X writes as in (1) we have the bound

$$\|p_{s,t_s+k_s}^{(\xi)}(X_t)\|_q \le \|X_t - \widetilde{X}_{t,s,k_s}\|_q,\tag{9}$$

where $\widetilde{X}_{t,s,k_s} = H\left(\left(\widetilde{\xi}_{t-j}\right)_{j\in\mathbb{Z}^d}\right)$, $\widetilde{\xi}_{t-j} = \xi_{t-j}$ if $j_s \neq -k_s$ and ξ'_{t-j} otherwise, ξ' being a copy of ξ . When d = 1, the right hand side of (9) is the coefficient denoted by $\theta_{q,k}$ used by Wu [13] to measure the dependence of a stationary process X. In this case, condition $A_q < \infty$ holds when

$$\sum_{k \in \mathbb{Z}} \| p_{1,k}^{(\xi)}(X_0) \|_q < \infty,$$

which is similar to the condition $\sum_{k\leq 0} \theta_{q,k} < \infty$ used by Liu and Lin [11] (see Lemma A1 and Lemma A2 of their paper) for a causal stationary process

$$X_t = H\left(\xi_{t-j}\right)_{j\geq 0}, \quad t\in\mathbb{Z}.$$

2. Note that when d = 1, A_q is finite if and only if the right hand side of inequality (8) is finite. For $d \ge 2$ this is no more true. For example, for the simple example of linear random fields (2), we have

$$A_q = \sum_{k \in \mathbb{Z}^d} |a_k| \times \|\xi_0\|_q.$$

In contrast, when q = 2 and ξ_0 is square integrable, the right hand side of (8) is finite provided for $s = 1, \ldots, d$

$$\sum_{n \in \mathbb{N}} n^{d-1} \sqrt{\sum_{|j_s|=n} a_j^2} < \infty,$$

which is more restrictive than the summability of coefficients (a_i) .

Proof of Corollary 1 From point 3 in Lemma 1, a straightforward induction leads to the following inequality:

$$\|p_{d,t_d+k_d}^{(\xi)} \circ \dots \circ p_{1,t_1+k_1}^{(\xi)}(X_t)\|_q \le 2^{d-1} \wedge_{s=1}^d \|p_{\ell,k_s+t_s}^{(\xi)}(X_t)\|_q.$$
(10)

Assume that the random field X satisfies the assumptions of Theorem 1. We have

$$\sum_{k \in \mathbb{Z}^d} \left(\sum_{t \in B} \wedge_{s=1}^d \| p_{s,t_s+k_s}^{(\xi)}(X_t) \|_q \right)^{1/q'}$$

$$\leq \sum_{s=1}^d \sum_{n \in \mathbb{N}} \sum_{k/\|k\|_{\infty} = |k_s| = n} c_{s,k_s}$$

$$\leq \sum_{s=1}^d \sum_{n \in \mathbb{N}} (2n+1)^{d-1} \sum_{|k_s| = n} c_{s,k_s}.$$

Then the conclusion of Corollary 1 follows. \Box

3 Examples

In the sequel ξ denotes an i.i.d random field satisfying $\mathbb{E}\xi_0 = 0$. For a random field X of the form (1), inequality (3) is interesting provided the constant A_q defined in (5) is finite. For the simple case of a linear random field (2), we have

$$A_q = \sum_{k \in \mathbb{Z}^d} |a_j| \|\xi_0\|_q.$$

Then provided that $\xi_0 \in \mathbb{L}^q$, we have $A_q < \infty$ if and only if $\sum_{j \in \mathbb{Z}^d} |a_j| < \infty$. The latter condition is a weak dependence condition since in this case the autocovariances of the field are summable.

A precise evaluation of the constant A_q may also be possible when the random field X has a polynomial expression with respect to the coordinates of ξ . As an example, we investigate a moment bound for the covariances of linear random fields.

Corollary 2 Let
$$Y_t = \sum_{j \in \mathbb{Z}^d} a_j \xi_{t-j}, t \in \mathbb{Z}^d$$
 with $\sum_{j \in \mathbb{Z}^d} |a_j| < \infty$, and for a given $h \in \mathbb{Z}^d$
$$X_t = Y_t Y_{t+h} - \mathbb{E}(Y_t Y_{t+h}).$$

Assume that $\xi_0 \in \mathbb{L}^{2m}$ with m > 1. Then for $1 < q \le m$, $A_q < \infty$.

Proof of Corollary 2 Let $k \in \mathbb{Z}^d$ and $B = p_{d,t_d+k_d}^{(\xi)} \circ \cdots \circ p_{1,t_1+k_1}^{(\xi)}(X_t)$. Then we are going to prove

$$B = \sum_{(\alpha,\beta)\in I} \left(\sum_{j\in A_{\alpha}} a_{j}\xi_{t-j} \right) \times \left(\sum_{j\in A_{\beta}} a_{j+h}\xi_{t-j} \right) + a_{-k}a_{-k+h} \left(\xi_{t+k}^{2} - \mathbb{E}\xi_{0}^{2} \right), \quad (11)$$

where

$$I = \bigcap_{s=1}^{d} \{ (\alpha, \beta) \in \{0, 1\}^d \times \{0, 1\}^d / (\alpha, \beta) \neq (0, 0) \text{ and } \alpha_s \beta_s = 0 \},$$

$$A_{\alpha} = \bigcap_{s=1}^{d} \{ j \in \mathbb{Z}^d / j_s = -k_s \text{ if } \alpha_s = 0 \text{ and } \alpha_s = 1 \text{ otherwise } \}.$$

One can easily see that if $(\alpha, \beta) \in I$, then $\alpha \neq \beta$ and since there exists $s \in \{1, \ldots, d\}$ such that $\alpha_s = 0$ and $\beta_s = 1$ (or the contrary) then the sums $\sum_{j \in A_\alpha} a_j \xi_{t-j}$ and $\sum_{j \in A_\beta} a_j \xi_{t-j}$ are independent. Before giving a proof of (11), we show why the conclusion of Corollary 2 holds. From (11) we obtain the following bound:

$$\|B\|_{q} \leq \sum_{(\alpha,\beta)\in I} \sum_{j\in A_{\alpha}} |a_{j}| \cdot \sum_{j\in A_{\beta}} |a_{j+h}| \|\xi_{0}\|_{q}^{2} + |a_{-k}a_{-k+h}(\xi_{0}^{2} - \sigma^{2})|,$$

where $\sigma^2 = \mathbb{E}\xi_0^2$. Let $q \in (1, m]$. Using assumptions on ξ and coefficients $(a_j), A_q < \infty$ if for $(\alpha, \beta) \in I$,

$$G = \sum_{k \in \mathbb{Z}^d} \left(\sum_{j \in A_\alpha} |a_j| \cdot \sum_{j \in A_\beta} |a_{j+h}| \right) < \infty.$$
(12)

If $\alpha = 0$, then $A_{\alpha} = -k$ and $G \leq \sum_{j} |a_{j}| \times \sum_{j} |a_{j+h}| < \infty$. The same holds if $\beta = 0$. Now suppose that $\ell = \sharp \{s \in \{1, \ldots, d\} / \alpha_{s} = 0\}$ satisfies $1 \leq \ell \leq d - 1$. Without loss of generality, we assume that $\alpha_{s} = 0$, $s \leq \ell$. Then by definition of I, we have $\beta_{s} = 0$ if $s \geq \ell + 1$. In this case, we obtain

$$G \leq \sum_{k \in \mathbb{Z}^d} \left(\sum_{j_{\ell+1}, \dots, j_d} |a_{-k_1, \dots, -k_\ell, j_{\ell+1}, \dots, j_d}| \right) \cdot \left(\sum_{j_1, \dots, j_\ell} |a_{j_1+h_1, \dots, j_\ell+h_\ell, -k_{\ell+1}+h_{\ell+1}, \dots, -k_d+h_d}| \right) \\ \leq \sum_{j \in \mathbb{Z}^d} |a_j| \cdot \sum_{j \in \mathbb{Z}^d} |a_{j+h}|.$$

Then $G < \infty$ and the conclusion of Corollary 2 follows.

Now we prove (11). We first introduce some notations. For $s \in \{1, \ldots, d\}$, let

$$I^{(s)} = \bigcap_{i=1}^{s} \{ (\alpha_1, \dots, \alpha_s, \beta_1, \dots, \beta_s) \in \{0, 1\}^{2s} \ \alpha_i \beta_i = 0 \},$$

$$A^{(s)}_{\alpha} = \bigcap_{i=1}^{s} \{ j \in \mathbb{Z}^d \ / \ j_i = -k_i \text{ if } \alpha_i = 0, \text{ and } j_i \neq -k_i \text{ otherwise } \},$$

$$B_s = p^{(\xi)}_{s, t_s + k_s} \circ \dots \circ p^{(\xi)}_{1, t_1 + k_1}(X_t).$$

Using a finite induction on the set $\{1, \ldots, d\}$, we are going to prove that for $s \in \{1, \ldots, d\}$,

$$B_s = \sum_{(\alpha,\beta)\in I^{(s)}} \left(\sum_{j\in A^{(s)}_{\alpha}} a_j \xi_{t-j}\right) \times \left(\sum_{j\in A^{(s)}_{\beta}} a_{j+h} \xi_{t-j}\right) - \sum_{j_i=-k_i, i=1,\dots,s} a_j a_{j+h} \sigma^2.$$
(13)

Observe that $B_d = B$. The proof of this induction uses the following remark: for two subsets C and D of \mathbb{Z}^d , we have for $s \in \{1, \ldots, d\}$ and $Z_t = \sum_{j \in C} a_j \xi_{t-j} \cdot \sum_{j \in D} a_{j+h} \xi_{t-j}$,

$$p_{s,t_s+k_s}(Z_t) = \sum_{j \in C, j_s = -k_s} a_j \xi_{t-j} \cdot \sum_{j \in D, j_s \neq -k_s} a_{j+h} \xi_{t-j}$$

$$+ \sum_{j \in C, j_s \neq -k_s} a_j \xi_{t-j} \cdot \sum_{j \in D, j_s = -k_s} a_{j+h} \xi_{t-j}$$

$$- \sigma^2 \sum_{j \in C \cap D, j_s = -k_s} a_j a_{j+h}.$$

- The case s = 1 is an easy consequence of the previous remark.
- Suppose that equality (13) holds for B_s . Then applying again the previous equality to $B_{s+1} = p_{s+1,k_{s+1}}(B_s)$, equality (13) easily follows for B_{s+1} .

The next corollary gives a sufficient condition in order to have $A_q < \infty$ for some locally Hölderian functionals of linear random fields. Here we apply Corollary 1.

Corollary 3 Let $h : \mathbb{R} \to \mathbb{R}$ be a function such that there exist $a \ge 0, b \in (0,1]$ and a positive constant K satisfying:

$$|h(x) - h(y)| \le K (1 + |x|^a + |y|^a) |x - y|^b, \quad x, y \in E.$$

For $t \in \mathbb{Z}^d$, let $Y_t = \sum_{j \in \mathbb{Z}^d} a_j \xi_{t-j}$, $t \in \mathbb{Z}^d$ and $X_t = h(Y_t) - \mathbb{E}h(Y_t)$. Assume that $\mathbb{E}|\xi_0|^m < \infty$ with m > 1, $m \ge (a+b)q$ and q > 1. If for $s \in \{1, \ldots, d\}$

$$\sum_{n \ge 0} (n+1)^{d-1} \left(\sum_{j/|j_s|=n} a_j^2 \right)^{b/2} < \infty,$$
(14)

then $A_q < \infty$.

In particular, if $a_j = O(||j||^{-\alpha})$ with $\alpha > \frac{(2+b)d-b}{2b}$, condition (14) is satisfied. Remarks

- 1. Corollary 3 shows that inequality $A_q < \infty$ holds with a loss for the decrease of coefficients (a_j) with respect to the condition $\sum_{j \in \mathbb{Z}^d} |a_j| < \infty$. For example, when $a_j =$ $O(||j||^{-\alpha})$ the summability of coefficients $(a_j)_j$ holds when $\alpha > d$, but $\frac{(2+b)d-b}{2b} > d$ when $d \ge 2$.
- 2. The result of Corollary 1 can also be applied if the functional h of Corollary 3 has some discontinuities. However, additional assumptions could be required in order to bound explicitly the quantity $\|p_{s,\ell}^{(\xi)}(X_0)\|_q$. For example if $h(x) = \mathbb{1}_{x>t}$, one can use an approximation of h by the Lipschitz function $h_{\epsilon}(x) = \mathbb{1}_{x>t} + \left(\frac{1}{\epsilon}x + 1 - \frac{t}{\epsilon}\right)\mathbb{1}_{t-\epsilon < x \leq t}$ We obtain

$$\|p_{s,\ell}^{(\xi)}(X_0)\|_q \le 2\|X_0 - h_{\epsilon}(Y_0)\|_q + \|p_{s,\ell}^{(\xi)}(h_{\epsilon}(Y_0))\|_q$$

Then we can bound the second term as in the proof of Corollary 3 and we get,

$$\|p_{s,\ell}^{(\xi)}(X_0)\|_q \le 2\mathbb{P}\left(t - \epsilon < Y_0 \le t\right) + \frac{2}{\epsilon} \|\xi_0\|_q \sqrt{\sum_{j_s = -\ell} a_j^2}.$$

Some regularity assumptions have be made for optimizing the previous bound in ϵ (e.g $\|p_{s,\ell}^{(\xi)}(X_0)\|_q \sim \left(\sum_{j_s=-\ell} a_j^2\right)^{1/4}$ if Y_0 has a locally bounded density).

Proof of Corollary 3 We first derive a bound for the quantities $||p_{s,t_s+k_s}(X_t)||_q$, s = $1, \ldots, d$ and $q \leq \frac{m}{a+b}$. For a copy ξ' of ξ , we set

$$Z = K \left(1 + \left| \sum_{j \in \mathbb{Z}^d} a_j \xi_{t-j} \right|^a + \left| \sum_{j/j_s \neq -k_s} a_j \xi_{t-j} + \sum_{j/j_s = -k_s} a_j \xi'_{t-j} \right|^a \right) \left| \sum_{j/j_s = -k_s} a_j (\xi_{t-j} - \xi'_{t-j}) \right|^b.$$

Then using the assumption on h, it is not difficult to prove that $||p_{s,t_s+k_s}(X_t)||_q \leq ||Z||_q$. If a > 0, we consider two positive numbers p and r satisfying $\frac{1}{q} = \frac{1}{p} + \frac{1}{r}$, $ar \le m$ and $m \ge bp > 1$. A such choice is possible from the assumptions on m. In the sequel C denotes a generic positive constant not depending on the index k or j. Then using the previous bound and Hölder inequality, we obtain

$$\|p_{s,t_s+k_s}(X_t)\|_q \le C\mathbb{E}^{1/p} |\sum_{j/j_s=-k_s} a_j(\xi_{t-j}-\xi'_{t-j})|^{bp}.$$

Note that the previous bound holds also when a = 0 if we set p = m/b. From Burkhölder inequality, we obtain

$$||p_{s,t_s+k_s}(X_t)||_q \le C\left(\sum_{j/j_s=-k_s} a_j^2\right)^{b/2}.$$
 (15)

The conclusion of the corollary follows easily from (15) and Corollary 1. Suppose now $a_j = O(||j||^{-\alpha})$ with $\alpha > \frac{(2+b)d-b}{2b}$. Then we have for $s \in \{1, \ldots, d\}$

$$\sum_{j/j_s=n} a_j^2 \le C \sum_{j/\|j\| \ge n, j_s=n} \|j\|^{-2\alpha} \le C n^{-2\alpha+d-1}.$$

Using the same bound for $\sum_{j/j_s=-n} a_j^2$, we deduce that condition (14) is satisfied and the

result follows. \Box

Acknowledgments: The author wish to thank an anonymous referee for its constructive suggestions which led to some improvements in some earlier versions of the manuscript.

References

- [1] Andrews, D. (1984) Non strong mixing autoregressive processes. Journal of Applied Probability 21, 930-934.
- [2] Bulinski, A., Shashkin, A. (2006) Strong invariance principle for dependent random fields IMS Lecture Notes-Monograph Series, Vol. 48, 128-143.
- [3] Dedecker, J. (2001) Exponential inequalities and functional central limit theorems for random fields. ESAIM Probab. Statist. 5 77-104.
- [4] Dedecker, J., Doukhan, P., Lang, G., León, J. R., Louhichi, and S., Prieur, C. (2007) Weak dependence, examples and applications, Lecture Notes in Statistics 190.
- [5] Delyon, B. (2009) Exponential inequalities for sums of weakly dependent variables. Electronic Journal of Probability, 752-779.

- [6] Doukhan, P. (1994) Mixing: Properties and Examples. Lecture Notes in Statist. 85. Springer, New York. MR1312160.
- [7] Doukhan P., Mayo N., Truquet L. (2009) Weak dependence, models and some applications. Metrika 69, 2-3, 199-225.
- [8] Doukhan, P., Truquet, L. A fixed point approach to model random fields. Alea 3, 111-132.
- [9] Guyon, X. (1995) Random fields on a network: modeling, statistics, and applications. Graduate Texts in Mathematics. Springer-Verlag, New-York.
- [10] Hall, P., Heyde, C.C. (1980) Martingal limit theory and its application. Academic Press.
- [11] Liu, W., Lin, Z. (2009) Strong approximation for stationary processes. Stochastic Processes and their Applications, Vol 119, 1, 249-280.
- [12] Móricz, F. (1983) A general moment inequality for the maximum of the rectangular partial sums of multiple series. Acta Math. Hung., 41, 3-4, 337-346.
- [13] Wu, W.B., (2005) Nonlinear system theory: Another look at dependence. Proc. Natl. Acad. Sci. USA. 102 (40), 14150-14154.
- [14] Wu, W.B. (2007) Strong invariance principles for dependent random variables. The Annals of Probability, Vol 35, 6, 2294-2320.

UMR 6625 CNRS Institut de Recherche Mathématique de Rennes (IRMAR) Université de Rennes I, Campus de Beaulieu, F-35042 Rennes Cedex, France, and CREST - ENSAI. Timbre J380. Campus de Ker Lann. rue Blaise Pascal. 35170 BRUZ. Email: lionel.truquet@univ-rennes1.fr