



# Segregation of calcium and magnesium into different substructures. Ca<sub>4</sub>Ag<sub>0.948</sub>Mg and other compounds with Gd<sub>4</sub>RhIn-type structure

Marcel Kersting, Samir F. Matar, Christian Schwickert, Rainer Pöttgen

## ► To cite this version:

Marcel Kersting, Samir F. Matar, Christian Schwickert, Rainer Pöttgen. Segregation of calcium and magnesium into different substructures. Ca<sub>4</sub>Ag<sub>0.948</sub>Mg and other compounds with Gd<sub>4</sub>RhIn-type structure. Zeitschrift fur Naturforschung B, 2012, 67 (1), pp.61-69. 10.1515/znb-2012-0111 . hal-00673265

HAL Id: hal-00673265

<https://hal.science/hal-00673265>

Submitted on 26 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Structure  
D 2000

21- 004

DOI: 10.1002/chin.201221004

**Segregation of Calcium and Magnesium into Different Substructures.**

**Ca<sub>4</sub>Ag<sub>0.948</sub>Mg and Other Compounds with Gd<sub>4</sub>RhIn-Type Structure.** — The compounds Ln<sub>4</sub>PdMg (Ln: Ca, Eu, Tb—Lu), Ln<sub>4</sub>AgMg (Ln: Ca, Yb), Ln<sub>4</sub>PtMg (Ln: Eu, Tb—Lu), and Ln<sub>4</sub>AuMg (Ln: Ca, Eu, Yb) are prepared by induction melting of the elements. The structures of Yb<sub>4</sub>PdMg and Ca<sub>4</sub>Ag<sub>0.948</sub>Mg are determined by single crystal XRD. They crystallize with the Gd<sub>4</sub>RhIn-type structure in the space group F43m with Z = 16. Ca<sub>4</sub>Ag<sub>0.948</sub>Mg exhibits segregation of Ca and Mg into different substructures. Calcium forms trigonal prisms around the silver atoms and magnesium forms covalently bonded tetrahedra, which are embedded in larger cavities of the adamantane-like three-dimensional network of edge- and corner-sharing AgCa<sub>6</sub> trigonal prisms. The electronic structure and chemical bonding in Ca<sub>4</sub>AgMg is analyzed by means of DFT calculations. Eu<sub>4</sub>PdMg and Eu<sub>4</sub>PtMg show Curie—Weiss behavior above 230 K. Ferromagnetic ordering occurs at 150.1 and 139.1 K for the Pd- and Pt-containing compound, respectively. — (KERSTING, M.; MATAR, S. F.; SCHWICKERT, C.; POETTGEN\*, R.; Z. Naturforsch., B: Chem. Sci. 67 (2012) 1, 61–69; Inst. Anorg. Anal. Chem., Westfael. Wilhelms-Univ., D-48149 Muenster, Germany; Eng.) — W. Pewestorf