



**HAL**  
open science

## **An Aetiological Mutation Causes Aberrant Striatal Activity and Alters Plasticity during Skill Learning**

Rui M Costa, Catherine Ann French, Xin Jin, Thomas G Campbell, Elissa Gerfen, Matthias Groszer, Simon Fisher

► **To cite this version:**

Rui M Costa, Catherine Ann French, Xin Jin, Thomas G Campbell, Elissa Gerfen, et al.. An Aetiological Mutation Causes Aberrant Striatal Activity and Alters Plasticity during Skill Learning. *Molecular Psychiatry*, 2011, 10.1038/mp.2011.105 . hal-00675395

**HAL Id: hal-00675395**

**<https://hal.science/hal-00675395>**

Submitted on 1 Mar 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **An Aetiological *Foxp2* Mutation Causes Aberrant Striatal Activity and Alters Plasticity during Skill Learning**

CA French<sup>1,4</sup>, X Jin<sup>2</sup>, TG Campbell<sup>1</sup>, E Gerfen<sup>2</sup>, M Groszer<sup>3</sup>, SE Fisher<sup>1,5</sup> and RM Costa<sup>2,4</sup>

<sup>1</sup>*Wellcome Trust Centre for Human Genetics, Roosevelt Drive, Oxford, OX3 7BN, United Kingdom;* <sup>2</sup>*Section on In Vivo Neural Function, Laboratory for Integrative Neuroscience, NIAAA/NIH, 5625 Fishers Lane, MSC 9411 Bethesda, MD 20852-9411, USA;* <sup>3</sup>*Institut du Fer à Moulin, INSERM U839, Université Pierre et Marie Curie (Paris 6), 17 rue du Fer à Moulin, Paris, 75005, France;* <sup>4</sup>*Champalimaud Neuroscience Programme at Instituto Gulbenkian de Ciência, Rua da Quinta Grande 6, 2780-901 Oeiras, Portugal;* <sup>5</sup>*Language and Genetics Department, Max Planck Institute for Psycholinguistics, Wundtlaan 1, 6525 XD Nijmegen, the Netherlands.*

Correspondence: Dr. RM Costa, Champalimaud Neuroscience Programme at Instituto Gulbenkian de Ciência, Rua da Quinta Grande 6, 2780-901 Oeiras, Portugal. E-mail: [ruicosta@fchampalimaud.org](mailto:ruicosta@fchampalimaud.org).

Running title: Aberrant striatal activity due to a *Foxp2* mutation

## Abstract

Mutations in the human *FOXP2* gene cause impaired speech development and linguistic deficits, which have been best characterized in a large pedigree called the KE family. The encoded protein is highly conserved in many vertebrates and is expressed in homologous brain regions required for sensorimotor integration and motor-skill learning, in particular cortico-striatal circuits. Independent studies in multiple species suggest that the striatum is a key site of FOXP2 action. Here we used *in vivo* recordings in awake behaving mice to investigate the effects of the KE-family mutation on the function of striatal circuits during motor-skill learning. We uncovered abnormally high ongoing striatal activity in mice carrying an identical mutation to that of the KE-family. Furthermore, there were dramatic alterations in striatal plasticity during the acquisition of a motor-skill, with most neurons in mutants showing negative modulation of firing rate, starkly contrasting with the predominantly positive modulation seen in control animals. We also observed striking changes in the temporal coordination of striatal firing during motor-skill learning in mutants. Our results indicate that FOXP2 is critical for the function of striatal circuits *in vivo*, which are important not only for speech but also for other striatal-dependent skills.

Keywords: speech and language, KE family, Foxp2, motor-skill learning, striatum, *in vivo* recording

## Introduction

Speech is one of the most complex and refined motor skills we perform. Developmental disorders of speech and language affect up to 7% of 5-to-7 year olds<sup>1</sup> and are known to be highly heritable<sup>2</sup>. However the inheritance patterns are usually complex, indicating that multiple genes are likely involved, a difficulty which has hindered investigations into the underlying molecular aetiology<sup>3</sup>. Identification of the genetic basis of a rare monogenic form of speech and language impairment has offered a unique opportunity to study these mechanisms<sup>4</sup>. In the large multigenerational KE family, around half the members carry a point mutation in one copy of the *FOXP2<sup>a</sup>* gene yielding an arginine-to-histidine substitution that disturbs the DNA-binding domain of the encoded transcription factor<sup>4, 5</sup>. These people have difficulty mastering the sequences of orofacial motor movements necessary for fluent speech<sup>6</sup>, a feature which is proposed to be a core deficit of the disorder<sup>7</sup>. The speech motor deficits are accompanied by other expressive and receptive problems in both oral and written language<sup>6</sup>. Additional families have been identified with speech and language problems caused by *FOXP2* mutations<sup>8</sup>, but the KE family remain the most well-studied example to date.

In the developing human brain, *FOXP2* is expressed in neuronal subpopulations of the cortex, basal ganglia, thalamus and cerebellum; regions which are known to be important for motor-related functions<sup>9</sup>. The striatum is the main recipient of cortical input to the basal ganglia, and neuroimaging studies of the KE family have implicated this structure in *FOXP2*-related speech and language deficits, uncovering changes in striatal grey-matter density<sup>10</sup>, as well as altered striatal activation during language based tasks<sup>11, 12</sup>. Importantly striatal lesions in humans, resulting from brain trauma or stroke, can result in verbal aphasia and amusia (a

---

<sup>a</sup> The standard nomenclature for FOX transcription factor proteins recommended by the winged helix/forkhead nomenclature committee (Kaestner KH, Knochel W, Martinez DE (2000) Unified nomenclature for the winged helix/forkhead transcription factors. *Genes Dev* 14: 142-146) is *FOXP2* for humans, *Foxp2* for mice and *FoxP2* for all other species, with genes and RNA displayed in italics. To maintain brevity we have used just *FoxP2* when referring to several species.

pitch processing deficit)<sup>13, 14</sup>. The FoxP2 protein sequence and neural expression pattern are highly conserved in a number of other vertebrates, including songbirds and rodents<sup>9, 15-18</sup>, suggesting some common ancestral functions. For basic studies investigating the role of FoxP2 in skill learning and striatal function, these species are more amenable to experimentation. We previously generated mutant mice (*Foxp2-R552H* line) which carry an identical mutation to that of the KE family<sup>19</sup>. Homozygous mutants show severe developmental delay and motor impairment, dying 3-4 weeks after birth. Heterozygous mutants have motor-skill deficits and slice recordings show impaired long-term depression (LTD) at glutamatergic inputs into the striatum<sup>19</sup>. Conversely, mice with a partially humanised version of Foxp2 have increased striatal LTD at cortico-striatal synapses<sup>20</sup>. Furthermore in the zebra finch, FoxP2 knockdown in Area X, a striatal nucleus essential for song development, impairs vocal learning<sup>21</sup>.

The evidence to date suggests that FoxP2 plays a role in striatal function with potential relevance for motor-skill learning. However, it is still unclear what impact the KE-family mutation has on striatal circuits during the learning of rapid motor sequences such as those required for speech. In this study we show the consequences of the KE-family mutation on the ongoing activity of striatal circuits *in vivo*, and on the *in-vivo* plasticity observed during skill learning, by performing multielectrode recordings in awake behaving *Foxp2-R552H/+* mice during training on the accelerating rotarod. This task engages the circuits required for the striatal-dependent learning of rapid motor sequences<sup>22</sup>, which are likely to be functioning abnormally in this mouse line<sup>19</sup>, and which have been proposed to be dysfunctional in affected members of the KE family<sup>7</sup>. We found that *Foxp2-R552H/+* mice displayed abnormally high ongoing striatal activity. Moreover during acquisition of a motor skill, most striatal neurons in *Foxp2-R552H/+* mice showed negative modulation of firing rate unlike the positive modulation observed in controls. Changes in the temporal coordination of striatal

firing were also evident during motor-skill learning in *Foxp2-R552H/+* mice. These results show that Foxp2 is critical for the normal functioning of striatal circuits during the acquisition of a motor-skills.

## **Materials and methods**

### *Animals*

The *Foxp2-R552H* and *Foxp2-S321X* lines were maintained on a C57BL/6J background and heterozygotes and wild type littermates, of between 2 and 6 months of age, were used for behavioral and recording experiments. The generation, marker-assisted backcrossing and genotyping of this strain are fully described in references<sup>19, 23, 24</sup>. All procedures were approved by the National Institute on Alcohol Abuse and Alcoholism Animal Care and Use Committee.

### *Accelerating rotarod*

A computer interfaced rotarod (ENV-575M, Med-Associates, St. Albans, VT) was set to accelerate from 6-60 rpm over a 300 s time period. Mice were trained for five consecutive days, with one daily session consisting of 10 trials separated by 300 s resting periods. Mice were placed on the rotarod and trials were deemed to have started when the rod began to turn. Trials ended when mice fell from the rod or after 300 s elapsed. Learning rate was calculated as follows (mean latency to fall trials 9 and 10) – (mean latency to fall trials 1 and 2)/ 9 (number of intertrial intervals)

### *In vivo extracellular recording*

Implantation of multielectrode arrays and *in vivo* recording of neural activity in mice training on the accelerating rotarod were performed essentially as described in reference<sup>22</sup>. Arrays had

2 rows of 8 polyamide-coated tungsten wires, which were 35  $\mu\text{m}$  in diameter with 90° blunt cut tips. Wires were separated by a 250  $\mu\text{m}$  gap, both within a row and across rows. The centre of each array was surgically positioned 2.2–2.4 mm below the surface of the brain, 0.5 mm anterior to and 1.7 mm lateral to Bregma. Final electrode positioning in the dorsomedial striatum (DMS) was determined by stereotaxic co-ordinates and by neural activity, which was monitored as the electrodes were lowered into the brain. Mice were allowed to recover for at least 10 days post surgery after which single-unit and multi-unit activity was recorded using the Plexon data acquisition system (Plexon, TX). The start and finish of each trial were signalled to the MAP recording system (Plexon, TX) as events. The recording cables were held by a custom-made pulley system allowing mice to run and fall relatively freely. Mice remained at the bottom of the apparatus during intertrial periods, usually in a fairly immobile state. Neural activity was initially sorted using an online sorting algorithm and then resorted, post-recording, using an offline sorting algorithm based on waveform, amplitude and interspike interval histogram (Plexon, TX). The total number of units recorded on days 1 to 5 was 56, 78, 80, 72 and 70 for wild-type mice and 55, 54, 64, 58 and 48 for *Foxp2-R552H/+* mice.

#### *Neural data analyses*

Data analyses were carried out using Neuroexplorer (Nex, TX) and custom written programs in Matlab (Mathworks, MA).

LFP power spectrum: The LFP power spectrum was estimated for 2-s sliding windows with 1-s steps via Welch's method; the parameters chosen allowed for a frequency resolution of 0.5 Hz. The percentage power for the following frequency bands was calculated; delta 1.5 - 4 Hz, theta 4.5 - 9 Hz and gamma 30 - 55 Hz.

Spike triggered average: STAs of the LFP were calculated for -3 s to 3 s intervals. An STA was considered significant when 20 or more 1 ms bins within the interval -0.1 s to 0.1 s lay outside the maximum or minimum values of the intervals -3 to -1 s and 1 s to 3 s.

### *Statistics*

Statistical analyses were carried out using SPSS (SPSS) and Neuroexplorer. Task related neurons were identified as having significant positive or negative modulation of firing rate during running. This was determined using a 99% confidence interval which was calculated from the firing rate during the intertrial intervals. All results were averaged per subject and subsequent analyses were performed on each subject's mean value. Data were analysed using repeated measures ANOVA, and *post hoc* tests (Fisher's LSD) were applied when appropriate. Planned comparisons were used to analyse the LFP data.

### *Histology*

Mice were perfused with 4% paraformaldehyde (PFA) and brains were removed and post-fixed overnight in 4% PFA. Electrode placement was verified by Nissl staining of 50  $\mu$ m vibrotome sections. For Foxp2 and interneuron immunohistochemistry (IHC), post-fixed brains were transferred to 30% sucrose and 40  $\mu$ m coronal cryosections were cut on a sliding microtome. IHC was carried out using the Vectastain ABC or Vector MOM kit (both Vector Laboratories). Free-floating sections were incubated with Foxp2 (1:2000, goat N16, Santa Cruz), ChAT (1:100, goat, Chemicon) or PV (1:5000, mouse, Swant) antibodies. Proteins were visualised using the Vector VIP substrate purple kit (Vector Laboratories) or the DAB substrate kit (Pierce). When looking at Foxp2 co-localisation with PV or ChAT, nuclei were marked using methyl green counterstain (Vector Laboratories). To ascertain interneuron numbers, serial sections were taken throughout the striatum and stained alternatively with PV

and ChAT antibodies. The top half of the striatum was considered to be dorsal, and this was subdivided into DM, DI and DL thirds. The area of each region of interest and the number of positive cells within it were determined using ImageJ. Cell density (cells/ mm<sup>2</sup>) was calculated by dividing the cell count by the area.

## Results

We first confirmed that mice carrying an aetiological *Foxp2* mutation display impaired motor skill learning<sup>19</sup>. Mice were trained for five consecutive days on an accelerating rotarod, with one daily session consisting of 10 trials (running periods), separated by 300 s intervals. This intensive schedule uses a higher acceleration rate (6-60 rpm) than previous experiments<sup>19, 22, 25</sup>, and robustly shows the motor-skill learning deficits of *Foxp2-R552H/+* animals. In line with previous findings<sup>19</sup>, latency to fall was reduced in *Foxp2-R552H/+* mice ( $F_{1,19} = 9.87$ ,  $P < 0.05$ ) (Figure 1a). A significant difference in performance was evident early in training (*post hoc*  $P < 0.05$ , comparison of third and fourth trials of session 1), and although *Foxp2-R552H/+* mice improved, they never reached the fall times of wild-type animals (*post hoc*  $P < 0.05$ , comparison of final two trials of session 5). The early deficits observed in *Foxp2-R552H/+* mice do not seem to stem from alterations in basal motor coordination, as performance during the first two trials of session 1 was not impaired (*post hoc*  $P > 0.05$ ). Latency to fall from a fixed speed rat rotarod, a task which addresses motor coordination, was also not significantly different between *Foxp2-R552H/+* and wild-type animals (Supplementary Figure 1a). Rather, the deficits appear to be due to a slower rate of learning during the first session (*post hoc*  $P < 0.05$ ) (Figure 1b). Notably, *Foxp2-R552H/+* mice do not have generalised striatal-dependent deficits in learning or motivation, because they learn to press a lever to earn a food reward in the same way as wild-type animals in an operant task which involves the striatum<sup>26</sup> (Supplementary Figure 2). Furthermore, the severity of the

deficits in rotarod skill learning appear to be related to the KE-family mutation, since the deficits observed in *Foxp2-S321X/+* mice, which carry a premature stop codon and have a half dosage of Foxp2 protein, were less severe than those observed in *Foxp2-R552H/+* mice (Supplementary Figure 3). *Foxp2-S321X/+* animals also have no impairment in basal motor coordination (Supplementary Figure 1b).

We therefore examined the impact of the KE-family mutation on striatal activity *in vivo* by performing multielectrode recordings in awake behaving *Foxp2-R552H/+* mice (n=5) and control wild-type littermates (n=6). Since skill learning deficits in *Foxp2-R552H/+* mice are apparent early in training, we targeted electrode arrays bilaterally to the dorsomedial striatum (DMS) (Supplementary Figure 4), which is preferentially engaged during this initial phase of learning<sup>25</sup>. After recovery from electrode implantation surgery, neuronal activity (Figure 2a) and local field potential oscillations (LFP) were measured during training on the accelerating rotarod.

We first investigated changes in ongoing striatal activity in *Foxp2-R552H/+* mice by measuring the firing rate of striatal neurons during a 5 minute period before the initial trial of each session, while mice were positioned at the bottom of the apparatus in a quiet but awake state (periods of no locomotion but occasional head movements). Intriguingly, we found that striatal neurons in *Foxp2-R552H/+* mice showed higher ongoing firing rates than those in wild-type controls across the five days of recording ( $F_{1,9} = 5.54, P < 0.05$ ) (Figure 2b). This is interesting because striatal medium spiny neurons (MSNs) usually display low *in vivo* firing rates<sup>27</sup>, a condition which is thought to be critical for normal action selection and movement<sup>28</sup>.

We next investigated the effects of the *Foxp2* mutation on the dynamic changes in striatal plasticity which are observed during motor-skill learning<sup>25</sup>. We determined whether the firing rate of striatal neurons increased or decreased during running compared to the intertrial intervals by calculating the modulation in firing rate  $[(\text{run-interval})/(\text{run+interval}) \times$

100], at the beginning and end of each session (first two trials of session – "first"; last two trials of session – "last")<sup>22</sup>. Consistent with previous experiments<sup>25</sup>, the average firing rate in wild-type DMS neurons increased during running compared to intertrial intervals in early training sessions, as revealed by positive modulation of firing rate (Figure 2c). However, striatal neurons in *Foxp2-R552H/+* mice were substantially different ( $F_{1,9} = 7.68$ ,  $P < 0.05$ ), showing predominantly negative modulation during running. This negative modulation was most evident at early stages and progressively diminished, with the last trials of day 5 being significantly different from the first trials of day 1 (*post hoc*  $P < 0.05$ ), which is consistent with the DMS becoming less engaged with training<sup>25</sup>. Furthermore, in wild-type mice the firing rate was significantly more modulated during the last trials of a session than the first ( $F_{1,5} = 27.88$ ,  $P < 0.05$ ), revealing within-session plasticity. This plasticity was not observed in *Foxp2-R552H/+* animals where no significant differences between the first and last trials of a session were found ( $F_{1,4} = 3.96$ ,  $P > 0.05$ ).

To examine how these differences in firing rate modulation emerge during skill learning, we determined the proportion of neurons whose firing rate changed significantly during running compared to the intertrial intervals (classified as task-related, see Materials and methods and Figure 3a). The percentage of task-related neurons recorded was similar in wild-type and *Foxp2-R552H/+* animals and remained relatively constant throughout training (Supplementary Figure 5). However, when task-related neurons were divided into those that increased or decreased firing rate during running, more neurons were found to increase firing rate in wild-type mice ( $F_{1,10} = 22.25$ ,  $P < 0.05$ ) and more to decrease firing rate in *Foxp2-R552H/+* mice ( $F_{1,8} = 7.12$ ,  $P < 0.05$ ) (Figure 3b). These results are consistent with the differences in firing rate modulation observed between *Foxp2-R552H/+* mutants and controls (Figure 2c).

We next investigated whether the negative modulation of firing rate observed in *Foxp2-R552H/+* during running on the rotarod was related to, or independent from, the higher ongoing activity of striatal neurons in these animals. It is plausible that neurons with high ongoing firing rates could be unable to increase firing further during running, and could even compensate by decreasing firing, hence resulting in the observed negative modulation of firing rate. However, we found that *Foxp2-R552H/+* neurons with high ongoing firing rates were not modulated differently during running compared to those with lower firing rates (Supplementary Figure 6). Furthermore, one would expect that if a significant proportion of *Foxp2-R552H/+* neurons were unable to modulate firing rate during running then there would be a corresponding reduction in the number of task-related neurons, which is not observed (Supplementary Figure 5). Finally, if *Foxp2-R552H/+* neurons with high baseline firing rates were not able to positively modulate firing during running, then the most prominent differences in rate modulation between genotypes would occur when the greatest levels of positive modulation are observed in wild-type mice, and vice versa. However, as previously discussed, firing rate is significantly more modulated in the last trials of a session compared to the first in wild-type mice but this within-session plasticity is clearly not evident in *Foxp2-R552H/+* mice (Figure 2c). Taken together these data indicate that the changes in striatal activity observed during skill learning in *Foxp2-R552H/+* mice are not simply a direct consequence of the increased ongoing firing.

In addition to changes in firing frequency, temporal coordination of striatal activity has also been shown to be important for striatal functions such as action selection and movement<sup>28, 29</sup>. We therefore measured LFP oscillations, which mostly reflect the coordinated input into an area, in *Foxp2-R552H/+* and control mice. We observed no significant differences in power at any of the frequencies examined, despite a tendency for higher power gamma oscillations in *Foxp2-R552H/+* mice on day 1 (Supplementary Figure 7). We then

investigated the temporal relationship between the oscillatory phase of the LFP (reflecting mostly input) and the firing of individual striatal neurons (reflecting output). We determined if a neuron fired preferentially during a particular phase of the LFP (was “entrained” to the LFP) during running and intertrial intervals (Supplementary Figure 8), using spike-triggered averages (STAs) of the LFP oscillations (see Materials and methods and References<sup>28, 29</sup>) (Figure 4a). We calculated how many neurons became entrained or lost entrainment to the LFP oscillations during running compared to the intertrial intervals. In wild-type mice more neurons lost entrainment to the LFP oscillations during running in the first trials of a session compared to the last ( $F_{1,5} = 9.00$ ,  $P < 0.05$ ), but in *Foxp2-R552H/+* animals this within-session plasticity was not evident ( $F_{1,4} = 0.80$ ,  $P > 0.05$ ) (Figure 4b). In addition, more neurons became entrained to the LFP during running in *Foxp2-R552H/+* than in wild-type mice ( $F_{1,8} = 6.21$ ,  $P < 0.05$ ), an effect that was particularly apparent during the late phase of training. These results show that the temporal coordination of striatal input and output differs between *Foxp2-R552H/+* and wild-type mice.

## Discussion

Previous neuroimaging studies of the KE family as well as experiments in mice and songbirds have highlighted possible roles for FoxP2 in the striatum, with potential relevance for motor-skill learning. However, the impact of the KE-family mutation on striatal function during the learning of rapid motor sequences was unclear. In this study we demonstrate *in vivo* that this mutation dramatically alters the functioning of striatal circuits during motor-skill learning. We found that in contrast to control animals, which showed positive modulation of firing rate during skill acquisition, *Foxp2-R552H/+* mice exhibited negative modulation of firing rate. We also found alterations in the temporal coordination of striatal activity in *Foxp2-R552H/+* mice compared to controls.

We found *Foxp2-R552H/+* mice to have abnormally high ongoing striatal activity. Striatal MSNs, which represent the vast majority of striatal neurons, usually display low *in vivo* firing rates<sup>27</sup>. This property, together with appropriate levels of concurrent activity between MSNs, is thought to be critical for normal action selection<sup>28</sup>. The high ongoing striatal firing rate in *Foxp2-R552H/+* mice could therefore result in increased collateral inhibition which may interfere with action selection and plasticity processes in this brain region. Our previous *ex vivo* data, which shows strong impairment of LTD at glutamatergic synapses onto MSNs in *Foxp2-R552H/+* mice<sup>19</sup>, is consistent with increased ongoing striatal activity. Nevertheless, it is important to note that besides glutamatergic input, MSNs receive dopaminergic input from the midbrain, and GABAergic input from interneurons and from other MSNs<sup>27</sup>. Hence elevated MSN activity could also result from changes in the strength of any of these other inputs (eg. increased dopaminergic input or decreased GABAergic input), as well as increased neuronal excitability. However, we found only modest co-localisation of Foxp2 with parvalbumin (PV) or cholineacetyltransferase (ChAT), markers for the main subtypes of striatal interneurons, and there were equivalent numbers of PV- or ChAT-positive cells in the striata of wild-type and *Foxp2-R552H/+* mice (Supplementary Figure 9), suggesting that striatal interneurons are unlikely to have a direct influence. We also showed that the increased ongoing striatal activity and the negative firing rate modulation during skill acquisition observed in *Foxp2-R552H/+* mice are not correlated. This suggests that the underlying mechanisms of the two phenomena are dissociable and raises the possibility that they are mechanistically distinct. Although any of the mechanisms outlined above could contribute to the firing rate modulation differences we observed, it seems unlikely that impaired LTD could directly give rise to an increase in negative modulation of firing rate. Selective deletion of Foxp2 will be a useful tool to investigate how specific cells types and brain regions contribute to changes in striatal activity as well as other phenotypes<sup>30</sup>.

*Foxp2-S321X/+* and *Foxp2-R552H/+* mice express a half dosage of normal wild-type Foxp2 protein, and so both could potentially display haploinsufficiency. However, *Foxp2-R552H/+* mice also simultaneously express a full-length mutant product with abnormal functional properties. *In vitro* studies indicate that the R552H mutation disrupts the DNA-binding and transactivation functions of Foxp2<sup>5</sup>, suggesting that the mutant protein could act in a dominant negative manner. Foxp2 has been shown to function as a homodimer, and also as a heterodimer with Foxp1 or Foxp4<sup>31</sup>, hence dominant negative effects could be exerted through protein-protein interactions or direct disturbance of transcription. In support of this proposal, we observed a less severe motor-skill learning phenotype in *Foxp2-S321X/+* mice than in *Foxp2-R552H/+* mice (Figure 1 and Supplementary Figure 3); further work will be necessary to determine if this is indeed the case. Studies using chromatin immunoprecipitation followed by microarray analysis have been used to identify direct neural targets of FOXP2 and have indicated roles in neurodevelopment, neurite outgrowth, axon guidance, neurotransmission and synaptic plasticity<sup>32, 33</sup>. The *Foxp2-R552H* mutation is likely to cause inappropriate or inefficient regulation of downstream targets mediating these processes<sup>34</sup>, resulting in the dramatic changes in striatal neural activity we observe.

In conclusion, the use of multielectrode recordings in awake behaving mice with an aetiological Foxp2 mutation has allowed for a detailed assessment of its impact on the neuronal dynamics of striatal circuits during the learning of rapid motor sequences, beyond the resolution of human neuroimaging. The resulting data gives novel insight into the mechanistic consequences of this mutation, which is important not only for speech but also for other striatal-dependent skills. It will be interesting to see whether the increased ongoing striatal activity and generalised motor-skill learning deficits observed in mice extend to people with disorders attributed to FOXP2 dysfunction. Although it is difficult to perform invasive electrophysiology studies on such people, our data suggest that resting state

functional imaging and relatively demanding tests of motor-skill learning ability could be highly informative.

### **Acknowledgments**

We would like to thank Sonali Darbar, Rathi Puliyadi, Terrell D. Holloway and Amber Luo for excellent technical assistance, John Burkhardt and Shih-Chieh Lin for help with Matlab, Emma Doran for running the operant learning task, and the staffs of the British Heart Foundation Functional Genetics Facility and the NIAAA vivarium for mouse husbandry. This work was supported by a Human Frontier Science Program short term fellowship to CAF; a Royal Society Research Fellowship to SEF; the Wellcome Trust [075491/Z/04], [080971]; the Division of Intramural Clinical and Basic Research of the National Institute on Alcohol Abuse and Alcoholism, US National Institutes of Health; the Champalimaud Neuroscience Programme; the European Research Council [STG 243393] and the Max Planck Society.

### **Conflict of interest**

The authors declare no conflict of interest.

### **References**

1. Tomblin JB, Records NL, Buckwalter P, Zhang X, Smith E, O'Brien M. Prevalence of specific language impairment in kindergarten children. *J Speech Lang Hear Res* 1997 Dec; **40**(6): 1245-1260.
2. Bishop DV. Genetic and environmental risks for specific language impairment in children. *Philos Trans R Soc Lond B Biol Sci* 2001 Mar 29; **356**(1407): 369-380.

3. Fisher SE, Lai CS, Monaco AP. Deciphering the genetic basis of speech and language disorders. *Annu Rev Neurosci* 2003; **26**: 57-80.
4. Lai CS, Fisher SE, Hurst JA, Vargha-Khadem F, Monaco AP. A forkhead-domain gene is mutated in a severe speech and language disorder. *Nature* 2001 Oct 4; **413**(6855): 519-523.
5. Vernes SC, Nicod J, Elahi FM, Coventry JA, Kenny N, Coupe AM *et al.* Functional genetic analysis of mutations implicated in a human speech and language disorder. *Hum Mol Genet* 2006 Nov 1; **15**(21): 3154-3167.
6. Watkins KE, Dronkers NF, Vargha-Khadem F. Behavioural analysis of an inherited speech and language disorder: comparison with acquired aphasia. *Brain* 2002 Mar; **125**(Pt 3): 452-464.
7. Vargha-Khadem F, Gadian DG, Copp A, Mishkin M. FOXP2 and the neuroanatomy of speech and language. *Nat Rev Neurosci* 2005 Feb; **6**(2): 131-138.
8. MacDermot KD, Bonora E, Sykes N, Coupe AM, Lai CS, Vernes SC *et al.* Identification of FOXP2 truncation as a novel cause of developmental speech and language deficits. *Am J Hum Genet* 2005 Jun; **76**(6): 1074-1080.
9. Lai CS, Gerrelli D, Monaco AP, Fisher SE, Copp AJ. FOXP2 expression during brain development coincides with adult sites of pathology in a severe speech and language disorder. *Brain* 2003 Nov; **126**(Pt 11): 2455-2462.

10. Watkins KE, Vargha-Khadem F, Ashburner J, Passingham RE, Connelly A, Friston KJ *et al.* MRI analysis of an inherited speech and language disorder: structural brain abnormalities. *Brain* 2002 Mar; **125**(Pt 3): 465-478.
11. Vargha-Khadem F, Watkins KE, Price CJ, Ashburner J, Alcock KJ, Connelly A *et al.* Neural basis of an inherited speech and language disorder. *Proc Natl Acad Sci U S A* 1998 Oct 13; **95**(21): 12695-12700.
12. Liegeois F, Baldeweg T, Connelly A, Gadian DG, Mishkin M, Vargha-Khadem F. Language fMRI abnormalities associated with FOXP2 gene mutation. *Nat Neurosci* 2003 Nov; **6**(11): 1230-1237.
13. Jarvis ED. Learned birdsong and the neurobiology of human language. *Ann N Y Acad Sci* 2004 Jun; **1016**: 749-777.
14. Lieberman P. On the nature and evolution of the neural bases of human language. *Am J Phys Anthropol* 2002; **Suppl 35**: 36-62.
15. Ferland RJ, Cherry TJ, Preware PO, Morrissey EE, Walsh CA. Characterization of Foxp2 and Foxp1 mRNA and protein in the developing and mature brain. *J Comp Neurol* 2003 May 26; **460**(2): 266-279.

16. Takahashi K, Liu FC, Hirokawa K, Takahashi H. Expression of Foxp2, a gene involved in speech and language, in the developing and adult striatum. *J Neurosci Res* 2003 Jul 1; **73**(1): 61-72.
17. Teramitsu I, Kudo LC, London SE, Geschwind DH, White SA. Parallel FoxP1 and FoxP2 expression in songbird and human brain predicts functional interaction. *J Neurosci* 2004 Mar 31; **24**(13): 3152-3163.
18. Haesler S, Wada K, Nshdejan A, Morrissey EE, Lints T, Jarvis ED *et al.* FoxP2 expression in avian vocal learners and non-learners. *J Neurosci* 2004 Mar 31; **24**(13): 3164-3175.
19. Groszer M, Keays DA, Deacon RM, de Bono JP, Prasad-Mulcare S, Gaub S *et al.* Impaired synaptic plasticity and motor learning in mice with a point mutation implicated in human speech deficits. *Curr Biol* 2008 Mar 11; **18**(5): 354-362.
20. Enard W, Gehre S, Hammerschmidt K, Holter SM, Blass T, Somel M *et al.* A humanized version of Foxp2 affects cortico-basal ganglia circuits in mice. *Cell* 2009 May 29; **137**(5): 961-971.
21. Haesler S, Rochefort C, Georgi B, Licznarski P, Osten P, Scharff C. Incomplete and inaccurate vocal imitation after knockdown of FoxP2 in songbird basal ganglia nucleus Area X. *PLoS Biol* 2007 Dec; **5**(12): e321.

22. Costa RM, Cohen D, Nicoletis MA. Differential corticostriatal plasticity during fast and slow motor skill learning in mice. *Curr Biol* 2004 Jul 13; **14**(13): 1124-1134.
23. Keays DA, Clark TG, Campbell TG, Broxholme J, Valdar W. Estimating the number of coding mutations in genotypic and phenotypic driven N-ethyl-N-nitrosourea (ENU) screens: revisited. *Mamm Genome* 2007 Feb; **18**(2): 123-124.
24. Coghill EL, Hugill A, Parkinson N, Davison C, Glenister P, Clements S *et al.* A gene-driven approach to the identification of ENU mutants in the mouse. *Nat Genet* 2002 Mar; **30**(3): 255-256.
25. Yin HH, Mulcare SP, Hilario MR, Clouse E, Holloway T, Davis MI *et al.* Dynamic reorganization of striatal circuits during the acquisition and consolidation of a skill. *Nat Neurosci* 2009 Mar; **12**(3): 333-341.
26. Yin HH, Knowlton BJ. The role of the basal ganglia in habit formation. *Nat Rev Neurosci* 2006 Jun; **7**(6): 464-476.
27. Kreitzer AC, Malenka RC. Striatal plasticity and basal ganglia circuit function. *Neuron* 2008 Nov 26; **60**(4): 543-554.
28. Costa RM, Lin SC, Sotnikova TD, Cyr M, Gainetdinov RR, Caron MG *et al.* Rapid alterations in corticostriatal ensemble coordination during acute dopamine-dependent motor dysfunction. *Neuron* 2006 Oct 19; **52**(2): 359-369.


29. Burkhardt JM, Jin X, Costa RM. Dissociable effects of dopamine on neuronal firing rate and synchrony in the dorsal striatum. *Front Integr Neurosci* 2009; **3**: 28.
30. French CA, Groszer M, Preece C, Coupe AM, Rajewsky K, Fisher SE. Generation of mice with a conditional Foxp2 null allele. *Genesis* 2007 Jul; **45**(7): 440-446.
31. Li S, Weidenfeld J, Morrisey EE. Transcriptional and DNA binding activity of the Foxp1/2/4 family is modulated by heterotypic and homotypic protein interactions. *Mol Cell Biol* 2004 Jan; **24**(2): 809-822.
32. Spiteri E, Konopka G, Coppola G, Bomar J, Oldham M, Ou J *et al.* Identification of the transcriptional targets of FOXP2, a gene linked to speech and language, in developing human brain. *Am J Hum Genet* 2007 Dec; **81**(6): 1144-1157.
33. Vernes SC, Spiteri E, Nicod J, Groszer M, Taylor JM, Davies KE *et al.* High-throughput analysis of promoter occupancy reveals direct neural targets of FOXP2, a gene mutated in speech and language disorders. *Am J Hum Genet* 2007 Dec; **81**(6): 1232-1250.
34. Fisher SE, Scharff C. FOXP2 as a molecular window into speech and language. *Trends Genet* 2009 Apr; **25**(4): 166-177.

**Figure 1** *Foxp2-R552H/+* mice show impaired motor-skill learning. **(a)** Latency to fall of wild-type and *Foxp2-R552H/+* mice during training on the accelerating rotarod. Mice received 10 trials a day for 5 consecutive days. **(b)** Rate of learning in wild-type and *Foxp2-R552H/+* mice during training on the accelerating rotarod. Error bars represent s.e.m.

**Figure 2** *Foxp2-R552H/+* mice have abnormally high ongoing striatal activity and show negative modulation of firing rate during skill learning. **(a)** Example of a single-unit waveform (top left). Units were distinguished from each other and from noise using interspike interval histograms (top right) and 2D cluster separation based on principal component analysis, where the circled yellow cluster represents the unit and the grey cluster noise (bottom right). The stability of a unit recording over the course of a session was assessed using 3D cluster displays with a time axis (bottom left). **(b)** Average ongoing firing rate in wild-type and *Foxp2-R552H/+* mice. **(c)** Average firing rate modulation during running compared with intertrial intervals, during the first and last trials of a session, in wild-type and *Foxp2-R552H/+* mice. Error bars represent s.e.m.

**Figure 3** A higher proportion of task-related neurons decrease firing rate during skill learning in *Foxp2-R552H/+* mice. **(a)** Examples of neurons from wild-type (top) and *Foxp2-R552H/+* (bottom) mice that increased (left) or decreased (right) firing rate (impulses per second) during running compared with intertrial intervals. Green triangles represent the start of a trial and red circles the end. **(b)** Percentage of task-related neurons that increased or decreased firing rate during running compared with intertrial intervals, during the first and last trials of a session, in wild-type (top panel) and *Foxp2-R552H/+* (bottom panel) mice. Error bars represent s.e.m.

**Figure 4** The temporal coordination of striatal activity is altered during skill learning in *Foxp2-R552H/+* mice. **(a)** STA histograms showing a neuron which is entrained to the LFP during intertrial intervals (left) but which loses entrainment during running periods (right). **(b)** Percentage of neurons that lost entrainment from (top panel) or gained entrainment to (bottom panel) the LFP between intertrial interval and running during the first and last trials of a session. Error bars represent s.e.m.

**a****b**