HAL open science

An elementary proof of Catalan-Mihailescu theorem

Jamel Ghannouchi

To cite this version:

Jamel Ghannouchi. An elementary proof of Catalan-Mihailescu theorem. 2011. hal-00677731v35

HAL Id: hal-00677731

https://hal.science/hal-00677731v35

Preprint submitted on 25 Oct 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An elementary proof of Catalan-Mihailescu theorem and generalization to Fermat-Catalan conjecture

Jamel Ghanouchi

jamel.ghanouchi@live.com

Abstract

(MSC=11D04) We begin with an equation, for example : $Y^p = X^q \pm Z^c$ and solve it.

(Keywords : Diophantine equations, Fermat-Catalan equation; Approach)

Resolution of Catalan equation

Let Catalan equation :

$$Y^p = X^q + 1$$

We have

$$X^{q-3}Y^2 - Y^{p-2}X^3 = A$$

And

$$Y^{p-2}Y^2 - X^{q-3}X^3 = Y^p - X^q = 1$$

If $A = 0$ then $X^{q-6} = Y^{p-4}$ leads, as $GCD(X, Y) = 1$, to $p = 4$ and $q = 6$. This case has been studied by Lebesgue in the XIX century, it has no solution. Thus $A \neq 0$.
And if $A = \pm 1$ then it means that both
$X^{q-4}Y^2 = \pm\frac{1}{X} + X^2Y^{p-2}$ and
$Y^{p-3}X^3 = \mp\frac{1}{X} + X^{q-3}Y$ are rationals

it means, as it is impossible, that $q = 3$ and $p = 2$.
We have

$$\frac{X^{q-3}}{A}Y^2 - \frac{Y^{p-2}}{A}X^3 = 1 = Y^{p-2}Y^2 - X^{q-3}X^3$$

And we have simultaneously

$$(Y^{p-2} - \frac{X^{q-3}}{A})Y^2 = (X^{q-3} - \frac{Y^{p-2}}{A})X^3$$

Or

$$(AY^{p-2} - X^{q-3})Y^2 = (AX^{q-3} - Y^{p-2})X^3$$

And

$$(Y^2 + \frac{X^3}{A})Y^{p-2} = (X^3 + \frac{Y^2}{A})X^{q-3}$$

Or

$$(AY^2 + X^3)Y^{p-2} = (AX^3 + Y^2)X^{q-3}$$

But X and Y are coprimes. It means that we have four cases with u and v integers

$$Y^2 = u(AX^{q-3} - Y^{p-2}); \quad X^3 = u(-X^{q-3} + AY^{p-2})$$
$$Y^{p-2} = v(AX^3 + Y^2); \quad X^{q-3} = v(X^3 + AY^2)$$

Or

$$uY^2 = AX^{q-3} - Y^{p-2}; \quad uX^3 = -X^{q-3} + AY^{p-2}$$
$$vY^{p-2} = AX^3 + Y^2; \quad vX^{q-3} = X^3 + AY^2$$

Or

$$Y^2 = u(AX^{q-3} - Y^{p-2}); \quad X^3 = u(-X^{q-3} + AY^{p-2})$$
$$vY^{p-2} = AX^3 + Y^2; \quad vX^{q-3} = X^3 + AY^2$$

Or

$$uY^2 = AX^{q-3} - Y^{p-2}; \quad uX^3 = -X^{q-3} + AY^{p-2}$$

$$Y^{p-2} = v(AX^3 + Y^2); \quad X^{q-3} = v(X^3 + AY^2)$$

First case

$$Y^p = uv(AX^q - Y^p + A(Y^2X^{q-3} - Y^{p-2}X^3))$$
$$= uv(A^2X^q - Y^p + A(A)) = uv(A^2X^q + A^2 - Y^p) = uv(A^2Y^p - Y^p)$$

Thus

$$uv = \frac{1}{A^2 - 1}$$

As uv is integer, it means that it is impossible thus $u = 0$ and $A^2 = 1$ or $A = \pm -1$ it means that $q = 3$ and $p = 2$.

Second case

$$uv\frac{Y^p}{A^2} = X^q - \frac{Y^p}{A^2} + \frac{Y^2X^{q-3} - Y^{p-2}X^3}{A}$$
$$= X^q - \frac{Y^p}{A^2} + 1 = X^q + 1 - \frac{Y^p}{A^2} = (\frac{A^2 - 1}{A^2})Y^p$$

Thus

$$uv = A^2 - 1$$

And

$$uv(Y^2X^{q-3} - X^3Y^{p-2}) = uvA = u(X^{2q-6} - Y^{2p-4})A = v(-X^6 + Y^4)A$$

Thus

$$u = -X^6 + Y^4; \quad v = X^{2q-6} - Y^{2p-4}$$
$$uv = A^2 - 1 = (-X^6 + Y^4)(X^{2q-6} - Y^{2p-4})$$

But

$$Y^p = X^q + 1 \Rightarrow Y^{p-2}X^3 = X^q\frac{X^3}{Y^2} + \frac{X^3}{Y^2}$$

And

$$Y^2X^{q-3} = X^q\frac{X^{q-3}}{Y^{p-2}} + \frac{X^{q-3}}{Y^{p-2}}$$

We deduce

$$A = Y^2X^{q-3} - X^3Y^{p-2} = (X^q + 1)(\frac{X^{q-3}}{Y^{p-2}} - \frac{X^3}{Y^2})$$

But

$$Y^p > X^q \Rightarrow A < 0 \Rightarrow \frac{X^{q-3}}{Y^{p-2}} < \frac{X^3}{Y^2}$$

And if

$$X^3 < Y^2 \Rightarrow X^{q-3} < Y^{p-2} \Rightarrow A^2 - 1 = (-X^6 + Y^4)(X^{2q-6} - Y^{2p-4} < 0$$

But

$$0 > Y^2(X^{q-3} - Y^{p-2}) = Y^2X^{q-3} - Y^p = Y^2X^{q-3} - X^q - 1 = X^{q-3}(Y^2 - X^3) - 1 > -1$$

And it is possible if $q - 3 = p - 2 = 0$.

And if

$$X^{q-3} > Y^{p-2} \Rightarrow X^3 > Y^2 \Rightarrow A^2 - 1 = (-X^6 + Y^4)(X^{2q-6} - Y^{2p-4} < 0$$

But

$$0 < Y^2(X^{q-3} - Y^{p-2}) = Y^2X^{q-3} - Y^p = Y^2X^{q-3} - X^q - 1 = X^{q-3}(Y^2 - X^3) - 1 < 0$$

And it is possible if $q - 3 = p - 2 = 0$

$$X^{q-3} > Y^{p-2} \Rightarrow X^{q-3}Y^2 > Y^p = X^q + 1 \Rightarrow Y^2 > X^3 + X^{3-q}$$

And

$$Y^2 - X^3 > X^{3-q} > 0$$

And $p-2=q-3=0$
Third case :
We have here

$$Y^2=u(AX^{q-3}-Y^{p-2});\quad X^3=u(-X^{q-3}+AY^{p-2})$$
$$vY^{p-2}=AX^3+Y^2;\quad vX^{q-3}=X^3+AY^2$$

And

$$vY^p=u(A^2X^q-Y^p+A^2)=u(A^2-1)Y^p$$
$$v=u(A^2-1)$$
$$v(Y^2X^{q-3}-X^3Y^{p-2})=vA=uvA(X^{2q-6}-Y^{2p-4})=A(-X^6+Y^4)$$

Thus

$$1=u(X^{2q-6}-Y^{2p-4})$$

With u and A^2-1 integers, it means $A^2=1$! Fourth case :

$$u\frac{Y^2}{A}=X^{q-3}-\frac{Y^{p-2}}{A};\quad u\frac{X^3}{A}=-\frac{X^{q-3}}{A}+Y^{p-2}$$
$$\frac{Y^{p-2}}{A}=v(X^3+\frac{Y^2}{A});\quad \frac{X^{q-3}}{A}=v(\frac{X^3}{A}+Y^2)$$

We have here

$$uY^2=AX^{q-3}-Y^{p-2};\quad uX^3-AY^{p-2}=-X^{q-3}$$

And

$$Y^{p-2}=AX^{q-3}-uY^2=(Y^2X^{q-3}-X^3Y^{p-2})X^{q-3}-uY^2$$

Hence

$$u\frac{Y^p}{A^2}=v(X^q-\frac{Y^p}{A^2}+1)=v(1-\frac{1}{A^2})Y^p$$

Thus

$$u=v(A^2-1)$$
$$u(Y^2X^{q-3}-X^3Y^{p-2})=uA=A(X^{2q-6}-Y^{2p-4})=uv(-X^6+Y^4)A$$

Thus

$$1=v(-X^6+Y^4)$$

With v and A^2-1 integers, it means $A^2-1=0$!
The only solution, in all cases, in $p=2$ and $q=3$.
And $Y^2=X^3+1$ whose solution is $(X,Y)=(2,\pm 3)$.

Resolution of Fermat equation

Let Fermat equation :

$$Y^n=X^n+Z^n$$

We have

$$X^{n-2}Y^2-Y^{n-2}X^2=A$$

And

$$Y^{n-2}Y^2-X^{n-2}X^2=Y^n-X^n=Z^n$$

If $A=0$ then $X^{n-4}=Y^{n-4}$ leads, as $GCD(X,Y)=1$, to $n=4$. This case has been studied by Fermat, it has no solution. Thus $A\neq 0$.
And if $A=\pm Z^n$ then it means that both
$X^{n-3}Y^2=\pm\frac{Z^n}{X}+XY^{n-2}$ and
$Y^{n-3}X^2=\mp\frac{Z^n}{Y}+X^{n-2}Y$ are rationals

it means, as it is impossible, that n is not greater than 2 or $n=2$.
We have

$$\frac{X^{n-2}Z^n}{A}Y^2-\frac{Y^{n-2}Z^n}{A}X^2=Z^n=Y^{n-2}Y^2-X^{n-2}X^2$$

And we have simultaneously

$$(Y^{n-2} - \frac{X^{n-2}Z^n}{A})Y^2 = (X^{n-2} - \frac{Y^{n-2}Z^n}{A})X^2$$

Or

$$(AY^{n-2} - X^{n-2}Z^n)Y^2 = (AX^{n-2} - Y^{p-2}Z^n)X^2$$

And

$$(Y^2 + \frac{X^2Z^n}{A})Y^{n-2} = (X^2 + \frac{Y^2Z^n}{A})X^{n-2}$$

Or

$$(AY^2 + X^2Z^n)Y^{p-2} = (AX^2 + Y^2Z^n)X^{n-2}$$

We have four cases with u and v integers

$$\frac{Y^2}{A} = u(X^{n-2} - \frac{Y^{n-2}Z^n}{A}); \quad \frac{X^2}{A} = u(-\frac{X^{n-2}Z^n}{A} + Y^{p-2})$$

$$\frac{Y^{n-2}}{A} = v(X^2 + \frac{Y^2Z^n}{A}); \quad \frac{X^{n-2}}{A} = v(\frac{X^2Z^n}{A} + Y^2)$$

Or

$$u\frac{Y^2}{A} = X^{n-2} - \frac{Y^{n-2}Z^n}{A}; \quad u\frac{X^2}{A} = -\frac{X^{n-2}Z^n}{A} + Y^{n-2}$$

$$v\frac{Y^{n-2}}{A} = X^2 + \frac{Y^2Z^n}{A}; \quad v\frac{X^{n-2}}{A} = \frac{X^2Z^n}{A} + Y^2$$

Or

$$\frac{Y^2}{A} = u(X^{n-2} - \frac{Y^{n-2}Z^n}{A}); \quad \frac{X^2}{A} = u(-\frac{X^{n-2}Z^n}{A} + Y^{n-2})$$

$$v\frac{Y^{n-2}}{A} = X^2 + \frac{Y^2Z^n}{A}; \quad v\frac{X^{n-2}}{A} = \frac{X^2Z^n}{A} + Y^2$$

Or

$$u\frac{Y^2}{A} = X^{n-2} - \frac{Y^{n-2}Z^n}{A}; \quad u\frac{X^2}{A} = -\frac{X^{n-2}Z^n}{A} + Y^{n-2}$$

$$\frac{Y^{n-2}}{A} = v(X^2 + \frac{Y^2Z^n}{A}); \quad \frac{X^{n-3}}{A} = v(\frac{X^2Z^n}{A} + Y^2)$$

First case

$$Y^n = uv(A^2X^n - Y^nZ^{2n} + AZ^n(Y^2X^{n-2} - Y^{n-2}X^2))$$

$$= uv(A^2X^n - Y^nZ^{2n} + A(AZ^n)) = uv(A^2X^n + A^2Z^n - Y^nZ^{2n}) = uv(A^2Y^n - Z^{2n}Y^n)$$

Thus

$$uv = \frac{1}{A^2 - Z^{2n}}$$

As uv is integer, it means that it is impossible thus $u = 0$ and $A^2 = Z^{2n}$
it means that $p = 2$.
Second case

$$uv\frac{Y^n}{A^2} = X^n - \frac{Y^nZ^{2n}}{A^2} + Z^n\frac{Y^2X^{n-2} - Y^{n-2}X^2}{A}$$

$$= X^n - \frac{Y^nZ^{2n}}{A^2} + Z^n = X^n + Z^n - \frac{Y^nZ^{2n}}{A^2} = (\frac{A^2 - Z^{2n}}{A^2})Y^n$$

Thus

$$uv = A^2 - Z^{2n}$$

And

$$uv(Y^2X^{n-2} - X^2Y^{n-2}) = uvA = u(X^{2n-4} - Y^{2n-4})A = v(-X^4 + Y^4)A$$

Thus

$$u = -X^4 + Y^4; \quad v = X^{2n-4} - Y^{2n-4}$$

$$uv = A^2 - Z^{2n} = (-X^4 + Y^4)(X^{2n-4} - Y^{2n-4})$$

But

$$Y^n = X^n + Z^n \Rightarrow Y^{n-2}X^2 = X^n\frac{X^2}{Y^2} + Z^n\frac{X^2}{Y^2}$$

And

$$Y^2X^{n-2} = X^n\frac{X^{n-2}}{Y^{n-2}} + Z^n\frac{X^{n-2}}{Y^{n-2}}$$

We deduce

$$A = Y^2X^{n-2} - X^2Y^{n-2} = (X^n + Z^n)(\frac{X^{n-2}}{Y^{n-2}} - \frac{X^2}{Y^2})$$

But

$$Y^n > X^n \Rightarrow A < 0 \Rightarrow \frac{X^{n-2}}{Y^{n-2}} < \frac{X^3}{Y^2}$$

And if

$$X^2 < Y^2 \Rightarrow X^{n-2} < Y^{n-2} \Rightarrow A^2 - 1 = (-X^4 + Y^4)(X^{2n-4} - Y^{2n-4} < 0$$

But

$$X^{n-2} < Y^{n-2} \Rightarrow X^{n-2}Y^2 < Y^n = X^n + Z^n \Rightarrow Y^2 < X^2 + Z^nX^{2-n}$$

And

$$0 < Y^2 - X^2 < Z^nX^{2-n}$$

and if $X^n > z^n$ it means $n = 2$. And if

$$X^{n-2} > Y^{n-2} \Rightarrow X^3 > Y^2 \Rightarrow A^2 - 1 = (-X^4 + Y^4)(X^{2n-4} - Y^{2n-4} > 0$$

But

$$X^{n-2} > Y^{n-2} \Rightarrow X^{n-2}Y^2 > Y^n = X^n + Z^n \Rightarrow Y^2 > X^2 + Z^nX^{2-n}$$

And

$$0 > Y^2 - X^2 > Z^nX^{2-n} > 0$$

And $n - 2 = 0$
Third case :
We have here

$$Y^2 = u(AX^{n-2} - Y^{n-2}Z^n); \quad X^2 = u(-X^{n-2}Z^n + AY^{n-2})$$

$$vY^{n-2} = AX^2 + Y^2Z^n; \quad vX^{n-2} = X^2Z^n + AY^2$$

And

$$vY^n = u(A^2X^n - Y^nZ^{2n} + A^2Z^n) = u(A^2 - Z^{2n})Y^n$$

$$v = u(A^2 - Z^{2n})$$

$$v(Y^2X^{n-2} - X^2Y^{p-2}) = vA = uvA(X^{2n-4} - Y^{2n-4}) = A(-X^4 + Y^4)$$

Thus

$$u(X^{2n-4} - Y^{2n-4}) = 1$$

With u and $A^2 - Z^{2n}$ integers, it means $A^2 = Z^{2n}$! Fourth case :

$$u\frac{Y^2}{A} = X^{n-2} - \frac{Y^{n-2}Z^n}{A}; \quad u\frac{X^2}{A} = -\frac{X^{n-2}Z^n}{A} + Y^{n-2}$$

$$\frac{Y^{n-2}}{A} = v(X^2 + \frac{Y^2Z^n}{A}); \quad \frac{X^{n-2}}{A} = v(\frac{X^2Z^n}{A} + Y^2)$$

Hence

$$u\frac{Y^n}{A^2} = v(X^n - \frac{Y^nZ2n}{A^2} + Z^n) = v(1 - \frac{Z^{2n}}{A^2})Y^n$$

Thus

$$u = v(A^2 - Z^{2n})$$

$$u(Y^2X^{n-2} - X^2Y^{n-2}) = uA = A(X^{2n-4} - Y^{2n-4}) = uv(-X^4 + Y^4)A$$

$$v(-X^4 + Y^4) = 1$$

With v and $A^2 - Z^{2n}$ integers, it means $A^2 - Z^{2n} = 0$!
The only solution, in all cases, in $n = 2$.

Resolution of Fermat- Catalan equation

Let Fermat-Catalan equation :

$$Y^p = X^q + aZ^c$$

$a = \pm 1$
We have

$$X^{q-w}Y^2 - Y^{p-2}X^w = aA$$

And

$$Y^{p-2}Y^2 - X^{q-w}X^w = Y^p - X^q = aZ^c$$

If $A = 0$ then $X^{q-2w} = Y^{p-4}$ leads, as $GCD(X, Y) = 1$, to $p = 4$ and $q = 2w$. Thus $A \neq 0$.
And if $A = \pm Z^c$ then it means that both
$X^{q-w-1}Y^2 = \pm\frac{aZ^c}{X} + X^{w-1}Y^{p-2}$ and
$Y^{p-3}X^w = \mp\frac{aZ^c}{Y} + X^{q-w}Y$ are rationals

it means, as i is imossble, that $q = w$ and $p = 2$.
We have

$$\frac{X^{q-w}Z^c}{A}Y^2 - \frac{Y^{p-2}Z^c}{A}X^w = aZ^c = Y^{p-2}Y^2 - X^{q-w}X^w$$

And we have simultaneously

$$(Y^{p-2} - \frac{X^{q-w}Z^c}{A})Y^2 = (X^{q-w} - \frac{Y^{p-2}Z^c}{A})X^w$$

Or

$$(AY^{p-2} - X^{q-w}Z^c)Y^2 = (AX^{q-w} - Y^{p-2}Z^c)X^w$$

And

$$(Y^2 + \frac{X^wZ^c}{A})Y^{p-2} = (X^w + \frac{Y^2Z^c}{A})X^{q-w}$$

Or

$$(AY^2 + X^wZ^c)Y^{p-2} = (AX^w + Y^2Z^c)X^{q-w}$$

We have four cases with u and v integers

$$\frac{Y^2}{A} = u(X^{q-w} - \frac{Y^{p-2}Z^c}{A}) \quad \frac{X^w}{A} = u(-\frac{X^{q-w}Z^c}{A} + Y^{p-2})$$

$$\frac{Y^{p-2}}{A} = v(X^w + \frac{Y^2Z^c}{A}) \quad \frac{X^{q-w}}{A} = v(\frac{X^wZ^c}{A} + Y^2)$$

Or

$$\frac{uY^2}{A}) = X^{q-w} - \frac{Y^{p-2}Z^c}{A} \quad \frac{uX^w}{A}) = -\frac{X^{q-w}Z^c}{A} + Y^{p-2}$$

$$\frac{vY^{p-2}}{A}) = X^w + \frac{Y^2Z^c}{A} \quad \frac{vX^{q-w}}{A} = \frac{X^wZ^c}{A} + Y^2$$

Or

$$\frac{Y^2}{A} = u(X^{q-w} - \frac{Y^{p-2}Z^c}{A}) \quad \frac{X^w}{A} = u(-\frac{X^{q-w}Z^c}{A} + Y^{p-2})$$

$$v\frac{Y^{p-2}}{A} = X^w + \frac{Y^2Z^c}{A} \quad v\frac{X^{q-w}}{A} = \frac{X^wZ^c}{A} + Y^2$$

Or

$$u\frac{Y^2}{A} = X^{q-w} - \frac{Y^{p-2}Z^c}{A} \quad u\frac{X^w}{A} = -\frac{X^{q-w}Z^c}{A} + Y^{p-2}$$

$$\frac{Y^{p-2}}{A} = v(X^w + \frac{Y^2Z^c}{A}) \quad \frac{X^{q-w}}{A} = v(\frac{X^wZ^c}{A} + Y^2)$$

First case

$$Y^p = uv(A^2X^q - Y^pZ^{2c} + AZ^c(Y^2X^{q-w} - Y^{p-2}X^w))$$

$$= uv(A^2X^q - Y^pZ^{2c} + aA(A)Z^c) = uv(A^2X^q + A^2Z^c - Y^pZ^{2c}) = uv(A^2Y^p - Y^pZ^{2c})$$

Thus

$$uv = \frac{1}{A^2 - Z^{2c}}$$

As uv is integer, it means that it is impossible thus $u = 0$ and $A^2 = Z^{2c}$ or $A = \pm -Z^c$ it means that $q = w$ and $p = 2$.
Second case

$$uv\frac{Y^p}{A^2} = X^q - \frac{Y^p Z^{2c}}{A^2} + Z^c \frac{Y^2 X^{q-w} - Y^{p-2} X^w}{A}$$
$$= X^q - \frac{Y^p Z^{2c}}{A^2} + aZ^c = X^q + aZ^c - \frac{Y^p Z^{2c}}{A^2} = (\frac{A^2 - Z^{2c}}{A^2})Y^p$$

Thus

$$uv = A^2 - Z^{2c}$$

And (with $a > 0$, the proof is similar for $a < 0$)

$$uv(Y^2 X^{q-w} - X^w Y^{p-2}) = uvaA = u(X^{2q-2w} - Y^{2p-4})A = v(-X^{2w} + Y^4)A$$

Thus

$$ua = -X^{2w} + Y^4; \quad va = X^{2q-2w} - Y^{2p-4}$$
$$uv = A^2 - Z^{2c} = (-X^{2w} + Y^4)(X^{2q-2w} - Y^{2p-4})$$

$$Y^b = X^a + Z^c \Rightarrow Y^{b-2} X^w = X^a \frac{X^w}{Y^2} + Z^n \frac{X^w}{Y^2}$$

And

$$Y^2 X^{a-w} = X^a \frac{X^{a-w}}{Y^{b-2}} + Z^c \frac{X^{a-w}}{Y^{b-2}}$$

We deduce

$$A = Y^2 X^{a-w} - X^w Y^{b-2} = (X^a + Z^c)(\frac{X^{a-w}}{Y^{b-2}} - \frac{X^w}{Y^2})$$

But

$$Y^b > X^a \Rightarrow A < 0 \Rightarrow \frac{X^{a-w}}{Y^{b-2}} < \frac{X^w}{Y^2}$$

And if

$$X^w < Y^2 \Rightarrow X^{a-w} < Y^{b-2} \Rightarrow A^2 - 1 = (-X^{2w} + Y^4)(X^{2a-2w} - Y^{2b-4} < 0$$

But

$$X^{a-w} < Y^{b-2} \Rightarrow X^{a-w} Y^2 < Y^n = X^a + Z^c \Rightarrow Y^2 < X^w + Z^c X^{w-a}$$

And

$$0 < Y^2 - X^w < Z^c X^{w-a}$$

and if $X^a > Z^c$ it means $b = 2$. And if

$$X^{a-w} > Y^{b-2} \Rightarrow X^w > Y^2 \Rightarrow A^2 - 1 = (-X^{2w} + Y^4)(X^{2a-2w} - Y^{2b-4} > 0$$

But

$$X^{a-w} > Y^{b-2} \Rightarrow X^{a-w} Y^2 > Y^b = X^a + Z^c \Rightarrow Y^2 > X^w + Z^c X^{w-a}$$

And

$$0 > Y^2 - X^w > Z^c X^{w-a} > 0$$

And $b - 2 = 0$
Third case :
We have here

$$Y^2 = u(AX^{q-w} - Y^{p-2} Z^c); \quad X^w = u(-X^{q-w} Z^c + AY^{p-2})$$
$$vY^{p-2} = AX^w + Y^2 Z^c; \quad vX^{q-w} = X^w Z^c + AY^2$$

And

$$vY^p = u(A^2 X^q - Y^p Z^{2c} + aA^2 Z^c) = u(A^2 - Z^{2c})Y^p$$
$$v = u(A^2 - Z^{2c})$$
$$v(Y^2 X^{q-w} - X^w Y^{p-2}) = vaA = uvA(X^{2q-2w} - Y^{2p-4}) = A(-X^{2w} + Y^4)$$

Thus $au(X^{2q-2w} - Y^{2p-4}) = 1$: it is possible only if $p = 2$ and $q = w$. Fourth case :

$$u\frac{Y^2}{A} = X^{q-w} - \frac{Y^{p-2}Z^c}{A}; \quad u\frac{X^w}{A} = -\frac{X^{q-w}Z^c}{A} + Y^{p-2}$$
$$\frac{Y^{p-2}}{A} = v(X^w + \frac{Y^2Z^c}{A}); \quad \frac{X^{q-w}}{A} = v(\frac{X^wZ^c}{A} + Y^2)$$

We have here

$$uY^2 = AX^{q-w} - Y^{p-2}Z^c; \quad uX^w - AY^{p-2} = -X^{q-w}Z^c$$

And

$$Y^{p-2} = AX^{q-w} - uY^2Z^c = (Y^2X^{q-w} - X^wY^{p-2})X^{q-w} - uY^2Z^c$$

Hence

$$u\frac{Y^p}{A^2} = v(X^q - \frac{Y^pZ^{2c}}{A^2} + aZ^c) = v(1 - \frac{Z^{2c}}{A^2})Y^p$$

Thus

$$u = v(A^2 - Z^{2c})$$
$$u(Y^2X^{q-w} - X^wY^{p-2}) = uaA = A(X^{2q-2w} - Y^{2p-4}) = uv(-X^{2w} + Y^4)A$$

Thus $-av(X^{2w} - Y^4) = 1$: it is possible only if $u = 0$ and $p - 2 = q - w = 0$
The only solution, in all cases, is $p = 2$.
And $Y^2 = X^q + aZ^c$.

Conclusion

Fermat-Catalan equation $Y^p = X^q \pm Z^c$ has solutions only for $p = 2$. We have shown a way to solve it.

References

[1] Paolo Ribenbo?m, The Catalan's conjecture *Academic press* , (1994).

[2] Robert Tijdeman, On the equation of Catalan *Acta Arith* , (1976).