



**HAL**  
open science

# Flavonoid conjugates interact with organic anion transporters (OATs) and attenuate cytotoxicity of adefovir mediated by organic anion transporter 1 (OAT1/SLC22A6)

Chi Chun Wong, Nigel P Botting, Caroline Orfila, Gary Williamson

## ► To cite this version:

Chi Chun Wong, Nigel P Botting, Caroline Orfila, Gary Williamson. Flavonoid conjugates interact with organic anion transporters (OATs) and attenuate cytotoxicity of adefovir mediated by organic anion transporter 1 (OAT1/SLC22A6). *Biochemical Pharmacology*, 2011, 81 (7), pp.942. <10.1016/j.bcp.2011.01.004>. <hal-00678041>

**HAL Id: hal-00678041**

**<https://hal.science/hal-00678041v1>**

Submitted on 12 Mar 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

## Accepted Manuscript

Title: Flavonoid conjugates interact with organic anion transporters (OATs) and attenuate cytotoxicity of adefovir mediated by organic anion transporter 1 (OAT1/SLC22A6)

Authors: Chi Chun Wong, Nigel P Botting, Caroline Orfila, Gary Williamson



PII: S0006-2952(11)00037-2  
DOI: doi:10.1016/j.bcp.2011.01.004  
Reference: BCP 10800

To appear in: *BCP*

Received date: 22-11-2010  
Revised date: 5-1-2011  
Accepted date: 6-1-2011

Please cite this article as: Wong CC, Botting NP, Orfila C, Williamson G, Flavonoid conjugates interact with organic anion transporters (OATs) and attenuate cytotoxicity of adefovir mediated by organic anion transporter 1 (OAT1/SLC22A6), *Biochemical Pharmacology* (2010), doi:10.1016/j.bcp.2011.01.004

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Flavonoid conjugates interact with organic anion transporters (OATs) and attenuate cytotoxicity of adefovir mediated by organic anion transporter 1 (OAT1/SLC22A6)**

Chi Chun Wong<sup>a</sup>, Nigel P Botting<sup>b</sup>, Caroline Orfila<sup>a</sup> and Gary Williamson<sup>a,c</sup>

<sup>a</sup> School of Food Science and Nutrition, University of Leeds, Leeds, LS2 9JT, UK

<sup>b</sup> School of Chemistry, University of St Andrews, North Haugh, St Andrews, Fife, KY16 9ST, Scotland.

<sup>c</sup> Nestle Research Center, Vers Chez les Blanc, 1000 Lausanne 26, Switzerland

Correspondence to: Professor Gary Williamson, School of Food Science and Nutrition, University of Leeds, Leeds, LS2 9JT, UK.

Tel: +44 (0) 113 343 8380

Fax: +44 (0) 113 343 2982

Email: g.williamson@leeds.ac.uk

Emails address of other authors: fscdw@leeds.ac.uk (Chi Chun Wong), npb@st-andrews.ac.uk (Nigel Botting), c.orfila@leeds.ac.uk (Caroline Orfila)

**Abstract**

1  
2  
3 Flavonoids are conjugated by phase II enzymes in humans to form glucuronidated and  
4  
5 sulfated metabolites that are excreted in urine via the kidney. In this study, we examined the  
6  
7 interaction between metabolites of quercetin and isoflavonoids found *in vivo* with human  
8  
9 organic anion transporters 1 (OAT1) and 3 (OAT3) and their potential in attenuating OATs-  
10  
11 induced cytotoxicity of adefovir. Accumulation of flavonoid conjugates was studied in  
12  
13 human embryonic kidney 293H cells overexpressing OAT1 or OAT3. OAT1-overexpressing  
14  
15 cells exhibited an increased uptake of the sulfated conjugates, genistein-4'-*O*-sulfate and  
16  
17 quercetin-3'-*O*-sulfate. OAT3-overexpressing cells demonstrated enhanced uptake of  
18  
19 glucuronide conjugates, such as daidzein-7-*O*-glucuronide, genistein-7-*O*-glucuronide,  
20  
21 glycitein-7-*O*-glucuronide and quercetin-3'-*O*-glucuronide. Position of conjugation was  
22  
23 important since quercetin-3-*O*-glucuronide and quercetin-7-*O*-glucuronide were poorly  
24  
25 transported. Kinetic analysis revealed high affinity uptake of quercetin-3'-*O*-sulfate by OAT1  
26  
27 ( $K_m = 1.73 \mu\text{M}$ ;  $V_{\text{max}} = 105 \text{ pmol}/(\text{min} \times \text{mg})$ ). OAT3 transported isoflavone glucuronides  
28  
29 with lower affinity ( $K_m = 7.9\text{-}19.1 \mu\text{M}$ ) but with higher  $V_{\text{max}}$  (171-420 pmol/(min x mg)).  
30  
31 Quercetin-3'-*O*-sulfate strongly inhibited OAT1-mediated *p*-aminohippuric acid uptake with  
32  
33 an  $\text{IC}_{50}$  of 1.22  $\mu\text{M}$ . Transport of 5-carboxyfluorescein by OAT3 was potently inhibited by  
34  
35 quercetin-3-*O*-glucuronide, quercetin-3'-*O*-glucuronide and quercetin-3'-sulfate ( $\text{IC}_{50} = 0.43\text{-}$   
36  
37 1.31  $\mu\text{M}$ ). In addition, quercetin-3'-*O*-sulfate was shown to effectively reduce OAT1-  
38  
39 mediated cytotoxicity of adefovir, an antiviral drug, in a dose-dependent manner. These data  
40  
41 suggest that OAT1 and OAT3 are responsible for basolateral uptake of flavonoid conjugates  
42  
43 in kidney, and flavonoid conjugates inhibit OAT1 and OAT3 activity at physiologically  
44  
45 relevant concentrations. Interaction with OATs limits systemic availability of flavonoids and  
46  
47 may be a mechanism of food-drug interaction via inhibition of renal uptake.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 **Keywords:** bioavailability, food-drug interaction, isoflavones, organic anion transporters,  
2 quercetin  
3

4  
5 **Abbreviations**  
6

7  
8  
9 BCRP: breast cancer resistance protein; MRP: multidrug resistance protein; OAT: organic  
10 anion transporter  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1. Introduction

1  
2  
3 Quercetin is a chemopreventive and anti-inflammatory flavonoid that interacts with  
4 nuclear factor E2-related factor 2 (Nrf2) and activates antioxidant response elements (ARE)  
5 [1, 2], contributing to enhanced cellular protection against carcinogens and oxidative stress  
6 [3]. Isoflavones are phytoestrogens that bind to the estrogen receptor [4] and exert preventive  
7 effects on hormone-dependent cancer [5], cardiovascular diseases [6], and post-menopausal  
8 osteoporosis [7]. Any biological activities of flavonoids, however, are influenced by their  
9 metabolism. Dietary flavonoids are metabolized extensively in the intestine and liver, and  
10 major forms found *in vivo* are glucuronide, sulfate and/or methyl-conjugates of the parent  
11 aglycone [8]. Quercetin is metabolized into quercetin-3-*O*-glucuronide, 3'-methylquercetin-3-  
12 *O*-glucuronide and quercetin-3'-*O*-sulfate following consumption of onions [9]. After soy  
13 ingestion, soy isoflavones such as daidzein and genistein are present in the circulation  
14 predominantly as their 7-*O*-glucuronide and 4'-*O*-glucuronide, and sulfates were minor  
15 metabolites [10]. Concentrations of these metabolites in human plasma is typically low after  
16 intake of 50 mg aglycone equivalent and it rarely exceeds 10  $\mu$ M even at much higher  
17 ingested doses [11].

18  
19  
20 Interaction with uptake and efflux transporters is considered to be one of the factors  
21 that limit the bioavailability of flavonoids, by facilitating the elimination of hydrophilic  
22 conjugates into bile and/or urine [12]. Urinary excretion of flavonoid conjugates has been  
23 well documented and varies according to the individual flavonoid aglycones. For the  
24 isoflavones daidzein and glycitein, about half of the intake could be recovered in the urine as  
25 conjugated metabolites [13]. A high urinary recovery of conjugates was also observed with  
26 catechins (~25 %) [14]; while for quercetin the urinary excretion is much less than 10 % of  
27 the intake [15]. Many drugs and their conjugated metabolites are eliminated in the urine via

1 tubular secretion [16]. Vectorial transport of hydrophilic substances across the renal proximal  
2 tubules involves the interplay of uptake transporters on the basolateral membrane and the  
3  
4 efflux transporters on the apical membrane. Active efflux is carried out by multidrug  
5  
6 resistance proteins (MRP), which transport glucuronide and sulfate conjugates of flavonoids  
7  
8 such as quercetin, baicalein and genistein [17-19]. The molecular mechanism underlying the  
9  
10 possible uptake of flavonoid conjugates into renal tubules via basolateral transporters is not  
11  
12 known.  
13  
14  
15

16  
17  
18       Organic anion transporters OAT1 (*SLC22A6*) and OAT3 (*SLC22A8*) are highly  
19  
20 expressed on the basolateral membrane of proximal tubules [20]. They play an important role  
21  
22 in the sodium-dependent renal uptake of structurally diverse organic anions from the blood  
23  
24 into proximal tubule cells. OAT1 and OAT3 are capable of mediating the uptake of these  
25  
26 organic anions against their electrochemical gradients by coupling to efflux of intracellular  $\alpha$ -  
27  
28 ketoglutarate. OAT1 interacts with endogenous metabolites, drugs including antivirals,  
29  
30 antibiotics, nonsteroidal anti-inflammatory agents and statins, as well as toxicants such as  
31  
32 mercury and ochratoxin [21]. OAT3 recognizes many OAT1 substrates; in addition, it has the  
33  
34 ability to transport corticosterone, estradiol-17 $\beta$ -glucuronide and taurocholate, that are not  
35  
36 substrates of OAT1 [22]. As toxins and cytotoxic drugs are taken up by OAT1 and OAT3,  
37  
38 these toxic substances may exert toxic effects on the proximal tubule cells and cause renal  
39  
40 injury. *In vitro* studies with OAT1-transfected cells showed that OAT1 over-expression  
41  
42 sensitizes the cytotoxicity caused by antiviral drugs, adefovir and cidofovir [23]. Co-  
43  
44 administration of the inhibitors of OATs may reduce the extent of OAT-induced  
45  
46 nephrotoxicity [24]. On the other hand, inhibition of OAT1 and OAT3 may also alter the  
47  
48 pharmacokinetics of a variety of drugs that are OAT substrates. OAT1 Inhibitors, such as  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

probenecid, could be administered with other drugs to prolong plasma half lives and improve therapeutic effects [25].

To date, only limited information is available concerning the interaction between OATs and flavonoids. Ellagic acid is a substrate and a potent inhibitor of OAT1 [26], while several flavonoids (naringenin, morin, silybin and quercetin) were also found to inhibit OAT1 and OAT3 [27]. However, the flavonoid aglycones are mostly absent in human plasma. The aim of this work is to investigate the transport of glucuronidated and sulfated metabolites of quercetin and isoflavones, major metabolites *in vivo*, in OAT1- and OAT3-transfected human embryonic kidney 293H cells. We also examined the inhibition of OATs by flavonoid conjugates and the potential of quercetin-3'-*O*-sulfate to attenuate OAT1-induced cytotoxicity of adefovir.

## 2. Methods and materials

### 2.1. Chemicals

Genistein and quercetin were obtained from Extrasynthese (Genay, France). Quercetin-7-*O*-glucuronide, quercetin-3-*O*-glucuronide and quercetin-3'-*O*-glucuronide were synthesized from quercetin using human liver S9 (Sigma-Aldrich, St Louis, MO) and purified by HPLC [28]. Quercetin-3'-*O*-sulfate was chemically synthesized as described and purified by gel filtration [9]. The identity of quercetin-3-*O*-glucuronide and quercetin-3'-*O*-sulfate was further confirmed by comparing the retention time and the absorption spectra of the authentic standards, kind gifts from Dr. Paul Kroon, Institute of Food Research, Norwich, UK. Purities were checked by HPLC to be over 95 %. Daidzein-7-*O*-glucuronide, genistein-7-*O*-glucuronide, glycitein-7-*O*-glucuronide, genistein-4'-*O*-sulfate and daidzein-7,4'-*O*-disulfate

1 were synthesized as previously described [10, 29]. Adefovir was obtained by the hydrolysis  
2 of adefovir dipivoxil (Sigma-Aldrich, St Louis, MO) using human intestinal S9 (Xenotech  
3 LLC, Kansas City, KS) and purified by gel filtration. Human embryonic kidney cells 293H  
4 were purchased from Invitrogen (Carlsbad, CA). OAT1 (SLC22A6 transcript variant 2,  
5 Genebank accession number: NM\_153276) and OAT3 (SLC22A8 transcript variant 1,  
6 Genebank accession number: NM\_004254) expression plasmids were obtained from Origene  
7 (Rockville, MD). Fugene HD was purchased from Roche (Basel, Switzerland). All other  
8 chemicals, unless otherwise stated, were purchased from Sigma-Aldrich (St Louis, MO).  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19

## 20 **2.2. HEK293 cell culture**

21  
22  
23 293H cells were routinely cultured in 75 cm<sup>2</sup> cell culture flasks (Corning Costar Corp.,  
24 Cambridge, MA) at 37 °C under a humidified 5 % CO<sub>2</sub> / O<sub>2</sub> atmosphere. Culture media  
25 consisted of Dulbecco's modified Minimum Essential Medium (DMEM) high-glucose media  
26 supplemented with 10% fetal bovine serum, 1 % nonessential amino acids and 50 U/ml  
27 penicillin-streptomycin (All from Sigma-Aldrich). Cells were split in a 1:4 ratio every 48 h.  
28  
29 Antibiotics were not added in transfection experiments. All experiments were performed with  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 66 **2.3. Transfection and uptake transport assay**

67 The cells were seeded into poly-L-lysine coated 24-well plates at a density of 1.2 x  
68 10<sup>5</sup> cells/per well the day prior to transient transfection. OAT1 or OAT3 plasmids, or the  
69 empty pCMV-XL6 vector, were transfected into 293H cells with the Fugene HD reagent  
70 according to the manufacturer's directions, using an optimized ratio of 3:2 (µl reagent: µg  
71 DNA). The transfection complexes were formed in Opti-MEM (Invitrogen, Carlsbad, CA)  
72 and then added to cells after incubation for 18 min. Uptake assays were performed 22-24 h  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

after transfection. The over-expression of the OATs was confirmed by quantitative RT-PCR (Taqman assays ID: Hs00537914\_m1 (SLC22A6) and Hs01056647\_m1 (SLC22A8), (Applied Biosystems, Foster City, CA), and by the uptake of model substrates *p*-aminohippuric acid (OAT1), and estrone-3-sulfate (OAT3). Uptake experiments were carried out using HBSS buffer containing 1.8 mM CaCl<sub>2</sub> and adjusted to pH 7.4 with 1 M HCl. The media were removed and the monolayers were washed twice with 0.25 mL transport buffer and incubated for 10 min. After that, the buffer was aspirated and 0.25 mL transport buffer containing the test compounds was added. Flavonoid and isoflavone conjugates were dissolved in DMSO (final concentration < 0.2 %), except quercetin-3'-*O*-sulfate which was dissolved in water. After the specified incubation time, the uptake was stopped by adding 1 mL ice-cold transport buffer containing 0.2 % bovine serum albumin (BSA). This was quickly aspirated and further washed twice with 0.5 mL ice-cold transport buffer with 0.2 % BSA. Finally, the cells were rinsed with 1 mL ice-cold transport buffer without BSA. Cells were collected with 0.4 mL lysis solution (50 % methanol) and stored at -20 °C. Extraction was performed by sonication for 5 min followed by the addition of 1 mL of ice-cold acetone. The samples were placed in a -20 °C freezer for 1 h and centrifuged at 17,000 g for 5 min. The supernatant was collected and evaporated to dryness *in vacuo* at 30 °C and stored at -20 °C until analysis. The protein pellet was re-dissolved in 0.1 N NaOH and the protein contents were determined by the Bradford assay. All the uptake values were standardized against protein content.

#### 2.4. Kinetics of uptake of flavonoid and isoflavonoid conjugates

Uptake assays were performed on the 293H cells 20-24 h after transfection. Uptake assays were performed for 10 min, during which uptake is linear with time. Kinetic analyses were performed for genistein-4'-*O*-sulfate and quercetin-3'-*O*-sulfate in OAT1-expressing

1 cells, with a substrate concentration range of 1 to 50  $\mu\text{M}$ . The uptake of glucuronides by  
2 OAT3 was linear over 30 min. The kinetics of uptake of daidzein-7-*O*-glucuronide, genistein-  
3 7-*O*-glucuronide, glycitein-7-*O*-glucuronide and quercetin-3'-*O*-glucuronide were measured  
4 in OAT3-expressing cells for 10 min, with substrate concentrations of 2.5 to 100  $\mu\text{M}$ . The  
5 apparent uptake of OATs was obtained by subtracting the uptake of OAT1 or OAT3 with  
6 uptake of control cells at each concentration.  
7  
8  
9  
10  
11  
12  
13

## 14 **2.5. HPLC analysis of flavonoid and isoflavonoid conjugates**

15 HPLC analyses were carried out on an Agilent 1200 series liquid chromatography  
16 system equipped with a diode array detector. The analyses were performed with a Zorbax  
17 XDB-C18 column (4.6 x 50 mm, 1.8  $\mu\text{m}$ ) with 20 mM ammonium formate, pH 4.5 (A) and  
18 methanol (B) as the mobile phase. For the analysis of quercetin and their conjugates, elution  
19 was performed at 0.75 mL/min and gradient started at 30 % (B) and increased linearly to 50  
20 % in 12 min, and then equilibrated at 30 % for 3 min. Quantification of quercetin metabolites  
21 was based on peak area of quercetin at 370 nm, assuming similar response factor for  
22 quercetin conjugates [30, 31]. For the analysis of isoflavones and their conjugates, elution  
23 started at 20 % (B), increasing to 30 % in 5 min and to 70 % in 8 min, after which it returned  
24 to 20 % for 3 min at a flow rate of 0.75 mL/min. The quantification of isoflavone metabolites  
25 was based on the peak area of the respective standard curves measured at 254 nm. Typically,  
26 50  $\mu\text{l}$  was injected for the analyses.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

## 48 **2.6. Inhibition of uptake of standard substrates**

49 *p*-aminohippuric acid and 5-carboxyfluorescein were used as the model substrates for  
50 OAT1 and OAT3, respectively. Uptake of *p*-aminohippuric acid (25  $\mu\text{M}$ ) and 5-  
51 carboxyfluorescein (100  $\mu\text{M}$ ) was measured for 10 min in the presence or absence of 10  $\mu\text{M}$   
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 test compounds. The intracellular *p*-aminohippuric acid was extracted as in section 2.3 and  
2 was analyzed by HPLC. The analyses were performed with a Zorbax XDB-C18 column (4.6  
3 x 50 mm, 1.8  $\mu$ m) with 20 mM ammonium formate, pH 4.5 as the mobile phase. It was run at  
4 a flow rate of 0.75 mL for 3 min. To quantify 5-carboxyfluorescein, the cells were extracted  
5 twice with 1 mL methanol in the dark, each for 30 min, and the supernatants were combined  
6 and analyzed by HPLC. The analyses were performed with Zorbax XDB-C18 column (4.6 x  
7 50 mm, 1.8  $\mu$ m) and eluted with 20 mM Tris-HCl (pH 8) and 20 % methanol, at 0.25 mL/min.  
8 5-carboxyfluorescein was detected using a fluorescence detector (Agilent) with excitation and  
9 emission of 492 and 517 nm, respectively.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

## 22 **2.7. Drug cytotoxicity assays**

23  
24  
25  
26 293H cells were plated onto 35-mm culture plates at with a density of  $5 \times 10^5$  cells per  
27 plate. After overnight incubation, the cells were transfected with OAT1, OAT3 or empty  
28 plasmids with fugene. Twenty-four hours after transfection, the cells were subcultivated into  
29 96 well plates at 20,000 cells per well and cultured in DMEM complete media. After  
30 overnight incubation, various concentrations of adefovir and/or quercetin-3'-*O*-sulfate were  
31 added (five replicates), after which the cells were further incubated for another 96 h. At the  
32 end of the incubation, cell viability was determined by a modified colorimetric assay using 3-  
33 [4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) [32]. Briefly, the culture  
34 medium was removed and replaced with 100  $\mu$ l phenol red-free complete media containing  
35 0.5 mg/mL MTT. Following 4 h incubation at 37  $^{\circ}$ C, 100  $\mu$ l of a solution containing 10%  
36 Triton X-100 and 0.1N HCL in isopropanol was added. The plate was incubated and gently  
37 mixed until MTT formazan crystals were dissolved. Absorbance at 570 nm was measured on  
38 a microplate reader and IC<sub>50</sub> calculated after subtraction of blank values.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 2.8 Data Analysis

1  
2  
3 Data are shown as mean  $\pm$  S.D. (as mean  $\pm$  S.E. for enzyme kinetics data). Raw data  
4  
5 from the enzyme kinetics studies were analyzed using GraphPad Prism 5.  $K_m$  and  $V_{max}$  were  
6  
7 derived from a nonlinear regression fit of the Michaelis–Menten model. Statistical differences  
8  
9 were determined using analysis of variance using the student's *t*-test. Differences were  
10  
11 considered significant when  $p \leq 0.05$ .  
12  
13  
14  
15  
16  
17  
18

## 3. Results

### 3.1. Uptake of flavonoid and isoflavone conjugates by OAT1

19  
20  
21  
22  
23  
24  
25  
26 To test whether flavonoid and isoflavone conjugates are transported by OAT1, 293H  
27  
28 cells transfected with empty or OAT1 plasmid were incubated with 25  $\mu$ M of the test  
29  
30 compounds for 30 min. The results are shown in Figure 1 and 2. Sulfated metabolites  
31  
32 quercetin-3'-*O*-sulfate and genistein-4'-*O*-sulfate, appeared to be the best substrates of OAT1  
33  
34 (> 10-fold over control), especially the former (Figure 1 and 2). However, daidzein-7,4'-*O*-  
35  
36 disulfate was not significantly transported. The glucuronide conjugates were rather poor  
37  
38 substrates of OAT1: daidzein-7-*O*-glucuronide, genistein-7-*O*-glucuronide, glycitein-7-*O*-  
39  
40 glucuronide and quercetin-3'-*O*-glucuronide were not transported; while uptake of quercetin-  
41  
42 7-*O*-glucuronide and quercetin-3-*O*-glucuronide was only 2- and 5-fold greater than the  
43  
44 control respectively. Subsequently, we further examined the uptake of the two sulfate  
45  
46 conjugates. Uptake of both sulfates was effectively inhibited by 1 mM probenecid, an OAT  
47  
48 inhibitor (Figure 3). It was found that OAT1 mediated a high-affinity transport of quercetin-  
49  
50 3'-*O*-sulfate, with  $K_m$  of  $1.73 \pm 0.38 \mu$ M (Table 1). Uptake of genistein-4'-*O*-sulfate was less  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 active compared to quercetin-3'-*O*-sulfate with a higher  $K_m$  and lower  $V_{max}$ . Quercetin and  
2 genistein, the aglycones with no attached sulfate, were not actively transported by OAT1.  
3  
4

### 5 **3.2. Uptake of flavonoid and isoflavone conjugates by OAT3**

6  
7

8  
9 The uptake of conjugates by OAT3 has a markedly different profile compared to  
10 OAT1 (Figure 1 and 2). OAT3 possessed a broader specificity since all the conjugates tested  
11 showed a significant increase in their uptake compared to the control transfected cells.  
12  
13 Glucuronide conjugates, in particular, were highly transported by OAT3. Uptake of daidzein-  
14 7-*O*-glucuronide and genistein-7-*O*-glucuronide in OAT3-expressing cells were in excess of  
15 300-fold over control cells; and for glycitein-7-*O*-glucuronide the transport into cells was  
16 100-fold greater. Uptake of the glucuronide conjugates was completely inhibited by co-  
17 incubation with 1 mM probenecid (Figure 3). Among quercetin conjugates, OAT3-mediated  
18 transport of quercetin-3'-*O*-glucuronide and quercetin-3'-*O*-sulfate were most active. Uptake  
19 of quercetin-7-*O*-glucuronide and quercetin-3-*O*-glucuronide were comparatively lower.  
20  
21 Genistein-4'-*O*-sulfate was also transported by OAT3. Interestingly, daidzein-7,4'-*O*-disulfate,  
22 not a substrate for OAT1, was significant transported by OAT3. However, the rate of  
23 transport was low for the tested isoflavone sulfates compared to glucuronide conjugates. We  
24 then determined the uptake kinetics of several glucuronide conjugates (Table 1). OAT3  
25 possessed a higher affinity (lower  $K_m$ ) towards quercetin-3'-*O*-glucuronide and genistein-7-  
26 *O*-glucuronide, with a lower affinity (higher  $K_m$ ) towards daidzein-7-*O*-glucuronide and  
27 glycitein-7-*O*-glucuronide. On the other hand, the maximal rate of transport was highest for  
28 daidzein-7-*O*-glucuronide, followed by genistein-7-*O*-glucuronide, glycitein-7-*O*-glucuronide  
29 and quercetin-3'-*O*-glucuronide. As seen for OAT1, the aglycones did not appear to be  
30 actively transported by OAT3.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### 3.3. Inhibition of OAT1-mediated uptake of *p*-aminohippuric acid by flavonoid and isoflavones conjugates

The flavonoid/isoflavone aglycones and conjugates were tested for their ability to inhibit OAT-mediated uptake of the model substrate for OAT1, *p*-aminohippuric acid. Uptake of *p*-aminohippuric acid in OAT1-expressing cells was sensitive to probenecid, an OAT1 inhibitor, with a complete inhibition at 1 mM. We tested the inhibitory effects of the aglycones and conjugates at 2 and 10  $\mu$ M. At 10  $\mu$ M, a concentration that is generally considered to be supraphysiological, all flavonoids conjugates tested showed a significant inhibition on OAT1-mediated uptake of *p*-aminohippuric acid (Figure 4). Quercetin-3'-*O*-sulfate is the most potent inhibitor and almost completely inhibited the uptake of *p*-aminohippuric acid. Quercetin-7'-*O*-glucuronide, quercetin-3'-*O*-glucuronide, genistein-4'-*O*-sulfate and the aglycones also resulted in over 50 % inhibition of uptake. Isoflavone glucuronides are comparatively weaker inhibitors of OAT1. At a physiologically achievable concentration (2  $\mu$ M), however, most of the conjugates tested show little inhibition of OAT1. Only the sulfated conjugates daidzein-7,4'-*O*-disulfate, genistein-4'-*O*-sulfate and quercetin-3'-*O*-sulfate demonstrated a statistically significant inhibition of *p*-aminohippuric acid uptake. In particular, quercetin-3'-sulfate demonstrated a strong inhibition (> 75 %) even at 2  $\mu$ M. The OAT1-inhibitory activity of quercetin-3'-*O*-sulfate was examined over a range of concentrations (0.1 to 10  $\mu$ M). IC<sub>50</sub> of the inhibition was 1.22  $\mu$ M, indicating that it is a potent inhibitor of OAT1 (Figure 5).

### 3.4. Inhibition of OAT3-mediated transport of 5-carboxyfluorescein by flavonoid and isoflavones conjugates

5-Carboxyfluorescein, a fluorescent marker, is a good model substrate of OAT3 [33]. Flavonoid/ isoflavone conjugates were tested for the inhibitory activity on OAT3-mediated

uptake of 5-carboxyfluorescein, at 2 and 10  $\mu\text{M}$  concentrations (Figure 6). We observed that the inhibition of OAT3 by the flavonoid conjugates tested was generally stronger compared to OAT1. All the flavonoids tested showed a considerable (40-90 %) inhibition of OAT3-mediated uptake at 10  $\mu\text{M}$ . Quercetin metabolites showed a stronger inhibition (> 60 %) of OAT3 compared to the isoflavone metabolites. Quercetin and genistein, non-substrates of OAT3, also showed a high inhibition of this transporter. In contrast to OAT1, many flavonoid metabolites tested showed considerable inhibition of OAT3 even at 2  $\mu\text{M}$ . Quercetin metabolites, quercetin-3-*O*-glucuronide, quercetin-3'-*O*-glucuronide and quercetin-3'-*O*-sulfate showed > 60 % inhibition of OAT3. Isoflavone glucuronides are poor inhibitors, despite being good substrates of OAT3. The  $\text{IC}_{50}$  values of inhibition of OAT3 for quercetin-3-*O*-glucuronide, quercetin-3'-*O*-glucuronide and quercetin-3'-*O*-sulfate was determined to be at 0.43  $\mu\text{M}$ , 1.31  $\mu\text{M}$  and 0.75  $\mu\text{M}$ , respectively (Figure 7). These data imply that physiologically relevant concentrations of quercetin metabolites have a significant impact on OAT3 activity.

### 3.5. Quercetin-3'-*O*-sulfate reduces OAT1-induced cytotoxicity of adefovir

Experiments with 293H control and OAT1-expressing cells demonstrated a marked enhancement of adefovir-induced cytotoxicity with OAT1 over-expression. In OAT1-expressing 293H cells, adefovir was highly cytotoxic with an  $\text{IC}_{50}$  value of 1.5  $\mu\text{M}$ ; while in control 293H cells, the  $\text{IC}_{50}$  value was 124  $\mu\text{M}$  (Figure 8). Quercetin-3'-*O*-sulfate, a good OAT1 substrate, may thus be protective against cytotoxicity of adefovir. To test this hypothesis, we evaluated the effect of co-incubation of adefovir (3 and 10  $\mu\text{M}$ ) with different concentrations of quercetin-3'-*O*-sulfate (2 to 50  $\mu\text{M}$ ) (Figure 8). The results showed that quercetin-3'-*O*-sulfate significantly reduced the cytotoxicity of adefovir dose-dependently at concentrations above 10  $\mu\text{M}$ . Maximal protection was observed at 50  $\mu\text{M}$ , with cell viability

1 increased from 37.3 % to 77.4 % and 16.2 % to 60.1 % for 3 and 10  $\mu$ M adefovir,  
2 respectively. Treatment of quercetin-3'-*O*-sulfate alone at concentrations tested (2 to 50  $\mu$ M)  
3  
4 had no effect on cell viability in OAT1-transfected cells.  
5  
6  
7  
8  
9

#### 10 11 **4. Discussion**

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Flavonoids are extensively metabolized in humans by the phase II enzyme families of sulfotransferases and UDP-glucuronosyltransferases following oral administration.

Conjugates of glucuronide and/or sulfate are the predominant forms in human plasma, while the parent aglycones are usually absent or present in minute amounts [11]. In the present study, we examined transport of nine quercetin and isoflavone metabolites found *in vivo* in OAT1- or OAT3- overexpressing human embryonic kidney 293H cells. The sulfated conjugates, quercetin-3'-*O*-sulfate and genistein-4'-*O*-sulfate, were efficiently transported by OAT1. On the other hand, glucuronide conjugates including daidzein-7-*O*-glucuronide, genistein-7-*O*-glucuronide, glycitein-7-*O*-glucuronide and quercetin-3'-*O*-glucuronide appeared to be preferential substrates for OAT3. Quercetin-3'-*O*-sulfate was also transported by OAT3, but the transport activity was much lower compared with OAT1. Although the aglycones quercetin and genistein were taken up into OAT1 and OAT3-expressing cells, uptake was not significantly different from the control cells. Due to the relative hydrophobicity of the aglycones, their uptake may be attributed to passive diffusion [34]. OAT1 and OAT3 are highly expressed on the basolateral membranes of renal proximal tubules where they mediate the rate-limiting uptake of xenobiotics and their metabolites. The relative specificity and activity of OATs may influence the extent to which flavonoid conjugates are eliminated via the urine. In humans, urinary excretion (20-60 % of an ingested dose) is a major pathway of elimination for the isoflavones [11]. Since isoflavone

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

glucuronides are the main metabolites in humans [10], OAT3 may contribute to the urinary excretion of isoflavones by mediating the uptake of glucuronide conjugates from blood into proximal tubules. Hence, OAT3-mediated transport could be one factor limiting the systemic availability of isoflavones. In contrast, a relatively small proportion (1-7 %) of ingested quercetin is excreted in urine. Although quercetin-3'-*O*-sulfate and quercetin-3'-*O*-glucuronide are good substrates of OAT1 and OAT3, respectively, quercetin-3-*O*-glucuronide and quercetin-7-*O*-glucuronide, the other major quercetin metabolites *in vivo*, were very poor substrates of OAT1 and OAT3. These data suggest that OAT1- and OAT3-mediated uptake into kidney may play an important role in determining disposition of flavonoid conjugates *in vivo*.

For the flavonoid metabolites, it was observed that OAT1 and OAT3 favored the transport of sulfate and glucuronide conjugates, respectively. OATs have been shown to transport various xenobiotics and their conjugated metabolites. In agreement with our data, OAT1 was reported to mediate the transport of sulfated conjugates such as indoxyl sulfate [35], edaravone sulfate [16] and 5-sulfooxymethylfurfural [36]. Transport of more bulky glucuronidated metabolites by OAT1 was not observed [16, 37]. Nonetheless, this study has also identified quercetin-3-*O*-glucuronide and quercetin-7-*O*-glucuronide as potential unique glucuronidated substrates of OAT1. OAT3, on the other hand, has been shown to transport the sulfate or glucuronide conjugates, such as estrone-3-sulfate, edaravone sulfate, edaravone glucuronide [16], and 7-*O*-mycophenolic acid glucuronide [37]. Indeed, all the flavonoid conjugates tested herein showed significantly enhanced uptake in the OAT3-overexpressing cells, suggesting that OAT3 possesses a broader specificity for the flavonoid conjugates compared to OAT1.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Vectorial transport of hydrophilic substances across the renal proximal tubules is achieved through co-operative transport of uptake transporters on the basolateral membrane and efflux transporters on the brush border membrane [38]. In the human kidney, multidrug resistance protein 2 (MRP2/ABCC2) and 4 (MRP4/ABCC4), p-glycoprotein (MDR1/ABCB1), breast cancer resistance protein (BCRP/ABCG2) and OAT4 are expressed on the brush border membrane [39]. Glucuronidated and sulfated conjugates of flavonoids are substrates of apical efflux transporters. Quercetin glucuronides are transported by MRP2 [17] and BCRP [40]; quercetin-3'-O-sulfate is transported by MRP2 [41]. BCRP, and to a lesser extent MRP2, were responsible for the apical efflux of isoflavone glucuronides and sulfates in the intestine [18]. In the human kidney, BCRP and MRP2 may play a similar role in the apical efflux of flavonoid conjugates from proximal tubule cells into urine. The coupled action of influx (OAT1 and OAT3) and efflux transporters (BCRP and MRP2) may result in effective elimination of flavonoid metabolites from the circulation, and contribute to the rapid elimination of flavonoids in humans.

Apart from being transported, the present study also shows that flavonoid conjugates inhibited the transport of model substrates by OAT1 and OAT3. Previous investigations showed that ellagic acid and morin are potent inhibitors of OAT1, with IC<sub>50</sub> values of 0.21 and 0.46 μM, respectively [26, 27]. However, the bioavailability of parent aglycones is often minimal due to extensive phase II metabolism. Herein, we demonstrated that some of the metabolites of flavonoids found *in vivo* retained the capability to inhibit OATs. Among the flavonoid conjugates tested, quercetin-3'-O-sulfate displayed the most effective inhibition of OAT1 (IC<sub>50</sub>: 1.7 μM). Indeed, the sulfation of this flavonoid was found to enhance its inhibitory activity on OAT1. Potent inhibition of OAT3-mediated uptake of 5-carboxyfluorescein was also found for quercetin-3-O-glucuronide, quercetin-7-O-glucuronide

1 and quercetin-3'-*O*-sulfate with IC<sub>50</sub> values of 0.43-1.31 μM. Isoflavone conjugates are  
2 generally weaker inhibitors of OATs. The low IC<sub>50</sub> of inhibition suggests that physiologically  
3 relevant concentrations of quercetin conjugates may have an impact on the OAT-mediated  
4 uptake of drugs. Such a flavonoid-drug interaction may be utilized to offer therapeutic  
5 benefits, for example, to improve the systemic availability of drugs by limiting the OAT1- or  
6 OAT3-mediated renal clearance.  
7  
8  
9  
10  
11  
12  
13

14  
15 Inhibitors of OATs are potential nephroprotective agents against drug-induced renal  
16 injury. The nephrotoxicity of some drugs is caused by the highly concentrative uptake of  
17 OATs, leading to excessive accumulation in the renal tubules and development of renal injury.  
18 OAT1-mediated transport has been implicated in the nephrotoxicity caused by adefovir,  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

OAT1-mediated transport has been implicated in the nephrotoxicity caused by adefovir, cidofovir and cephalosporin antibiotics [42, 43]. It has been suggested that probenecid and non-steroidal anti-inflammatory drugs (NSAIDs), at clinically relevant concentrations, are effective in reducing OAT1-induced nephrotoxicity of drugs [24, 42]. Probenecid and NSAIDs, however, are not without adverse effects. Flavonoids may thus be useful alternatives in the management of OAT1-induced nephrotoxicity of drugs. Adefovir, a substrate of OAT1, was used to investigate the potential of quercetin-3'-*O*-sulfate to reduce OAT1-mediated toxicity of drugs. OAT1-transfected cells were shown to be more sensitive to the cytotoxicity of adefovir compared to control cells [24, 42]. On the other hand, we showed that quercetin-3'-*O*-sulfate ( $\geq 10 \mu\text{M}$ ) effectively reduced the cytotoxicity of adefovir in OAT1-overexpressing cells. This suggests that flavonoids have the potential to limit the severity of drug-induced renal injury via the inhibition of OAT1. Lin et al. [44] also reported that the flavonoid morin protected against imipenem-induced nephrotoxicity, via the inhibition of OAT3-mediated accumulation of this antibiotic. Alternatively, flavonoids may attenuate drug-induced renal damage via free radical scavenging [45, 46] or anti-apoptotic

1 mechanisms [47]. Nevertheless, OAT1 and OAT3 may play an active role in these protective  
2 effects by mediating the entry of flavonoids and their metabolites into the proximal tubule  
3  
4 cells.  
5  
6

7  
8 In conclusion, our results indicate that both OAT1 and OAT3 are likely to be  
9 responsible for uptake of flavonoid sulfates and glucuronides across the basolateral  
10 membrane of the human kidney, respectively. OAT-mediated transport may be one factor  
11 limiting the systemic availability of flavonoids *in vivo*. Our study also showed that quercetin  
12 conjugates are able to inhibit OAT1 and OAT3 at physiologically achievable concentrations,  
13 and quercetin-3'-*O*-sulfate reduces OAT1-mediated cytotoxicity of adefovir, thereby  
14  
15 indicating a potential for flavonoid-drug interactions.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Acknowledgements

We thank the Nestle Research Center for funding.

## 5. References

- [1] Murakami A, Ashida H, Terao J. Multitargeted cancer prevention by quercetin. *Cancer Letters* 2008;269:315-25.
- [2] Tanigawa S, Fujii M, Hou DX. Action of Nrf2 and Keap1 in ARE-mediated NQO1 expression by quercetin. *Free Radic Biol Med* 2007;42:1690-703.
- [3] Kwak MK, Kensler TW. Targeting NRF2 signaling for cancer chemoprevention. *Toxicol Appl Pharmacol* 2010;244:66-76.
- [4] Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT, et al. Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* 1998;139:4252-63.
- [5] Hilakivi-Clarke L, Onojafe I, Raygada M, Cho E, Skaar T, Russo I, et al. Prepubertal exposure to zearalenone or genistein reduces mammary tumorigenesis. *Br J Cancer* 1999;80:1682-8.
- [6] Nestel PJ, Yamashita T, Sasahara T, Pomeroy S, Dart A, Komesaroff P, et al. Soy isoflavones improve systemic arterial compliance but not plasma lipids in menopausal and perimenopausal women. *Arterioscler Thromb Vasc Biol* 1997;17:3392-8.
- [7] Strauss L, Santti R, Saarinen N, Streng T, Joshi S, Makela S. Dietary phytoestrogens and their role in hormonally dependent disease. *Toxicol Lett* 1998;102-103:349-54.
- [8] Williamson G, Barron D, Shimoi K, Terao J. In vitro biological properties of flavonoid conjugates found in vivo. *Free Radic Res* 2005;39:457-69.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [9] Day AJ, Mellon F, Barron D, Sarrazin G, Morgan MRA, Williamson G. Human metabolism of dietary flavonoids: Identification of plasma metabolites of quercetin. *Free Radical Res* 2001;35:941-52.
- [10] Clarke DB, Lloyd AS, Botting NP, Oldfield MF, Needs PW, Wiseman H. Measurement of intact sulfate and glucuronide phytoestrogen conjugates in human urine using isotope dilution liquid chromatography-tandem mass spectrometry with [<sup>13</sup>C(3)]isoflavone internal standards. *Anal Biochem* 2002;309:158-72.
- [11] Manach C, Williamson G, Morand C, Scalbert A, Remesy C. Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am J Clin Nutr* 2005;81:230S-42S.
- [12] Silberberg M, Morand C, Mathevon T, Besson C, Manach C, Scalbert A, et al. The bioavailability of polyphenols is highly governed by the capacity of the intestine and of the liver to secrete conjugated metabolites. *Eur J Nutr* 2006;45:88-96.
- [13] Watanabe S, Yamaguchi M, Sobue T, Takahashi T, Miura T, Arai Y, et al. Pharmacokinetics of soybean isoflavones in plasma, urine and feces of men after ingestion of 60 g baked soybean powder (kinako). *J Nutr* 1998;128:1710-5.
- [14] Goldberg DM, Yan J, Soleas GJ. Absorption of three wine-related polyphenols in three different matrices by healthy subjects. *Clin Biochem* 2003;36:79-87.
- [15] Mullen W, Edwards CA, Crozier A. Absorption, excretion and metabolite profiling of methyl-, glucuronyl-, glucosyl- and sulpho-conjugates of quercetin in human plasma and urine after ingestion of onions. *Brit J Nutr* 2006;96:107-16.
- [16] Mizuno N, Takahashi T, Iwase Y, Kusuhara H, Niwa T, Sugiyama Y. Human organic anion transporters 1 (hOAT1/SLC22A6) and 3 (hOAT3/SLC22A8) transport edaravone (MCI-186; 3-methyl-1-phenyl-2-pyrazolin-5-one) and its sulfate conjugate. *Drug Metab Dispos* 2007;35:1429-34.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [17] Williamson G, Aeberli I, Miguet L, Zhang Z, Sanchez MB, Crespy V, et al. Interaction of positional isomers of quercetin glucuronides with the transporter ABCC2 (cMOAT, MRP2). *Drug Metab Dispos* 2007;35:1262-8.
- [18] Chen J, Lin H, Hu M. Absorption and metabolism of genistein and its five isoflavone analogs in the human intestinal Caco-2 model. *Cancer Chemother Pharmacol* 2005;55:159-69.
- [19] Zhang L, Lin G, Kovacs B, Jani M, Krajcsi P, Zuo Z. Mechanistic study on the intestinal absorption and disposition of baicalein. *Eur J Pharm Sci* 2007;31:221-31.
- [20] Kojima R, Sekine T, Kawachi M, Cha SH, Suzuki Y, Endou H. Immunolocalization of multispecific organic anion transporters, OAT1, OAT2, and OAT3, in rat kidney. *J Am Soc Nephrol* 2002;13:848-57.
- [21] VanWert AL, Gionfriddo MR, Sweet DH. Organic anion transporters: discovery, pharmacology, regulation and roles in pathophysiology. *Biopharm Drug Dispos* 2010;31:1-71.
- [22] Cha SH, Sekine T, Fukushima JI, Kanai Y, Kobayashi Y, Goya T, et al. Identification and characterization of human organic anion transporter 3 expressing predominantly in the kidney. *Mol Pharmacol* 2001;59:1277-86.
- [23] Cihlar T, Lin DC, Pritchard JB, Fuller MD, Mendel DB, Sweet DH. The antiviral nucleotide analogs cidofovir and adefovir are novel substrates for human and rat renal organic anion transporter 1. *Mol Pharmacol* 1999;56:570-80.
- [24] Mulato AS, Ho ES, Cihlar T. Nonsteroidal anti-inflammatory drugs efficiently reduce the transport and cytotoxicity of adefovir mediated by the human renal organic anion transporter 1. *J Pharmacol Exp Ther* 2000;295:10-5.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [25] Dattwyler RJ, Volkman DJ, Conaty SM, Platkin SP, Luft BJ. Amoxicillin plus probenecid versus doxycycline for treatment of erythema migrans borreliosis. *Lancet* 1990;336:1404-6.
- [26] Whitley AC, Sweet DH, Walle T. The dietary polyphenol ellagic acid is a potent inhibitor of hOAT1. *Drug Metab Dispos* 2005;33:1097-100.
- [27] Hong SS, Seo K, Lim SC, Han HK. Interaction characteristics of flavonoids with human organic anion transporter 1 (hOAT1) and 3 (hOAT3). *Pharmacol Res* 2007;56:468-73.
- [28] Day AJ, Bao YP, Morgan MRA, Williamson G. Conjugation position of quercetin glucuronides and effect on biological activity. *Free Radical Bio Med* 2000;29:1234-43.
- [29] Fairley B, Botting NP, Cassidy A. The synthesis of daidzein sulfates. *Tetrahedron* 2003;59:5407-10.
- [30] Boersma MG, van der Woude H, Bogaards J, Boeren S, Vervoort J, Cnubben NH, et al. Regioselectivity of phase II metabolism of luteolin and quercetin by UDP-glucuronosyl transferases. *Chem Res Toxicol* 2002;15:662-70.
- [31] Day AJ, Bao Y, Morgan MR, Williamson G. Conjugation position of quercetin glucuronides and effect on biological activity. *Free Radic Biol Med* 2000;29:1234-43.
- [32] Carmichael J, DeGraff WG, Gazdar AF, Minna JD, Mitchell JB. Evaluation of a tetrazolium-based semiautomated colorimetric assay: assessment of chemosensitivity testing. *Cancer Res* 1987;47:936-42.
- [33] Truong DM, Kaler G, Khandelwal A, Swaan PW, Nigam SK. Multi-level analysis of organic anion transporters 1, 3, and 6 reveals major differences in structural determinants of antiviral discrimination. *J Biol Chem* 2008;283:8654-63.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [34] Day AJ, Gee JM, DuPont MS, Johnson IT, Williamson G. Absorption of quercetin-3-glucoside and quercetin-4'-glucoside in the rat small intestine: the role of lactase phlorizin hydrolase and the sodium-dependent glucose transporter. *Biochem Pharmacol* 2003;65:1199-206.
- [35] Deguchi T, Kusuhara H, Takadate A, Endou H, Otagiri M, Sugiyama Y. Characterization of uremic toxin transport by organic anion transporters in the kidney. *Kidney Int* 2004;65:162-74.
- [36] Bakhiya N, Stephani M, Bahn A, Ugele B, Seidel A, Burckhardt G, et al. Uptake of chemically reactive, DNA-damaging sulfuric acid esters into renal cells by human organic anion transporters. *J Am Soc Nephrol* 2006;17:1414-21.
- [37] Uwai Y, Motohashi H, Tsuji Y, Ueo H, Katsura T, Inui K. Interaction and transport characteristics of mycophenolic acid and its glucuronide via human organic anion transporters hOAT1 and hOAT3. *Biochem Pharmacol* 2007;74:161-8.
- [38] Sekine T, Miyazaki H, Endou H. Molecular physiology of renal organic anion transporters. *Am J Physiol-Renal* 2006;290:F251-F61.
- [39] Leslie EM, Deeley RG, Cole SP. Multidrug resistance proteins: role of P-glycoprotein, MRP1, MRP2, and BCRP (ABCG2) in tissue defense. *Toxicol Appl Pharmacol* 2005;204:216-37.
- [40] Sesink AL, Arts IC, de Boer VC, Breedveld P, Schellens JH, Hollman PC, et al. Breast cancer resistance protein (Bcrp1/Abcg2) limits net intestinal uptake of quercetin in rats by facilitating apical efflux of glucuronides. *Mol Pharmacol* 2005;67:1999-2006.
- [41] O'Leary KA, Day AJ, Needs PW, Mellon FA, O'Brien NM, Williamson G. Metabolism of quercetin-7- and quercetin-3-glucuronides by an in vitro hepatic model: the role of human beta-glucuronidase, sulfotransferase, catechol-O-methyltransferase

and multi-resistant protein 2 (MRP2) in flavonoid metabolism. *Biochem Pharmacol* 2003;65:479-91.

- [42] Ho ES, Lin DC, Mendel DB, Cihlar T. Cytotoxicity of antiviral nucleotides adefovir and cidofovir is induced by the expression of human renal organic anion transporter 1. *J Am Soc Nephrol* 2000;11:383-93.
- [43] Takeda M, Tojo A, Sekine T, Hosoyamada M, Kanai Y, Endou H. Role of organic anion transporter 1 (OAT1) in cephaloridine (CER)-induced nephrotoxicity. *Kidney Int* 1999;56:2128-36.
- [44] Lim SC, Im YB, Bae CS, Han SI, Kim SE, Han HK. Protective effect of morin on the imipenem-induced nephrotoxicity in rabbits. *Arch Pharm Res* 2008;31:1060-5.
- [45] Sung MJ, Kim DH, Jung YJ, Kang KP, Lee AS, Lee S, et al. Genistein protects the kidney from cisplatin-induced injury. *Kidney Int* 2008;74:1538-47.
- [46] Zhong Z, Connor HD, Li X, Mason RP, Forman DT, Lemasters JJ, et al. Reduction of ciclosporin and tacrolimus nephrotoxicity by plant polyphenols. *J Pharm Pharmacol* 2006;58:1533-43.
- [47] Hisamura F, Kojima-Yuasa A, Kennedy DO, Matsui-Yuasa I. Protective effect of green tea extract and tea polyphenols against FK506-induced cytotoxicity in renal cells. *Basic Clin Pharmacol Toxicol* 2006;98:192-6.

**Figure legends:**

1  
2  
3 **Fig. 1.** Uptake of quercetin and its conjugates by OAT1 and OAT3. 293H-control (white bar),  
4  
5 293H-OAT1 (hatched bar) and 293H-OAT3 (black bar) cells were incubated with 25  $\mu\text{M}$   
6  
7 substrate for 30 min. \*,  $p < 0.05$  compared to control.  
8  
9

10  
11 **Fig. 2.** Uptake of genistein and isoflavone conjugates by OAT1 and OAT3. 293H-control  
12  
13 (white bar), 293H-OAT1 (hatched bar) and 293H-OAT3 (black bar) cells were incubated  
14  
15 with 25  $\mu\text{M}$  substrate for 30 min. \*,  $p < 0.05$  compared to control.  
16  
17  
18

19 **Fig. 3.** Effect of probenecid on OAT1 or OAT3-mediated uptake of flavonoid conjugates.  
20  
21 293H-OAT1 (left panel) or 293H-OAT3 (right panel) cells were incubated with 25  $\mu\text{M}$  of  
22  
23 flavonoid conjugates without (hatched bar) or with 1 mM probenecid (black bar) for 10 min.  
24  
25  
26  
27 \*,  $p < 0.05$  compared to vector control (white bar); \*\*  $p < 0.05$  compared to uptake without  
28  
29 probenecid.  
30  
31

32 **Fig. 4.** Inhibition of OAT1-mediated uptake of *p*-aminohippuric acid (PAH) by flavonoid  
33  
34 conjugates. OAT1-expressing cells were incubated for 10 min with 25  $\mu\text{M}$  PAH in the  
35  
36 presence of 2 (grey bar) and 10  $\mu\text{M}$  (white bar) of flavonoid conjugates. \*,  $p < 0.05$  compared  
37  
38 to control (black bar).  
39  
40  
41

42  
43 **Fig. 5.** Concentration-dependent inhibition of OAT1-mediated uptake of *p*-aminohippuric  
44  
45 acid (PAH) by quercetin-3'-*O*-sulfate. OAT1-expressing cells were incubated for 10 min with  
46  
47 25  $\mu\text{M}$  PAH in the presence of 0.1-10  $\mu\text{M}$  of quercetin-3'-*O*-sulfate.  
48  
49  
50

51 **Fig. 6.** Inhibition of OAT3-mediated uptake of 5-carboxyfluorescein (FAM) by flavonoid  
52  
53 conjugates. OAT3-expressing cells were incubated for 10 min with 25  $\mu\text{M}$  FAM in the  
54  
55 presence of 2 (grey bar) and 10  $\mu\text{M}$  (white bar) of flavonoid conjugates. \*,  $p < 0.05$  compared  
56  
57 to control (black bar).  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
**Fig. 7.** Concentration-dependent inhibition of OAT3-mediated uptake of 5-carboxyfluorescein (FAM) by quercetin-3-*O*-glucuronide, quercetin-3'-*O*-glucuronide and quercetin-3'-*O*-sulfate. OAT3-expressing cells were incubated for 10 min with 25  $\mu$ M PAH in the presence of 0.1-10  $\mu$ M of the quercetin conjugates.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
**Fig. 8.** Quercetin-3'-*O*-sulfate attenuated OAT1-induced cytotoxicity of adefovir. (A). The effect of OAT1 expression on the cytotoxicity of adefovir. 293H-control and 293H-OAT1 cells were incubated with various concentrations of adefovir for 96 h; (B). The effect of quercetin-3'-*O*-sulfate (2-50  $\mu$ M) on the cytotoxicity of 3 (white bar) and 10  $\mu$ M (grey bar) adefovir. \*,  $p < 0.05$  compared to control.

**Table 1.** Kinetics of OAT1 and OAT3-mediated uptake of selected flavonoid conjugates

Substrate	OAT1			OAT3		
	$K_m$ ( $\mu\text{M}$ )	$V_{\text{max}}$ (pmol/min per mg)	CL ( $\mu\text{L}/\text{min}$ per mg)	$K_m$ ( $\mu\text{M}$ )	$V_{\text{max}}$ (pmol/min per mg)	CL ( $\mu\text{L}/\text{min}$ per mg)
Daidzein-7- <i>O</i> -glucuronide				$19.1 \pm 1.9$	$420 \pm 15$	22.0
Genistein-4'- <i>O</i> -sulfate	$5.07 \pm 1.81$	$28.3 \pm 2.9$	5.58			
Genistein-7- <i>O</i> -glucuronide				$7.94 \pm 1.42$	$210 \pm 10$	26.4
Glycitein-7- <i>O</i> -glucuronide				$17.6 \pm 3.9$	$171 \pm 13$	9.72
Quercetin-3'- <i>O</i> -sulfate	$1.73 \pm 0.38$	$105 \pm 7$	60.7			
Quercetin-3'- <i>O</i> -glucuronide				$5.25 \pm 0.86$	$73.0 \pm 3.5$	13.9

Figure 1  
[Click here to download high resolution image](#)




Figure 2  
[Click here to download high resolution image](#)




Figure 3  
[Click here to download high resolution image](#)




Figure 4  
[Click here to download high resolution image](#)




Figure 5  
[Click here to download high resolution image](#)





Figure 6  
[Click here to download high resolution image](#)






Figure 8  
[Click here to download high resolution image](#)

