



**HAL**  
open science

**catena-Poly[bis(dibenzylammonium)  
[[dichloridomercurate(II)]-[mu]-sulfato-[kappa]2O:O']]**

Mouhamadou Sembene Boye, Aminata Diasse-Sarr, Arnaud Grosjean,  
Philippe Guionneau

► **To cite this version:**

Mouhamadou Sembene Boye, Aminata Diasse-Sarr, Arnaud Grosjean, Philippe Guionneau. catena-Poly[bis(dibenzylammonium) [[dichloridomercurate(II)]-[mu]-sulfato-[kappa]2O:O']]. Acta Crystallographica Section E: Structure Reports Online [2001-2014], 2012, 68 (3), pp.m340. 10.1107/S1600536812006927 . hal-00679526

**HAL Id: hal-00679526**

**<https://hal.science/hal-00679526>**

Submitted on 26 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## catena-Poly[bis(dibenzylammonium) [[dichloridomercurate(II)]- $\mu$ -sulfato- $\kappa^2$ O:O']]

Mouhamadou Sembene Boye,<sup>a\*</sup> Aminata Diasse-Sarr,<sup>a</sup> Arnaud Grosjean<sup>b</sup> and Philippe Guionneau<sup>b</sup>

<sup>a</sup>Département de Chimie, Faculté des Sciences et Techniques, Université Cheikh Anta Diop, Dakar, Senegal, and <sup>b</sup>CNRS, Université de Bordeaux, ICMCB, 87 Avenue du Docteur A. Schweitzer, F-33608 Pessac, France  
Correspondence e-mail: mouhasboye@hotmail.com


Received 5 January 2012; accepted 15 February 2012

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.017$  Å;  $R$  factor = 0.046;  $wR$  factor = 0.121; data-to-parameter ratio = 17.3.

The structure of the title compound,  $(\text{C}_{14}\text{H}_{16}\text{N})_2[\text{HgCl}_2(\text{SO}_4)]$ , consists of an infinite chain propagating along the  $c$  direction, containing  $\text{Hg}^{\text{II}}$  ions tetraordinated by two bridging O atoms of bis-monodentate sulfate anions and two chloride ligands. In the the crystal,  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonding between the cations and the anionic chains consolidates the packing. The crystal structure was determined from an inversion twin with approximately equal twin domains.

### Related literature

For the behavior of sulfate as a ligand, see: Sall *et al.* (1992); Diop *et al.* (2000); Boye *et al.* (2007). For the IR vibrational frequencies of sulfate, see: Nakamoto (1978).



### Experimental

#### Crystal data

$(\text{C}_{14}\text{H}_{16}\text{N})_2[\text{HgCl}_2(\text{SO}_4)]$   
 $M_r = 764.14$

Monoclinic,  $Cc$   
 $a = 22.8275$  (5) Å

$b = 12.9547$  (3) Å  
 $c = 10.1512$  (3) Å  
 $\beta = 92.095$  (2)°  
 $V = 2999.94$  (13) Å<sup>3</sup>  
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 5.41$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.40 \times 0.25 \times 0.25$  mm

#### Data collection

Nonius Kappa CCD diffractometer  
Absorption correction: empirical (using intensity measurements) (*SCALEPACK*; Otwinowski & Minor, 1997)  
 $T_{\min} = 0.221$ ,  $T_{\max} = 0.345$

9151 measured reflections  
5464 independent reflections  
5304 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.020$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$   
 $wR(F^2) = 0.121$   
 $S = 1.02$   
5464 reflections  
315 parameters

2 restraints  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 1.12$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -2.40$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O1}^{\text{i}}$	0.90	2.44	2.920 (9)	114
$\text{N1}-\text{H1A}\cdots\text{O3}^{\text{i}}$	0.90	2.29	3.037 (10)	141
$\text{N1}-\text{H1B}\cdots\text{O3}^{\text{ii}}$	0.90	1.90	2.766 (9)	161
$\text{N2}-\text{H2C}\cdots\text{O4}^{\text{iii}}$	0.90	2.32	3.043 (10)	137
$\text{N2}-\text{H2C}\cdots\text{O1}^{\text{iii}}$	0.90	2.12	2.857 (9)	139

Symmetry codes: (i)  $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (ii)  $x + \frac{1}{2}, y - \frac{1}{2}, z$ ; (iii)  $x, -y + 1, z + \frac{1}{2}$ .

Data collection: *COLLECT* (Nonius, 2003); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS86* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FI2124).

### References

- Boye, M. S., Diassé-Sarr, A., Diop, L., Russo, U., Biesemans, M. & Willem, R. (2007). *C. R. Chim.* **10**, 466–468.  
Diop, C. A. K., Kochikpa, Y. A. O., Diop, L. & Mahieu, B. (2000). *Main Group Met. Chem.* **23**, 9, 493–495.  
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
Nakamoto, K. (1978). *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, 3rd ed., p. 142. New York: John Wiley & Sons.  
Nonius (2003). *COLLECT*. Nonius BV, Delft, The Netherlands.  
Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography, Part A*, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.  
Sall, A. S., Diassé, A., Sarr, O. & Diop, L. (1992). *Main Group Met. Chem.* **15**, 265–268.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supplementary materials

*Acta Cryst.* (2012). E68, m340 [doi:10.1107/S1600536812006927]

**catena-Poly[bis(dibenzylammonium) [[dichloridomercurate(II)]- $\mu$ -sulfato- $\kappa^2$ O:O']]**

**Mouhamadou Sembene Boye, Aminata Diasse-Sarr, Arnaud Grosjean and Philippe Guionneau**

**Comment**

In the framework of our research work for understanding the behavior of sulfate acting as ligand (Sall *et al.*, 1992; Diop *et al.*, 2000; Boye *et al.*, 2007), we report the crystallographic study of  $2[(C_6H_5CH_2)_2NH_2]^+[HgSO_4Cl_2]^{2-}$ .

The structure obtained by single-crystal XRD (Fig. 1) indicate an infinite chain in which each Hg atom is tetracoordinated by two O atoms of two sulfates and two chloride atoms in a distorted tetrahedral geometry. The tetrahedral angles are in the range 80.8 (2)–152.42 (10). The sulfate behaves as a bidentate anion with disparate Hg—O distances [Hg—O(1) = 2.433 (6) and Hg—O(4) = 2.533 (7) Å]. The S—O distances vary from 1.446 (6) to 1.492 (7) Å, the S—O distances for two O atoms linked to Hg atoms [1.474 (7)–1.492 (5) Å] are longer than those non-bonding [1.446 (6)–1.465 (6) Å].


The behavior of bidentate sulfate ( $C_{2v}$  symmetry) is confirmed by the infrared data,  $\nu_s(SO_4^{2-})$  appears at 988  $cm^{-1}$  (Nakamoto, 1978) and  $\nu_{as}(SO_4^{2-})$  splits into three bands (1115, 1082, 1041  $cm^{-1}$ ). The crystal packing of the title compound is shown in Fig.2.

**Experimental**

$(C_6H_5CH_2)_2NH$ ,  $H_2SO_4$  and  $HgCl_2$  (Aldrich chemicals) were used without further purification.

The title compound was obtained by mixing ethanolic solutions of  $(C_6H_5CH_2)_2NH$  (17.66 mmol),  $H_2SO_4$  (8.83 mmol) and  $HgCl_2$  (4.41 mmol) in a 8–4–1 ratio. The mixture was stirred for around two hours at room temperature. Suitable crystals for X-ray diffraction were obtained after slow solvent evaporation. (m.p. 459 K).

The title compound was isolated according to the following reaction:



- **Infrared data ( $cm^{-1}$ )** [ $\nu_s$  = very strong; s = strong]

988 s  $\nu_s(SO_4^{2-})$ ; 1115 s, 1082 s, 1041 s  $\nu_{as}(SO_4^{2-})$ ; 454 s  $\delta_s(SO_4^{2-})$ ; 610 vs  $\delta_{as}(SO_4^{2-})$ .

**Refinement**


Inversion twin matrix instruction was used during refinement. The twin components were 0.498 (9) and 0.502 (9), respectively.

All H atoms were placed in geometrically calculated positions (C—H = 0.93 Å for phenyl H and 0.97 Å for methylene H, N—H = 0.90 Å) and refined using a riding model with  $U_{iso}(H) = 1.2U_{eq}$  of the respective carrier atom.


**Computing details**

Data collection: *COLLECT* (Nonius, 2003); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS86* (Sheldrick, 2008); program(s)


used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *publCIF* (Westrip, 2010).



**Figure 1**  
Molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.



**Figure 2**  
Molecular packing viewed along the *c* axis. H atoms have been omitted for clarity.


**Figure 3**

Chains propagate along the *c* direction. H atoms have been omitted for clarity.

**catena-Poly[bis(dibenzylammonium) [[dichloridomercurate(II)]- $\mu$ -sulfato- $\kappa^2$ O:O']**
*Crystal data*

(C<sub>14</sub>H<sub>16</sub>N)<sub>2</sub>[HgCl<sub>2</sub>(SO<sub>4</sub>)]

*M<sub>r</sub>* = 764.14

Monoclinic, *Cc*

Hall symbol: C-2yc

*a* = 22.8275 (5) Å

*b* = 12.9547 (3) Å

*c* = 10.1512 (3) Å

$\beta$  = 92.095 (2)°

*V* = 2999.94 (13) Å<sup>3</sup>

*Z* = 4

*F*(000) = 1504

*D<sub>x</sub>* = 1.692 Mg m<sup>-3</sup>

Mo *K* $\alpha$  radiation,  $\lambda$  = 0.71073 Å

Cell parameters from 10249 reflections

$\theta$  = 0.4–26.0°

$\mu$  = 5.41 mm<sup>-1</sup>

*T* = 293 K

Prism, colourless

0.40 × 0.25 × 0.25 mm

*Data collection*

Nonius Kappa CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: empirical (using intensity measurements)

(*SCALEPACK*; Otwinowski & Minor, 1997)

*T<sub>min</sub>* = 0.221, *T<sub>max</sub>* = 0.345

9151 measured reflections

5464 independent reflections

5304 reflections with *I* > 2 $\sigma$ (*I*)

*R<sub>int</sub>* = 0.020

$\theta_{\max}$  = 26.0°,  $\theta_{\min}$  = 2.7°

*h* = -27→28

*k* = -15→15

*l* = -12→12

*Refinement*

Refinement on *F*<sup>2</sup>

Least-squares matrix: full

*R*[*F*<sup>2</sup> > 2 $\sigma$ (*F*<sup>2</sup>)] = 0.046

*wR*(*F*<sup>2</sup>) = 0.121

*S* = 1.02

5464 reflections

315 parameters

2 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.099P)^2 + 6.3836P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.005$$

$$\Delta\rho_{\max} = 1.12 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -2.40 \text{ e } \text{\AA}^{-3}$$

Extinction correction: *SHELXL97* (Sheldrick, 2008),  $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0063 (3)

### Special details

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.47859 (7)	0.47550 (14)	0.25406 (15)	0.0294 (3)
O1	0.4797 (2)	0.4792 (5)	0.1074 (4)	0.0377 (11)
O3	0.4174 (2)	0.4818 (5)	0.2924 (6)	0.0427 (12)
O2	0.5059 (3)	0.3804 (5)	0.2988 (6)	0.0460 (13)
N1	0.8594 (3)	-0.0322 (6)	0.5261 (7)	0.0454 (15)
H1A	0.8729	0.0132	0.5870	0.054*
H1B	0.8851	-0.0336	0.4610	0.054*
C2	0.8012 (4)	0.0059 (8)	0.4703 (8)	0.0452 (18)
H2A	0.7860	-0.0443	0.4070	0.054*
H2B	0.8076	0.0698	0.4231	0.054*
C4	0.7555 (4)	0.0246 (7)	0.5720 (8)	0.0428 (17)
C6	0.7115 (5)	-0.0450 (9)	0.5881 (11)	0.057 (2)
H6	0.7110	-0.1064	0.5406	0.068*
C5	0.7568 (5)	0.1163 (8)	0.6417 (9)	0.051 (2)
H5	0.7867	0.1639	0.6307	0.061*
C7	0.7123 (5)	0.1359 (8)	0.7291 (11)	0.057 (2)
H7	0.7122	0.1980	0.7750	0.068*
C8	0.6683 (5)	0.0643 (9)	0.7486 (12)	0.063 (3)
H8	0.6400	0.0764	0.8102	0.075*
C11	0.5329 (10)	0.1734 (10)	0.580 (2)	0.0868 (16)
C17	0.6597 (4)	0.4041 (7)	0.4804 (10)	0.0476 (19)
C16	0.6563 (4)	0.3853 (10)	0.3472 (10)	0.059 (3)
H16	0.6204	0.3923	0.3015	0.071*
C13	0.7640 (4)	0.3654 (9)	0.4791 (12)	0.055 (2)
H13	0.8000	0.3616	0.5244	0.067*
C14	0.7590 (5)	0.3438 (12)	0.3478 (13)	0.067 (3)
H14	0.7915	0.3210	0.3035	0.081*
C15	0.7054 (5)	0.3559 (16)	0.2797 (13)	0.086 (4)
H15	0.7024	0.3443	0.1892	0.103*
C12	0.7138 (4)	0.3934 (8)	0.5433 (10)	0.050 (2)

H12	0.7168	0.4055	0.6336	0.060*
C18	0.6675 (5)	-0.0243 (10)	0.6755 (13)	0.067 (3)
H18	0.6371	-0.0713	0.6843	0.081*
C29	0.8848 (4)	-0.2461 (7)	0.3960 (9)	0.0486 (19)
H29	0.9186	-0.2067	0.3883	0.058*
C30	0.8736 (8)	-0.3307 (8)	0.3096 (13)	0.069 (4)
H30	0.9000	-0.3463	0.2449	0.083*
C31	0.8241 (7)	-0.3898 (10)	0.3207 (14)	0.074 (3)
H31	0.8170	-0.4461	0.2656	0.088*
C33	0.7948 (5)	-0.2806 (11)	0.4996 (12)	0.064 (3)
H33	0.7671	-0.2645	0.5613	0.076*
C32	0.7861 (6)	-0.3641 (9)	0.4130 (17)	0.078 (4)
H32	0.7524	-0.4037	0.4201	0.094*
C34	0.8456 (4)	-0.2227 (7)	0.4910 (7)	0.0407 (16)
C35	0.8576 (5)	-0.1388 (8)	0.5890 (9)	0.051 (2)
H35A	0.8949	-0.1522	0.6348	0.061*
H35B	0.8274	-0.1397	0.6538	0.061*
N2	0.5594 (3)	0.3587 (6)	0.5478 (7)	0.0403 (14)
H2C	0.5291	0.3831	0.5927	0.048*
H2D	0.5470	0.3490	0.4636	0.048*
C37	0.6092 (4)	0.4390 (7)	0.5523 (10)	0.0467 (19)
H37A	0.5948	0.5033	0.5144	0.056*
H37B	0.6212	0.4521	0.6434	0.056*
C40	0.4857 (9)	0.1670 (12)	0.6625 (18)	0.0868 (16)
H40	0.4829	0.2140	0.7312	0.104*
C42	0.5402 (8)	0.1009 (11)	0.4852 (15)	0.0868 (16)
H42	0.5723	0.1030	0.4312	0.104*
C38	0.4515 (8)	0.0193 (11)	0.5571 (16)	0.0868 (16)
H38	0.4247	-0.0347	0.5558	0.104*
C39	0.4449 (8)	0.0952 (12)	0.6454 (15)	0.0868 (16)
H39	0.4114	0.0973	0.6948	0.104*
C41	0.4946 (8)	0.0166 (11)	0.4705 (14)	0.0868 (16)
H41	0.4963	-0.0340	0.4057	0.104*
O4	0.5118 (3)	0.5651 (6)	0.3063 (7)	0.0540 (17)
C3	0.5785 (5)	0.2590 (8)	0.6051 (10)	0.052 (2)
H3A	0.5854	0.2670	0.6993	0.063*
H3B	0.6152	0.2387	0.5673	0.063*
Cl1	0.40170 (12)	0.2706 (3)	-0.0297 (3)	0.0596 (7)
Cl2	0.59979 (13)	0.2769 (3)	0.0059 (4)	0.0672 (8)
Hg1	0.50000 (3)	0.315360 (17)	0.00102 (5)	0.04119 (15)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0266 (7)	0.0391 (8)	0.0226 (7)	-0.0018 (6)	0.0036 (6)	0.0009 (6)
O1	0.041 (3)	0.054 (3)	0.018 (2)	-0.003 (2)	0.0051 (19)	-0.002 (2)
O3	0.033 (3)	0.055 (3)	0.041 (3)	-0.003 (2)	0.011 (2)	0.005 (2)
O2	0.048 (3)	0.042 (3)	0.048 (3)	0.001 (2)	-0.001 (2)	0.009 (2)
N1	0.045 (4)	0.054 (4)	0.036 (3)	-0.011 (3)	0.000 (3)	-0.004 (3)
C2	0.040 (4)	0.059 (5)	0.036 (4)	-0.006 (4)	0.001 (3)	-0.001 (4)

C4	0.044 (4)	0.052 (5)	0.032 (3)	-0.001 (3)	-0.006 (3)	0.005 (3)
C6	0.048 (5)	0.066 (6)	0.057 (5)	-0.016 (4)	0.005 (4)	-0.009 (5)
C5	0.059 (5)	0.048 (5)	0.045 (4)	0.001 (4)	-0.010 (4)	0.003 (4)
C7	0.057 (5)	0.056 (5)	0.056 (5)	0.025 (4)	-0.007 (4)	-0.009 (4)
C8	0.052 (5)	0.075 (7)	0.061 (6)	0.012 (5)	-0.002 (4)	0.004 (5)
C11	0.109 (5)	0.068 (3)	0.082 (4)	-0.007 (3)	-0.009 (3)	0.008 (3)
C17	0.044 (4)	0.046 (5)	0.053 (5)	0.007 (4)	0.010 (4)	0.007 (4)
C16	0.036 (4)	0.093 (8)	0.048 (5)	0.013 (4)	-0.004 (4)	0.010 (5)
C13	0.039 (4)	0.063 (6)	0.064 (6)	0.007 (4)	0.002 (4)	-0.002 (5)
C14	0.041 (5)	0.098 (8)	0.065 (7)	0.015 (5)	0.018 (5)	-0.004 (6)
C15	0.043 (5)	0.162 (13)	0.053 (6)	0.023 (7)	0.012 (5)	-0.015 (8)
C12	0.038 (4)	0.058 (5)	0.052 (5)	0.002 (4)	-0.003 (4)	0.004 (4)
C18	0.049 (5)	0.070 (7)	0.084 (8)	0.000 (5)	0.023 (5)	-0.013 (6)
C29	0.055 (5)	0.038 (4)	0.054 (5)	-0.007 (4)	0.013 (4)	0.003 (3)
C30	0.116 (11)	0.040 (5)	0.051 (6)	0.003 (6)	0.010 (6)	0.001 (4)
C31	0.094 (9)	0.056 (6)	0.069 (7)	-0.008 (6)	-0.017 (7)	-0.006 (6)
C33	0.063 (6)	0.069 (7)	0.060 (6)	-0.018 (5)	0.016 (5)	0.013 (6)
C32	0.067 (7)	0.045 (6)	0.121 (12)	-0.028 (5)	-0.010 (7)	0.006 (7)
C34	0.048 (4)	0.042 (4)	0.032 (4)	-0.011 (3)	-0.002 (3)	0.004 (3)
C35	0.065 (6)	0.054 (5)	0.033 (4)	-0.006 (4)	-0.006 (4)	-0.006 (4)
N2	0.032 (3)	0.047 (4)	0.042 (4)	0.006 (3)	0.009 (3)	0.001 (3)
C37	0.042 (4)	0.041 (4)	0.058 (5)	0.009 (3)	0.004 (4)	0.000 (4)
C40	0.109 (5)	0.068 (3)	0.082 (4)	-0.007 (3)	-0.009 (3)	0.008 (3)
C42	0.109 (5)	0.068 (3)	0.082 (4)	-0.007 (3)	-0.009 (3)	0.008 (3)
C38	0.109 (5)	0.068 (3)	0.082 (4)	-0.007 (3)	-0.009 (3)	0.008 (3)
C39	0.109 (5)	0.068 (3)	0.082 (4)	-0.007 (3)	-0.009 (3)	0.008 (3)
C41	0.109 (5)	0.068 (3)	0.082 (4)	-0.007 (3)	-0.009 (3)	0.008 (3)
O4	0.046 (3)	0.078 (5)	0.038 (3)	-0.025 (3)	0.016 (3)	-0.022 (3)
C3	0.048 (5)	0.054 (5)	0.054 (5)	-0.002 (4)	-0.008 (4)	0.016 (4)
Cl1	0.0447 (13)	0.0551 (16)	0.0792 (18)	-0.0145 (12)	0.0057 (12)	0.0028 (13)
Cl2	0.0498 (15)	0.075 (2)	0.0761 (19)	0.0215 (15)	-0.0071 (13)	-0.0090 (16)
Hg1	0.04012 (18)	0.04296 (19)	0.04054 (18)	0.00013 (15)	0.00241 (10)	-0.00068 (16)

*Geometric parameters (Å, °)*

S1—O2	1.446 (6)	C18—H18	0.9300
S1—O3	1.465 (5)	C29—C34	1.374 (13)
S1—O4	1.473 (7)	C29—C30	1.422 (15)
S1—O1	1.491 (5)	C29—H29	0.9300
O1—Hg1	2.433 (6)	C30—C31	1.37 (2)
N1—C2	1.508 (12)	C30—H30	0.9300
N1—C35	1.523 (12)	C31—C32	1.34 (2)
N1—H1A	0.9000	C31—H31	0.9300
N1—H1B	0.9000	C33—C34	1.386 (13)
C2—C4	1.514 (12)	C33—C32	1.40 (2)
C2—H2A	0.9700	C33—H33	0.9300
C2—H2B	0.9700	C32—H32	0.9300
C4—C6	1.364 (13)	C34—C35	1.492 (12)
C4—C5	1.382 (13)	C35—H35A	0.9700
C6—C18	1.390 (14)	C35—H35B	0.9700



C6—H6	0.9300	N2—C3	1.477 (12)
C5—C7	1.396 (14)	N2—C37	1.539 (11)
C5—H5	0.9300	N2—H2C	0.9000
C7—C8	1.385 (17)	N2—H2D	0.9000
C7—H7	0.9300	C37—H37A	0.9700
C8—C18	1.367 (17)	C37—H37B	0.9700
C8—H8	0.9300	C40—C39	1.32 (2)
C11—C42	1.36 (2)	C40—H40	0.9300
C11—C40	1.39 (3)	C42—C41	1.51 (2)
C11—C3	1.536 (19)	C42—H42	0.9300
C17—C16	1.374 (15)	C38—C41	1.34 (2)
C17—C12	1.375 (13)	C38—C39	1.34 (2)
C17—C37	1.459 (12)	C38—H38	0.9300
C16—C15	1.388 (14)	C39—H39	0.9300
C16—H16	0.9300	C41—H41	0.9300
C13—C14	1.363 (18)	O4—Hg1 <sup>i</sup>	2.533 (7)
C13—C12	1.387 (14)	C3—H3A	0.9700
C13—H13	0.9300	C3—H3B	0.9700
C14—C15	1.392 (18)	Cl1—Hg1	2.327 (3)
C14—H14	0.9300	Cl2—Hg1	2.331 (3)
C15—H15	0.9300	Hg1—O4 <sup>ii</sup>	2.533 (7)
C12—H12	0.9300		
O2—S1—O3	111.6 (4)	C31—C30—C29	120.6 (13)
O2—S1—O4	110.3 (4)	C31—C30—H30	119.7
O3—S1—O4	110.0 (4)	C29—C30—H30	119.7
O2—S1—O1	108.5 (4)	C32—C31—C30	118.3 (12)
O3—S1—O1	108.3 (3)	C32—C31—H31	120.8
O4—S1—O1	107.9 (4)	C30—C31—H31	120.8
S1—O1—Hg1	115.2 (3)	C34—C33—C32	118.5 (11)
C2—N1—C35	114.7 (7)	C34—C33—H33	120.7
C2—N1—H1A	108.6	C32—C33—H33	120.7
C35—N1—H1A	108.6	C31—C32—C33	123.2 (11)
C2—N1—H1B	108.6	C31—C32—H32	118.4
C35—N1—H1B	108.6	C33—C32—H32	118.4
H1A—N1—H1B	107.6	C29—C34—C33	119.5 (9)
N1—C2—C4	114.5 (7)	C29—C34—C35	121.3 (8)
N1—C2—H2A	108.6	C33—C34—C35	119.1 (9)
C4—C2—H2A	108.6	C34—C35—N1	112.8 (7)
N1—C2—H2B	108.6	C34—C35—H35A	109.0
C4—C2—H2B	108.6	N1—C35—H35A	109.0
H2A—C2—H2B	107.6	C34—C35—H35B	109.0
C6—C4—C5	120.5 (9)	N1—C35—H35B	109.0
C6—C4—C2	120.3 (9)	H35A—C35—H35B	107.8
C5—C4—C2	119.0 (8)	C3—N2—C37	111.9 (7)
C4—C6—C18	120.1 (10)	C3—N2—H2C	109.2
C4—C6—H6	120.0	C37—N2—H2C	109.2
C18—C6—H6	120.0	C3—N2—H2D	109.2
C4—C5—C7	118.7 (9)	C37—N2—H2D	109.2

C4—C5—H5	120.7	H2C—N2—H2D	107.9
C7—C5—H5	120.7	C17—C37—N2	111.8 (8)
C8—C7—C5	121.1 (10)	C17—C37—H37A	109.2
C8—C7—H7	119.4	N2—C37—H37A	109.2
C5—C7—H7	119.4	C17—C37—H37B	109.2
C18—C8—C7	118.7 (11)	N2—C37—H37B	109.2
C18—C8—H8	120.7	H37A—C37—H37B	107.9
C7—C8—H8	120.7	C39—C40—C11	121.3 (18)
C42—C11—C40	120.2 (16)	C39—C40—H40	119.4
C42—C11—C3	121.2 (18)	C11—C40—H40	119.4
C40—C11—C3	118.4 (15)	C11—C42—C41	117.8 (17)
C16—C17—C12	117.2 (8)	C11—C42—H42	121.1
C16—C17—C37	122.0 (9)	C41—C42—H42	121.1
C12—C17—C37	120.7 (9)	C41—C38—C39	124.0 (15)
C17—C16—C15	121.1 (9)	C41—C38—H38	118.0
C17—C16—H16	119.5	C39—C38—H38	118.0
C15—C16—H16	119.5	C40—C39—C38	120.3 (19)
C14—C13—C12	118.2 (10)	C40—C39—H39	119.8
C14—C13—H13	120.9	C38—C39—H39	119.8
C12—C13—H13	120.9	C38—C41—C42	115.8 (14)
C13—C14—C15	120.2 (10)	C38—C41—H41	122.1
C13—C14—H14	119.9	C42—C41—H41	122.1
C15—C14—H14	119.9	S1—O4—Hg1 <sup>i</sup>	134.0 (4)
C16—C15—C14	119.8 (11)	N2—C3—C11	112.1 (10)
C16—C15—H15	120.1	N2—C3—H3A	109.2
C14—C15—H15	120.1	C11—C3—H3A	109.2
C17—C12—C13	123.4 (10)	N2—C3—H3B	109.2
C17—C12—H12	118.3	C11—C3—H3B	109.2
C13—C12—H12	118.3	H3A—C3—H3B	107.9
C8—C18—C6	120.8 (11)	C11—Hg1—C12	152.43 (10)
C8—C18—H18	119.6	C11—Hg1—O1	94.47 (15)
C6—C18—H18	119.6	C12—Hg1—O1	112.27 (16)
C34—C29—C30	119.7 (10)	C11—Hg1—O4 <sup>ii</sup>	100.19 (18)
C34—C29—H29	120.2	C12—Hg1—O4 <sup>ii</sup>	90.91 (17)
C30—C29—H29	120.2	O1—Hg1—O4 <sup>ii</sup>	80.8 (2)
O2—S1—O1—Hg1	15.7 (4)	C30—C29—C34—C35	176.0 (10)
O3—S1—O1—Hg1	-105.6 (4)	C32—C33—C34—C29	2.1 (17)
O4—S1—O1—Hg1	135.3 (4)	C32—C33—C34—C35	-175.3 (11)
C35—N1—C2—C4	65.4 (10)	C29—C34—C35—N1	65.7 (12)
N1—C2—C4—C6	-102.6 (10)	C33—C34—C35—N1	-116.9 (10)
N1—C2—C4—C5	81.8 (10)	C2—N1—C35—C34	68.1 (10)
C5—C4—C6—C18	-0.2 (16)	C16—C17—C37—N2	62.6 (13)
C2—C4—C6—C18	-175.8 (10)	C12—C17—C37—N2	-120.4 (10)
C6—C4—C5—C7	0.2 (14)	C3—N2—C37—C17	59.3 (10)
C2—C4—C5—C7	175.9 (8)	C42—C11—C40—C39	-5 (3)
C4—C5—C7—C8	1.5 (14)	C3—C11—C40—C39	179.8 (14)
C5—C7—C8—C18	-3.3 (16)	C40—C11—C42—C41	2 (2)
C12—C17—C16—C15	0.5 (19)	C3—C11—C42—C41	177.3 (13)

C37—C17—C16—C15	177.5 (13)	C11—C40—C39—C38	8 (2)
C12—C13—C14—C15	4 (2)	C41—C38—C39—C40	-8 (3)
C17—C16—C15—C14	1 (3)	C39—C38—C41—C42	5 (2)
C13—C14—C15—C16	-3 (3)	C11—C42—C41—C38	-2 (2)
C16—C17—C12—C13	0.3 (16)	O2—S1—O4—Hg1 <sup>i</sup>	-97.7 (6)
C37—C17—C12—C13	-176.8 (9)	O3—S1—O4—Hg1 <sup>i</sup>	25.8 (7)
C14—C13—C12—C17	-2.5 (17)	O1—S1—O4—Hg1 <sup>i</sup>	143.8 (5)
C7—C8—C18—C6	3.3 (18)	C37—N2—C3—C11	-170.1 (11)
C4—C6—C18—C8	-1.6 (19)	C42—C11—C3—N2	102.8 (16)
C34—C29—C30—C31	-0.5 (18)	C40—C11—C3—N2	-82.1 (16)
C29—C30—C31—C32	1 (2)	S1—O1—Hg1—C11	92.6 (3)
C30—C31—C32—C33	-1 (2)	S1—O1—Hg1—C12	-80.6 (3)
C34—C33—C32—C31	-1 (2)	S1—O1—Hg1—O4 <sup>ii</sup>	-167.8 (4)
C30—C29—C34—C33	-1.3 (15)		

Symmetry codes: (i)  $x, -y+1, z+1/2$ ; (ii)  $x, -y+1, z-1/2$ .

*Hydrogen-bond geometry* ( $\text{\AA}, ^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A $\cdots$ O1 <sup>iii</sup>	0.90	2.44	2.920 (9)	114
N1—H1A $\cdots$ O3 <sup>iii</sup>	0.90	2.29	3.037 (10)	141
N1—H1B $\cdots$ O3 <sup>iv</sup>	0.90	1.90	2.766 (9)	161
N2—H2C $\cdots$ O4 <sup>i</sup>	0.90	2.32	3.043 (10)	137
N2—H2C $\cdots$ O1 <sup>i</sup>	0.90	2.12	2.857 (9)	139
C37—H37A $\cdots$ Cl2 <sup>i</sup>	0.97	2.85	3.716 (10)	149

Symmetry codes: (i)  $x, -y+1, z+1/2$ ; (iii)  $x+1/2, -y+1/2, z+1/2$ ; (iv)  $x+1/2, y-1/2, z$ .