



**HAL**  
open science

## Windings of planar stable processes

Ron A. Doney, Stavros Vakeroudis

► **To cite this version:**

| Ron A. Doney, Stavros Vakeroudis. Windings of planar stable processes. 2012. hal-00679710v3

**HAL Id: hal-00679710**

**<https://hal.science/hal-00679710v3>**

Preprint submitted on 22 Dec 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Windings of planar stable processes

R.A. Doney<sup>\*</sup> and S. Vakeroudis<sup>\*†</sup>

December 22, 2012

## Abstract

Using a generalization of the skew-product representation of planar Brownian motion and the analogue of Spitzer's celebrated asymptotic Theorem for stable processes due to Bertoin and Werner, for which we provide a new easy proof, we obtain some limit Theorems for the exit time from a cone of stable processes of index  $\alpha \in (0, 2)$ . We also study the case  $t \rightarrow 0$  and we prove some Laws of the Iterated Logarithm (LIL) for the (well-defined) winding process associated to our planar stable process.

**AMS 2010 subject classification:** Primary: 60G52, 60G51, 60F05, 60J65; secondary: 60E07, 60B12, 60G18.

**Key words:** Stable processes, Lévy processes, Brownian motion, windings, exit time from a cone, Spitzer's Theorem, skew-product representation, Lamperti's relation, Law of the Iterated Logarithm (LIL) for small times.

## 1 Introduction

In this paper, we study the windings of planar isotropic stable processes. More precisely, having as a starting point a work of Bertoin and Werner [BeW96] concerning this subject (following their previous work on windings of planar Brownian motion<sup>§</sup> [BeW94a]) and motivated by some works of Shi [Shi98], we attempt to generalize some results obtained recently for the case of planar Brownian motion (see e.g. [Vak11, Vakth11, VaY12] and the references therein). In particular, we are interested in the behaviour of stable processes for small time, an aspect which has already been investigated e.g. by Doney [Don04]

---

<sup>\*</sup>Probability and Statistics Group, School of Mathematics, University of Manchester, Alan Turing Building, Oxford Road, Manchester M13 9PL, United Kingdom. E-mail: ron.doney@manchester.ac.uk

<sup>†</sup>Laboratoire de Probabilités et Modèles Aléatoires (LPMA) CNRS : UMR7599, Université Pierre et Marie Curie - Paris VI, Université Paris-Diderot - Paris VII, 4 Place Jussieu, 75252 Paris Cedex 05, France. E-mail: stavros.vakeroudis@ulb.ac.be

<sup>§</sup>When we simply write: Brownian motion, we always mean real-valued Brownian motion, starting from 0. For 2-dimensional Brownian motion, we indicate planar or complex BM.

in terms of Spitzer's condition for stable processes (see e.g. [BeD97] and the references therein).

In Section 2, we recall some facts about standard isotropic stable processes of index  $\alpha \in (0, 2)$  taking values in the complex plane. Then, we follow Bertoin and Werner [BeW96] to define the process of its winding number, we generalize the skew-product representation of planar BM (see e.g. [Kiu80, ReY99, Chy06]) and we present two Lemmas for the winding process of isotropic stable Lévy processes obtained in [BeW96]. Finally, we mention some properties of the positive and the negative moments of the exit times from a cone of this process.

In Section 3, we use some continuity arguments of the composition function due to Whitt [Whi80] and we obtain a new simple proof of the analogue of Spitzer's Theorem for isotropic stable Lévy processes, initially proven by Bertoin and Werner [BeW96]. We reformulate and we extend this result in terms of the exit times from a cone. More precisely, Spitzer's asymptotic Theorem says that, if  $(\vartheta_t, t \geq 0)$  denotes the continuous determination of the argument of a planar BM starting away from the origin, then:

$$\frac{2}{\log t} \vartheta_t \xrightarrow[t \rightarrow \infty]{(law)} C_1, \quad (1)$$

where  $C_1$  is a standard Cauchy variable. For other proofs of (1), see e.g. [Wil74, Dur82, MeY82, PiY86, BeW94a, Yor97, Vak11, VaY12]. Bertoin and Werner state that because an isotropic stable Lévy process is transient, we expect that it winds more slowly than planar Brownian motion and prove that, with  $\theta$  now denoting the process of its winding number (appropriately defined, see e.g. Section 2),  $\theta_t/\sqrt{\log t}$  converges in distribution to some centered Gaussian law as  $t \rightarrow \infty$  (Theorem 1 in Bertoin and Werner [BeW96], stated here as Theorem 3.2).

In Section 4, and more precisely in Theorems 4.1 and 4.4, we study the asymptotics of a symmetric Lévy process and of the winding process of isotropic stable Lévy processes for  $t \rightarrow 0$ , respectively, which are the main results of this article. In particular, we show that  $t^{-1/\alpha}\theta_t$  converges in distribution to an  $\alpha$ -stable law as  $t \rightarrow 0$ . Using this result, in Proposition 4.5 we obtain the (weak) limit in distribution of the process of the exit times from a cone with narrow amplitude and we further obtain several generalizations. We also study the windings of planar stable processes in  $(t, 1]$ , for  $t \rightarrow 0$  and we note that, with obvious notation, Spitzer's law is still valid for  $\theta_{(t,1]}$  (see Remark 4.7). Section 5 deals with the Law of the Iterated Logarithm LIL for Lévy processes for small times, in the spirit of some well-known (LIL) for Brownian motion for  $t \rightarrow \infty$  from Bertoin and Werner [BeW94a, BeW94b] and from Shi [Shi94, Shi98], and for stable subordinators with index  $\alpha \in (0, 1)$  for  $t \rightarrow 0$  from Fristedt [Fri64, Fri67] and Khintchine [Khi39] (see also [Ber96]). Moreover, we prove a LIL for the winding number process of stable processes, for  $t \rightarrow 0$ .

Finally, in Section 6 we discuss the planar Brownian motion case and in Theorem 6.1 we obtain the asymptotic behaviour of the winding process as  $t \rightarrow 0$ . More precisely, the process  $(c^{-1/2}\vartheta_{ct}, t \geq 0)$  converges in law to a 1-dimensional Brownian motion as  $c \rightarrow 0$ .

**Notation:** In the following text, with the symbol " $\implies$ " we shall denote the weak convergence in distribution on the appropriate space, endowed with the Skorohod topology.

## 2 Preliminaries

Following Lamperti [Lam72], a Markov process  $X$  with values in  $\mathbb{R}^d$ ,  $d \geq 2$  is called isotropic or  $O(d)$ -invariant ( $O(d)$  stands for the group of orthogonal transformations on  $\mathbb{R}^d$ ) if its transition satisfies:

$$P_t(\phi(x), \phi(\mathcal{B})) = P_t(x, \mathcal{B}), \quad (2)$$

for any  $\phi \in O(d)$ ,  $x \in \mathbb{R}^d$  and Borel subset  $\mathcal{B} \subset \mathbb{R}^d$ .

Moreover,  $X$  is said to be  $\alpha$ -self-similar if, for  $\alpha > 0$ ,

$$P_{\lambda t}(x, \mathcal{B}) = P_t(\lambda^{-\alpha}x, \lambda^{-\alpha}\mathcal{B}), \quad (3)$$

for any  $\lambda > 0$ ,  $x \in \mathbb{R}^d$  and  $\mathcal{B} \subset \mathbb{R}^d$ .

We focus now our study on the 2-dimensional case ( $d = 2$ ), where (3) holds, and we denote by  $(Z_t, t \geq 0)$  a standard isotropic stable process of index  $\alpha \in (0, 2)$  taking values in the complex plane and starting from  $z_0 + i0$ ,  $z_0 > 0$ . A scaling argument shows that we may assume  $z_0 = 1$ , without loss of generality, since, with obvious notation:

$$\left( Z_t^{(z_0)}, t \geq 0 \right) \stackrel{(law)}{=} \left( z_0 Z_{(t/z_0^\alpha)}^{(1)}, t \geq 0 \right). \quad (4)$$

Thus, from now on, we shall take  $z_0 = 1$ . More precisely,  $Z$  has stationary independent increments, its sample path is right continuous and has left limits (cadlag) and, with  $\langle \cdot, \cdot \rangle$  standing for the Euclidean inner product,  $E[\exp(i\langle \lambda, Z_t \rangle)] = \exp(-t|\lambda|^\alpha)$ , for all  $t \geq 0$  and  $\lambda \in \mathbb{C}$ .  $Z$  is transient,  $\lim_{t \rightarrow \infty} |Z_t| = \infty$  a.s. and it a.s. never visits single points. We remark that for  $\alpha = 2$ , we are in the Brownian motion case.

We are now going to recall some properties of stable processes and Lévy processes (for more details see e.g. [Ber96] or [Kyp06]).

To start with, if  $\mathcal{Z} = (\mathcal{Z}_t, t \geq 0)$  denotes a planar Brownian motion starting from 1 and  $S = (S(t), t \geq 0)$  an independent stable subordinator with index  $\alpha/2$  starting from 0, i.e.:

$$E[\exp(-\mu S(t))] = \exp(-t\mu^{\alpha/2}), \quad (5)$$

for all  $t \geq 0$  and  $\mu \geq 0$ , then the subordinated planar BM  $(\mathcal{Z}_{2S(t)}, t \geq 0)$  is a standard isotropic stable process of index  $\alpha$ . The Lévy measure of  $S$  is:

$$\frac{\alpha}{2\Gamma(1 - \alpha/2)} s^{-1-\alpha/2} 1_{\{s>0\}} ds$$

thus, the Lévy measure  $\nu$  of  $Z$  is:

$$\begin{aligned} \nu(dx) &= \frac{\alpha}{2\Gamma(1 - \alpha/2)} \int_0^\infty s^{-1-\alpha/2} P(\mathcal{Z}_{2s} - 1 \in dx) ds \\ &= \frac{\alpha}{8\pi\Gamma(1 - \alpha/2)} \left( \int_0^\infty s^{-2-\alpha/2} \exp(-|x|^2/(4s)) ds \right) dx \\ &= \frac{\alpha 2^{-1+\alpha/2} \Gamma(1 + \alpha/2)}{\pi\Gamma(1 - \alpha/2)} |x|^{-2-\alpha} dx. \end{aligned} \quad (6)$$

Contrary to planar Brownian motion, as  $Z$  is discontinuous, we cannot define its winding number (recall that, as is well known [ItMK65], for planar BM, since it starts away from the origin, it does not visit a.s. the point 0 but keeps winding around it infinitely often. In particular, the winding process is well defined, for further details see also e.g. [PiY86]). However, following [BeW96], we can consider a path on a finite time interval  $[0, t]$  and "fill in" the gaps with line segments in order to obtain the curve of a continuous function  $f : [0, 1] \rightarrow \mathbb{C}$  with  $f(0) = 1$ . Now, since 0 is polar and  $Z$  has no jumps across 0 a.s., we have  $f(u) \neq 0$  for every  $u \in [0, 1]$ . Hence, we can define the process of the winding number of  $Z$  around 0, which we denote by  $\theta = (\theta_t, t \geq 0)$ . It has cadlag paths of absolute length greater than  $\pi$  and, for all  $t \geq 0$ ,

$$\exp(i\theta_t) = \frac{Z_t}{|Z_t|} . \quad (7)$$

We also introduce the clock:

$$H(t) \equiv \int_0^t \frac{ds}{|Z_s|^\alpha} , \quad (8)$$

and its inverse:

$$A(u) \equiv \inf\{t \geq 0, H(t) > u\} . \quad (9)$$

Bertoin and Werner following [GVA86] obtained these two Lemmas for  $\alpha \in (0, 2)$  (for the proofs see [BeW94a]):

**Lemma 2.1.** *The time-changed process  $(\theta_{A(u)}, u \geq 0)$  is a real-valued symmetric Lévy process. It has no Gaussian component and its Lévy measure has support in  $[-\pi, \pi]$ .*

We now denote by  $dz$  the Lebesgue measure on  $\mathbb{C}$ . Then, for every complex number  $z \neq 0$ ,  $\phi(z)$  denotes the determination of its argument valued in  $(-\pi, \pi]$ .

**Lemma 2.2.** *The Lévy measure of  $\theta_{A(\cdot)}$  is the image of the Lévy measure  $\nu$  of  $Z$  by the mapping  $z \rightarrow \phi(1+z)$ . As a consequence,  $E[(\theta_{A(u)})^2] = uk(\alpha)$ , where*

$$k(\alpha) = \frac{\alpha 2^{-1+\alpha/2} \Gamma(1+\alpha/2)}{\pi \Gamma(1-\alpha/2)} \int_{\mathbb{C}} |z|^{-2-\alpha} |\phi(1+z)|^2 dz . \quad (10)$$

Using Lemma 2.1, we can obtain the analogue of the skew product representation for planar BM which is the Lamperti correspondence for stable processes. Indeed, following [GVA86] and using Lamperti's relation (see e.g. [Lam72, Kiu80, Chy06, CPP11] or [ReY99]) and Lemma 2.1, there exist two real-valued Lévy processes  $(\xi_u, u \geq 0)$  and  $(\rho_u, u \geq 0)$ , the first one non-symmetric whereas the second one symmetric, both starting from 0, such that:

$$\log |Z_t| + i\theta_t = (\xi_u + i\rho_u) \Big|_{u=H_t=\int_0^t \frac{ds}{|Z_s|^\alpha}} . \quad (11)$$

We remark here that  $|Z|$  and  $Z_{A(\cdot)}/|Z_{A(\cdot)}|$  are NOT independent. Indeed, the processes  $|Z_{A(\cdot)}|$  and  $Z_{A(\cdot)}/|Z_{A(\cdot)}|$  jump at the same times hence they cannot be independent. Moreover,  $A(\cdot)$  depends only upon  $|Z|$ , hence  $|Z|$  and  $Z_{A(\cdot)}/|Z_{A(\cdot)}|$  are not independent. For

further discussion on the independence, see e.g. [LiW11], where is shown that an isotropic  $\alpha$ -self-similar Markov process has a skew-product structure if and only if its radial and its angular part do not jump at the same time.

We also remark that

$$H^{-1}(u) \equiv A(u) \equiv \inf\{t \geq 0 : H(t) > u\} = \int_0^u \exp\{\alpha\xi_s\} ds . \quad (12)$$

Hence, (11) may be equivalently written as:

$$\begin{cases} |Z_t| = \exp(\xi(H_t)) \Leftrightarrow |Z_{A(t)}| = \exp(\xi_t), & \text{(extension of Lamperti's identity)} \\ \theta_t = \rho(H_t) \Leftrightarrow \theta(A(t)) = \rho(t) . \end{cases} \quad (13)$$

We also define the random times  $T_c^{|\theta|} \equiv \inf\{t : |\theta_t| \geq c\}$  and  $T_c^{|\rho|} \equiv \inf\{t : |\rho_t| \geq c\}$ , ( $c > 0$ ). Using the "generalized" skew-product representation (11) (or (13)), we obtain:

$$T_c^{|\theta|} = H_u^{-1} \Big|_{u=T_c^{|\rho|}} = \int_0^{T_c^{|\rho|}} ds \exp(\alpha\xi_s) \equiv A_{T_c^{|\rho|}} . \quad (14)$$

Following [VaY12], for the random times  $T_{-d,c}^\theta \equiv \inf\{t : \theta_t \notin (-d, c)\}$ ,  $d, c > 0$ , and  $T_c^\theta \equiv \inf\{t : \theta_t \geq c\}$ , we have:

**Remark 2.3.** For  $0 < c < d$ , the random times  $T_{-d,c}^\theta$ ,  $T_c^{|\theta|}$  and  $T_c^\theta$  satisfy the trivial inequality:

$$T_c^{|\theta|} \leq T_{-d,c}^\theta \leq T_c^\theta . \quad (15)$$

Hence, with  $p > 0$ :

$$E \left[ (T_c^{|\theta|})^p \right] \leq E \left[ (T_{-d,c}^\theta)^p \right] \leq E \left[ (T_c^\theta)^p \right] , \quad (16)$$

and for the negative moments:

$$E \left[ (T_c^\theta)^{-p} \right] \leq E \left[ (T_{-d,c}^\theta)^{-p} \right] \leq E \left[ (T_c^{|\theta|})^{-p} \right] . \quad (17)$$

**Remark 2.4.** For further details concerning the finiteness of the positive moments of  $T_c^{|\theta|}$ , see e.g. [DeB90, BaB04]. Recall also that for the positive moments of the exit time from a cone of planar Brownian motion, Spitzer showed that (with obvious notation) [Spi58, Bur77]:

$$E \left[ (T_c^{|\theta|})^p \right] < \infty \Leftrightarrow p < \frac{\pi}{4c} , \quad (18)$$

whereas all the negative moments  $E \left[ (T_c^{|\theta|})^{-p} \right]$  are finite [VaY12].

We denote now by  $\Psi(u)$  the exponent of the symmetric Lévy process  $\rho$ , hence (Lévy-Khintchine formula)  $E[e^{iu\rho t}] = e^{-t\Psi(u)}$ , with:

$$\Psi(u) = \int_{(-\infty, \infty)} (1 - e^{iux} + iux1_{\{|x| \leq 1\}}) \mu(dx), \quad u \in \mathbb{R}, \quad (19)$$

where  $\mu$  is a Radon measure on  $\mathbb{R} \setminus \{0\}$  such that:

$$\int_{(-\infty, \infty)} (x^2 \wedge 1) \mu(dx) < \infty .$$

$\mu$  is the Lévy measure of  $\rho$  and is symmetric.

### 3 Large time asymptotics

Concerning the clock  $H$ , we have the almost sure convergence (see Corollary 1 in Bertoin and Werner [BeW96]):

$$\frac{H(e^u)}{u} \xrightarrow[u \rightarrow \infty]{a.s.} 2^{-\alpha} \frac{\Gamma(1 - \alpha/2)}{\Gamma(1 + \alpha/2)} \equiv K(\alpha) = E [|Z_1|^{-\alpha}] . \quad (20)$$

Moreover, we have the following:

**Proposition 3.1.** *The family of processes*

$$H_x^{(u)} \equiv \left( \frac{H(e^{ux})}{u}, x \geq 0 \right)$$

is tight, as  $u \rightarrow \infty$ .

**Proof of Proposition 3.1:**

To prove this, we could repeat some arguments of Pitman and Yor [PiY89] (see the estimates in their proof of Theorem 6.4), however, we give here a straightforward proof, using the definition of tightness:

for every  $\varepsilon, \eta > 0$ , there exist  $\delta > 0$  and  $C_\delta > 0$  such that, for every  $0 < x < y$ :

$$P \left( \sup_{|x-y| \leq \delta} |H(e^{uy}) - H(e^{ux})| \geq u\varepsilon \right) \leq \eta , \quad \text{for } u \geq C_\delta , \quad (21)$$

or equivalently:

$$P \left( \frac{1}{u} |H(e^{u(x+\delta)}) - H(e^{ux})| \geq \varepsilon \right) \leq \eta , \quad \text{for } u \geq C_\delta . \quad (22)$$

First, following Bertoin and Werner [BeW96], we introduce the "Ornstein-Uhlenbeck type" process:

$$\tilde{Z}_u = \exp(-u/\alpha) Z_{\exp(u)}, \quad u \geq 0 , \quad (23)$$

which is a stationary Markov process under  $P_0$  (see e.g. [Bre68]). We denote by  $p_t(\cdot)$  the semigroup of  $Z$ :

$$p_t(\bar{z}) = P_0(Z_t \in d\bar{z})/d\bar{z}, \quad \bar{z} \in \mathbb{C}.$$

We denote by  $Z^{(0)}$  another stable process starting at 0. Then, using the scaling property, given that  $\tilde{Z}_0 \equiv Z_1 \equiv 1 + Z_1^{(0)} = \bar{x}$ , the semigroup  $q_u(\cdot)$  of  $\tilde{Z}$  is given by:

$$\begin{aligned} q_u(\bar{x}, \bar{y}) &= p_{\exp(u)-1}(e^{u/\alpha}\bar{y} - \bar{x}) e^{2u/\alpha} \\ &= (e^u - 1)^{-2/\alpha} e^{2u/\alpha} p_1((e^u - 1)^{-1/\alpha}(e^{u/\alpha}\bar{y} - \bar{x})) \\ &= (l(u))^2 p_1(l(u)(\bar{y} - e^{-u/\alpha}\bar{x})) , \end{aligned} \quad (24)$$

where  $l(v) = e^{v/\alpha}(e^v - 1)^{-1/\alpha}$ . For every  $\delta > 0$  and changing variables  $s = \exp(v)$ , with obvious notation, we have:

$$E [|H(e^{u(x+\delta)}) - H(e^{ux})|] = \int_{e^{ux}}^{e^{u(x+\delta)}} E [|Z_s|^{-\alpha}] ds = \int_{ux}^{u(x+\delta)} E_{\tilde{Z}_0} [| \tilde{Z}_v |^{-\alpha}] dv . \quad (25)$$

We also define  $\varepsilon(v) \equiv l(v)e^{-v/\alpha} = (e^v - 1)^{-1/\alpha}$ . From (23), using the stability of  $Z$ , we have:

$$\begin{aligned}\tilde{Z}_v &= e^{-v/\alpha} Z_{\exp(v)} = e^{-v/\alpha} (Z_{\exp(v)-1} + Z_1) \stackrel{(law)}{=} e^{-v/\alpha} \left( (e^v - 1)^{1/\alpha} Z_1^{(0)} + Z_1 \right) \\ &= (l(v))^{-1} \left( Z_1^{(0)} + \varepsilon(v) Z_1 \right).\end{aligned}\tag{26}$$

Hence (for simplicity, we use  $E \equiv E_0$ ):

$$E_{\tilde{Z}_0} \left[ |\tilde{Z}_v|^{-\alpha} \right] = (l(v))^\alpha E \left[ |Z_1^{(0)} + \varepsilon(v) \tilde{Z}_0|^{-\alpha} \right] \equiv (l(v))^\alpha (E_1 + E_2),\tag{27}$$

where, with  $\delta' > 0$ ,

$$\begin{aligned}E_1 &= E \left[ |Z_1^{(0)} + \varepsilon(v) \tilde{Z}_0|^{-\alpha} : |Z_1^{(0)} + \varepsilon(v) \tilde{Z}_0| \geq \delta' \right], \\ E_2 &= E \left[ |Z_1^{(0)} + \varepsilon(v) \tilde{Z}_0|^{-\alpha} : |Z_1^{(0)} + \varepsilon(v) \tilde{Z}_0| \leq \delta' \right].\end{aligned}$$

We have:  $l(v) \xrightarrow{v \rightarrow \infty} 1$  and  $\varepsilon(v) \xrightarrow{v \rightarrow \infty} 0$ , thus, by Dominated Convergence Theorem:

$$E_1 \xrightarrow{v \rightarrow \infty} E \left[ |Z_1^{(0)}|^{-\alpha} : |Z_1^{(0)}| \geq \delta' \right] \xrightarrow{\delta' \rightarrow 0} E \left[ |Z_1^{(0)}|^{-\alpha} \right].\tag{28}$$

Moreover, changing the variables:  $\bar{w} = \bar{z} + \varepsilon(v)\bar{x}$ , we have:

$$\begin{aligned}E_2 &= \int_{\bar{x}, \bar{z}: |\bar{z} + \varepsilon(v)\bar{x}| \leq \delta'} P(\tilde{Z}_0 \in d\bar{x}) P(Z_1^{(0)} \in d\bar{z}) |\bar{z} + \varepsilon(v)\bar{x}|^{-\alpha} \\ &= \int_{\bar{x}, \bar{w}: |\bar{w}| \leq \delta'} P(\tilde{Z}_0 \in d\bar{x}) P(Z_1^{(0)} \in d\bar{w}) |\bar{w}|^{-\alpha}.\end{aligned}$$

Remarking now that for stable processes:  $P(Z_1^{(0)} \in d\bar{y}) \leq C' d\bar{y}$ , where  $C'$  stands for a positive constant and using  $w = (w_1, w_2)$ , we have:

$$E_2 \leq C' \int_{\bar{x}, \bar{w}: |\bar{w}| \leq \delta'} P(\tilde{Z}_0 \in d\bar{x}) \frac{dw_1 dw_2}{|\bar{w}|^\alpha} = C' \int_{\bar{z}: |\bar{w}| \leq \delta'} \frac{dw_1 dw_2}{|\bar{w}|^\alpha} \xrightarrow{\delta' \rightarrow 0} 0.\tag{29}$$

Thus, from (27), (28) and (29), invoking again the Dominated Convergence Theorem, we deduce:

$$\lim_{u \rightarrow \infty} E_{\tilde{Z}_0} \left[ |\tilde{Z}_v|^{-\alpha} \right] = E \left[ |Z_1^{(0)}|^{-\alpha} \right],\tag{30}$$

which is a constant. Hence, for every  $\varepsilon, \eta > 0$ , there exist  $\delta > 0$  and  $C_\delta > 0$  such that (22) is satisfied for  $u \geq C_\delta$ .  $\blacksquare$

Bertoin and Werner in [BeW96] obtained the analogue of Spitzer's asymptotic Theorem [Spi58] for isotropic stable Lévy processes of index  $\alpha \in (0, 2)$ :

**Theorem 3.2.** *The family of processes*

$$\Theta_t^{(c)} \equiv (c^{-1/2} \theta_{\exp(ct)}, t \geq 0)$$



converges in distribution on  $D([0, \infty), \mathbb{R})$  endowed with the Skorohod topology, as  $c \rightarrow \infty$ , to  $(\sqrt{r(\alpha)}B_t, t \geq 0)$ , where  $(B_s, s \geq 0)$  is a real valued Brownian motion and

$$r(\alpha) = \frac{\alpha 2^{-1-\alpha/2}}{\pi} \int_{\mathbb{C}} |z|^{-2-\alpha} |\phi(1+z)|^2 dz . \quad (31)$$

Using some results due to Whitt [Whi80], we can obtain a simple proof of this Theorem.

**Proof of Theorem 3.2: (new proof)**

Essentially, an argument of continuity of the composition function (Theorem 3.1 in [Whi80]) may replace the martingale argument in the lines of the proof for  $t \rightarrow \infty$  from Bertoin and Werner. We split the proof in three parts:

i) Concerning the clock  $H$ , we have the almost sure convergence (20):

$$\frac{H(e^u)}{u} \xrightarrow[u \rightarrow \infty]{a.s.} K(\alpha) = E [|Z_1|^{-\alpha}] .$$

From this result follows the convergence of the finite dimensional distributions of  $v^{-1}H(\exp(vt))$ , for  $v \rightarrow \infty$  and every  $t > 0$ .

Moreover, from Proposition 3.1, the family of processes

$$H_x^{(u)} \equiv \left( \frac{H(e^{ux})}{u}, x \geq 0 \right)$$

is tight as  $u \rightarrow \infty$ . Hence, from (20) and (21), finally,  $H^{(u)}(t) \equiv (u^{-1}H(\exp(ut)), t \geq 0)$  converges weakly to  $(tK(\alpha), t \geq 0)$  as  $u \rightarrow \infty$ , i.e.:

$$\left( \frac{H(e^{ut})}{u}, t \geq 0 \right) \xrightarrow[u \rightarrow \infty]{(d)} (tK(\alpha), t \geq 0), \quad (32)$$

where the convergence in distribution is viewed on  $D([0, \infty), \mathbb{R})$  endowed with the Skorohod topology.

ii) Using the skew product representation analogue (11) and Lemma 2.2, we have:

$$\left( \frac{\rho_{tu}}{\sqrt{u}}, t \geq 0 \right) \xrightarrow[u \rightarrow \infty]{(d)} \left( \sqrt{k(\alpha)} B_t, t \geq 0 \right) , \quad (33)$$

where the convergence in distribution is viewed again on  $D([0, \infty), \mathbb{R})$  endowed with the Skorohod topology and  $k(\alpha)$  is given by (10). This follows from the convergence of the finite dimensional distributions:

$$\frac{\rho_u}{\sqrt{u}} = \frac{\theta_{A(u)}}{\sqrt{u}} \xrightarrow[u \rightarrow \infty]{(d)} \sqrt{k(\alpha)} B_1 , \quad (34)$$

a condition which is sufficient for the weak convergence (33), since Lévy processes are semimartingales with stationary independent increments; for further details see e.g. [Sko91] or [JaS03] (Corollary 3.6, Chapter VII, p. 415).

iii) Theorem 3.1 in [Whi80] states that the composition function on  $D([0, \infty), \mathbb{R}) \times D([0, \infty), [0, \infty))$  is continuous at each  $(\rho, H) \in (C([0, \infty), \mathbb{R}) \times D_0([0, \infty), [0, \infty)))$ , with  $C$  denoting the set of continuous functions and  $D_0$  the subset of increasing cadlag functions in  $D$  (hence the subset of non-decreasing cadlag functions in  $D$ ). Hence, from (32) and (33), we have: for every  $t > 0$ ,

$$\frac{\theta_{\exp(ct)}}{\sqrt{c}} = \frac{\rho_{H(\exp(ct))}}{\sqrt{c}} = \frac{\rho_{c(H(e^{ct})/c)}}{\sqrt{c}}. \quad (35)$$

The result now follows from the continuity of the composition function together with (35) and the weak convergence of  $H^{(c)}(\cdot)$  and  $c^{-1/2}\rho_c$ , as  $c \rightarrow \infty$ . ■

From Theorem 3.2, we can obtain the asymptotic behaviour of the exit times from a cone for isotropic stable processes which generalizes a recent result in [VaY12]:

**Proposition 3.3.** *For  $c \rightarrow \infty$ , for every  $x > 0$ , we have the weak convergence:*

$$\left( \frac{1}{c} \log \left( T_{x\sqrt{c}}^\theta \right), x \geq 0 \right) \xrightarrow[c \rightarrow \infty]{(d)} \left( \tau_{\sqrt{1/r(\alpha)}}^{(1/2)}, x \geq 0 \right), \quad (36)$$

where for every  $y > 0$ ,  $\tau_y^{(1/2)}$  stands for the  $\frac{1}{2}$ -stable process defined by:  $\tau_y^{(1/2)} \equiv \inf\{t : B_t = y\}$ .

**Proof of Proposition 3.3:**

We rely now upon Theorem 3.2, the analogue of Spitzer's Theorem for stable processes by Bertoin and Werner:

$$\Theta_t^{(c)} \equiv (c^{-1/2}\theta_{\exp(ct)}, t \geq 0) \xrightarrow[c \rightarrow \infty]{(d)} (B_{r(\alpha)t}, t \geq 0). \quad (37)$$

Hence, for every  $x > 0$ ,

$$\begin{aligned} \frac{1}{c} \log \left( T_{x\sqrt{c}}^\theta \right) &= \frac{1}{c} \log \left( \inf \{ t : \theta_t > x\sqrt{c} \} \right) \\ &\stackrel{t=\exp(cs)}{=} \frac{1}{c} \log \left( \inf \left\{ e^{cs} : \frac{1}{\sqrt{c}} \theta_{\exp(cs)} > x \right\} \right) \\ &= \inf \left\{ s : \frac{1}{\sqrt{c}} \theta_{\exp(cs)} > x \right\} \\ &\xrightarrow{c \rightarrow \infty} \inf \{ s : B_{r(\alpha)s} > x \} \\ &= \inf \left\{ s : \sqrt{r(\alpha)} B_s > x \right\} \equiv \tau_{x/\sqrt{r(\alpha)}}^{(1/2)}. \end{aligned} \quad (38)$$

Moreover, from Theorem 7.1 in [Whi80], we know that the first passage time function mapping is continuous, thus, we deduce (36). ■

If we replace  $c$  by  $ac$ , we can obtain several variants of Proposition 3.3 for the random times  $T_{-bc, ac}^\theta$ ,  $0 < a, b \leq \infty$ , for  $c \rightarrow \infty$ , and  $a, b > 0$  fixed:

**Corollary 3.4.** *The following asymptotic results hold:*

$$\frac{1}{c} \log \left( T_{\sqrt{ac}}^\theta \right) \xrightarrow[c \rightarrow \infty]{(law)} \tau_{\sqrt{a/r(\alpha)}}^{(1/2)}, \quad (39)$$

$$\frac{1}{c} \log \left( T_{\sqrt{ac}}^{|\theta|} \right) \xrightarrow[c \rightarrow \infty]{(law)} \tau_{\sqrt{a/r(\alpha)}}^{|\theta|}, \quad (40)$$

$$\frac{1}{c} \log \left( T_{-\sqrt{bc}, \sqrt{ac}}^\theta \right) \xrightarrow[c \rightarrow \infty]{(law)} \tau_{-\sqrt{b/r(\alpha)}, \sqrt{a/r(\alpha)}}^B, \quad (41)$$

where for every  $x, y > 0$ ,  $\tau_x^{|\theta|} \equiv \inf\{t : |B_t| = x\}$  and  $\tau_{-y,x}^B \equiv \inf\{t : B_t \notin (-y, x)\}$ .

**Proposition 3.5.** *The following asymptotic result for  $\alpha \in (0, 2)$ , holds: for every  $b > 0$ ,*

$$P \left( T_{b\sqrt{\log t}}^\theta > t \right) \xrightarrow[t \rightarrow \infty]{} \operatorname{erf} \left( \frac{b}{\sqrt{2r(\alpha)}} \right), \quad (42)$$

where  $\operatorname{erf}(x) \equiv \frac{2}{\sqrt{\pi}} \int_0^x e^{-y^2} dy$  is the error function.

**Proof of Proposition 3.5:**

Using the notation of Theorem 3.2, for every  $b > 0$ , we have:

$$\begin{aligned} P \left( T_{b\sqrt{\log t}}^\theta > t \right) &= P \left( \sup_{u \leq t} \theta_u < b\sqrt{\log t} \right) \stackrel{u=tv}{=} P \left( \sup_{v \leq 1} (\log t)^{-1/2} \theta_{tv} < b \right) \\ &\stackrel{t=e^c}{=} P \left( \sup_{v \leq 1} c^{-1/2} \theta_{\exp(cv)} < b \right) \end{aligned}$$

Hence, using Theorem 3.2 for  $t \rightarrow \infty$ , we deduce:

$$\begin{aligned} P \left( T_{b\sqrt{\log t}}^\theta > t \right) &\xrightarrow[t \rightarrow \infty]{} P \left( \sup_{v \leq 1} \sqrt{r(\alpha)} B_v < b \right) = P \left( |B_1| < \frac{b}{\sqrt{r(\alpha)}} \right) \\ &= 2 \int_0^{b/\sqrt{r(\alpha)}} \frac{dw}{\sqrt{2\pi}} e^{-w^2/2}, \end{aligned}$$

and changing the variables  $w = y\sqrt{2}$ , we obtain (42). ■

As mentioned in [BeW96], because an isotropic stable Lévy process  $Z$  is transient, the difference between  $\theta$  and the winding number around an arbitrary fixed  $z \neq 1$  is bounded and converges as  $t \rightarrow \infty$ . Hence, with  $(\theta_t^i, t > 0)$ ,  $1 \leq i \leq n$  denoting the continuous total angle wound of  $Z$  of index  $\alpha \in (0, 2)$  around  $z^i$  ( $z^1, \dots, z^n$  are  $n$  distinct points in the complex plane  $\mathbb{C}$ ) up to time  $t$ , we obtain the following concerning the finite dimensional distributions (windings around several points):

**Proposition 3.6.** *For isotropic stable Lévy processes of index  $\alpha \in (0, 2)$ , we have:*

$$\left( \frac{\theta_t^i}{\sqrt{\log t}}, 1 \leq i \leq n \right) \xrightarrow[t \rightarrow \infty]{(d)} \left( \sqrt{r(\alpha)} B_1^i, 1 \leq i \leq n \right), \quad (43)$$

where  $(B_s^i, 1 \leq i \leq n, s \geq 0)$  is an  $n$ -dimensional Brownian motion and  $r(\alpha)$  is given by (31).

## 4 Small time asymptotics

We turn now our study to the behaviour of  $\theta_t$  for  $t \rightarrow 0$ .

**Theorem 4.1.** *For  $\alpha \in (0, 2)$ , the following convergence in law holds:*

$$(t^{-1/\alpha} \rho_{ts}, s \geq 0) \xrightarrow[t \rightarrow 0]{(d)} (\zeta_s, s \geq 0) , \quad (44)$$

where  $(\zeta_s, s \geq 0)$  is a symmetric 1-dimensional  $\alpha$ -stable process and the convergence in distribution is considered on  $D([0, \infty), \mathbb{R})$  endowed with the Skorohod topology.

**Proof of Theorem 4.1:**

From Lemma 2.2, we use the Lévy measure, say  $\tilde{\pi}$ , of  $\theta_{A(\cdot)}$  (thus the Lévy measure of  $\rho$ ) and we prove that for  $t \rightarrow 0$  it converges to the Lévy measure of a 1-dimensional  $\alpha$ -stable process. Indeed, with

$$L \equiv \frac{\alpha 2^{-1+\alpha/2} \Gamma(1 + \alpha/2)}{\pi \Gamma(1 - \alpha/2)},$$

and  $z$  denoting a number in  $\mathbb{C}$ , using polar coordinates, we have:

$$\phi(1+z) = \int_{\mathbb{C}} \frac{dz}{1+z} = 2L \int_0^\pi \int_0^\infty \frac{r dr d\varphi}{(1+r^2-2r \cos \varphi)^{1+\alpha/2}} .$$

We remark that:

$$1+r^2-2r \cos \varphi = (r - \cos \varphi)^2 + \sin^2 \varphi ,$$

hence, changing the variables  $(r - \cos \varphi)^2 = t^{-1} \sin^2 \varphi$  and denoting by:

$$B(y; a, b) = \int_0^y u^{a-1} (1-u)^{b-1} du ,$$

the incomplete Beta function, for  $\varphi > 0$  (we can repeat the same arguments for  $\varphi < 0$ ) we have:

$$\begin{aligned} \tilde{\pi}(d\varphi) &= d\varphi 2L \int_0^\infty \frac{r dr}{((r - \cos \varphi)^2 + \sin^2 \varphi)^{1+\alpha/2}} \\ &= d\varphi \frac{2L}{2} \left( \frac{2}{\alpha} + \cos \varphi (1 - \cos^2 \varphi)^{-\frac{1}{2}-\frac{\alpha}{2}} \int_0^{1-\frac{1}{\cos^2 \varphi}} t^{-\frac{1}{2}+\frac{\alpha}{2}} (1+t)^{-\frac{\alpha}{2}-1} dt \right) \\ &\stackrel{u=-t}{=} d\varphi L \left( \frac{2}{\alpha} + \cos \varphi (-1 + \cos^2 \varphi)^{-\frac{1}{2}-\frac{\alpha}{2}} \int_0^{1-\frac{1}{\cos^2 \varphi}} u^{-\frac{1}{2}+\frac{\alpha}{2}} (1-u)^{-\frac{\alpha}{2}-1} du \right) \\ &= d\varphi L \left( \frac{2}{\alpha} + \cos \varphi (-1 + \cos^2 \varphi)^{-\frac{1}{2}-\frac{\alpha}{2}} B \left( 1 - \frac{1}{\cos^2 \varphi}; \frac{1}{2} + \frac{\alpha}{2}, -\frac{\alpha}{2} \right) \right) \\ &\stackrel{\varphi \sim 0}{\sim} \tilde{L} \varphi^{-1-\alpha} d\varphi , \end{aligned} \quad (45)$$

which is the Lévy measure of an  $\alpha$ -stable process. The result now follows by standard arguments. ■

Concerning the clock  $H$  and its increments, we have:

**Theorem 4.2.** *The following a.s. convergence holds:*

$$\left( \frac{H(ux)}{u}, x \geq 0 \right) \xrightarrow[u \rightarrow 0]{a.s.} (x, x \geq 0) . \quad (46)$$

**Proof of Theorem 4.2:**

From the definition of the clock  $H$  we have:

$$\frac{H(ux)}{u} = \frac{1}{u} \int_0^{ux} \frac{ds}{|Z_s|^\alpha} .$$

Hence, for every  $x_0 > 0$ , we have:

$$\begin{aligned} \sup_{x \leq x_0} \left| \frac{H(ux) - ux}{u} \right| &= \sup_{x \leq x_0} \frac{1}{u} \left| \int_0^{ux} \left( \frac{1}{|Z_s|^\alpha} - 1 \right) ds \right| \leq \frac{1}{u} \int_0^{ux_0} \left| \frac{1}{|Z_s|^\alpha} - 1 \right| ds \\ &\stackrel{s=uw}{=} \int_0^{x_0} \left| \frac{1}{|Z_{uw}|^\alpha} - 1 \right| dw \xrightarrow[u \rightarrow 0]{a.s.} 0 . \end{aligned} \quad (47)$$

because:

$$|Z_u|^\alpha \xrightarrow[u \rightarrow 0]{a.s.} 1 . \quad (48)$$

Thus, as (47) is true for every  $x_0 > 0$ , we obtain (46). ■

**Remark 4.3.** *We remark that this behaviour of the clock is different for the case  $t \rightarrow \infty$ , where (32) can be equivalently stated as:*

$$\left( \frac{H(ux)}{\log u}, x \geq 0 \right) \xrightarrow[u \rightarrow \infty]{(d)} \left( 2^{-\alpha} \frac{\Gamma(1 - \alpha/2)}{\Gamma(1 + \alpha/2)} x, x \geq 0 \right) . \quad (49)$$

Using Theorems 4.1 and 4.2, we obtain:

**Theorem 4.4.** *With  $\alpha \in (0, 2)$ , the family of processes*

$$(c^{-1/\alpha} \theta_{ct}, t \geq 0)$$

*converges in distribution on  $D([0, \infty), \mathbb{R})$  endowed with the Skorohod topology, as  $c \rightarrow 0$ , to a symmetric 1-dimensional  $\alpha$ -stable process  $(\zeta_t, t \geq 0)$ .*

**Proof of Theorem 4.4:**

We will use Theorems 4.1 and 4.2. More precisely, we shall rely again upon the continuity of the composition function as studied in Theorem 3.1 in [Whi80].

i) First, concerning the clock  $H$ , for every  $t > 0$ , we have the almost sure convergence (46), which yields the weak convergence (on  $D([0, \infty), \mathbb{R})$  endowed with the Skorohod topology) of the family of processes  $\tilde{H}^{(u)}(t) \equiv (u^{-1}H(ut), t \geq 0)$  to  $(t, t \geq 0)$  as  $u \rightarrow 0$ .

ii) We use another result of Whitt [Whi80] which states that the composition function on  $D([0, \infty), \mathbb{R}) \times D([0, \infty), [0, \infty))$  is continuous at each  $(\rho, H) \in (D([0, \infty), \mathbb{R}) \times C_0([0, \infty), [0, \infty)))$ , with  $D$  denoting the set of cadlag functions and  $C_0$  the subset of

strictly-increasing functions in  $C$ . Hence, from Theorem 4.1 and (46), using the weak convergence of  $\tilde{H}^{(c)}(\cdot)$  and of  $c^{-1/\alpha}\rho_c$ , as  $c \rightarrow 0$ , we deduce: for every  $t > 0$ ,

$$\frac{\theta_{ct}}{c^{1/\alpha}} = \frac{\rho_{H(ct)}}{c^{1/\alpha}} = \frac{\rho_{c(H(ct)/c)}}{c^{1/\alpha}} \xrightarrow[c \rightarrow 0]{(d)} \zeta_t, \quad (50)$$

where the convergence in distribution is viewed on  $D([0, \infty), \mathbb{R})$  endowed with the Skorohod topology.  $\blacksquare$

From the previous results, we deduce the asymptotic behaviour, for  $c \rightarrow 0$ , of the first exit times from a cone for isotropic stable processes of index  $\alpha \in (0, 2)$  taking values in the complex plane:

**Proposition 4.5.** *For  $c \rightarrow 0$ , we have the weak convergence:*

$$\left( \frac{1}{c} T_{c^{1/\alpha}x}^\theta, x \geq 0 \right) \xrightarrow[c \rightarrow 0]{(d)} (T_x^\zeta, x \geq 0), \quad (51)$$

where for every  $x$ ,  $T_x^\zeta$  is the first hitting time defined by:  $T_x^\zeta \equiv \inf\{t : \zeta_t = x\}$ .

**Proof of Proposition 4.5:**

Using Theorem 4.4, we have:

$$\begin{aligned} \frac{1}{c} T_{c^{1/\alpha}x}^\theta &= \frac{1}{c} \inf\{t : \theta_t > c^{1/\alpha}x\} \stackrel{t=cs}{=} \frac{1}{c} \inf\{cs : c^{-1/\alpha}\theta_{cs} > x\} = \inf\{s : c^{-1/\alpha}\theta_{cs} > x\} \\ &\xrightarrow[c \rightarrow 0]{} \inf\{s : \zeta_s > x\}, \end{aligned}$$

which, using again the continuity of the first passage time function mapping (see Theorem 7.1 in [Whi80]), yields (51).  $\blacksquare$

Finally, we can obtain several variants of Proposition 4.5 for the random times  $T_{-bc,ac}^\theta$ ,  $0 < a, b \leq \infty$  fixed, for  $c \rightarrow 0$ :

**Corollary 4.6.** *The following asymptotic results hold:*

$$\frac{1}{c} T_{ac^{1/\alpha}}^\theta \xrightarrow[c \rightarrow 0]{(law)} T_a^\zeta, \quad (52)$$

$$\frac{1}{c} T_{ac^{1/\alpha}}^{|\theta|} \xrightarrow[c \rightarrow 0]{(law)} T_a^{|\zeta|}, \quad (53)$$

$$\frac{1}{c} T_{-bc^{1/\alpha}, ac^{1/\alpha}}^\theta \xrightarrow[c \rightarrow 0]{(law)} T_{-b,a}^\zeta, \quad (54)$$

where, for every  $x, y > 0$ ,  $T_x^{|\zeta|} \equiv \inf\{t : |\zeta_t| = x\}$  and  $T_{-y,x}^\zeta \equiv \inf\{t : \zeta_t \notin (-y, x)\}$ .

**Remark 4.7. (Windings of planar stable processes in  $(t, 1]$  for  $t \rightarrow 0$ )**

*i) We consider now our stable process  $Z$  starting from 0 and we want to investigate its windings in  $(t, 1]$  for  $t \rightarrow 0$ . We know that it doesn't visit again the origin but it winds a.s. infinitely often around it, hence, its winding process  $\theta$  in  $(t, 1]$  is well-defined. With*

obvious notation, concerning now the clock  $H_{(t,1]} = \int_t^1 du |Z_u|^{-\alpha}$ , the change of variables  $u = tv$  and the stability property, i.e.:  $Z_{tv} \stackrel{(law)}{=} t^{1/\alpha} Z_v$ , yield:

$$H_{(t,1]} = \int_1^{1/t} \frac{t dv}{|Z_{tv}|^\alpha} \stackrel{(law)}{=} \int_1^{1/t} \frac{dv}{|Z_v|^\alpha} = H_{(1,1/t]} .$$

Hence, as before, using Whitt's result [Whi80] on the continuity of the composition function on  $D([0, \infty), \mathbb{R}) \times D([0, \infty), [0, \infty))$  at each  $(\rho, H) \in (D([0, \infty), \mathbb{R}) \times C_0([0, \infty), [0, \infty)))$ , we have (with obvious notation):

$$\theta_{(t,1]} = \rho_{H_{(t,1]}} \stackrel{(law)}{=} \rho_{H_{(1,1/t]}} = \theta_{(1,1/t]} . \quad (55)$$

The only difference with respect to the "normal" stable case is that the winding process is considered from 1 and not from 0, but this doesn't provoke any problem.

Hence, Bertoin and Werner's Theorem 3.2 is still valid for  $Z$  in  $(t, 1]$ ,  $t \rightarrow 0$ : for  $\alpha \in (0, 2)$ :

$$\frac{1}{\sqrt{\log(1/t)}} \theta_{(t,1]} \stackrel{(law)}{=} \frac{1}{\sqrt{\log(1/t)}} \theta_{(1,1/t]} \xrightarrow[t \rightarrow 0]{(d)} \sqrt{r(\alpha)} N, \quad (56)$$

with  $r(\alpha)$  defined in (31) and  $N \sim \mathcal{N}(0, 1)$ .

ii) We note that this study is also valid for a planar Brownian motion starting from 0 in  $(t, 1]$  for  $t \rightarrow 0$  and for planar stable processes or planar Brownian motion starting both from a point different from 0 (in order to have an well-defined winding number) in  $[0, 1]$ . In particular, for planar Brownian motion  $\mathcal{Z}$  with associated winding number  $\vartheta$ , we obtain that Spitzer's law is still valid for  $t \rightarrow 0$  (see e.g. [ReY99, LeG92]):

$$\frac{2}{\log(1/t)} \vartheta_{(t,1]} \xrightarrow[t \rightarrow 0]{(law)} C_1, \quad (57)$$

where  $C_1$  is a standard Cauchy variable. We also remark that this result could also be obtained from a time inversion argument, that is: with  $\mathcal{Z}'$  denoting another planar Brownian motion starting from 0, with winding number  $\vartheta'$ , by time inversion we have:  $\mathcal{Z}_u = u\mathcal{Z}'_{1/u}$ . Changing now the variables  $u = 1/v$ , we obtain:

$$\begin{aligned} \vartheta_{(t,1]} &\equiv \operatorname{Im} \left( \int_t^1 \frac{d\mathcal{Z}_u}{\mathcal{Z}_u} \right) = \operatorname{Im} \left( \int_t^1 \frac{d(u\mathcal{Z}'_{1/u})}{u\mathcal{Z}'_{1/u}} \right) = \operatorname{Im} \left( \int_t^1 \frac{d\mathcal{Z}'_{1/u}}{\mathcal{Z}'_{1/u}} \right) \\ &= \operatorname{Im} \left( \int_1^{1/t} \frac{d\mathcal{Z}'_v}{\mathcal{Z}'_v} \right) \equiv \vartheta'_{(1,1/t]} , \end{aligned}$$

and we continue as before.

Note that this time inversion argument is NOT valid for planar stable processes.

## 5 The Law of the Iterated Logarithm (LIL)

In this Section, we shall use some notation introduced in [DoM02a, DoM02b]. Recall (19); then, for all  $x > 0$ , because  $\rho$  is symmetric, we define:

$$L(x) = 2\hat{\mu}(x) = 2\mu(x, +\infty), \quad U(x) = 2 \int_0^x yL(y)dy .$$

We remark that  $U$  plays essentially the role of the truncated variance in the random walk case (see e.g. [DoM00]).

Hence, from (44), for  $t \rightarrow 0$ , we have ( $\tilde{K}(\alpha)$  is a constant depending on  $\alpha$ ):

$$U(x) \stackrel{x \rightarrow 0}{\sim} \tilde{K}(\alpha)x^{2-\alpha}. \quad (58)$$

Then, we obtain the following Law of the Iterated Logarithm (LIL) for Lévy processes for small times:

### Theorem 5.1. (LIL for Lévy processes for small times)

For any non-decreasing function  $f > 0$ ,

$$\limsup_{t \rightarrow 0} \frac{\rho_t}{f(t)} = \begin{cases} 0 ; \\ \infty \end{cases} \quad a.s. \Leftrightarrow \int_1^\infty (f(t))^{-\alpha} dt \begin{cases} < \infty ; \\ = \infty . \end{cases} \quad (59)$$

We can reformulate Theorem 5.1 by using the skew-product representation (13) stating:  $\rho_t = \theta_{A(t)}$ , in order to deduce a LIL for the winding process  $\theta_{A(\cdot)}$  for small times.

**Corollary 5.2.** For any non-decreasing function  $f > 0$ ,

$$\limsup_{t \rightarrow 0} \frac{\theta_{A(t)}}{f(t)} = \begin{cases} 0 ; \\ \infty \end{cases} \quad a.s. \Leftrightarrow \int_1^\infty (f(t))^{-\alpha} dt \begin{cases} < \infty ; \\ = \infty . \end{cases} \quad (60)$$

### Proof of Theorem 5.1:

First, we define:

$$h(y) = y^{-2}U(y), \quad y > 0. \quad (61)$$

Then, we consider  $t_n = 2^{-n}$  and we note that (Cauchy's test):

$$I(f) \equiv \int_1^\infty dt h(f(t)) < \infty \iff \sum_{n=1}^\infty t_n h(f(t_n)) < \infty .$$

Using now Lemma 2 from Doney and Maller [DoM02a], because  $\rho$  is symmetric, there exists a positive constant  $c_1$  such that for every  $x > 0$ ,  $t > 0$ ,

$$P(\rho_t \geq x) \leq P\left(\sup_{0 \leq u \leq t} \rho_u \geq x\right) \leq c_1 t h(x) = c_1 t \frac{U(x)}{x^2} . \quad (62)$$

Thus:

$$\sum_{n=1}^\infty P(\rho_{t_{n-1}} \geq f(t_n)) \leq c_1 \sum_{n=1}^\infty t_n \frac{U(f(t_n))}{(f(t_n))^2} .$$



From (58), we have that,

$$\frac{U(f(t_n))}{(f(t_n))^2} \stackrel{t_n \sim 0}{\sim} (f(t_n))^{-\alpha}.$$

Hence, when  $I(f) < \infty$ , from Borel-Cantelli Lemma we have that with probability 1,  $\rho_{t_{n-1}} \leq f(t_n)$  for all  $n$ 's, except for a finite number of them. Now, from a monotonicity argument for  $f$ , if  $t \in [t_n, t_{n-1}]$ , we have that:  $\rho_{t_{n-1}} \leq f(t_n) \leq f(t)$  for every  $t$  sufficiently small. It follows now that  $\lim_{t \rightarrow 0} (\rho_t/f(t)) \leq 1$  a.s. Finally, we remark that as  $I(f) < \infty$ , we also have that  $I(\varepsilon f) < \infty$ , for arbitrarily small  $\varepsilon > 0$  and follows that  $\rho_t/f(t) \rightarrow 0$  a.s.

The proof of the second statement follows from the same kind of arguments. Indeed, using Lemma 2 from Doney and Maller [DoM02a] and the fact that  $\rho$  is symmetric, there exists a positive constant  $c_2$  such that for every  $x > 0$ ,  $t > 0$ ,

$$P\left(\sup_{0 \leq u \leq t} \rho_u \leq x\right) \leq \frac{c_2}{t h(x)}. \quad (63)$$

Hence:

$$\begin{aligned} \sum_{n=1}^{\infty} P(\rho_{t_{n-1}} \leq f(t_{n-1})) &\leq \sum_{n=1}^{\infty} P\left(\sup_{0 \leq u \leq t_{n-1}} \rho_u \leq f(t_{n-1})\right) \\ &\leq \sum_{n=1}^{\infty} \frac{c_2}{t_{n-1} h(f(t_{n-1}))}. \end{aligned}$$

Thus, for  $I(f) = \infty$  (or equivalently  $\sum h(f(t_{n-1})) = \infty$ ), Borel-Cantelli Lemma yields that for every  $n$ , a.s.  $\rho_{t_{n-1}} > f(t_{n-1})$  infinitely often, which finishes the proof.  $\blacksquare$

**Remark 5.3.** For other kinds of LIL for Lévy processes for small times e.g. of the Chung type, see [ADS10] and the references therein.

**Theorem 5.4. (LIL for the angular part of planar stable processes for small times)**

For any non-decreasing function  $f > 0$ ,

$$\limsup_{t \rightarrow 0} \frac{\theta_t}{f(t)} = \begin{cases} 0; \\ \infty \end{cases} \quad a.s. \Leftrightarrow \int_1^{\infty} (f(t))^{-\alpha} dt \begin{cases} < \infty; \\ = \infty. \end{cases} \quad (64)$$

**Proof of Theorem 5.4:**

We use the skew-product representation (13) together with (46), which essentially writes:  $t^{-1}H(t) \xrightarrow[t \rightarrow 0]{a.s.} 1$ . Thus, for every  $\varepsilon, \delta > 0$ , there exists  $t_0 > 0$  such that:

$$P\left(\frac{H(t)}{t} \leq 1 + \varepsilon\right) \geq 1 - \delta, \quad \text{for } t \leq t_0. \quad (65)$$

We define now the setting:

$$\mathcal{K} \equiv \mathcal{K}(\omega) \equiv \left\{ \omega : \frac{H(t)}{t} \leq 1 + \varepsilon \right\}, \quad \text{thus : } \bar{\mathcal{K}} \equiv \bar{\mathcal{K}}(\omega) \equiv \left\{ \omega : \frac{H(t)}{t} \geq 1 + \varepsilon \right\},$$

hence, there exists  $t_0 > 0$  such that: for every  $t \leq t_0$ ,

$$P(\mathcal{K}) \geq 1 - \delta \quad \text{and} \quad P(\overline{\mathcal{K}}) \leq \delta .$$

Hence, choosing  $\delta > 0$  small enough, it suffices to restrict our study in the set  $\mathcal{K}$  and it follows that:

$$\begin{aligned} P\left(\sup_{0 \leq u \leq t} \theta_u > x\right) &= P\left(\sup_{0 \leq u \leq t} \rho_{H(u)} > x\right) = P\left(\left\{\sup_{0 \leq u \leq t} \rho_{H(u)} > x\right\} \cap \mathcal{K}\right) \\ &\leq P\left(\sup_{0 \leq u \leq t} \rho_{u(1+\varepsilon)} > x\right) \end{aligned}$$

Changing now the variables  $\tilde{u} = u(1 + \varepsilon)$ , and invoking (62), there exists another positive constant  $c_3$  such that, for every  $x > 0$  and  $t > 0$ :

$$P\left(\sup_{0 \leq u \leq t} \theta_u > x\right) \leq P\left(\sup_{0 \leq \tilde{u} \leq t(1+\varepsilon)} \rho_{\tilde{u}} > x\right) \leq c_3 t(1 + \varepsilon) \frac{U(x)}{x^2} . \quad (66)$$

Mimicking now the proof of Theorem 5.1, we obtain the first statement.

For the second statement, we use the settings:

$$\mathcal{K}' \equiv \mathcal{K}'(\omega) \equiv \left\{ \omega : \frac{H(t)}{t} \geq 1 - \varepsilon \right\} , \quad \text{thus} : \quad \overline{\mathcal{K}'} \equiv \overline{\mathcal{K}'}(\omega) \equiv \left\{ \omega : \frac{H(t)}{t} \leq 1 - \varepsilon \right\} .$$

Hence, for every  $\varepsilon, \delta > 0$ , there exists  $t_0 > 0$  such that: for every  $t \leq t_0$ ,

$$P(\mathcal{K}') \geq 1 - \delta \quad \text{and} \quad P(\overline{\mathcal{K}'}) \leq \delta .$$

As before, we choose  $\delta > 0$  small enough and we restrict our study in the set  $\mathcal{K}'$ . The proof finishes by repeating the arguments of the proof of Theorem 5.1.  $\blacksquare$

## 6 The planar Brownian motion case

Before starting, we remark that the notations used in this Section are independent from the ones used in the text up to now.

In this Section, we state and give a new proof of the analogue of Theorem 4.4 for the planar Brownian motion case, which is equivalent to a result obtained in [VaY12]. For this purpose, and in order to avoid complexity, we will use the same notation as in the "stable" case. Hence, for a planar BM  $\mathcal{Z}$  starting from a point different  $z_0$  from 0 (without loss of generality, let  $z_0 = 1$ ) and with  $\vartheta = (\vartheta_t, t \geq 0)$  denoting now the (well defined - see eg. [ItMK65]) continuous winding process, we have the skew product representation (see e.g. [ReY99]):

$$\log |\mathcal{Z}_t| + i\vartheta_t \equiv \int_0^t \frac{d\mathcal{Z}_s}{\mathcal{Z}_s} = (\beta_u + i\gamma_u) \Big|_{u=\mathcal{H}_t=\int_0^t \frac{ds}{|\mathcal{Z}_s|^2}} , \quad (67)$$

where  $(\beta_u + i\gamma_u, u \geq 0)$  is another planar Brownian motion starting from  $\log 1 + i0 = 0$ . The Bessel clock  $\mathcal{H}$  plays a key role in many aspects of the study of the winding number

process  $(\vartheta_t, t \geq 0)$  (see e.g. [Yor80]). We shall also make use of the inverse of  $\mathcal{H}$ , which is given by:

$$\mathcal{H}_u^{-1} = \inf\{t \geq 0 : \mathcal{H}(t) > u\} = \int_0^u ds \exp(2\beta_s) = \mathcal{A}_u. \quad (68)$$

Rewriting (67) as:

$$\log |\mathcal{Z}_t| = \beta_{\mathcal{H}_t}; \quad \vartheta_t = \gamma_{\mathcal{H}_t}, \quad (69)$$

we easily obtain that the two  $\sigma$ -fields  $\sigma\{|\mathcal{Z}_t|, t \geq 0\}$  and  $\sigma\{\beta_u, u \geq 0\}$  are identical, whereas  $(\gamma_u, u \geq 0)$  is independent from  $(|\mathcal{Z}_t|, t \geq 0)$ , a fact that is in contrast to what happens in the "stable" case.

**Theorem 6.1.** *The family of processes*

$$(c^{-1/2}\vartheta_{ct}, t \geq 0)$$

*converges in distribution, as  $c \rightarrow 0$ , to a 1-dimensional Brownian motion  $(\gamma_t, t \geq 0)$ .*

*Proof.* We split the proof in two parts:

i) First, repeating the arguments in the proof of Theorem 4.2 with  $\alpha = 2$ , we obtain:

$$\left( \frac{\mathcal{H}(ux)}{u}, x \geq 0 \right) \xrightarrow[u \rightarrow 0]{a.s.} (x, x \geq 0). \quad (70)$$

which also implies the weak convergence:

$$\left( \frac{\mathcal{H}(ux)}{u}, x \geq 0 \right) \xrightarrow[u \rightarrow 0]{(d)} (x, x \geq 0). \quad (71)$$

ii) Using the skew product representation (69) and the scaling property of BM together with (70), we have that for every  $s > 0$ :

$$t^{-1/2}\vartheta_{st} = t^{-1/2}\gamma_{\mathcal{H}(st)} \stackrel{(law)}{=} \sqrt{\frac{\mathcal{H}(st)}{t}}\gamma_1 \xrightarrow[t \rightarrow 0]{a.s.} \sqrt{s}\gamma_1 \stackrel{(law)}{=} \gamma_s, \quad (72)$$

which finishes the proof.

We remark that for part (ii) of the proof, we could also invoke Whitt's Theorem 3.1 concerning the composition function [Whi80], however, the independence in the planar Brownian motion case simplifies the proof.  $\square$   $\square$

From Theorem 6.1 now, with  $T_c^{|\vartheta|} \equiv \inf\{t : |\vartheta_t| = c\}$  and  $T_c^{|\gamma|} \equiv \inf\{t : |\gamma_t| = c\}$ , ( $c > 0$ ), we deduce for the exit time from a cone of planar BM (this result has already been obtained in [VaY12], where one can also find several variants):

**Corollary 6.2.** *The following convergence in law holds:*

$$\left( \frac{1}{c^2} T_c^{|\vartheta|}, x \geq 0 \right) \xrightarrow[c \rightarrow 0]{(law)} (T_x^{|\gamma|}, x \geq 0). \quad (73)$$

**Remark 6.3.** We highlight the different behaviour of the clock  $\mathcal{H}$  for  $t \rightarrow 0$  and for  $t \rightarrow \infty$  (for the second see e.g. [PiY84], followed by [PiY86, LGY86, LGY87], a result which is equivalent to Spitzer's Theorem [Spi58] stated in (1) ), that is:

$$\frac{\mathcal{H}(t)}{t} \xrightarrow[t \rightarrow 0]{a.s.} 1, \quad (74)$$

$$\frac{4\mathcal{H}(t)}{(\log t)^2} \xrightarrow[t \rightarrow \infty]{(law)} T_1 \equiv \inf\{t : \beta_t = 1\}, \quad (75)$$

where the latter follows essentially from the classical Laplace argument:

$$\|\cdot\|_p \xrightarrow{p \rightarrow \infty} \|\cdot\|_\infty.$$

We also remark that, from Remark 4.3, the behaviour of the clock for  $t \rightarrow 0$  is a.s. the same for Brownian motion and for stable processes, whereas it is different for  $t \rightarrow \infty$ . In particular, for  $t \rightarrow \infty$ , compare (49) to (75).

### Acknowledgements

The author S. Vakeroudis is very grateful to Professor M. Yor for the financial support during his stay at the University of Manchester as a Post Doc fellow invited by Professor R.A. Doney.

## References

- [ADS10] F. Aurzada, L. Döring and M. Savov. Small time Chung type LIL for Lévy processes. To appear in *Bernoulli*. ArXiv:1002.0675.
- [BaB04] R. Banuelos and K. Bogdan (2004). Symmetric stable processes in cones. *Potential Anal.*, **21**, p. 263-288.
- [Ber96] J. Bertoin (1996). Lévy Processes. Cambridge University Press, Cambridge.
- [BeD97] J. Bertoin and R.A. Doney (1997). Spitzer's condition for random walks and Lévy Processes. *Ann. Inst. Henri Poincaré*, **33**, p. 167-178.
- [BeW94a] J. Bertoin and W. Werner (1994). Asymptotic windings of planar Brownian motion revisited via the Ornstein-Uhlenbeck process. *Sém. Prob. XXVIII, Lect. Notes in Math.*, **1583**, Springer, Berlin Heidelberg New York, p. 138-152.
- [BeW94b] J. Bertoin and W. Werner (1994). Comportement asymptotique du nombre de tours effectués par la trajectoire brownienne plane. *Séminaire de Probabilités XXVIII, Lecture Notes in Math.*, **1583**, Springer, Berlin Heidelberg New York, p. 164-171.
- [BeW96] J. Bertoin and W. Werner (1996). Stable windings. *Ann. Probab.*, **24**, p. 1269-1279.
- [Bre68] L. Breiman (1968). A delicate law of the iterated logarithm for non-decreasing stable processes. *Ann. Math. Statist.*, **39**, p. 1818-1824; correction **41**, p. 1126.

- [Bur77] D. Burkholder (1977). Exit times of Brownian Motion, Harmonic Majorization and Hardy Spaces. *Adv. in Math.*, **26**, p. 182-205.
- [CPP11] M. E. Caballero, J. C. Pardo, and J. L. Pérez (2011). Explicit identities for Lévy processes associated to symmetric stable processes. *Bernoulli*, **17** (1), p. 34-59.
- [Chy06] O. Chybiryakov (2006). The Lamperti correspondence extended to Lévy processes and semi-stable Markov processes in locally compact groups. *Stochastic Processes and their Applications*, **116**, p. 857-872.
- [DeB90] R.D. De Blassie (1990). The first exit time of a two-dimensional symmetric stable process from a wedge. *Ann. Prob.*, **18**, p. 1034-1070.
- [Don04] R.A. Doney (2004). Small time behaviour of Lévy processes. *Electron. J. Probab.*, **9**, p. 209-229.
- [DoM00] R.A. Doney and R.A. Maller (2000). Random walks crossing curved boundaries: a functional limit theorem, stability and asymptotic distributions for exit positions. *Adv. Appl. Probab.*, **32**, p. 1117-1149.
- [DoM02a] R.A. Doney and R.A. Maller (2002). Stability of the overshoot for Lévy processes. *Ann. Prob.*, **30**, p. 188-212.
- [DoM02b] R.A. Doney and R.A. Maller (2002). Stability and Attraction to Normality for Lévy processes at zero and infinity. *J. Theoretical Probab.*, **15**, p. 751-792.
- [Dur82] R. Durrett (1982). A new proof of Spitzer's result on the winding of 2-dimensional Brownian motion. *Ann. Prob.*, **10**, p. 244-246.
- [Fri64] B.E. Fristedt (1964). The behavior of increasing stable processes for both small and large times. *J. Math. Mech.*, **13**, p. 849-856.
- [Fri67] B.E. Fristedt (1967). Sample function behavior of increasing processes with stationary, independent increments. *Pacific J. Math.*, vol. **21**, n. 1, p. 21-33.
- [GVA86] S.E. Graversen and J. Vuolle-Apiala (1986).  $\alpha$ -self-similar Markov Processes. *Probab. Theory Related Fields*, **71**, p. 149-158.
- [ItMK65] K. Itô and H.P. McKean (1965). Diffusion Processes and their Sample Paths. Springer, Berlin Heidelberg New York.
- [JaS03] J. Jacod and A. N. Shiryaev (2003). Limit theorems for stochastic processes. 2nd ed., Springer-Verlag, Berlin.
- [Khi39] A. Khintchine (1939). Sur la croissance locale des processus stochastiques homogènes à accroissements indépendants. (Russian article and French resume) *Akad. Nauk. SSSR, Izvestia, Ser. Math.*, **3** (5-6), p. 487-508.
- [Kiu80] S.W. Kiu (1980). Semi-stable Markov processes in  $\mathbb{R}^n$ . *Stochastic Processes and their Applications*, **10**, n. 2, p. 183-191.

- [Kyp06] A.E. Kyprianou (2006). *Introductory Lectures on Fluctuations of Lévy Processes with Applications*. Springer, Berlin.
- [Lam72] J. Lamperti (1972). Semi-stable Markov processes I. *Z. Wahr. Verw. Gebiete*, **22**, p. 205-225.
- [LeG92] J.F. Le Gall (1992). Some properties of planar Brownian motion. Cours de l'école d'été de St-Flour XX. *Lecture Notes in Math.*, Vol. **1527**. Springer, Berlin. p. 111-235.
- [LGY86] J.F. Le Gall and M. Yor (1986). Etude asymptotique de certains mouvements browniens complexes avec drift . *Probab. Th. Rel. Fields*, **71** (2), p. 183-229.
- [LGY87] J.F. Le Gall and M. Yor (1987). Etude asymptotique des enlacements du mouvement brownien autour des droites de l'espace. *Probab. Th. Rel. Fields*, **74** (4), p. 617-635.
- [LiW11] M. Liao and L. Wang (2011). Isotropic self-similar Markov processes. *Stochastic Processes and their Applications*, **121**, n. 9, p. 2064-2071.
- [MeY82] P. Messulam and M. Yor (1982). On D. Williams' "pinching method" and some applications. *J. London Math. Soc.*, **26**, p. 348-364.
- [PiY84] J.W. Pitman and M. Yor (1984). The asymptotic joint distribution of windings of planar Brownian motion. *Bull. Amer. Math. Soc.*, **10**, p. 109-111.
- [PiY86] J.W. Pitman and M. Yor (1986). Asymptotic Laws of planar Brownian Motion. *Ann. Prob.*, **14**, p. 733-779.
- [PiY89] J.W. Pitman and M. Yor (1989). Further Asymptotic Laws of planar Brownian Motion. *Ann. Prob.*, **17** (3), p. 965-1011.
- [ReY99] D. Revuz and M. Yor (1999). *Continuous Martingales and Brownian Motion*. 3rd ed., Springer, Berlin.
- [Shi94] Z. Shi (1994). Liminf behaviours of the windings and Lévy's stochastic areas of planar Brownian motion. *Séminaire de Probabilités XXVIII, Lecture Notes in Math.*, **1583**, Springer, Berlin Heidelberg New York, p. 122-137.
- [Shi98] Z. Shi (1998). Windings of Brownian motion and random walks in the plane. *Ann. Probab.*, **26**, n. 1, p. 112-131.
- [Sko91] A.V. Skorohod (1991). *Random Processes with Independent Increments*. Kluwer, Dordrecht.
- [Spi58] F. Spitzer (1958). Some Theorems concerning two-dimensional Brownian Motion. *Trans. Amer. Math. Soc.*, **87**, p. 187-197.
- [Vakth11] S. Vakeroudis (2011). Nombres de tours de certains processus stochastiques plans et applications à la rotation d'un polymère. (Windings of some planar Stochastic Processes and applications to the rotation of a polymer). PhD Dissertation, Université Pierre et Marie Curie (Paris VI), April 2011.

- [Vak11] S. Vakeroudis (2012). On hitting times of the winding processes of planar Brownian motion and of Ornstein-Uhlenbeck processes, via Bougerol's identity. *SIAM Theory of Probability and its Applications*, Vol. **56** (3), pp. 485-507 (originally published in 2011 in *Teor. Veroyatnost. i Primenen.*, Vol. **56** (3), pp. 566-591).
- [VaY12] S. Vakeroudis and M. Yor (2012). Integrability properties and Limit Theorems for the exit time from a cone of planar Brownian motion. To appear in *Bernoulli*. ArXiv:1201.2716.
- [Whi80] W. Whitt (1980). Some useful functions for functional limit Theorems. *Math. Operat. Res.*, **5**, p. 67-85.
- [Wil74] D. Williams (1974). A simple geometric proof of Spitzer's winding number formula for 2-dimensional Brownian motion. University College, Swansea. Unpublished.
- [Yor80] M. Yor (1980). Loi de l'indice du lacet Brownien et Distribution de Hartman-Watson. *Z. Wahrsch. verw. Gebiete*, **53**, p. 71-95.
- [Yor97] M. Yor (1997). Generalized meanders as limits of weighted Bessel processes, and an elementary proof of Spitzer's asymptotic result on Brownian windings. *Studia Scient. Math. Hung.*, **33**, p. 339-343.