



HAL
open science

Limiting laws of supercritical branching random walks

Julien Barral, Rémi Rhodes, Vincent Vargas

► **To cite this version:**

Julien Barral, Rémi Rhodes, Vincent Vargas. Limiting laws of supercritical branching random walks. Comptes rendus de l'Académie des sciences. Série I, Mathématique, 2012. hal-00682212v2

HAL Id: hal-00682212

<https://hal.science/hal-00682212v2>

Submitted on 26 May 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Limiting laws of supercritical branching random walks

Julien Barral ^{*}, Rémi Rhodes [†], Vincent Vargas ^{‡§}

May 26, 2012

Abstract

In this note, we make explicit the limit law of the renormalized supercritical branching random walk, giving credit to a conjecture formulated in [5] for a continuous analogue of the branching random walk. Also, in the case of a branching random walk on a homogeneous tree, we express the law of the corresponding limiting renormalized Gibbs measures, confirming, in this discrete model, conjectures formulated by physicists about the Poisson-Dirichlet nature of the jumps in the limit, and precisising the conjecture by giving the spatial distribution of these jumps.

1. Introduction and notations

We consider a branching random walk on the real line \mathbb{R} . Initially, a particle sits at the origin. Its children form the first generation; their displacements from the origin correspond to a point process on the line. These children have children of their own (who form the second generation), and behave relative to their respective positions like independent copies of the same point process L . And so on. We write $|u| = n$ if an individual u is in the n -th generation, and denote its position $V(u)$ (and $V(e) = 0$ for the initial ancestor). We define the (logarithmic) moment generating function

$$\forall t \geq 0, \quad \psi(t) = \ln \mathbb{E} \left[\sum_{|u|=1} e^{-tV(u)} \right] \in]-\infty; +\infty],$$

^{*}Université Paris 13, Institut Galilée, LAGA, UMR CNRS 7539, 99 rue Jean-Baptiste Clément 93430 Villetaneuse, France.

[†]Université Paris-Dauphine, Ceremade, UMR 7534, Place du marechal de Lattre de Tassigny, 75775 Paris Cedex 16, France.

[‡]Université Paris-Dauphine, Ceremade, UMR 7534, Place du marechal de Lattre de Tassigny, 75775 Paris Cedex 16, France.

[§]The last two authors are partially supported by the CHAMU project (ANR-11-JCJC).

where \mathbb{E} denotes expectation with respect to \mathbb{P} , the law of the branching random walk.

The numbers of particles N_n , $n \geq 1$, in successive generations form a Galton-Watson process, which is supercritical if and only if $\ln \mathbb{E}(N_1) = \psi(0) > 0$, a condition we assume throughout, together with normalizing condition $\psi(1) = 0$. Then the sequence

$$W_n = \sum_{|u|=n} e^{-V(u)}, \quad n = 0, 1, 2, \dots \quad \left(\sum_{\emptyset} := 0 \right) \quad (1)$$

is a nonnegative martingale, which converges almost surely to a limit W_∞ . Moreover, $\mathbb{P}(W_\infty > 0) > 0$ if and only if $\psi'(1^-) < 0$ and $\mathbb{E}[(W_1) \ln^+(W_1)] < \infty$. In that case, the convergence holds in L^1 norm too and $W_\infty > 0$ almost surely on the set of non-extinction of the Galton-Watson process. When $\psi'(1^-) \geq 0$, $W_\infty = 0$ almost surely, and this gives rise to the issue of finding natural renormalizations of $(W_n)_{n \geq 1}$. It is straightforward to see that this reduces to considering the renormalization of

$$W_{n,\beta} = \sum_{|u|=n} e^{-\beta V(u)} \quad (\beta \geq 1), \quad (2)$$

where now $\psi'(1^-) = 0$. We further introduce the derivative martingale:

$$Z_n = \sum_{|u|=n} V(u) e^{-V(u)} \quad (3)$$

We reinforce our assumptions. Let $\widetilde{W}_1 = W_1 + \sum_{|u|=1} V(u) e^{-V(u)}$. We assume

$$a) \quad \psi''(1) < \infty, \quad b) \quad \mathbb{E} \left[\widetilde{W}_1 \ln_+(\widetilde{W}_1)^3 \right] < +\infty. \quad (4)$$

It is known that under (4) the martingale $(Z_n)_{n \geq 1}$ converges almost surely towards a random variable Z_∞ , which is strictly positive conditionally on the survival of the system (see [6, 1]). In what follows, we investigate the limiting laws of the renormalized sequence $(W_{n,\beta})_n$ for $\beta > 1$. In the following theorem, a,b) of (4) are needed for the convergence in law as it is shown in [13]. We identify the limit under additional assumptions.

Theorem 1. *Assume that L is non-lattice, $\psi(0) > 0$, $\psi(1) = \psi'(1^-) = 0$ and (4). Let $\beta > 1$. The random variables $(n^{\frac{3}{2}\beta} W_{n,\beta})_n$ converge in law towards a non trivial random variable $W_{\infty,\beta}$. Moreover, if there exists $\delta > 1$ such that $\mathbb{E} \left[(W_{1,\beta} + N_1)^\delta \right] < +\infty$, the law of $W_{\infty,\beta}$ is, up to a multiplicative constant c , that of a stable subordinator T_β of index $1/\beta$ taken at an independent time Z_∞ :*

$$W_{\infty,\beta} \stackrel{\text{law}}{=} cT_\beta(Z_\infty).$$

We notice that this result is obtained in [16] in the special case where the Galton-Watson-tree is homogeneous and the variables $V(u)$, $|u| = 1$, are Gaussian and independent.

2. The case of Random multiplicative cascades

We point out that Theorem 1 has direct applications to random multiplicative cascades since random multiplicative cascades can be seen as branching random walks where the number of children for each particle has been set to two. Random multiplicative cascades were introduced by B. Mandelbrot in [14] as a toy model for energy dissipation in a turbulent flow. More precisely, the closed dyadic subintervals of $[0, 1]$ are naturally encoded by the nodes of the binary tree, i.e., the finite words over the alphabet $\{0, 1\}$ via the mapping $u = u_1 \cdots u_n \mapsto I_u = [\sum_{k=1}^n u_k 2^{-k}, 2^{-n} + \sum_{k=1}^n u_k 2^{-k}]$.

In the notations of the previous section (recall that $\psi(1) = 0$ and notice that $\psi(0) = \log(2) > 0$), let us assume that $V(0)$ and $V(1)$ are i.i.d., and define a sequence of random measures on $[0, 1]$ as follows. Let dx denote the Lebesgue measure on $[0, 1]$. We define for $n \geq 1$ the random measure

$$\mu_n(dx) = 2^n e^{-V(u(x))} dx, \quad \text{where}$$

where $u(x)$ is the unique element of $\{0, 1\}^n$ such that $x \in I_{u(x)}$ when x is not dyadic. We recall the following theorem of [11] first conjectured in the seminal work [14]:

Theorem 2 (Kahane). *The weak-star limit $\mu(dx) = \lim_{n \rightarrow \infty} \mu_n(dx)$ exists almost surely. Moreover, μ is almost surely non trivial if and only if $\psi'(1^-) < 0$.*

We are interested in the situation where $\psi'(1^-) \geq 0$ in which case the limiting measure vanishes. More precisely, as suggested in [14], we want to find a renormalization sequence $(\lambda_n)_{n \geq 1}$ such that $\nu_n(dx) = \lambda_n \mu_n(dx)$ converges weakly, at least in law. In accordance with the previous section we assume now that (4) holds and $\psi'(1) = 0$. We define for $\beta > 0$:

$$\mu_{n,\beta}(dx) = 2^n e^{-\beta V(u(x))} dx.$$

When $0 < \beta < 1$, the situation boils down to the non-degenerate (classical) situation. For $\beta = 1$, this corresponds to the so-called boundary case which was studied in [2]. One can deduce from [2] and [6] that $n^{1/2} \mu_{n,1}$ converges in probability to a positive multiple of a positive statistically self-similar measure obtained as the almost sure weak limit of $-\frac{d\mu_{n,\beta}}{d\beta}|_{\beta=1}(dx)$, call it $\tilde{\mu}_\infty$ (the same convergence in law when $V(0)$ is Gaussian can be deduced from [16]).

We are especially interested in the low temperature case $\beta > 1$. From Theorem 1, we deduce:

Corollary 3. *Assume $\psi(1) = \psi'(1) = 0$ and (4). Let $\beta > 1$. Suppose there exists $\delta > 1$ such that $\psi(\delta\beta) < \infty$. The random measures $(n^{\frac{3}{2}\beta} \mu_{n,\beta}(dx))_n$ weakly converge in law towards a random Borel measure $\mu_{\infty,\beta}(dx)$ on $[0, 1]$. The law of the measure $\mu_{\infty,\beta}(dx)$ is given by the derivative (in the sense of distributions) of the function $t \rightarrow cT_\beta(\tilde{\mu}_\infty[0, t])$ where $c > 0$ is some positive constant and T_β is a stable subordinator of index $1/\beta$ independent from $\tilde{\mu}_\infty(dx)$.*

We will denote this limiting law $cT_\beta(\tilde{\mu}_\infty(dx))$. We stress that this implies the following result on renormalized Gibbs measures:

Corollary 4. *Under the assumptions of Corollary 3, we have the following convergence in law (for the weak convergence of measures):*

$$\lim_{n \rightarrow \infty} \frac{\mu_{n,\beta}(dx)}{\mu_{n,\beta}([0, 1])} = \frac{T_\beta(\tilde{\mu}_\infty(dx))}{T_\beta(\tilde{\mu}_\infty[0, 1])}.$$

3. The continuous case: Gaussian multiplicative Chaos

In this section, we discuss the continuous analogue of the Random multiplicative cascades with lognormal weights: Gaussian multiplicative Chaos. One can work with other weights than lognormal but we stick to this case for the sake of simplicity. Gaussian multiplicative Chaos was introduced by J.P. Kahane in [10]. Gaussian multiplicative Chaos and lognormal Random multiplicative cascades bear striking similarities; in particular, one can introduce a continuous analogue to the discrete star equation (cf. [3, 15]) which plays a key role in the identification of the limit in the above theorems. In some sense, there is a correspondance between both models in the sense that roughly all theorems in the discrete case have continuous analogues. In the paper [5] (section 6 perspectives), the authors conjectured an analogue to corollary 4 for Gaussian multiplicative Chaos. In fact, theorem 1 leads to an even stronger version of the conjecture in [5].

The above result on renormalized Gibbs measures, i.e. corollary 4, and the conjecture in [5] on the limit of a continuous renormalized Gibbs measure share a common feature which is proved or conjectured to hold for many supercritical Gibbs measures (also called the glassy phase in the physics litterature) : the limiting object is a purely atomic measure whose jumps are distributed according to a Poisson-Dirichlet distribution, i.e. the ordered jumps of a renormalized stable subordinator. This universal feature (proved for instance in the case of the REM, etc.) has been conjectured by physicists (see [8]). Corollary 4 and the corresponding conjecture in [5] are stronger than the conjecture of physicists since it also provides the spatial distribution of the jumps.

4. Proofs

4.1 Proof of Theorem 1

It is proved in [13] that the sequence $(n^{\frac{3}{2}\beta} W_{n,\beta})_n$ converges in law towards a random variable $W_{\infty,\beta}$. This random variable is non trivial since

$$n^{\frac{3}{2}\beta} W_{n,\beta} \geq \left(n^{3/2} \max_{|u|} e^{-V(u)} \right)^\beta \geq \left(n^{3/2} e^{-\min_{|u|} V(u)} \right)^\beta = \left(e^{-\left(\min_{|u|} V(u) - \frac{3}{2} \ln n \right)} \right)^\beta.$$

It is proved in [1] that the last quantity $\min_{|u|} V(u) - \frac{3}{2} \ln n$ converges in law as $n \rightarrow \infty$ towards a non trivial limit. So $W_{\infty, \beta}$ cannot be 0. It just remains to identify the limiting law $W_{\infty, \beta}$. To that purpose, it is straightforward to check that $W_{\infty, \beta}$ satisfies the star equation ([14]):

$$W_{\infty, \beta} \stackrel{\text{law}}{=} \sum_{|u|=1} e^{-\beta V(u)} W_{\infty, \beta}^{(u)},$$

where conditionally on $(u, V(u), |u| = 1)$ the family $(W_{\infty, \beta}^{(u)})_{|u|=1}$ are independent copies of $W_{\infty, \beta}$. This equation is carefully studied in [12]. Since $t \mapsto \psi_\beta(t) = \ln \mathbb{E}[\sum_{|u|=1} e^{-\beta t V(u)}]$ vanishes for $t = \frac{1}{\beta}$ with $\psi'_\beta(\frac{1}{\beta}) = 0$, the result follows from [12] (see also [9] for the case of homogenous trees). \square

4.2 Convergence of the random measures of section 2

It is convenient to first consider measures on the symbolic space $\{0, 1\}^{\mathbb{N}^+}$. For each finite dyadic word u we denote by $[u]$ the cylinder consisting of infinite words with u as prefix and define $Q(u) = e^{-V(u)}$. $\{0, 1\}^{\mathbb{N}^+}$ is endowed with the Borel σ -field generated by the cylinders taken as a basis of open sets. Denote the dyadic coding by $\pi : \{0, 1\}^{\mathbb{N}^+} \rightarrow [0, 1]$: $\pi(x_1 x_2 \dots) = \sum_{k \geq 1} x_k 2^{-k}$. Also denote by λ the uniform measure on $\{0, 1\}^{\mathbb{N}^+}$.

For $\beta \geq 1$ and $n \geq 1$, let $\nu_{n, \beta}$ stand for the measure whose density with respect to λ is given by $2^n Q(u)^\beta$; we have $\mu_{n, \beta} = \nu_{\beta, n} \circ \pi^{-1}$. Also, let $\tilde{\nu}_n$ be the signed measure whose density with respect to λ is given by $-2^n Q(u) \log Q(u)$ over each cylinder $[u]$ of generation n ; we have $\tilde{\nu}_n \circ \pi^{-1}(dx) = -\frac{d\mu_{n, \beta}}{d\beta}|_{\beta=1}(dx)$. It follows from [7] that, almost surely, $\tilde{\nu}_n$ is ultimately positive and converges weakly to a positive measure $\tilde{\nu}_\infty$; consequently $-\frac{d\mu_{n, \beta}}{d\beta}|_{\beta=1}(dx)$ weakly converges to $\tilde{\mu}_\infty(dx) = \tilde{\nu}_\infty \circ \pi^{-1}(dx)$.

Let $N_{\tilde{\nu}_\infty}^{(\beta)}$ be a positive Borel random measure on $\{0, 1\}^{\mathbb{N}^+} \times \mathbb{R}_+^*$, whose law conditionally on $\tilde{\nu}_\infty$ is that of a Poisson point measure with intensity $\beta \Gamma(1/\beta) \frac{\tilde{\nu}_\infty(dx) dz}{z^{1+1/\beta}}$.

Then define the random measure $\tilde{\nu}_{\infty, \beta}(A) = \int_A \int_{\mathbb{R}_+^*} z N_{\tilde{\nu}_\infty}^{(\beta)}(dx, dz)$ and assume that T_β is normalized so that its Laplace exponent is $-\theta^{1/\beta}$. For all continuous function $f : \{0, 1\}^{\mathbb{N}^+} \rightarrow \mathbb{R}$, $|n^{3\beta/2} \nu_{n, \beta}(f)| \leq n^{3\beta/2} \|\nu_{n, \beta}\| \|f\|_\infty$ ($W_{n, \beta} = \|\nu_{n, \beta}\|$). Since the laws of the variables $n^{3\beta/2} \nu_{n, \beta}(\{0, 1\}^{\mathbb{N}^+})$, $n \geq 1$ form a tight sequence (Theorem 1), so do the laws of the random measures $n^{3\beta/2} \nu_{n, \beta}$, $n \geq 1$, on the compact set $\{0, 1\}^{\mathbb{N}^+}$ for the weak convergence of measures. Let us show that the unique limit point of this sequence is the law of $c\tilde{\nu}_{\infty, \beta}$, for some $c > 0$. Since the linear combinations of the indicator functions of cylinders generate the space of real-valued continuous functions on $\{0, 1\}^{\mathbb{N}^+}$, it is enough to show that for some $c > 0$, for each $p \geq 1$, $(n^{3\beta/2} \nu_{n, \beta}([u]))_{u \in \{0, 1\}^p}$ converges in law to $c(\tilde{\nu}_{\infty, \beta}([u]))_{u \in \{0, 1\}^p}$ as $n \rightarrow \infty$. Using the self-similarity of the construction and Theorem 1, it is not difficult to see that $(n^{3\beta/2} \nu_{n, \beta}([u]))_{u \in \{0, 1\}^p}$ converges in law to $(Q(u)^\beta W_{\infty, \beta}(u))_{u \in \{0, 1\}^p}$, where

$(W_{\infty,\beta}(u))_{u \in \{0,1\}^p}$ is independent of $(Q(u))_{u \in \{0,1\}^p}$ and its components are independent copies of $W_{\infty,\beta}$. On the other hand, a calculation combining standard properties of Poisson measures with the fact that $(\tilde{\nu}_{\infty}([u]))_{u \in \{0,1\}^p} = (Q(u)Z_{\infty}(u))_{u \in \{0,1\}^p}$, where $(Z_{\infty}(u))_{u \in \{0,1\}^p}$ is independent of $(Q(u))_{u \in \{0,1\}^p}$ and its components are independent copies of Z_{∞} , shows that the Laplace transform of $(\nu_{\infty,\beta}([u]))_{u \in \{0,1\}^p}$ equals that of $(Q(u)^{\beta}T_{\beta}^{(u)}(Z_{\infty}(u)))_{u \in \{0,1\}^p}$, where the $T_{\beta}^{(u)}$ are independent copies of T_{β} , and also are independent of the $(V(u), Z_{\infty}(u))_{u \in \{0,1\}^p}$. Due to Theorem 1, this yields the desired equality in law. It follows that $n^{3\beta/2}\mu_{n,\beta}$ converges weakly in law to $c\tilde{\mu}_{\infty,\beta} := c\tilde{\nu}_{\infty,\beta} \circ \pi^{-1}$. Moreover, the measure $\tilde{\nu}_{\infty}$ assigns no mass to the countable set of points of $\{0,1\}^{\mathbb{N}^+}$ encoding dyadic points (see [4]), hence so does $\tilde{\nu}_{\infty,\beta}$, and after projection so do $\tilde{\mu}_{\infty}$ and $\tilde{\mu}_{\infty,\beta}$ with dyadic numbers. The previous information are enough to get the equality in law, for each $p \geq 1$, of $(\tilde{\mu}_{\infty,\beta}(I_u))_{u \in \{0,1\}^p}$ and that of $(T_{\beta}(\tilde{\mu}_{\infty})(I_u))_{u \in \{0,1\}^p}$, and then the equality in law of $\tilde{\mu}_{\infty,\beta}(dx)$ and $T_{\beta}(\tilde{\mu}_{\infty}(dx))$, hence the desired conclusion. \square

Acknowledgements: The authors are thankful to J.P. Bouchaud and Alberto Rosso for useful discussions and to Yueyun Hu for orienting them to Thomas Madaule's paper.

References

- [1] E. Aidekon, Convergence of the minimum of a branching random walk, arXiv:1107.2543.
- [2] E. Aidekon, Z. Shi, The Seneta-Heyde scaling for the branching random walk, arXiv:1102.0217.
- [3] R. Allez, R. Rhodes, V. Vargas, Lognormal scale invariant random measures, to appear in *Probability Theory and Related Fields*.
- [4] J. Barral, Continuity of the multifractal spectrum of a random statistically self-similar measure, *J. Theoretic. Probab.*, **13** (2000), 1027–1060.
- [5] J. Barral, X. Jin, R. Rhodes, V. Vargas, Gaussian multiplicative chaos and KPZ duality, arXiv:1202.5296.
- [6] J.D. Biggins, A.E. Kyprianou, Measure change in multitype branching, *Adv. in Appl. Proba.*, **36**(2) (2004), 544-581.
- [7] J.D. Biggins, A.E. Kyprianou, Seneta-Heyde norming in the branching random walk, *Ann. Proba.* **25** (1997), 337-360.
- [8] D. Carpentier, P. Le Doussal, Glass transition for a particle in a random potential, front selection in nonlinear renormalization group, and entropic phenomena in Liouville and Sinh-Gordon models, *Phys. Rev. E* **63** (2001), 026110.

- [9] R. Durrett, T.M. Liggett, Fixed points of the smoothing transformation, *Probability Theory and Related Fields*, **64** (1983), n. 3, 275-301.
- [10] J.-P. Kahane, Sur le chaos multiplicatif, *Ann. Sci. Math. Québec*, **9** no.2 (1985), 105-150.
- [11] J.-P. Kahane, J. Peyrière, Sur certaines martingales de Benoit Mandelbrot, *Advances in Math.* **22** (1976), no. 2, 131-145.
- [12] Q.S. Liu, Fixed points of a generalized smoothing transformation and applications to the branching random walk, *Adv. Appl. Probab.* **30** (1998), 85-112.
- [13] T. Madaule, Convergence in law for the branching random walk seen from its tip, arXiv:1107.2543.
- [14] B.B. Mandelbrot, Multiplications aléatoires itérées et distributions invariantes par moyenne pondérée aléatoire, I and II. *Comptes Rendus* (Paris): **278A** (1974), 289-292 and 355-358.
- [15] R. Rhodes, J. Sohier, V. Vargas, \star -scale invariant random measures, arXiv:1201.5219.
- [16] C. Webb, Exact asymptotics of the freezing transition of a logarithmically correlated random energy model, *J. Stat. Phys.*, **145** (2011), 1595–1619.