



HAL
open science

A three-species model explaining cyclic dominance of pacific salmon

Christian Guill, Barbara Drossel, Wolfram Just, Eddy Carmack

► **To cite this version:**

Christian Guill, Barbara Drossel, Wolfram Just, Eddy Carmack. A three-species model explaining cyclic dominance of pacific salmon. *Journal of Theoretical Biology*, 2011, 276 (1), pp.16. 10.1016/j.jtbi.2011.01.036 . hal-00682411

HAL Id: hal-00682411

<https://hal.science/hal-00682411>

Submitted on 26 Mar 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author's Accepted Manuscript

A three-species model explaining cyclic dominance of pacific salmon

Christian Guill, Barbara Drossel, Wolfram Just, Eddy Carmack

PII: S0022-5193(11)00062-2
DOI: doi:10.1016/j.jtbi.2011.01.036
Reference: YJTBI6348



www.elsevier.com/locate/jtbi

To appear in: *Journal of Theoretical Biology*

Received date: 29 July 2010
Revised date: 21 January 2011
Accepted date: 25 January 2011

Cite this article as: Christian Guill, Barbara Drossel, Wolfram Just and Eddy Carmack, A three-species model explaining cyclic dominance of pacific salmon, *Journal of Theoretical Biology*, doi:[10.1016/j.jtbi.2011.01.036](https://doi.org/10.1016/j.jtbi.2011.01.036)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A three-species model explaining cyclic dominance of pacific salmon

Christian Guill^a, Barbara Drossel^a, Wolfram Just^b, Eddy Carmack^c^a*Institut für Festkörperphysik, TU Darmstadt, Hochschulstrasse 6, 64289 Darmstadt, Germany*^b*School of Mathematical Sciences, Queen Mary University of London, Mile End Road, London E14NS, UK*^c*Institute of Ocean Sciences, 9860 West Saanich Road Sidney B.C. V8L 4B2 Canada*

Abstract

The four-year oscillations of the number of spawning sockeye salmon (*Oncorhynchus nerka*) that return to their native stream within the Fraser River basin in Canada are a striking example of population oscillations. The period of the oscillation corresponds to the dominant generation time of these fish. Various - not fully convincing - explanations for these oscillations have been proposed, including stochastic influences, depensatory fishing, or genetic effects. Here, we show that the oscillations can be explained as an attractor of the population dynamics, resulting from a strong resonance near a Neimark Sacker bifurcation. This explains not only the long-term persistence of these oscillations, but also reproduces correctly the empirical sequence of salmon abundance within one period of the oscillations. Furthermore, it explains the observation that these oscillations occur only in sockeye stocks originating from large oligotrophic lakes, and that they are usually not observed in salmon species that have a longer generation time.

Key words: cyclic dominance, population oscillations, single-generation cycles, Neimark Sacker bifurcation, strong resonance

*Corresponding author

Email addresses: guill@fkp.tu-darmstadt.de (Christian Guill), drossel@fkp.tu-darmstadt.de (Barbara Drossel), w.just@qmul.ac.uk (Wolfram Just), Eddy.Carmack@dfo-mpo.gc.ca (Eddy Carmack)

1. Introduction

2 In many ecological systems distinct population oscillations are known, with their specific dynamical
 4 characteristics often captured by simple generic models. Among these are the spatial synchronisa-
 6 tion of the lynx-hare oscillations in Canada [1, 2, 3], the chaotic oscillations of boreal rodents in
 8 Fennoscandia [4], or the cyclic outbreak dynamics of the spruce budworm [5, 6]. The four-year os-
 10 cillations of sockeye salmon (*Oncorhynchus nerka*) in the Fraser River basin in Canada are another
 12 well-documented example of such large-scale population oscillations [7, 8, 9]. They are a typical
 14 example of single-generation cycles [10, 11]. Every fourth year (which is the dominant generation
 16 time) the abundance of these fish is at very high levels, reaching several million fish in some spawning
 populations, but drops to numbers between several hundred and a few ten thousand individuals in
 the following years (hence the term *cyclic dominance*). Different stocks can have their population
 maximum in different years (Fig. 1 and Appendix). The oscillations were reported as early as the
 19th century and are evident for instance in the extremely high catches by fisheries every fourth year
 [12]. This both economically and conservationally important phenomenon has been ascribed either to
 transient effects or to stochastic influences [13], to depensatory predation [14], to fishing [15], or to
 genetic effects [16, 17], but all of these explanations are still not fully convincing [16, 9].




Figure 1: **Empirical time series of sockeye salmon abundance.** Total escapement (number of male and female adults that escape the fisheries) of the late Shuswap Lake run. Shuswap lake is one of the most important rearing lakes of juvenile sockeye salmon. Data for 5 other lakes are provided in the Appendix.

The sockeye salmon return to spawn in their native stream or lake in late summer and then die,
 18 which means that the salmon generations do not overlap. The hatched fry migrate downstream in the
 20 following spring and feed for one season in large freshwater lakes, before they migrate to the ocean,
 22 where they spend the next two and a half years. The carcasses of the adult salmon are decomposed
 and provide an important phosphorus input into the rearing lakes of the fry. Since the oscillations
 of sockeye salmon originating from different lakes are not in phase, we have clear evidence that the
 24 relevant processes causing the phenomenon of cyclic dominance take place in the rearing lakes rather
 than in the ocean.

2. Model and results of computer simulations

26 It is our aim to develop a generic model capturing only the essential mechanisms required for the
 occurrence of cyclic dominance. As such, the model is kept very simple and is not designed to
 28 quantitatively predict the population dynamics of all species in the corresponding lake ecosystems.

The model uses standard continuous population dynamics equations for sockeye fry, $s_n(t)$, their predator (e.g. rainbow trout), $p_n(t)$, and their zooplankton food (mainly daphnia), $z_n(t)$, during the growth season from spring ($t = 0$) to fall ($t = T$) in year n , combined with a rule for calculating the three population sizes at the beginning of the next season as a function of the population sizes at the end of the previous season(s). The sockeye fry population at the end of a season gives rise to the number of spawning adults 3 years later, which in turn determines the number of sockeye fry in the following spring, and represents a nutrient input that affects the carrying capacity of the zooplankton. A small fraction of the salmon stays in the ocean for one more year and mature at age 5, thus causing a mixing between the four brood lines of a spawning population. An even smaller fraction matures at age 3, but since these fish are predominantly small-sized males [9] that do not influence the number of fertilised eggs, they are neglected in this study.

The general structure of the model, which combines continuous population dynamics during the growth season with a discrete update scheme capturing the seasonally driven migration and reproduction of the species of interest, has also been applied in a modelling study of the dynamics of fish preying on zooplankton [18]. Here, however, we have chosen a much simpler form of the continuous dynamics which is sufficient to generate cyclic dominance. It is given by the following equations of motion:

$$\begin{aligned}
 \frac{d}{dt}s_n(t) &= \lambda \cdot a_{sz} \frac{z_n(t) \cdot s_n(t)}{1 + c_s \cdot s_n(t) + z_n(t)} \\
 &\quad - a_{ps} \frac{s_n(t) \cdot p_n(t)}{1 + c_p \cdot p_n(t) + s_n(t)} - d_s \cdot s_n(t) \\
 \frac{d}{dt}z_n(t) &= z_n(t) \cdot \left(1 - \frac{z_n(t)}{K_n}\right) \\
 &\quad - a_{sz} \frac{z_n(t) \cdot s_n(t)}{1 + c_s \cdot s_n(t) + z_n(t)} \\
 \frac{d}{dt}p_n(t) &= \lambda \cdot a_{ps} \frac{s_n(t) \cdot p_n(t)}{1 + c_p \cdot p_n(t) + s_n(t)} - d_p \cdot p_n(t)
 \end{aligned} \tag{1}$$

The variables s_n , z_n , and p_n are biomass densities that are made dimensionless. The parameters in these equations and their numerical values used in the computer simulations are as follows. $\lambda = 0.85$ denotes the assimilation efficiency of ingested prey biomass of carnivores [19]. The maximal per unit mass ingestion rates of salmon, $a_{sz} = 10$, and predators, $a_{ps} = 1.6$, scale allometrically with bodymass according to a power law with exponent $-1/4$ [20]. We assume the predators to be approximately 1500 times heavier than the sockeye fry, which fixes the ratio a_{sz}/a_{ps} . The parameters $d_s = 1$ and $d_p = 0.16$ represent biomass loss due to respiration and mortality; they also scale allometrically with body mass. The feeding terms include saturation at high prey densities, and a predator interference term in the denominator with interference strengths set to $c_s = 1$ and $c_p = 0.2$. This Beddington functional response [21] has been chosen since it describes consumer-resource interactions more accurately than the more frequently used Holling type II form [22]. Similar predator interference terms are also used by other modellers [23].

The matching conditions used to determine the biomasses of the species at the beginning of the next

season from their values at the end of the previous season(s) are given by

$$\begin{aligned} s_{n+1}(0) &= \gamma((1 - \epsilon)s_{n-3}(T) + \epsilon s_{n-4}(T)) \\ z_{n+1}(0) &= K_{n+1} \\ p_{n+1}(0) &= p_n(T), \end{aligned} \tag{2}$$

52 with ϵ the proportion of surviving sockeye that return to their native lakes at the age of 5 to spawn
and die. The parameter γ summarises survival from smolt to adult fish (including ocean survivability
54 and loss to fisheries), spawning success, and egg to fry survival. The various factors are estimated
from empirical data [15, 24, 25] and yield $\gamma = 0.4$.

According to empirical observations [16], the zooplankton level at the end of one year has no effect
on the following year. However, its carrying capacity K_{n+1} is a function of the nutrients provided by
the adult salmon and thus a function of the initial fry biomass of that year [26]:

$$K_{n+1} = K_0 + \left(\kappa \frac{s_{n+1}(0)}{\kappa_0 + s_{n+1}(0)} \right). \tag{3}$$

56 (For a review on the importance of salmon-derived nutrients for freshwater ecosystems, see [27] and
citations therein.) Consumer dependent resource productivity is certainly a rather uncommon phe-
58 nomenon. In the lake ecosystems under consideration, it provides an important positive feedback
to the system, but it does not seem to be crucial for the mathematical mechanism underlying the
60 oscillations [28]. The three parameters determining the carrying capacity K_n of the zooplankton in
year n are set to $K_0 = 15$, $\kappa = 10$, and $\kappa_0 = 2$. The numerical values of these parameters are difficult
62 to determine from empirical data. However, since the nutrient concentration in the brood lakes nearly
doubles in years following dominant sockeye returns compared to years following non-dominant sock-
64 eye returns [26], it seems reasonable that the constant fraction of the carrying capacity, K_0 , and the
maximum of the variable fraction, κ , are of the same order of magnitude.

66 Fig. 2a shows a time series of the biomass of the sockeye fry at the end of the growth season. The first
300 years are cut off to show only the stable periodic oscillation with one strong year followed by one
68 intermediate year and two weak years, just as in the empirical data of most sockeye stocks showing
cyclic dominance (Fig. 1 and Appendix). When the simulation parameters are chosen differently (e.g.,
70 $c_s = 0.2$), the difference between the strong and the intermediate year may become less pronounced.
By additionally increasing K_0 and c_p , their order may even become reversed. This is also observed in
72 the empirical data of some stocks.

When a parameter is varied, for instance the constant fraction of the zooplankton carrying capacity,
74 K_0 , the dynamical pattern may change. Fig. 3 shows the biomass $s_n(T)$ of the sockeye fry at the
end of the season for different values of K_0 , from year $t = 5000$ to year 5200. For small K_0 , all
76 sockeye lines are equally strong (only one point is visible for each value of K_0). This means that
the dynamics reaches a fixed point and that there is no cyclic dominance. With increasing K_0 , the
78 fixed point eventually becomes unstable. First a bifurcation to quasiperiodic behaviour occurs (the
attractor contains infinitely many points for each value of K_0), and then the frequency of the oscillation
80 becomes locked at 4. Now only four points are visible for each value of the bifurcation parameter,
indicating periodic oscillations.




Figure 2: **Simulated time series.** Attractor of sockeye salmon dynamics generated by computation of the time-continuous model, showing oscillations with period 4 and the same pattern in the sequence of salmon abundance as in the empirical data. Only the biomasses at the end of each growth season are shown. a: deterministic model, b: dynamics with up to 50 percent fluctuations in the survivability of the sockeye salmon in the ocean (proportional to γ), see *Discussion*. Parameter values for these simulations are as explained in section 2.




Figure 3: **Bifurcation diagram of the biomass of the sockeye fry at the end of the season.** The bifurcation parameter is the constant fraction of the zooplankton carrying capacity K_0 . For each value of K_0 , 200 consecutive data points are plotted.

82 **3. Bifurcation analysis**

In order to understand and interpret these observations, we first note that the continuous population
 84 dynamics during the season, together with the matching conditions applied between two seasons, can
 be viewed as a discrete map (a so-called Poincaré map), giving the sockeye and predator biomasses
 86 at the end of one year as function of the biomasses at the end of the previous years.

To obtain this map, one first has to integrate the dynamical equations over one season, giving $s_n(T)$
 88 and $p_n(T)$ as a function of $s_n(0)$ and $p_n(0)$. The zooplankton can be eliminated since the initial
 condition of the zooplankton depends on $s_n(0)$ only. Next, one expresses $s_{n+1}(0)$ and $p_{n+1}(0)$ as a
 90 function of $s_{n-3}(T)$, $s_{n-4}(T)$ and $p_n(T)$, using the matching conditions.

The mechanism which generates the population oscillation is based on the nature of the instability of
 the stationary state $s_n(T) = s^*(T)$, $p_n(T) = p^*(T)$ of our system. The corresponding bifurcation can
 be investigated in terms of a linear stability analysis. Close to the fixed point, the dynamics can be
 approximated by linear terms. Denoting the distance of the biomasses from their fixed point value by
 $\delta s_n = s_n(T) - s^*(T)$ etc, the linear approximation of this map has the form

$$\begin{pmatrix} \delta s_{n+1} \\ \delta s_n \\ \delta s_{n-1} \\ \delta s_{n-2} \\ \delta s_{n-3} \\ \delta p_{n+1} \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & m_s & \epsilon_1 & -a \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & b & c & m_p \end{pmatrix} \begin{pmatrix} \delta s_n \\ \delta s_{n-1} \\ \delta s_{n-2} \\ \delta s_{n-3} \\ \delta s_{n-4} \\ \delta p_n \end{pmatrix} \quad (4)$$

with positive parameters $m_s, m_p, a, b, c, \epsilon_1$. The general structure of the matrix in Eq. (4) is determined
 92 only by the matching conditions and is valid for any time-continuous model applied during the seasons.
 The equations of motion of the latter only determine the numerical values of the parameters in Eq.
 94 (4).

The first line of the matrix describes the influence on the sockeye fry of year $n + 1$ of the sockeye fry
 96 of year $(n - 3)$ and $(n - 4)$ (which are the parents of the fry in year $(n + 1)$), and of the predator
 population. m_s and ϵ_1 are positive, since more parents imply more offspring. $-a$ is negative, since
 98 more predators imply less fry. The other nontrivial line of this matrix, the last line, describes the
 influence on the predators of year $(n + 1)$ of the sockeye fry of year $(n - 3)$ and $(n - 4)$ (which are
 100 the parents of the predator's food), and of the predator population in the previous year. All three
 parameters are positive, since more food implies more predator growth and since more predators in
 102 one year give rise to more predators in the next year.

The eigenvalues of this matrix determine the nature of the dynamics near the bifurcation. When
 104 all eigenvalues have an absolute value smaller than 1, the fixed point is stable, and the dynamics
 converges to this fixed point. When the absolute value of one or more eigenvalues is larger than 1,
 106 the fixed point is unstable, and the dynamics approaches a different attractor.

In order to understand the dynamics resulting from this matrix, we first consider the case that the
 108 matrix elements ϵ_1, b , and c vanish. This means that all salmon return at age 4, and that the trout
 have a good choice of other food so that their dynamics is independent of that of the salmon fry. In

110 this case the eigenvalues of the matrix are m_p , 0, and the four fourth roots of m_s . Since the four
 112 salmon lines are independent from each other in this case, the sequence δs_n has trivially the period
 4 and simply iterates the initial four values, with an amplitude decreasing for $m_s < 1$ and increasing
 114 otherwise (and with the trout being independent of the salmon). When m_s is increased from a value
 smaller than 1 to a value larger than 1, all four eigenvalues $m_s^{1/4}$ cross the unit circle simultaneously,
 116 and the fixed point becomes unstable. This degeneracy is lifted when the parameters ϵ_1 , b , and c are
 made nonzero. As long as these parameters are not large, one can expect the four main eigenvalues
 118 to remain close to the real and imaginary axis, respectively, implying a (possibly damped) oscillation
 with a period close to 4.

The type of bifurcation that occurs when the fixed point becomes unstable depends on which eigenvalue
 120 crosses first the unit circle as a control parameter is increased. In order to determine the type of the
 bifurcation, we evaluated the parameters of the matrix numerically from our computer simulations
 122 of the time continuous model when going through the bifurcation (i. e., when K_0 increases from
 6.4 to 7.5), and we calculated the eigenvalues λ of the matrix. Figure 4 shows that the bifurcation is
 124 caused by a pair of complex conjugate eigenvalues crossing the unit circle, indicating a Neimark Sacker
 bifurcation (the discrete version of the Hopf bifurcation). Since the dominant eigenvalues (depicted
 126 with the symbol +) are close to $\pm i$, the period of the resulting oscillation is close to 4. In the linear,
 time discrete model (Eq. (4)), the bifurcation is mainly driven by the parameters m_s and, to a minor
 128 extent, a , which correspond to reproduction of the four-year old sockeyes and predation by the rainbow
 trouts, respectively. The fixed point becomes unstable when either of these parameters increases.




Figure 4: **Eigenvalues of the matrix from Eq. (4), as obtained from a linear stability analysis of the time-continuous model.** K_0 increases from 6.4 to 7.5 as the eigenvalues move outwards. The zoom shows that the eigenvalues close to the imaginary axis (+) are the first ones to cross the unit cycle.

130 Now, it is known from the theory of bifurcations that if the period at such a bifurcation is close to
 4, there occurs a strong resonance, which means that the period becomes locked exactly at the value
 132 4 not far beyond the bifurcation point. In contrast to conventional mode locking that gives rise to
 Arnold tongues, a strong resonance is due to nonlinearities that are of the same order as the leading
 134 nonlinearity (which ensures the stability of the quasiperiodic orbit in the first place), and frequency
 locking therefore occurs over a much wider range of parameters [29].

136 The two complex conjugate eigenvalues are the first ones to cross the unit circle when a and ϵ_1 are

small non-negative numbers and when m_p is well below 1. Since ϵ_1 , b , and c are small compared to m_s , the period of the resulting oscillation remains close to 4, and the locking at period 4 therefore occurs not far beyond the bifurcation. When these conditions are not met, the unit circle is typically first crossed at -1, and a period-doubling (or *flip*) bifurcation occurs.

4. Discussion

The four-year oscillations of sockeye salmon in the Fraser River are a typical example of single-generation cycles (as opposed to predator-prey cycles [10]), a phenomenon that has also been extensively studied in the context of, e.g., host-parasitoid [30] or daphnia-zooplankton [31] systems. In the latter case, individuals of different age or size of a consumer species directly compete with each other for a resource, while the biological situation in the present study is somewhat different: The salmon fry of different brood lines do not compete with one another directly, since they populate the rearing lakes at distinct, non-overlapping times. The interaction between the brood lines is mediated by the long-living predators, which have a net effect on the salmon that is similar to competition (apparent competition). In contrast to other models that produce single-generation cycles, the cycle period 4 observed in our system is not identical to the mean generation time, which is larger than 4 years, and not all salmon mature and reproduce at the same age. The locking of the period at 4 is due to a strong resonance, a phenomenon which plays no role in the cited models.

The three-species model analysed in this study is in principle also capable of showing conventional predator-prey cycles. If the parameters were chosen such that the complex conjugate pair of eigenvalues of the matrix in Eq. (4) that is close to the real axis at +1 is the first to cross the unit circle, oscillations with a period much larger than four years occur and the year-to-year variation between brood lines is rather small (results not shown). However, this type of oscillations is not seen in the sockeye populations rearing in the Fraser River basin.

The considerations in the previous section lead to two basic conditions under which strong resonance and thus period-four generation cycles can be observed: First, an increase in the number of spawning salmon must lead to a sufficiently strong increase in this number four years later (i.e. m_s must be large enough). If the number of fry migrating to the ocean is dominated by other factors, such as a strongly limited carrying capacity for the fry, the population will be at a fixed point rather than on the oscillating side of the bifurcation. Second, the four salmon lines must be coupled in order for the Neimark Sacker bifurcation to occur, rather than a period doubling bifurcation. In our model, this coupling is due to a fraction of sockeye returning at age 5 instead of age 4, and, more importantly, due to the predator being sufficiently strongly coupled to sockeye dynamics.

Previous studies of salmon dynamics, used in salmon management, are based on the Ricker model [7, 24] or the Larkin model [14, 32, 33]. Because both models have a strongly limited carrying capacity for the fry, and because they do not include explicitly the predator dynamics, the only bifurcation occurring in those models is the flip bifurcation. Building on both modelling approaches, Myers et al. [13] have demonstrated that an unstable (decaying) period-4 oscillation can be excited by stochastic driving. However, it remains unclear over how long the oscillations can be maintained in that study without a change in the phase. Furthermore, that stochastic model produces only episodically the

176 typical sequence of a dominant, a subdominant and two weak years described in section 2 and seen
in our simulations, Fig. 2).

178 In principle, the coupling between the four salmon lines can also occur through the food of the sockeye
fry. However, there is no empirical evidence that there is a negative effect of a strong sockeye year
180 on the daphnia population in the following year that limits the growth of the sockeye fry in that year
[16].

182 The two conditions for a strong resonance of period 4 fit together with the empirical observation that
it occurs only in large oligotrophic lakes, such as those of the Fraser River basin. The smaller and
184 ultra-oligotrophic lakes in the outer coast regions of British Columbia do not show these oscillations.
Their nutrient content is neither large enough to raise large fry numbers, nor to allow for salmon
186 predators to become strong. On the other hand, cyclic dominance cannot be expected in nutrient-
rich lakes, because the spawning adults would not be an important nutrient input, and because the
188 coupling to the predator would not be strong enough in a situation where there were more predator
species and more prey species for these predators.

190 Since there are large fluctuations in the proportion of fry that survive to return to their nursery lake,
we included noise in the matching condition for the sockeyes, Eq. (2), in order to determine with
192 how much noise superimposed on the deterministic dynamics the period-4 oscillation can persist. Fig.
2b shows a data series generated with 50 percent noise in ocean survivability superimposed to the
194 deterministic dynamics. The oscillation is still clearly visible, although the system has a phase shift
every 300 years on average. With less noise, the phase shift occurs less often, and with 100 percent
196 noise, the cyclic dominance vanishes in our simulations. When a large perturbation acts only for a
limited time, the oscillation quickly recovers afterwards. In fact, the recovery following the blocking
198 of the Fraser River migration routes early in the last century, can be seen in the non-dominant brood
lines of the Shuswap stock in Fig. 1 and in the data of some of the stocks presented in the Appendix.

200 Unfortunately, the year 2009 has seen another large perturbation, with most Fraser sockeye expected
to return that year not surviving in the ocean, so that the expected strong escapement did not occur.

202 The age composition of sockeye stocks was also found to strongly influence the potential of the system
to show a strong resonance. The fraction of adult sockeyes that returns at the age of 5 years instead
204 of 4 years, ϵ , was set to 0.1 in the simulations and we could show that the resonance appears for
 $0 < \epsilon \leq 0.3$, but for very small values of ϵ (< 0.02) the non-dominant lines disappear completely
206 with increasing K_0 . The resonance condition was best met at $\epsilon \approx 0.2$, where the resonance occurred
nearly directly after the Neimark Sacker bifurcation. The parameter ϵ is also measured in the real
208 populations [34]. For the sockeye salmon populations of the Fraser River it is approximately 0.08,
while for the less productive outer coast lakes of British Columbia (where cyclic dominance is not
210 observed) it is between 0.56 and 0.76.

Some of the sockeye populations of the Bristol Bay area (Alaska), most notably the Kvichak River
212 stocks, also exhibit strong oscillations [35, 36] despite a broader distribution of the age at spawning
than in the Fraser River stocks [37, 38]. However, the oscillations are not as regular, with maxima of
214 the populations occurring every fourth or fifth year. This is consistent with our model, as it indicates

a quasiperiodic oscillation rather than a fixed oscillation period associated with a strong resonance.

216 Our results do not rule out additional mechanisms such as compensatory fishing [15] or genetic effects
[17], which would reduce the population sizes of weak lines to values smaller than those resulting from
218 our model. However, these additional assumptions are not needed to explain the occurrence of cyclic
dominance in the first place.

220 Acknowledgements

This work was supported by the German Research Foundation (DFG) under contract number Dr300/7
222 and Br2315/9-1. Kim Hyatt, Jeremy Hume, and Carl Walters commented on an earlier version of
this manuscript. Empirical data were kindly provided by Tracy Cone (escapement data) and by Kim
224 Hyatt (age composition of the sockeye stocks). This collaboration started at the Complex Systems
Workshop in Fairbanks, Alaska, 2007. The 2008 Cultus Lake Modelling Workshop, which had an
226 important impact on this project, was hosted by Fisheries and Oceans Canada.

References

- 228 [1] C. Elton, M. Nicholson, The ten-year cycle in numbers of lynx in Canada, *J. Anim. Ecol.* 11
(1942) 215–244.
- 230 [2] E.P. Odum, *Fundamentals of Ecology*, W.B. Saunders, Philadelphia, 1953.
- [3] B. Blasius, A. Huppert, L. Stone, Complex dynamics and phase synchronization in spatially
232 extended ecological systems, *Nature* 399 (1999) 354–359.
- [4] I. Hanski, P. Turchin, E. Korpimäki, H. Henttonen, Population oscillations of boreal rodents –
234 Regulation by mustelid predators leads to chaos, *Nature* 364 (1993) 232–235.
- [5] D. Ludwig, D.D. Jones, C.S. Holling, Qualitative analysis of insect outbreak systems – spruce
236 budworm and forest, *J. Anim. Ecol.* 47 (1978) 315–332.
- [6] T. Royama, Population dynamics of the spruce budworm *choristoneura-fumiferana*, *Ecol. Mon.*
238 54 (1984) 429–462.
- [7] W.E. Ricker, Cycle dominance among the Fraser sockeye, *Ecology* 31 (1950) 6–26.
- 240 [8] C.R. Townsend, Population cycles in freshwater fish, *J. Fish Biol.* 35 (Supplement A) (1989)
125–131.
- 242 [9] W.E. Ricker, Cycles of abundance among Fraser River sockeye salmon (*Oncorhynchus nerka*),
Can. J. Fish. Aquat. Sci. 54 (1997) 950–968.
- 244 [10] W.S.C. Gurney, R.M. Nisbet, Fluctuation periodicity, generation separation, and the expression
of larval competition, *Theor. Pop. Biol.* 28 (1985) 150–180.
- 246 [11] W.W. Murdoch, B.E. Kendall, R.M. Nisbet, C.J. Briggs, E. McCauley, R. Boiser, Single-species
models for many-species food webs, *Nature* 417 (2002) 541–543.

- 248 [12] G.A. Rounsefell, G.B. Kelez, The salmon and salmon fisheries of Swiftsure Bank, Puget Sound,
and the Fraser River, *Bull. Bur. Fish.* 49 (1938) 692–823.
- 250 [13] R.A. Myers, G. Mertz, J.M. Bridson, M.J. Bradford, Simple dynamics underlie sockeye salmon
(*Oncorhynchus nerka*) cycles, *Can. J. Fish. Aquat. Sci.* 55 (1998) 2355–2364.
- 252 [14] P.A. Larkin, Simulation studies of the Adams River sockeye salmon, *Oncorhynchus nerka*, *J. Fish.*
Res. Board Can. 18 (1971) 1493–1502.
- 254 [15] C.J. Walters, M.J. Staley, Evidence against the existence of cyclic dominance in Fraser River
sockeye salmon (*Oncorhynchus nerka*), in: *Sockeye salmon (Oncorhynchus nerka) population*
256 *biology and future management*, *Can. Spec. Publ. Fish. Aquat. Sci.* 96 (1987) 375–384.
- [16] D.A. Levy, C.C. Wood, Review of proposed mechanisms for sockeye salmon population cycles in
258 the Fraser River, *Bull. Math. Biol.* 54 (1992) 241–261.
- [17] C.J. Walters, J.C. Woodey, Genetic models for cyclic dominance in sockeye salmon (*Oncorhynchus*
260 *nerka*), *Can. J. Fish. Aquat. Sci.* 49 (1992) 281–292.
- [18] L. Persson, K. Leonardsson, A.M. de Roos, M. Gyllenberg, B. Christensen, Ontogenetic scaling
262 of foraging rates and the dynamics of a size-structured consumer-resource model, *Theor. Pop.*
Biol. 54 (1998) 270–293.
- 264 [19] P. Yodzis, S. Innes, Body size and consumer-resource dynamics, *Am. Nat.* 139 (1992) 1151–1175
- [20] J.H. Brown, J.F. Gillooly, A.P. Allen, V.M. Savage, G.B. West, Toward a metabolic theory of
266 ecology, *Ecology* 85 (2004) 1771–1789.
- [21] J.R. Beddington, Mutual interference between parasites or predators and its effects on searching
268 efficiency, *J. Anim. Ecol.* 44 (1975) 331–340.
- [22] G.T. Skalski, J.F. Gilliam, Functional responses with predator interference: viable alternatives
270 to the Holling type II model, *Ecology* 82 (2001) 3083–3092.
- [23] C.J. Walters, V. Christensen, Adding realism to foraging arena predictions of trophic flows in
272 Ecosim ecosystem models: shared foraging arenas and bout feeding, *Ecol. Mod.* 209 (2007) 342–
350.
- 274 [24] J.M.B. Hume, K.S. Shortreed, K.F. Morton, Juvenile sockeye rearing capacities of three lakes in
the Fraser River system, *Can. J. Fish. Aquat. Sci.* 53 (1996) 719–733.
- 276 [25] G.B. Pauley, R. Risher, G.L. Thomas, Species profiles: life histories and environmental require-
ments of coastal fishes and invertebrates (Pacific Northwest) – Sockeye salmon. US Dep. Int.,
278 Fish and Wildlife Service, Biological Report 82(11.116) (1989).
- [26] J. Hume, K. Shortreed, T. Whitehouse, Sockeye fry, smolt, and nursery lake monitoring of Quesnel
280 and Shuswap lakes in 2004 (2005), available at http://www.unbc.ca/qrrc/historical_research.html

- 282 [27] S.M. Gende, R.T. Edwards, M.F. Willson, M.S. Wipfli, Pacific salmon in aquatic and terrestrial
ecosystems, *BioScience* 52 (2002) 917–928.
- 284 [28] C. Guill, B. Reichhardt, B. Drossel, W. Just, Coexistence of two periodic attractors: the degen-
erate Neimark Sacker bifurcation as a generic mechanism, submitted, arXiv:1007.4981
- 286 [29] Y.A. Kuznetsov, *Elements of Applied Bifurcation Theory*, third ed., Springer Verlag, New York,
2004.
- 288 [30] H.J.C. Godfray, M.P. Hassell, Discrete and continuous insect populations in tropical environ-
ments. *J. Anim. Ecol.* 58 (1989) 153–174.
- 290 [31] A.M. de Roos, J.A.J. Metz, E. Evers, A. Leipoldt, A size dependent predator-prey interaction:
who pursues whom? *J. Math. Biol.* 28 (1990) 609–643.
- 292 [32] S.J.D. Martell, C.J. Walters, R. Hilborn, Retrospective analysis of harvest management perfor-
mance for Bristol Bay and Fraser River sockeye salmon, *Can. J. Fish. Aquat. Sci.* 65 (2008)
409–424.
- 294 [33] A.D. Marsden, S.J.D. Martell, U.R. Sumaila, Retrospective bioeconomic analysis of Fraser River
sockeye salmon fishery management, *Fish. Res.* 95 (2009) 32–41.
- 296 [34] M. Healey, The adaptive significance of age and size at maturity in female sockeye salmon
(*Oncorhynchus nerka*), in: H.D. Smith, L. Margolis, C.C. Wood (Eds.), *Sockeye Salmon (On-*
298 *corhynchus nerka) Population Biology and Future Management*. Can. Spec. Publ. Fish. Aquat.
Sci. 96, Ottawa: Fisheries and Oceans Canada, 1987, pp. 110–117.
- 300 [35] L.F. Fair, Critical elements of Kvichak River sockeye salmon management, *Alaska Fish. Res.*
Bull. 10 (2003) 95–103.
- 302 [36] L.A. Rogers, D.E. Schindler, Asynchrony in population dynamics of sockeye salmon in southwest
Alaska, *Oikos* 117 (2008) 1578–1586.
- 304 [37] F.W. West, L.F. Fair, Abundance, age, sex, and size statistics for pacific salmon in Bristol Bay,
2003, Fishery Data Series No. 06-47, Alaska Department of Fish and Game, Division of Sport
306 Fish and Commercial Fisheries, Anchorage, 2006.
- 308 [38] F.W. West, L.F. Fair, T. Baker, S. Morstad, K. Weiland, T. Sands, C. Westings, Abundance,
age, sex, and size statistics for pacific salmon in Bristol Bay, 2004, Fishery Data Series No. 09-
51, Alaska Department of Fish and Game, Division of Sport Fish and Commercial Fisheries,
310 Anchorage, 2009.

Appendix: Escapement data for sockeye stocks showing cyclic dominance

312 Many spawning stocks of sockeye salmon in the upper Fraser River basin are not stationary but have
 314 been dramatically increasing in size over the last decades (e.g. Figure A1 a, b, and d), following a
 massive disturbance of the stocks at the beginning of the 20th century. Nevertheless, the oscillatory
 pattern has clearly emerged again.

316 In some stocks (most notably the Bowron Lake stock, Figure A1 c), cyclic dominance is observed only
 episodically. This dynamical pattern may appear when the system is close to the bifurcation point.

318 The sockeye salmon stock of Seton Lake (Figure A1 e) also displayed cyclic dominance, but for nearly
 two decades (five cycles), the dominant line is followed by the weakest line instead of the sub-dominant
 320 line. In this stock, cyclic dominance broke down in the early 1990's.




Figure A1: Escapement data of for sockeye spawning stocks in the Fraser River basin. a: Quesnel Lake, b: Stuart Lake, c: Bowron Lake, d: Seton Lake, e: Chilko Lake.