



**HAL**  
open science

# On the zero-field orbital magnetic susceptibility of Bloch electrons in graphene-like solids: Some rigorous results.

Baptiste Savoie

► **To cite this version:**

Baptiste Savoie. On the zero-field orbital magnetic susceptibility of Bloch electrons in graphene-like solids: Some rigorous results.. 2012. hal-00684309v1

**HAL Id: hal-00684309**

**<https://hal.science/hal-00684309v1>**

Preprint submitted on 1 Apr 2012 (v1), last revised 4 Jul 2012 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On the zero-field orbital magnetic susceptibility of Bloch electrons in graphene-like solids: Some rigorous results.

April 1, 2012

Baptiste Savoie\*

## Abstract

Starting with a nearest-neighbors tight-binding model, we rigorously investigate the bulk zero-field orbital susceptibility of a non-interacting Bloch electrons gas in graphene-like solids at fixed temperature and density of particles. In the zero-temperature limit and in the semi-conducting situation, we derive a complete expression which holds for an arbitrary number of bands with possible degeneracies. In the particular case of a two-bands gapped model, all involved quantities are exactly written down. Besides the formula we obtain have the special feature to be suitable for numerical computations since it only involves the eigenvalues and associated eigenfunctions of the Bloch Hamiltonian, together with the derivatives (up to the second order) w.r.t. the quasi-momentum of the matrix-elements of the Bloch Hamiltonian. Finally we give a simple application for the two-bands gapped model by considering the case of a dispersion law which is linear w.r.t. the quasi-momentum in the gapless limit. Through this instance, the origin of the singularity, which express as a Dirac delta function of the Fermi energy, implied by the McClure's formula in purely monolayer graphene is discussed.

## Contents

<b>1</b>	<b>Introduction and the main results.</b>	<b>1</b>
1.1	Introduction. . . . .	1
1.2	The setting and the main results. . . . .	3
1.3	A simple application for the two-bands gapped model - Discussions. . . . .	7
1.4	The content of the paper. . . . .	9
<b>2</b>	<b>Proof of Lemma ??.</b>	<b>9</b>
2.1	Proof of (i) – (ii). . . . .	9
2.2	Proof of (iii). . . . .	11
2.3	Appendix: proofs of intermediate results. . . . .	13
<b>3</b>	<b>Proof of Theorem ??.</b>	<b>14</b>
3.1	Proof of (i). . . . .	14
3.2	Proof of (ii) – (iii). . . . .	17
3.2.1	Proof of (ii). . . . .	18
3.2.2	Proof of (iii). . . . .	20
3.3	Appendix: proofs of the intermediate results. . . . .	22
<b>4</b>	<b>Acknowledgments.</b>	<b>26</b>

PACS-2010 number: 51.60.+a, 73.20.-r, 73.22.Pr, 75.20.Ck, 75.30.Cr.

MSC-2010 number: 81Q10, 82B10, 82D20, 82D37.

Keywords: Diamagnetism, zero-field susceptibility, two-dimensional electrons gas, graphene, semi-conductors.

---

\*Institute of Mathematics 'Simion Stoilow' of the Romanian Academy, P.O. Box 1-764, RO-70700 Bucharest, Romania; e-mail: baptiste.savoie@gmail.com .

# 1 Introduction and the main results.

## 1.1 Introduction.

Isolated for the first time in 2004 by Novoselov *et al.* [30], graphene is one of the solids the most fascinating by virtue of its exceptional electronic properties which, in great part, originate from its purely two-dimensional structure; see [9] for a review of these properties. Regarded to be a semimetal since the works of Wallace in 1947 [40] (he was the first one who investigated the band structure of graphene through a nearest-neighbors tight-binding model), graphene has the feature to exhibit a very high diamagnetic susceptibility. Note that this characteristic is shared by the most semimetals, and to a lesser degree, by the most narrow-gap intrinsic semiconductors.

From a theoretical point of view, the first attempts to account for this large diamagnetic behavior go back at least to the 50's with the works of Smoluchowski [39] and Hove [19]. Even though these papers only were concerned with graphite (viewed as a piling up of graphene sheets) which shows the same feature than graphene (graphite is known as one of the strongest diamagnetic solids), they nevertheless are interesting in regards to the method. The starting point in [39, 19] is the formula of Peierls [33, Eq. (62c)] obtained in the tight-binding approximation (it amounts to the same thing to consider a single-band model) for the zero-field orbital susceptibility of conduction electrons at non-zero temperature, together with the Wallace band structure [40] (two bands touching at the Brillouin zone corners: the lower one completely filled, the upper completely empty at zero temperature). However in 1953, the Peierls formula threw back into doubt by Adams [2]. This latter pointed out the existence of other contributions coming from the interband effects induced by the magnetic field; but apart from particular cases in which these contributions are far from being negligible, no general formulas were derived. Note that the quest to a complete formula (including the interband contributions) of the zero-field susceptibility of Bloch electrons in solids generated an intense activity up to the middle of 70's, see e.g. [26] for a review. Note also that the first rigorous investigations in this direction came as late as 1990, see [17]. Recently, Briet *et al.* [5] rigorously derived a complete formula for the zero-field susceptibility of Bloch electrons at fixed temperature and density for both metals and semiconductors/insulators. They also gave a rigorous justification to the Landau-Peierls approximation in low temperature and density limit.

In 1956, McClure wrote down in the high temperature regime a formula for the zero-field orbital susceptibility of Bloch electrons in a *two-dimensional honeycomb lattice* (modeling a single-layer graphite) taking into account the interband effects, see [24, Eq. (3.15)]. To achieve that, he first calculated the energy levels induced by the magnetic field for the two-dimensional Wallace band structure (this band structure yields a linear dispersion law near the corners of the Brillouin zone). Subsequently he used these calculations to work out the Helmholtz free energy from which he derived the zero-field orbital susceptibility. He stressed that '*all the conduction electrons diamagnetism of the two-dimensional model is due to band-to-band transitions*'.

Following [24], the most papers were mainly concentrated either on 'real' graphite, or on compounds involving graphite-like structure, see e.g. [25, 38, 36, 37]. All the same we mention that when dealing with (infinite) single-layer graphite model, Safran *et al.* [36], and later on [37], recovered the McClure formula in a regime of small values of the chemical potential (they do not tend to show some restrictions on the temperature regime). However the question of the behavior of the zero-field orbital susceptibility at zero temperature is not addressed. It should be point out that the method used in [25, 38, 36, 37] basically is different from the one implemented in [24]. Indeed their starting point is a given formula for the zero-field susceptibility of a 3-D Bloch electrons gas, in this case the one derived by Fukuyama [16]. Afterwards they carry out some transformations in that formula to take into account the dispersion law induced by the band structure of materials.

In 1987, Levintovitch *et al.* [23] interested in the zero-field orbital susceptibility of a *two-dimensional Bloch electrons gas in narrow-gap semiconductors* at zero temperature and fixed density. Their method consisted in expressing a (quite) modified version of the zero-field susceptibility formula derived by Wannier *et al.* [41] only in terms of a dispersion law corresponding to a two-bands gapped model (the origin of energy is taken at the middle of the gap, and when the gap width is zero, the two-bands model used by McClure [24] is recovered). In the zero-temperature

limit and in the semiconducting situation, they get a simple formula involving only the gap width and the Fermi energy, see [23, Eqs. (22)-(23)]. In the gapless limit, they found that only the interband terms contribute to the zero-field orbital susceptibility, and moreover, it has a singularity expressed as a Dirac delta-like dependence on the Fermi energy (i.e. the zero-field orbital susceptibility is minus infinity at band-touching point, and vanishes otherwise). They claimed that their result is in good agreement with the McClure's formula in the zero-temperature limit.

After the experimental discovery of Novoselov *et al.* [30], many investigations arose to account for the large diamagnetic behavior of monolayer graphene. Their approach substantially is different from the previous authors: it is based on the observation that the linear dispersion law near the band-touching points in the Brillouin zone (derived from the 2-D Wallace band structure [40]) is analogous to the energy of massless relativistic particles. Since that suggests a massless Dirac fermions-like behavior of quasi-particles around the band-touching points, most of papers are focused on the diamagnetism of massless Dirac fermions, see e.g. [27, 20, 21]. We do not give details about their method since this present paper will not deal with Dirac fermions. However we emphasize that for a gapless Dirac fermions system, Koshino *et al.* obtained that the zero-field orbital susceptibility at zero temperature is expressed as a delta function of the Fermi energy (the origin of energy is taken at band-touching point), see [20, Eq. (37)]-[21, Eq. (33)]. Besides they argued that it corresponds to the zero-temperature limit in the McClure's formula. In [27], Nakamura interested in gapped Dirac fermions systems. In the semiconducting situation, he found that the zero-field orbital susceptibility at zero temperature is in inverse proportion to the energy gap, see [27, Eq. (7)]. In the zero-energy gap limit, the results of [20, 21] are recovered.

In the light of the above mentioned results, the delta-like singularity contained in the zero-field orbital susceptibility at zero temperature of Bloch electrons in graphene seems to be regularized when adding a band gap in the band structure model of graphene. This leads one to be interested in the zero-field orbital susceptibility of gapped graphene to shed light on the possible origins of this singularity, other than the semimetallic nature of graphene at zero temperature. Furthermore, the study of fundamental properties of gapped graphene has recently found a renewal of interest due to new technical advances allowing to open the gap in graphene materials, see e.g. [31, 32].

In this present paper, we rigorously investigate the zero-field orbital magnetic susceptibility of a non-interacting Bloch electrons gas in graphene-like solids. By a graphene-like solid, we mean a purely two-dimensional crystal which is an intrinsic semiconductor (possibly with a narrow gap) at zero temperature. Our starting point is a nearest-neighbors tight-binding model for the Hamiltonian of a Bloch electron in a finite-size graphene-like solid plunged into a constant magnetic field applied perpendicular to the solid plane. Our model of tight-binding Hamiltonian is a Harper-like operator (see e.g. [11]), and is analogous to the one of Brynildsen *et al.* [8] who lately have rigorously investigated the Faraday rotation in graphene-like solids. From the thermodynamic limit of the grand-canonical pressure (the question of its existence is there fully addressed, see Lemma 1.1 below), we derive a general formula for the bulk zero-field orbital susceptibility at fixed positive temperature and density of electrons, see (1.11). This is made possible by the use of the so-called gauge invariant perturbation theory (see e.g. [10, 28] and [3, 12, 13, 7, 8] for further applications), followed by the Bloch-Floquet theory. After carrying out some convenient transformations needed to perform the zero-temperature limit, then in the semiconducting situation, we get from (1.11) a complete formula for the zero-field orbital susceptibility at zero temperature and fixed density which holds for an arbitrary number of bands with possible degeneracies, see (1.15). It consists merely of (1.16) when considering the particular case of a two-bands gapped model. The whole of terms involved in (1.16) are explicitly written down and are expressed only in terms of eigenvalues and associated eigenfunctions of the Bloch Hamiltonian, together with the derivatives (up to the second order) w.r.t. the quasi-momentum of its matrix-elements. The terms involved in (1.15) have the same special feature what makes our formulas suitable for numerical computations. Finally, we give an application for the two-bands gapped model by treating the case of a dispersion law which is linear w.r.t. the quasi-momentum in the gapless limit (this is a simple 'textbook model' for monolayer graphene). The singular behavior of the bulk zero-field orbital susceptibility at zero temperature obtained in the gapless limit is construed from a mathematical point of view.

## 1.2 The setting and the main results.

Consider a two-dimensional finite-size crystal subjected to a uniform external magnetic field applied orthogonal to the solid. By crystal, we mean an ideal crystalline ordered solid composed of a regular lattice of identical atoms whose nuclei are fixed at their equilibrium positions. Due to the magnetic field, the Bloch electrons possess an orbital magnetic moment together with a spin magnetic moment. Since we are only interested in orbital diamagnetic effects, we disregard the electron spin. Furthermore we neglect the self-interactions: the Bloch electrons gas is supposed strongly diluted. Besides the finite-size crystal is at equilibrium with a thermal and particles bath.

Let us precise our assumptions. Let  $\Upsilon := \{\mathbf{v} : \mathbf{v} = \sum_{k=1}^2 n_k \mathbf{a}_k; n_k \in \mathbb{Z}\}$  be a Bravais lattice generated by two independent  $\mathbb{R}^2$ -vectors  $\mathbf{a}_k$ ,  $k = 1, 2$ . Let  $\Omega := \{\mathbf{x} : |\mathbf{x}| \leq \inf_{\mathbf{v} \in \Upsilon \setminus \{\mathbf{0}\}} |\mathbf{x} - \mathbf{v}|\}$  be the Wigner-Seitz (W-S) cell of  $\Upsilon$ , and denote by  $\Theta$  the finite set of vectors which models the basis (i.e. the positions of the ions at equilibrium in  $\Omega$ ). Hereafter, we suppose that  $\text{Card}(\Theta) \geq 2$ . The finite-size ion-core mesh is given by  $\Lambda_N := \{\underline{\mathbf{x}} + \mathbf{v} : \underline{\mathbf{x}} \in \Theta, \mathbf{v} = \sum_{k=1}^2 n_k \mathbf{a}_k; |n_k| \leq N\}$ ,  $N \in \mathbb{N}^*$ . Note that it basically boils down to the same thing to consider a central region of  $\Lambda := \Theta + \Upsilon = \{\underline{\mathbf{x}} + \mathbf{v} : \underline{\mathbf{x}} \in \Theta, \mathbf{v} \in \Upsilon\}$  which stands for the ion-core mesh of the corresponding infinite crystal. The magnetic field is given by the 3-dimensional vector  $\mathbf{B} := (0, 0, B)$ ,  $B \geq 0$ . By seeing a  $\mathbb{R}^2$ -vector  $\mathbf{x} = (x_1, x_2)$  as the 3-dimensional vector  $(x_1, x_2, 0)$ , the magnetic potential vector which reigns in the crystal is defined by  $B\mathbf{a}(\mathbf{x}) := \frac{1}{2}(0, 0, B) \wedge (x_1, x_2, 0) = \frac{B}{2}(-x_2, x_1, 0)$  (we have used the symmetric transverse gauge). In the following, we denote by  $b := eB/c \in \mathbb{R}^+$  the electron cyclotron frequency where we set  $m_{\text{electron}} = 1$ . Note that below we set  $\hbar = 1$  too.

Under the above conditions, let us introduce the one-particle Hamiltonian. To fix notation and give some additional assumptions, we firstly start with the situation in which the crystal is infinite. In view of our discrete approach, let  $l^2(\Lambda)$  be the one-particle Hilbert space and denote by  $\{\delta_{\mathbf{x}}\}_{\mathbf{x} \in \Lambda}$  the canonical basis, where  $\delta_{\mathbf{x}}(\mathbf{y}) = 1$  when  $\mathbf{y} = \mathbf{x}$  and 0 otherwise. Recall that the kernel of any bounded operators  $A$  on  $l^2(\Lambda)$  is given by  $A(\mathbf{x}, \mathbf{y}) := \langle \delta_{\mathbf{x}}, A\delta_{\mathbf{y}} \rangle$  ( $\mathbf{x}, \mathbf{y} \in \Lambda^2$ ), where  $\langle \cdot, \cdot \rangle$  denotes the usual scalar product on  $l^2(\Lambda)$ . In the one-body approximation, let  $H_0$  be the Hamiltonian on  $l^2(\Lambda)$  which determines the dynamics of each Bloch electron in the absence of the magnetic field. We denote by  $H_0(\cdot, \cdot) : \Lambda \times \Lambda \rightarrow \mathbb{C}$  the kernel of  $H_0$ . In the whole of this paper, we assume that  $H_0$  satisfies the following three properties (given through its kernel):

(P1)  $H_0$  is a self-adjoint operator, i.e.:

$$\forall (\mathbf{x}, \mathbf{y}) \in \Lambda^2, \quad H_0(\mathbf{x}, \mathbf{y}) = \overline{H_0(\mathbf{y}, \mathbf{x})}.$$

(P2)  $H_0$  is a nearest-neighbour tight-binding operator, that is:

$$\exists C > 0 \quad : \quad H_0(\mathbf{x}, \mathbf{y}) = 0 \quad \text{when } |\mathbf{x} - \mathbf{y}| \geq C. \quad (1.1)$$

(P3)  $H_0$  commutes with the translations of the  $\Upsilon$  Bravais lattice, i.e.:

$$\forall \mathbf{v} \in \Upsilon, \quad H_0(\mathbf{x} + \mathbf{v}, \mathbf{y} + \mathbf{v}) = H_0(\mathbf{x}, \mathbf{y}) \quad (\mathbf{x}, \mathbf{y}) \in \Lambda^2.$$

Usually the magnitude of the non-zero hopping matrix-elements for  $H_0(\cdot, \cdot)$  is fixed to some finite constant (typically, fixed to 1). In this case, and we suppose that hereafter, the kernel  $H_0(\cdot, \cdot)$  is *EAD* (=Exponentially Almost Diagonal), i.e. there exists two constants  $c_1, c_2 > 0$  s.t.

$$\forall (\mathbf{x}, \mathbf{y}) \in \Lambda^2, \quad |H_0(\mathbf{x}, \mathbf{y})| \leq c_1 e^{-c_2 |\mathbf{x} - \mathbf{y}|}. \quad (1.2)$$

In the presence of the magnetic field, the Hamiltonian  $H_b$  on  $l^2(\Lambda)$  is defined through its kernel  $H_b(\cdot, \cdot) : \Lambda \times \Lambda \rightarrow \mathbb{C}$  which is obtained from  $H_0(\cdot, \cdot)$  by using the so-called 'Peierls substitution':

$$\forall (\mathbf{x}, \mathbf{y}) \in \Lambda^2, \quad H_b(\mathbf{x}, \mathbf{y}) := e^{ib\phi(\mathbf{x}, \mathbf{y})} H_0(\mathbf{x}, \mathbf{y}) \quad b \in \mathbb{R}^+; \quad (1.3)$$

here  $\phi(\cdot, \cdot)$  stands for the usual magnetic phase defined by  $\phi(\mathbf{x}, \mathbf{y}) := \mathbf{x} \cdot \mathbf{a}(\mathbf{y}) = \frac{1}{2}(y_1 x_2 - x_1 y_2)$ . Since the phase is antisymmetric, then  $H_b$  is a self-adjoint operator by virtue of (P1). Moreover (1.1) still holds true for  $H_b(\cdot, \cdot)$ , and due to (P3),  $H_b$  commutes with the magnetic translations:

$$\forall \mathbf{v} \in \Upsilon, \quad H_b(\mathbf{x} + \mathbf{v}, \mathbf{y} + \mathbf{v}) e^{ib\phi(\mathbf{y}, \mathbf{v})} = e^{ib\phi(\mathbf{x}, \mathbf{v})} H_b(\mathbf{x}, \mathbf{y}) \quad (\mathbf{x}, \mathbf{y}) \in \Lambda^2. \quad (1.4)$$

Defined via (1.3),  $H_b$  eventually is a Harper-like operator; see e.g. [11] for its spectral properties.

Let us get back to our initial problem dealing with the Bloch electrons gas in the finite-size crystal. The one-particle Hamiltonian in the presence of the magnetic field is defined from  $H_b$  by:

$$H_{N,b} := \chi_N H_b \chi_N \quad b \in \mathbb{R}^+, \quad (1.5)$$

where  $\chi_N$  denotes the characteristic function of the ion-core mesh  $\Lambda_N$ . For any  $b \in \mathbb{R}^+$ , (1.5) defined a family of bounded and self-adjoint operators on  $l^2(\Lambda)$  for any  $N \in \mathbb{N}^*$ . Obviously this definition corresponds to choose Dirichlet boundary conditions on the edges of  $\Lambda_N$ . Hereafter we denote by  $\{e_b^{(j)}\}_{j=1}^N$  the set of eigenvalues of  $H_{N,b}$  counting multiplicities and in increasing order.

Now define the pressure of the Bloch electrons gas in the finite-size crystal under the grand-canonical conditions. Recall that in the grand-canonical ensemble, the fixed external parameters are  $(\beta, z, \text{Card}(\Lambda_N))$ , where:  $\beta := (k_B T)^{-1} > 0$  is the 'inverse' temperature ( $k_B$  stands for the Boltzmann constant) and  $z := e^{\beta\mu} > 0$  is the fugacity ( $\mu \in \mathbb{R}$  stands for the chemical potential). For any  $\beta > 0$ ,  $z > 0$  and  $b \geq 0$ , the grand-canonical pressure is given by (see e.g. [34]):

$$P_N(\beta, z, b) := \frac{1}{\beta} \frac{1}{\text{Card}(\Lambda_N)} \sum_{j=1}^N \ln(1 + ze^{-\beta e_b^{(j)}}). \quad (1.6)$$

Let us introduce some notation again. Let  $R_b(\cdot, \cdot; \xi) : \Lambda \times \Lambda \rightarrow \mathbb{C}$  be the kernel of the resolvent operator  $R_b(\xi) := (H_b - \xi)^{-1} \forall \xi \in \rho(H_b)$ . Let  $\Gamma$  be a simple positively oriented closed contour around the spectrum of  $H_b$  defined as in (2.1). We set  $\mathfrak{f}(\beta, z; \cdot) := \ln(1 + ze^{-\beta \cdot})$  for any  $\beta, z > 0$ .

Our main technical result is given in the following lemma:

**Lemma 1.1.** (i). *For any  $\beta > 0$ ,  $z > 0$  and  $b \geq 0$ , the thermodynamic limit of the grand-canonical pressure exists. More precisely:*

$$P(\beta, z, b) := \lim_{N \rightarrow \infty} P_N(\beta, z, b) = \frac{1}{\beta} \frac{1}{\text{Card}(\Theta)} \frac{i}{2\pi} \int_{\Gamma} d\xi \mathfrak{f}(\beta, z; \xi) \sum_{\mathbf{x} \in \Theta} R_b(\mathbf{x}, \mathbf{x}; \xi), \quad (1.7)$$

uniformly in  $(\beta, z) \in [\beta_1, \beta_2] \times [z_1, z_2]$ ,  $0 < \beta_1 < \beta_2 < \infty$  and  $0 < z_1 < z_2 < \infty$ .

(ii). *For any  $\beta > 0$  and  $b \geq 0$ ,  $P(\beta, \cdot, b)$  is a  $C^\infty$ -function on  $\mathbb{R}_+^*$ .*

(iii). *For any  $\beta > 0$  and  $z > 0$ ,  $P(\beta, z, \cdot)$  is a  $C^2$ -function on  $\mathbb{R}_+$ .*

By virtue of Lemma 1.1, introduce the bulk density of particles defined as the following:

$$\rho(\beta, z, b) := \beta z \frac{\partial P}{\partial z}(\beta, z, b), \quad \beta > 0, z > 0, b \geq 0, \quad (1.8)$$

as well as the bulk orbital susceptibility defined as the second derivative of the bulk pressure w.r.t. the intensity  $B$  of the magnetic field (see e.g. [1, 3, 5, 7] and references therein):

$$\mathcal{X}(\beta, z, b) := \left(\frac{e}{c}\right)^2 \frac{\partial^2 P}{\partial b^2}(\beta, z, b), \quad \beta > 0, z > 0, b \geq 0. \quad (1.9)$$

From now on, the density of particles  $\rho_0 > 0$  becomes our fixed external parameter. Seeing the bulk density  $\rho$  as a function of the  $\mu$ -variable (instead of the  $z$ -variable), we denote by  $\mu^{(0)}(\beta, \rho_0, b) \in \mathbb{R}$  the unique solution of the equation  $\rho(\beta, e^{\beta\mu}, b) = \rho_0$ . Let us note that the inversion of the relation between the bulk density and the chemical potential is ensured by the fact that  $\mu \mapsto \rho(\beta, e^{\beta\mu}, b)$  is a strictly increasing function on  $(0, \infty)$ . Under these conditions, the bulk zero-field orbital susceptibility at fixed positive 'temperature'  $\beta > 0$  and density  $\rho_0 > 0$  is defined by:

$$\mathcal{X}(\beta, \rho_0) = \mathcal{X}(\beta, \rho_0, 0) := \mathcal{X}(\beta, e^{\beta\mu^{(0)}(\beta, \rho_0, b=0)}, b=0).$$

Before formulating the main results of this paper, let us introduce some more notation.

Let  $\Upsilon^*$  be the dual lattice of  $\Upsilon$  generated by the vectors  $\mathbf{b}_l$ ,  $l = 1, 2$  defined from the  $\mathbf{a}_k$ 's by

$\mathbf{a}_k \cdot \mathbf{b}_l = 2\pi\delta_{k,l}$ . Denote by  $\Omega^*$  the Brillouin zone of  $\Upsilon^*$ . By virtue of (P3), the Floquet theory for periodic operators allows us to use the bands structure of the spectrum of  $H_0$ , see e.g. [22]. Under the Bloch-Floquet unitary transformation,  $l^2(\Lambda) \cong \int_{\Omega^*}^{\oplus} d\mathbf{k} l^2(\Theta)$  and  $H_0$  can be seen as the direct integral  $\int_{\Omega^*}^{\oplus} d\mathbf{k} H(\mathbf{k})$ , see e.g. [35, Sect. XIII.16]. For each  $\mathbf{k} \in \Omega^*$ , the fiber Hamiltonian  $H(\mathbf{k})$  lives on  $l^2(\Theta)$  and it is defined via its kernel which reads as (see e.g. [6, Eq. (2.4)]):

$$\forall(\underline{\mathbf{x}}, \underline{\mathbf{y}}) \in \Theta^2, \quad H(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) := \sum_{\mathbf{v} \in \Upsilon} e^{-i\mathbf{k} \cdot (\underline{\mathbf{x}} + \mathbf{v} - \underline{\mathbf{y}})} H_0(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}}) \quad \mathbf{k} \in \Omega^*.$$

Let us note that in the canonical basis of  $l^2(\Theta)$ , the matrix of  $H(\mathbf{k})$  is nothing but  $[H(\underline{\mathbf{x}}_l, \underline{\mathbf{x}}_m; \mathbf{k})]_{l,m}$ ,  $1 \leq l, m \leq \text{Card}(\Theta)$ . Introduce as well on  $l^2(\Theta)$  the operators  $(\partial_{k_\alpha} H)(\mathbf{k})$  and  $(\partial_{k_\gamma} \partial_{k_\alpha} H)(\mathbf{k})$  with  $\alpha, \gamma = 1, 2$  and  $\mathbf{k} \in \Omega^*$ , generated via their kernels respectively defined on  $\Theta^2$  as:

$$\begin{aligned} (\partial_{k_\alpha} H)(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) &:= \frac{\partial}{\partial k_\alpha} H(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) = -i \sum_{\mathbf{v} \in \Upsilon} (\underline{x}_\alpha + v_\alpha - \underline{y}_\alpha) e^{-i\mathbf{k} \cdot (\underline{\mathbf{x}} + \mathbf{v} - \underline{\mathbf{y}})} H_0(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}}), \quad (1.10) \\ (\partial_{k_\gamma} \partial_{k_\alpha} H)(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) &:= \frac{\partial^2}{\partial k_\gamma \partial k_\alpha} H(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) = \\ &= - \sum_{\mathbf{v} \in \Upsilon} (\underline{x}_\gamma + v_\gamma - \underline{y}_\gamma) (\underline{x}_\alpha + v_\alpha - \underline{y}_\alpha) e^{-i\mathbf{k} \cdot (\underline{\mathbf{x}} + \mathbf{v} - \underline{\mathbf{y}})} H_0(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}}) = \frac{\partial^2}{\partial k_\alpha \partial k_\gamma} H(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}). \end{aligned}$$

Since  $H_0(\cdot, \cdot)$  is *EAD*, the kernels  $(\partial_{k_\alpha} H)(\cdot, \cdot; \mathbf{k})$  and  $(\partial_{k_\gamma} \partial_{k_\alpha} H)(\cdot, \cdot; \mathbf{k})$  are well-defined on  $\Theta^2$ . Getting back to the Bloch Hamiltonian, we denote by  $\{E_j(\mathbf{k})\}_{j=1}^{\text{Card}(\Theta)}$  the set of its eigenvalues, counting multiplicities and *in increasing order*. Due to this choice of labelling, the Bloch energies  $E_j(\cdot)$  are  $\Upsilon^*$ -periodic and continuous functions but they are not differentiable at (possible) crossing-points; for a review see e.g. [29, Sect. III]. Hereafter we will consider the both situations:

- (A1) The  $E_j(\cdot)$ 's,  $j = 1, \dots, \text{Card}(\Theta)$  are non-degenerate for Lebesgue-almost all  $\mathbf{k} \in \Omega^*$ .  
(A2) The  $E_j(\cdot)$ 's,  $j = 1, \dots, \text{Card}(\Theta)$  are non-degenerate everywhere on  $\Omega^*$ .

In other words, (A1) means that the set formed by the crossing-points has a zero Lebesgue-measure. As for (A2), it amounts to the same thing to assume that the Bloch bands are simple; the  $j$ -th Bloch band is defined by  $\mathcal{E}_j := \bigcup_{\mathbf{k} \in \Omega^*} E_j(\mathbf{k})$ . Under either assumption, the spectrum of  $H_0$  is absolutely continuous and given by  $\sigma(H_0) = \bigcup_{j=1}^{\text{Card}(\Theta)} \mathcal{E}_j$ . The energy bands of  $H_0$  corresponds to the disjoint union of the  $\mathcal{E}_j$ 's. Under the condition (A1), the  $\mathcal{E}_j$ 's may overlap, be contained in each other, or even coincide. Generally speaking if  $\max \mathcal{E}_j < \mathcal{E}_{j+1}$  for some  $j = 1, \dots, \text{Card}(\Theta) - 1$ , then we have a non-trivial spectral gap. If  $\max \mathcal{E}_j = \mathcal{E}_{j+1}$ , then we have a trivial spectral gap.

Lastly, denote by  $\mathcal{E}_F(\rho_0)$  the Fermi energy which is defined by  $\mathcal{E}_F(\rho_0) := \lim_{\beta \rightarrow \infty} \mu^{(0)}(\beta, \rho_0, 0)$ . This limit always exists and eventually defines an increasing function of  $\rho_0$ , see [5, Thm. 1.1].

The main results of this paper are collected in the following theorem:

**Theorem 1.2.** *Let  $\rho_0 > 0$  be fixed.*

(i). *For any  $\beta > 0$ , let  $\mu^{(0)} = \mu^{(0)}(\beta, \rho_0, 0) \in \mathbb{R}$  be the unique solution of the equation  $\rho(\beta, e^{\beta\mu}, 0) = \rho_0$ . Then the bulk zero-field orbital susceptibility at fixed positive temperature and density can be written as:*

$$\mathcal{X}(\beta, \rho_0) = \frac{1}{4} \left( \frac{e}{c} \right)^2 \frac{1}{\text{Card}(\Theta) |\Omega^*|} \frac{1}{\beta} \frac{i}{2\pi} \int_{\Gamma} d\xi \mathfrak{f}(\beta, \mu^{(0)}; \xi) \int_{\Omega^*} d\mathbf{k} \text{Tr}_{l^2(\Theta)} \left( \sum_{l=3}^5 \mathcal{P}_l(\mathbf{k}; \xi) \right), \quad (1.11)$$

where for each  $\mathbf{k} \in \Omega^*$ , denoting by  $R(\mathbf{k}; \xi) := (H(\mathbf{k}) - \xi)^{-1}$  the fiber of the resolvent:

$$\begin{aligned} \mathcal{P}_5(\mathbf{k}; \xi) &:= R(\mathbf{k}; \xi) \left\{ (\partial_{k_1} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_2} H)(\mathbf{k}) - (\partial_{k_2} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_1} H)(\mathbf{k}) \right\} \times \\ &\times R(\mathbf{k}; \xi) \left\{ (\partial_{k_2} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_1} H)(\mathbf{k}) - (\partial_{k_1} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_2} H)(\mathbf{k}) \right\} R(\mathbf{k}; \xi), \quad (1.12) \end{aligned}$$

$$\begin{aligned}
\mathcal{P}_4(\mathbf{k}; \xi) := & R(\mathbf{k}; \xi) \left\{ (\partial_{k_1}^2 H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_2} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_2} H)(\mathbf{k}) + \right. \\
& - (\partial_{k_1} \partial_{k_2} H)(\mathbf{k}) R(\mathbf{k}; \xi) [(\partial_{k_1} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_2} H)(\mathbf{k}) + (\partial_{k_2} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_1} H)(\mathbf{k})] + \\
& \left. + (\partial_{k_2}^2 H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_1} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_1} H)(\mathbf{k}) \right\} R(\mathbf{k}; \xi), \quad (1.13)
\end{aligned}$$

$$\begin{aligned}
\mathcal{P}_3(\mathbf{k}; \xi) := & -R(\mathbf{k}; \xi) \left\{ \frac{1}{2} (\partial_{k_1}^2 H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_2}^2 H)(\mathbf{k}) + \frac{1}{2} (\partial_{k_2}^2 H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_1}^2 H)(\mathbf{k}) + \right. \\
& \left. - (\partial_{k_1} \partial_{k_2} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_1} \partial_{k_2} H)(\mathbf{k}) \right\} R(\mathbf{k}; \xi). \quad (1.14)
\end{aligned}$$

(ii). Suppose (A1). Assume the semiconducting situation, i.e. the Fermi energy lies in the middle of a non-trivial gap, that is there exists  $M \in \{1, \dots, \text{Card}(\Theta) - 1\}$  s.t.  $\mathcal{E}_M < \mathcal{E}_{M+1}$  and  $\mathcal{E}_F(\rho_0) = (\max \mathcal{E}_M + \min \mathcal{E}_{M+1})/2$ . Then there exists  $2M$  functions  $\mathfrak{d}_{j,1}(\cdot)$  and  $\mathfrak{d}_{j,0}(\cdot)$ ,  $j = 1, \dots, M$  defined on  $\Omega^*$  outside a set of Lebesgue-measure zero s.t.

$$\begin{aligned}
\mathcal{X}(\rho_0) &:= \lim_{\beta \rightarrow \infty} \mathcal{X}(\beta, \rho_0) \\
&= \left(\frac{e}{c}\right)^2 \frac{1}{4} \frac{1}{\text{Card}(\Theta) |\Omega^*|} \int_{\Omega^*} d\mathbf{k} \sum_{j=1}^M \left\{ \mathfrak{d}_{j,1}(\mathbf{k}) + \{E_j(\mathbf{k}) - \mathcal{E}_F(\rho_0)\} \mathfrak{d}_{j,0}(\mathbf{k}) \right\}, \quad (1.15)
\end{aligned}$$

and the above integrand can be extended by continuity to the whole of  $\Omega^*$ .

(iii). Suppose (A2) together with  $\text{Card}(\Theta) = 2$ . Assume the semiconducting situation, i.e. the Fermi energy lies in the middle of the spectral gap separating the band  $\mathcal{E}_1$  from  $\mathcal{E}_2$ . Then in the zero-temperature limit, the bulk zero-field orbital susceptibility at fixed density is given by:

$$\mathcal{X}(\rho_0) = \left(\frac{e}{c}\right)^2 \frac{1}{8} \frac{1}{|\Omega^*|} \int_{\Omega^*} d\mathbf{k} \sum_{l=0}^2 \left\{ \frac{\mathbf{u}_{1,l}(\mathbf{k})}{(E_2(\mathbf{k}) - E_1(\mathbf{k}))^{l+1}} + \frac{\{E_1(\mathbf{k}) - \mathcal{E}_F(\rho_0)\} \mathbf{v}_{1,l}(\mathbf{k})}{(E_2(\mathbf{k}) - E_1(\mathbf{k}))^{l+2}} \right\}, \quad (1.16)$$

where the functions  $\mathbf{u}_{1,l}(\cdot)$  and  $\mathbf{v}_{1,l}(\cdot)$  are respectively defined in (3.29)-(3.31) and (3.32)-(3.34).

**Remark 1.** Each and every one functions  $\mathfrak{d}_{j,1}(\cdot)$  and  $\mathfrak{d}_{j,0}(\cdot)$ ,  $j = 1, \dots, M$  appearing in (1.15) can be explicitly written down. They only involve the eigenvalues and associated eigenfunctions of the Bloch Hamiltonian  $H(\mathbf{k})$ , together with the derivatives (up to the second order) w.r.t. the  $k_\alpha$ -variables ( $\alpha = 1, 2$ ) of the  $\text{Card}(\Theta) \times \text{Card}(\Theta)$ -matrix elements of  $H(\mathbf{k})$ , see Section 3.2.1. As for the functions  $\mathbf{u}_{1,l}(\cdot)$  and  $\mathbf{v}_{1,l}(\cdot)$ ,  $l = 0, 1, 2$ , they have the same characteristics, see page 21 for their explicit expressions. This special feature makes our formulas suitable for computations.

**Remark 2.** We mention that the assumption (A1) in (ii) is artificial in the following sense. One eventually can prove that (1.15) still holds true when the  $E_j$ 's are degenerate on a subset of  $\Omega^*$  with full Lebesgue-measure. However in this case the functions  $\mathfrak{d}_{j,1}(\cdot)$  and  $\mathfrak{d}_{j,0}(\cdot)$ ,  $j = 1, \dots, M$  can not be expressed as it is mentioned in Remark 1: their expressions are more complicated and require the use of the orthogonal projection corresponding to  $E_j$ . For further details, see [5].

**Remark 3.** When (A1) is replaced with (A2) in (ii), then (1.15) still holds true with the same functions  $\mathfrak{d}_{j,l}(\cdot)$ ,  $j = 1, \dots, M$  and  $l = 0, 1$ . But in this instance these functions are smooth in the  $\mathbf{k}$ -variable by virtue of the analytic perturbation theory, see e.g. [35, Sect. XII]. In the particular case of  $\text{Card}(\Theta) = 2$ , one has the following identities:

$$\forall \mathbf{k} \in \Omega^*, \quad \mathfrak{d}_{1,1}(\mathbf{k}) = \sum_{l=0}^2 \frac{\mathbf{u}_{1,l}(\mathbf{k})}{(E_2(\mathbf{k}) - E_1(\mathbf{k}))^{l+1}}, \quad \mathfrak{d}_{1,0}(\mathbf{k}) = \sum_{l=0}^2 \frac{\mathbf{v}_{1,l}(\mathbf{k})}{(E_2(\mathbf{k}) - E_1(\mathbf{k}))^{l+2}}.$$

**Remark 4.** As it is common practice in physics, the starting point in this kind of problem is a given matrix for the Bloch Hamiltonian  $H(\mathbf{k})$ . Since the trace in the formula (1.11) only involves



the derivatives (up to the second order) w.r.t. the  $k_\alpha$ -variables ( $\alpha = 1, 2$ ) of the  $\text{Card}(\Theta) \times \text{Card}(\Theta)$ -matrix elements of the Bloch Hamiltonian together with its 'inverse matrix', then a such formula allows 'hand'-calculations when dealing notably with  $2 \times 2$  matrices. More precisely, the calculations from (1.11) are quite easy when considering dispersion laws which are polynomial in  $\mathbf{k}$  in the gapless limit. Under these conditions, the zero-temperature limit simply follows from the formula (1.11) by the application of the residue theorem without resorting to the formula (1.15) or (1.16), see the following paragraph for some examples. Besides we mention that under the assumption (A1), we give another formula for the zero-field orbital susceptibility at fixed positive temperature and density which have the same special feature than the one mentioned for the formula (1.15), see (3.12) in Proposition 3.2. In the particular case of  $\text{Card}(\Theta) = 2$ , all the terms involved in (3.12) are explicitly written down, see Lemma 3.8. Furthermore the assumption (A2) makes possible the rewriting of (3.12) as a sum of two terms: the first one is the so-called Peierls contribution, the second one stands for the interband contributions, see (3.38) in Proposition 3.9. In the zero-temperature limit and in the semiconducting situation, only a large number of the terms involved in the interband contributions will give rise to (1.16).

**Remark 5.** The regularity properties announced in (ii)-(iii) of Lemma 1.1 are far from being optimum. On the one hand, one can prove that  $z \mapsto P(\beta, z, b)$  can be analytically extended to the complex domain  $\mathbb{C} \setminus (-\infty, -e^{\beta E_b^{(0)}}]$  where  $E_b^{(0)} := \inf \sigma(H_b)$ , see e.g. [3, 4, 7]. On the other hand, the use of the gauge invariant magnetic perturbation theory to prove (iii) allows us eventually to get that  $b \mapsto P(\beta, z, b)$  is a  $\mathcal{C}^\infty$ -function. For further details, see the proof of Proposition 2.3.

### 1.3 A simple application for the two-bands gapped model - Discussions.

The purpose of this paragraph is to present and to discuss some results on the bulk zero-field orbital susceptibility of Bloch electrons at zero temperature and fixed density in the framework of a two-bands gapped model for which the dispersion law is linear w.r.t. the quasi-momentum in the gapless limit. For monolayer graphene, this model is a case in point as a first approximation.

We first give a series of calculations from two distinct Bloch Hamiltonian, see (1.17) and (1.21) below, which both have the same eigenvalues. Subsequently, we discuss the results.

**Calculation (1).** The starting point is the following Bloch Hamiltonian:

$$H^{(l)}(\mathbf{k}) := \begin{pmatrix} \delta & k_1 + ik_2 \\ k_1 - ik_2 & -\delta \end{pmatrix} \quad \delta > 0, \mathbf{k} \in \Omega^*. \quad (1.17)$$

Let  $E_1(\mathbf{k}) := -\sqrt{\delta^2 + |\mathbf{k}|^2}$  and  $E_2(\mathbf{k}) := -E_1(\mathbf{k})$  be its two eigenvalues. The matrix-resolvent is:

$$R^{(l)}(\mathbf{k}; \xi) := \frac{-1}{(E_1(\mathbf{k}) - \xi)(E_2(\mathbf{k}) - \xi)} \begin{pmatrix} \delta + \xi & k_1 + ik_2 \\ k_1 - ik_2 & \xi - \delta \end{pmatrix} \quad \mathbf{k} \in \Omega^*.$$

Note that the second derivatives w.r.t. the  $k_\alpha$ -variable,  $\alpha = 1, 2$  of the matrix elements in (1.17) are all zero. Therefore, each quantity in (1.13) and (1.14) is reduced to the  $2 \times 2$ -zero matrix. After carrying out some calculations involving products of  $2 \times 2$ -matrices, one gets for the quantity defined in (1.12) (which we denote by  $\mathcal{P}_5^{(l)}(\mathbf{k}; \xi)$  to express the dependence on the model in (1.17)):

$$\mathcal{P}_5^{(l)}(\mathbf{k}; \xi) := \frac{4(\delta^2 - \xi^2)}{(E_1(\mathbf{k}) - \xi)^4 (E_2(\mathbf{k}) - \xi)^4} \begin{pmatrix} \delta + \xi & k_1 + ik_2 \\ k_1 - ik_2 & \xi - \delta \end{pmatrix} \quad \mathbf{k} \in \Omega^*. \quad (1.18)$$

Let  $\beta > 0$  and  $\rho_0 > 0$  be fixed. From (1.18), the formula (1.11) leads to:

$$\mathcal{X}(\beta, \rho_0) = \left(\frac{e}{c}\right)^2 \frac{1}{|\Omega^*|} \frac{1}{\beta} \int_{\Omega^*} d\mathbf{k} \left(\frac{1}{2i\pi}\right) \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi) g(\xi)}{(E_1(\mathbf{k}) - \xi)^4 (E_2(\mathbf{k}) - \xi)^4}, \quad g(\xi) := \xi(\xi^2 - \delta^2).$$

Suppose the semiconducting situation, i.e. the Fermi energy lies in the middle of the spectral gap (the width of the gap is  $2\delta$ ) separating the band  $\mathcal{E}_1$  from  $\mathcal{E}_2$ . By performing first the integration

w.r.t. the  $\xi$ -variable by the residue theorem, and subsequently the zero-temperature limit (for further details in regards to the method, see the proof of Theorem 1.2 (ii) – (iii)), one has:

$$\mathcal{X}(\rho_0) := \lim_{\beta \rightarrow \infty} \mathcal{X}(\beta, \rho_0) = -\frac{1}{32} \left( \frac{e}{c} \right)^2 \frac{1}{|\Omega^*|} \int_{\Omega^*} d\mathbf{k} \{ \mathfrak{d}_{1,1}(\mathbf{k}) + (E_1(\mathbf{k}) - \mathcal{E}_F(\rho_0)) \mathfrak{d}_{1,0} \},$$

$$\text{where: } \mathfrak{d}_{1,1}(\mathbf{k}) = \frac{1}{(\delta^2 + |\mathbf{k}|^2)^{\frac{3}{2}}} + \frac{\delta^2}{(\delta^2 + |\mathbf{k}|^2)^{\frac{5}{2}}} \quad \text{and} \quad \mathfrak{d}_{1,0}(\mathbf{k}) = 0. \quad (1.19)$$

Finally, by performing first the integration w.r.t. the  $\mathbf{k}$ -variable (note that the presence of the  $\delta$ 's makes integrable the function  $\mathfrak{d}_{1,1}(\cdot)$  near  $\mathbf{k} = \mathbf{0}$ ) and only after the gapless limit, one obtains:

$$\lim_{\delta \rightarrow 0} \mathcal{X}(\rho_0) = -\infty. \quad (1.20)$$

**Calculation (2).** The starting point is the following diagonal Bloch Hamiltonian:

$$H^{(d)}(\mathbf{k}) := \begin{pmatrix} E_1(\mathbf{k}) & 0 \\ 0 & E_2(\mathbf{k}) \end{pmatrix}, \quad \text{with: } \begin{cases} E_1(\mathbf{k}) := -\sqrt{\delta^2 + |\mathbf{k}|^2}, \\ E_2(\mathbf{k}) := -E_1(\mathbf{k}) \end{cases} \quad \delta > 0, \mathbf{k} \in \Omega^*. \quad (1.21)$$

The matrix-resolvent is:

$$R^{(d)}(\mathbf{k}; \xi) := \frac{1}{(E_1(\mathbf{k}) - \xi)(E_2(\mathbf{k}) - \xi)} \begin{pmatrix} E_2(\mathbf{k}) - \xi & 0 \\ 0 & E_1(\mathbf{k}) - \xi \end{pmatrix} \quad \mathbf{k} \in \Omega^*.$$

Let us remark that:

$$(\partial_{k_j} H^{(d)})(\mathbf{k}) := \frac{k_j}{\sqrt{\delta^2 + |\mathbf{k}|^2}} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \quad j = 1, 2.$$

Due to this feature, it is easy to see that the quantity in (1.12) is reduced to the  $2 \times 2$ -zero matrix. After working out the trace of the quantities in (1.13)-(1.14), (1.11) yields:

$$\mathcal{X}(\beta, \rho_0) = \frac{1}{8} \left( \frac{e}{c} \right)^2 \frac{1}{|\Omega^*|} \frac{1}{\beta} \sum_{j=1}^2 \int_{\Omega^*} d\mathbf{k} \left\{ \frac{(-1)^j |\mathbf{k}|^2}{(\delta^2 + |\mathbf{k}|^2)^{\frac{3}{2}}} \left( \frac{1}{2i\pi} \right) \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_j(\mathbf{k}) - \xi)^4} + \right.$$

$$\left. - \frac{\delta^2}{(\delta^2 + |\mathbf{k}|^2)^2} \left( \frac{1}{2i\pi} \right) \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_j(\mathbf{k}) - \xi)^3} \right\}.$$

Let us mention that the application of the residue theorem for each complex integration (inside the braces) provides a derivative of  $\mathfrak{f}(\beta, \mu^{(0)}, \cdot)$  of order three and two respectively. This feature leads, in the zero-temperature limit and in the semiconducting situation, to the following:

$$\mathcal{X}(\rho_0) := \lim_{\beta \rightarrow \infty} \mathcal{X}(\beta, \rho_0) = 0. \quad (1.22)$$

Let us now discuss these results. Starting with the Bloch Hamiltonian in (1.17), the singularity expressed as a delta function of the Fermi energy is recovered, see (1.20) and Section 1.1. Through the calculations leading to (1.22), this singularity can be accounted for by the following.

The Bloch Hamiltonian-matrix in (1.17) is related to the one in (1.21) by:

$$H^{(l)}(\mathbf{k}) = P(\mathbf{k}) H^{(d)}(\mathbf{k}) P^{-1}(\mathbf{k}) \quad \mathbf{k} \in \Omega^*, \quad (1.23)$$

where the change-of-basis matrix together with its inverse read for instance as:

$$P(\mathbf{k}) := \begin{pmatrix} \frac{-(k_1 + ik_2)}{\delta + \sqrt{\delta^2 + |\mathbf{k}|^2}} & \frac{-(k_1 + ik_2)}{\delta - \sqrt{\delta^2 + |\mathbf{k}|^2}} \\ 1 & 1 \end{pmatrix}, \quad P^{-1}(\mathbf{k}) := \begin{pmatrix} \frac{-(k_1 - ik_2)}{2\sqrt{\delta^2 + |\mathbf{k}|^2}} & \frac{\delta + \sqrt{\delta^2 + |\mathbf{k}|^2}}{2\sqrt{\delta^2 + |\mathbf{k}|^2}} \\ \frac{k_1 - ik_2}{2\sqrt{\delta^2 + |\mathbf{k}|^2}} & \frac{-\delta + \sqrt{\delta^2 + |\mathbf{k}|^2}}{2\sqrt{\delta^2 + |\mathbf{k}|^2}} \end{pmatrix}. \quad (1.24)$$

Let  $\mathcal{P}_5^{(d)}(\mathbf{k}; \xi)$  be the quantity defined in (1.12) but with  $H^{(d)}(\mathbf{k})$ . By replacing  $H^{(l)}(\mathbf{k})$  with the r.h.s. of (1.23) in  $\mathcal{P}_5^{(l)}(\mathbf{k}; \xi)$  which only survives in the calculation (1), one gets:

$$\mathcal{P}_5^{(l)}(\mathbf{k}; \xi) = P(\mathbf{k})\mathcal{P}_5^{(d)}(\mathbf{k}; \xi)P^{-1}(\mathbf{k}) + W_5^{(l,d)}(\mathbf{k}; \xi) \quad \mathbf{k} \in \Omega^*, \quad (1.25)$$

where  $W_5^{(l,d)}(\mathbf{k}; \xi)$  consists of a sum of terms which each of them contains at least a 'derivative w.r.t. the  $k_\alpha$ -variable of the change-of-basis matrix' (or its inverse); a generical term is for example:

$$P(\mathbf{k})R^{(d)}(\mathbf{k}; \xi)P^{-1}(\mathbf{k})(\partial_{k_j}P)(\mathbf{k})H^{(d)}(\mathbf{k})R^{(d)}(\mathbf{k}; \xi)(\partial_{k_i}H^{(d)})(\mathbf{k})R^{(d)}(\mathbf{k}; \xi)(\partial_{k_i}H^{(d)})(\mathbf{k}) \times \\ \times R^{(d)}(\mathbf{k}; \xi)H^{(d)}(\mathbf{k})(\partial_{k_i}P^{-1})(\mathbf{k})P(\mathbf{k})R^{(d)}(\mathbf{k}; \xi)P^{-1}(\mathbf{k}) \quad i, j = 1, 2, i \neq j.$$

As mentioned in calculation (2),  $\mathcal{P}_5^{(d)}(\mathbf{k}; \xi) = 0$  (in the matrices-sense). In view of (1.25) and (1.18), this means that only the term  $W_5^{(l,d)}(\mathbf{k}; \xi)$  (through its trace) provides a non-zero contribution to the zero-field orbital susceptibility at zero-temperature, see (1.19). Therefore, the singular behavior appearing in the gapless limit (1.20) is closely related to the fact that the maps  $\mathbf{k} \mapsto P(\mathbf{k})$  and  $\mathbf{k} \mapsto P^{-1}(\mathbf{k})$  are non-differentiable in  $\mathbf{k} = \mathbf{0}$  when  $\delta = 0$ , see (1.24).

Finally, the identically zero result obtained in (1.22) tends to show that our 'linear model' as a first approximation for monolayer graphene is not good enough. Some numerical computations based on a more 'realistic' dispersion law derived from a 'gapped two-dimensional Wallace'-type band structure is needed.

## 1.4 The content of the paper.

Our current paper is organized as follows. Section 2 is dedicated to the proof of Lemma 1.1. The proof of (i) essentially is based on a simplified version of the geometric perturbation theory recently developed in [13]. Since the method for the discrete case is fully detailed in [8, Sect. 3.3], the proof of (1.7) just is outlined. The crucial ingredient involved in the proof of (iii) is the gauge invariant magnetic perturbation theory applied on the resolvent operator  $R_b(\xi)$ . It allows us to keep a good control over the linear growth induced by the magnetic vector potential when dealing with the diagonal part of the kernel  $R_b(\cdot, \cdot; \xi)$ . Section 3 is devoted to the proof of Theorem 1.2. In Section 3.1 we focus on bulk the zero-field orbital susceptibility at fixed positive temperature and density. In particular, we prove Theorem 1.2 (i) from the results of Lemma 1.1 (iii) together with the Bloch-Floquet decomposition. In Sections 3.2 and 3.3 we prove Theorem 1.2 (ii) and (iii) respectively. It essentially follows the outline of the proof of [5, Thm. 1.2 (i)]. From the formula (1.11), we separately consider the both situations corresponding to each one of the assumptions (A1)-(A2) and we perform the zero-temperature limit in the semiconducting situation.

## 2 Proof of Lemma 1.1.

### 2.1 Proof of (i) – (ii).

Under the grand-canonical conditions, let  $\beta := (k_B T)^{-1} > 0$  and  $z := e^{\beta\mu} > 0$ . For any  $b \geq 0$ , let  $\Gamma$  be the counter-clockwise oriented simple contour defined as the following:

$$\Gamma := \{\Re\xi \in [\delta_-, \delta_+], \Im\xi = \pm \frac{\pi}{2\beta}\} \cup \{\Re\xi = \delta_\pm, \Im\xi \in [-\frac{\pi}{2\beta}, \frac{\pi}{2\beta}]\}, \quad (2.1)$$

where  $\delta_- := \inf \sigma(H_b) - 1$ ,  $\delta_+ := \sup \sigma(H_b) + 1$ . By construction  $\Gamma$  surrounds the spectrum of  $H_b$ , and the closed subset surrounding by  $\Gamma$  is a strict subset of  $\mathfrak{D} := \{\zeta \in \mathbb{C} : \Im\zeta \in (-\pi/\beta, \pi/\beta)\}$  which is the holomorphic domain of the function  $\xi \mapsto \mathfrak{f}(\beta, z; \xi) := \ln(1 + ze^{-\beta\xi})$  for any  $z > 0$ . We point out the fact that the reals  $\delta_\pm$  in (2.1) can eventually be chosen  $b$ -independent since the  $l^2$ -norm of  $H_b$  is bounded from above by some constants uniformly in  $b$ . This follows from

the Schur-Holmgren criterion (see [11, Eq. (1.1)]) applied to the kernel (1.3), combined with the estimate (1.2). Note also that under our conditions, there exists a constant  $c = c(\beta, z) > 0$  s.t.

$$\forall \xi \in \Gamma, \quad |\mathfrak{f}(\beta, z; \xi)| \leq ce^{-\beta \Re \xi}. \quad (2.2)$$

Now we use the Dunford functional calculus [15, Sect. VI.3] to write down a more convenient formula for the grand-canonical pressure of the Bloch electrons gas in the finite-size crystal.

First of all we need give some properties on the resolvent operator of  $H_{N,b}$  defined in (1.5). Let  $N \in \mathbb{N}^*$  and  $b \geq 0$ . For any  $\xi \in \rho(H_{N,b})$ , denote by  $R_{N,b}(\xi) := (H_{N,b} - \xi)^{-1}$ . Obviously  $R_{N,b}(\xi)$  lives on the Hilbert space  $l^2(\Lambda_N)$ , and moreover it owns a kernel denoted by  $R_{N,b}(\cdot, \cdot; \xi) : \Lambda_N \times \Lambda_N \rightarrow \mathbb{C}$ . This kernel is *EAD* (=Exponentially Almost Diagonal) in the following sense: for any compact subset  $K$  of  $\rho(H_{N,b})$ , there exists two constants  $c_1, c_2 > 0$  independent of  $b, N$  s.t.

$$\forall (\mathbf{x}, \mathbf{y}) \in \Lambda_N^2, \quad \sup_{\xi \in K} |R_{N,b}(\mathbf{x}, \mathbf{y}; \xi)| \leq c_1 e^{-c_2 |\mathbf{x} - \mathbf{y}|}. \quad (2.3)$$

The estimate (2.3) simply follows from the fact that the kernel of  $H_{N,b}$  is *EAD*, see [8, Prop. 3.5]. An important ingredient for the following is the uniform estimate on the trace of  $R_{N,b}$ :

$$\sup_{\xi \in K} |\text{Tr}_{l^2(\Lambda_N)}(R_{N,b}(\xi))| \leq c, \quad (2.4)$$

for some  $K$ -dependent constants  $c > 0$ . To obtain (2.4), all we have to do is write the trace as the sum over  $\Lambda_N$  of the diagonal matrix-elements of  $R_{N,b}(\xi)$ , and afterwards use the estimate (2.3).

Let  $\beta > 0$ ,  $z > 0$ ,  $b \geq 0$  and  $N \in \mathbb{N}^*$ . With  $\Gamma$  the contour as in (2.1), introduce on  $l^2(\Lambda_N)$ :

$$\mathcal{L}_N(\beta, z, b) := \frac{i}{2\pi} \int_{\Gamma} d\xi \mathfrak{f}(\beta, z; \xi) R_{N,b}(\xi).$$

The Dunford functional calculus provides us the identification  $\mathcal{L}_N(\beta, z, b) = \ln(\mathbb{I}_N + ze^{-\beta H_{N,b}})$  which holds on  $l^2(\Lambda_N)$ . In view of (1.6), this allows us to define the grand-canonical pressure as:

$$P_N(\beta, z, b) = \frac{1}{\beta} \frac{1}{\text{Card}(\Lambda_N)} \text{Tr}(\mathcal{L}_N(\beta, z, b)) = \frac{1}{\beta} \frac{1}{\text{Card}(\Lambda_N)} \frac{i}{2\pi} \int_{\Gamma} d\xi \mathfrak{f}(\beta, z; \xi) \text{Tr}(R_{N,b}(\xi)). \quad (2.5)$$

Obviously the trace in the r.h.s. of the first equality makes sense (do not forget that we in fact deal with  $N \times N$ -matrices). The commutation of the complex integral w.r.t.  $\xi$  with the trace in the r.h.s. of the second equality is ensured by the Tonneli's theorem, see estimates (2.4) and (2.2). We draw the attention to the fact that if the following limit holds:

$$\lim_{N \rightarrow \infty} \frac{1}{\text{Card}(\Lambda_N)} \text{Tr}(R_{N,b}(\xi)) = \frac{1}{\text{Card}(\Theta)} \sum_{\mathbf{x} \in \Theta} R_b(\mathbf{x}, \mathbf{x}; \xi), \quad (2.6)$$

then (1.7) (only in the pointwise-sense) will follow from the dominated convergence theorem.

Let us outline the proof of (2.6) which uses a simplified version of the geometric perturbation theory [13, Sect. 4]. All the details about this method in the discrete case are in [8, Sect. 3.3.1]. Below  $\lfloor \cdot \rfloor$  denotes the floor function. Pick  $\epsilon \in (0, 1)$ , and assume that  $N$  is large enough. The first step consists in dividing  $\Lambda_N$  into a 'core part'  $\tilde{\Lambda}_N$  and an 'edge part'  $\tilde{\tilde{\Lambda}}_N := \Lambda_N \setminus \tilde{\Lambda}_N$ , where:

$$\tilde{\Lambda}_N := \{\mathbf{x} + \mathbf{v} : \mathbf{x} \in \Theta, \mathbf{v} = \sum_{k=1}^2 n_k \mathbf{a}_k; |n_k| \leq N - \lfloor N^\epsilon \rfloor\}.$$

From a geometrical argument, one can see that  $\tilde{\Lambda}_N$  contains  $(2(N - \lfloor N^\epsilon \rfloor) + 1)^2$  W-S cells. This implies that the number of W-S cells in  $\tilde{\tilde{\Lambda}}_N$  is of order  $\mathcal{O}(N^{1+\epsilon})$  for  $N$  sufficiently large. Owing to this splitting of  $\Lambda_N$ , we expect only the contribution to  $\text{Tr}(R_{N,b}(\xi))$  coming from the core

region  $\tilde{\Lambda}_N$  to give the announced limit in (2.6). With this aim in view, the geometric perturbation method consists in approximating the resolvent  $R_{N,b}(\xi)$  with the operator  $U_{N,b}(\xi)$  defined by:

$$U_{N,b}(\xi) := \chi_{\tilde{\Lambda}_N} R_b(\xi) \chi_{\tilde{\Lambda}_N} + \chi_{\tilde{\Lambda}_N^c} R_{N,b}(\xi) \chi_{\tilde{\Lambda}_N^c}, \quad (2.7)$$

where  $\chi_{\tilde{\Lambda}_N}$  and  $\chi_{\tilde{\Lambda}_N^c}$  are the characteristic functions of  $\tilde{\Lambda}_N$  and  $\tilde{\Lambda}_N^c$  respectively. Afterwards, it remains to control the behavior when  $N \rightarrow \infty$  of the trace of the r.h.s. of (2.7). On the one hand:

$$\frac{1}{\text{Card}(\Lambda_N)} \text{Tr}(\chi_{\tilde{\Lambda}_N} R_{N,b}(\xi) \chi_{\tilde{\Lambda}_N}) = \frac{1}{(2N+1)^2 \text{Card}(\Theta)} \sum_{\mathbf{x} \in \tilde{\Lambda}_N} R_{N,b}(\mathbf{x}, \mathbf{x}; \xi) = \mathcal{O}\left(\frac{1}{N^{1-\epsilon}}\right),$$

where we used (2.3) together with the fact that the number of W-S cells in  $\tilde{\Lambda}_N$  is of order  $\mathcal{O}(N^{1+\epsilon})$ . On the other hand, by using that  $R_b(\xi)$  commutes with the magnetic translations, see (1.4):

$$\frac{\text{Tr}(\chi_{\tilde{\Lambda}_N} R_b(\xi) \chi_{\tilde{\Lambda}_N})}{\text{Card}(\Lambda_N)} = \frac{(2(N - \lfloor N^\epsilon \rfloor) + 1)^2}{(2N+1)^2 \text{Card}(\Theta)} \sum_{\mathbf{x} \in \Theta} R_b(\mathbf{x}, \mathbf{x}; \xi) = \frac{1}{\text{Card}(\Theta)} \sum_{\mathbf{x} \in \Theta} R_b(\mathbf{x}, \mathbf{x}; \xi) + \mathcal{O}\left(\frac{1}{N^{1-\epsilon}}\right).$$

To complete the proof of (2.6), we need control the behavior of the trace of  $R_{N,b}(\xi) - U_{N,b}(\xi)$ . One can prove that  $\text{Tr}(R_{N,b}(\xi) - U_{N,b}(\xi))/\text{Card}(\Lambda_N) = \mathcal{O}(N^{-1})$ . We do not give further details.

In order to complete the proof of Lemma 1.1 (i), consider for any  $(\beta, z) \in [\beta_1, \beta_2] \times [z_1, z_2]$ , with  $0 < \beta_1 < \beta_2 < \infty$  and  $0 < z_1 < z_2 < \infty$ , the following quantity:

$$\mathcal{Q}(\beta, z, b) := \frac{i}{2\pi} \int_{\Gamma} d\xi f(\beta, z; \xi) \left( \frac{1}{\text{Card}(\Lambda_N)} \sum_{\mathbf{x} \in \Lambda_N} R_{N,b}(\mathbf{x}, \mathbf{x}; \xi) - \frac{1}{\text{Card}(\Theta)} \sum_{\mathbf{x} \in \Theta} R_b(\mathbf{x}, \mathbf{x}; \xi) \right),$$

where the  $\Gamma$ -contour is defined as in (2.1) but with  $\beta_2$  instead of  $\beta$ . Then  $\forall (\beta, z) \in [\beta_1, \beta_2] \times [z_1, z_2]$ :

$$|\mathcal{Q}(\beta, z, b)| \leq c \int_{\Gamma} |d\xi| e^{-\beta_1 \Re \xi} \left| \frac{1}{\text{Card}(\Lambda_N)} \sum_{\mathbf{x} \in \Lambda_N} R_{N,b}(\mathbf{x}, \mathbf{x}; \xi) - \frac{1}{\text{Card}(\Theta)} \sum_{\mathbf{x} \in \Theta} R_b(\mathbf{x}, \mathbf{x}; \xi) \right|,$$

for some constants  $c = c(\beta_1, z_2) > 0$ . By virtue of (2.6), the proof of Lemma 1.1 (i) now is over.

Let us conclude this paragraph by proving (ii) of Lemma 1.1. Let  $\beta > 0$  be fixed. For any  $\xi \in \mathfrak{D}$  (the holomorphic domain of  $f$ ),  $z \mapsto f(\beta, z; \xi)$  is a  $\mathcal{C}^\infty$ -function on  $(0, \infty)$ . In particular:

$$(\partial_z^m f)(\beta, z; \xi) = (m-1)! (-1)^{m+1} \frac{e^{-m\beta\xi}}{(1 + ze^{-\beta\xi})^m} \quad m \in \mathbb{N}^*,$$

and  $\xi \mapsto (\partial_z^m f)(\beta, z; \xi)$ ,  $m \in \mathbb{N}^*$  is holomorphic on  $\mathfrak{D}$ . Moreover it obeys the following estimate:

$$\forall \xi \in \Gamma, \quad |(\partial_z^m f)(\beta, z; \xi)| \leq ce^{-m\beta\Re\xi} \quad m \in \mathbb{N}^*,$$

for some constants  $c = c(m) > 0$ . The proof of Lemma 1.1 (ii) follows by standard arguments.

## 2.2 Proof of (iii).

The proof of (iii) essentially is based on the application of the so-called gauge invariant magnetic perturbation theory to the resolvent operator  $R_b(\xi)$ , see Lemma 2.2 below. This method makes it possible not only to get the regularity of the kernel  $R_b(\cdot, \cdot; \xi)$  w.r.t. the  $b$ -variable, but also to write down formulas for its partial derivatives w.r.t.  $b$ , see Proposition 2.3. For reader's convenience we collect in the appendix of this section all the proofs of intermediary results needed to prove (iii).

For the sake of simplicity, let us see  $b$  as a parameter on the whole real line. Pick a  $b_0 \in \mathbb{R}$ .

For any  $\xi \in \rho(H_{b_0})$  and  $b \in \mathbb{R}$ , introduce on  $l^2(\Lambda)$  the operators  $\tilde{R}_b(\xi)$  and  $\tilde{T}_b(\xi)$  through their kernel which respectively are defined by:

$$\forall(\mathbf{x}, \mathbf{y}) \in \Lambda^2, \quad \tilde{R}_b(\mathbf{x}, \mathbf{y}; \xi) := e^{i\delta b \phi(\mathbf{x}, \mathbf{y})} R_{b_0}(\mathbf{x}, \mathbf{y}; \xi) \quad \delta b := b - b_0, \quad (2.8)$$

$$\tilde{T}_b(\mathbf{x}, \mathbf{y}; \xi) := e^{i\delta b \phi(\mathbf{x}, \mathbf{y})} \sum_{\mathbf{z} \in \Lambda} \{e^{i\delta b \text{fl}(\mathbf{x}, \mathbf{z}, \mathbf{y})} - 1\} H_{b_0}(\mathbf{x}, \mathbf{z}) R_{b_0}(\mathbf{z}, \mathbf{y}; \xi), \quad (2.9)$$

where, for any arbitrary vectors  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \Lambda$ ,  $\text{fl}(\mathbf{u}, \mathbf{v}, \mathbf{w}) := \phi(\mathbf{u}, \mathbf{v}) + \phi(\mathbf{v}, \mathbf{w}) + \phi(\mathbf{w}, \mathbf{u})$ . Note that  $\text{fl}(\mathbf{u}, \mathbf{v}, \mathbf{w})$  stands for the flux of a unit magnetic field through the triangle generated by  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ . From (2.8),  $\tilde{R}_b(\xi)$  is clearly bounded on  $l^2(\Lambda)$  since its kernel is  $EAD$ . Ditto for  $\tilde{T}_b(\xi)$ :

**Lemma 2.1.** *Let  $b_0 \in \mathbb{R}$  be fixed. For each compact subset  $K \subset \rho(H_{b_0})$  there exists  $c_K > 0$  s.t.:*

$$\forall b \in \mathbb{R}, \quad \sup_{\xi \in K} \|\tilde{T}_b(\xi)\| \leq c_K |\delta b|. \quad (2.10)$$

The main point of the gauge invariance magnetic perturbation theory is the below identity (2.11). It asserts that for any  $b$  sufficiently close to  $b_0$  kept fixed, the resolvent  $R_b(\xi)$  can be approximated on  $l^2(\Lambda)$  by  $\tilde{R}_b(\xi)$ ; the  $l^2$ -norm of the 'corrective term' being of order  $\mathcal{O}(|b - b_0|)$ :

**Lemma 2.2.** *Let  $b_0 \in \mathbb{R}$  be fixed. Then for each compact subset  $K \subset \rho(H_{b_0})$  there exists  $\varsigma_K > 0$  s.t. in the bounded operators sense on  $l^2(\Lambda)$ :*

$$\forall b \in [b_0 - \varsigma_K, b_0 + \varsigma_K], \quad \forall \xi \in K, \quad R_b(\xi) = \tilde{R}_b(\xi) - R_b(\xi) \tilde{T}_b(\xi). \quad (2.11)$$

The above identity, twice iterated and written in the kernel sense, allows us to establish:

**Proposition 2.3.** *Let  $b_0 \in \mathbb{R}$  and  $K \subset \rho(H_{b_0})$  be a compact subset. Then there exists  $\varsigma_K > 0$  s.t.  $b \mapsto R_b(\mathbf{x}, \mathbf{y}; \xi)$  is a  $\mathcal{C}^2$ -function on  $(b_0 - \varsigma_K, b_0 + \varsigma_K) \forall \xi \in K$  and  $\forall(\mathbf{x}, \mathbf{y}) \in \Lambda^2$ . In particular, its second partial derivative at  $b_0$  reads as:*

$$\begin{aligned} \forall(\mathbf{x}, \mathbf{y}) \in \Lambda^2, \quad (\partial_b^2 R)_{b_0}(\mathbf{x}, \mathbf{y}; \xi) = & -\frac{(\phi(\mathbf{x}, \mathbf{y}))^2}{2} R_{b_0}(\mathbf{x}, \mathbf{y}; \xi) + \\ & + \sum_{\mathbf{z}_1, \mathbf{z}_2 \in \Lambda} R_{b_0}(\mathbf{x}, \mathbf{z}_1; \xi) \{ \phi(\mathbf{x}, \mathbf{z}_1) + \phi(\mathbf{z}_1, \mathbf{y}) \} \text{fl}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{y}) H_{b_0}(\mathbf{z}_1, \mathbf{z}_2) R_{b_0}(\mathbf{z}_2, \mathbf{y}; \xi) + \\ - \sum_{\mathbf{z}_1, \dots, \mathbf{z}_4 \in \Lambda} & R_{b_0}(\mathbf{x}, \mathbf{z}_1; \xi) \text{fl}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3) H_{b_0}(\mathbf{z}_1, \mathbf{z}_2) R_{b_0}(\mathbf{z}_2, \mathbf{z}_3; \xi) \text{fl}(\mathbf{z}_3, \mathbf{z}_4, \mathbf{y}) H_{b_0}(\mathbf{z}_3, \mathbf{z}_4) R_{b_0}(\mathbf{z}_4, \mathbf{y}; \xi) + \\ & + \frac{1}{2} \sum_{\mathbf{z}_1, \mathbf{z}_2 \in \Lambda} R_{b_0}(\mathbf{x}, \mathbf{z}_1; \xi) \{ \text{fl}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{y}) \}^2 H_{b_0}(\mathbf{z}_1, \mathbf{z}_2) R_{b_0}(\mathbf{z}_2, \mathbf{y}; \xi), \quad (2.12) \end{aligned}$$

and the first both terms in the r.h.s. of (2.12) vanish when  $\mathbf{x} = \mathbf{y}$  by antisymmetry of the phase.

Let us prove (iii) of Lemma 1.1. Let  $\beta > 0$ ,  $z > 0$  and  $b_0 \in \mathbb{R}$ . With the contour  $\Gamma$  defined in (2.1) (with  $b_0$  in place of  $b$ ), recall that the bulk pressure of the Bloch electrons gas reads as:

$$P(\beta, z, b_0) = \frac{1}{\beta} \frac{1}{\text{Card}(\Theta)} \frac{i}{2\pi} \int_{\Gamma} d\xi \mathfrak{f}(\beta, z; \xi) \sum_{\mathbf{x} \in \Theta} R_{b_0}(\mathbf{x}, \mathbf{x}; \xi).$$

From Proposition 2.3, we know that there exists  $\varsigma_{\Gamma} > 0$  such that  $\Gamma$  encloses the spectrum of  $H_b$  for all  $b$  satisfying  $|b - b_0| < \varsigma_{\Gamma}$ , and moreover  $b \mapsto R_b(\mathbf{x}, \mathbf{x}; \xi)$  is a  $\mathcal{C}^2$ -function on  $(b_0 - \varsigma_{\Gamma}, b_0 + \varsigma_{\Gamma}) \forall \xi \in \Gamma$  and  $\forall \mathbf{x} \in \Lambda$ . On the other hand, from the expressions (2.12) and (2.15) (with  $\mathbf{y} = \mathbf{x}$ ) and by using the same arguments as the ones used in the proof of Lemma 2.1 below, then one has:

$$\forall \mathbf{x} \in \Lambda, \quad \forall b \in (b_0 - \varsigma_{\Gamma}, b_0 + \varsigma_{\Gamma}), \quad \sup_{\xi \in \Gamma} |(\partial_b^l R)_b(\mathbf{x}, \mathbf{x}; \xi)| \leq c \quad l = 0, 1, 2,$$

for some  $\Gamma$ -dependent constants  $c > 0$ . Lemma 1.1 (iii) now follows by standard arguments.

### 2.3 Appendix: proofs of intermediate results.

**Proof of Lemma 2.1.** From the definition (2.9), it holds on  $\Lambda^2$ :

$$|\tilde{T}_b(\mathbf{x}, \mathbf{y}; \xi)| \leq |\delta b| \sum_{\mathbf{z} \in \Lambda} |\mathbf{x} - \mathbf{z}| |H_{b_0}(\mathbf{x}, \mathbf{z})| |\mathbf{z} - \mathbf{y}| |R_{b_0}(\mathbf{z}, \mathbf{y}; \xi)|,$$

where we used that  $|e^{ix} - 1| \leq |x| \forall x \in \mathbb{R}$ , followed by the estimate  $|\text{fl}(\mathbf{x}, \mathbf{z}, \mathbf{y})| \leq |\mathbf{x} - \mathbf{z}| |\mathbf{z} - \mathbf{y}|$ . Now use that  $H_{b_0}(\cdot, \cdot)$  is *EAD*, and therefore  $R_{b_0}(\cdot, \cdot; \xi)$  too by [8, Prop. 3.5], to get ride of the factor  $|\mathbf{x} - \mathbf{z}|$  and  $|\mathbf{z} - \mathbf{y}|$  respectively. Ergo there exists two  $K$ -dependent constants  $c_1, c_2 > 0$  s.t.:

$$\forall \xi \in K, \quad |\tilde{T}_b(\mathbf{x}, \mathbf{y}; \xi)| \leq c_1 |\delta b| e^{-c_2 |\mathbf{x} - \mathbf{y}|}.$$

The lemma follows by the Shur-Holmgren criterion in [11, Eq. (1.1)]. ■

**Proof of Lemma 2.2.** Let  $K \subset \rho(H_{b_0})$  be a compact subset. Pick a  $\xi \in K$ . From (2.8), one has:

$$\forall (\mathbf{x}, \mathbf{y}) \in \Lambda^2, \quad (H_b \tilde{R}_b(\xi))(\mathbf{x}, \mathbf{y}) = \sum_{\mathbf{z} \in \Lambda} e^{i\delta b(\phi(\mathbf{x}, \mathbf{z}) + \phi(\mathbf{z}, \mathbf{y}))} e^{ib_0 \phi(\mathbf{x}, \mathbf{z})} H_0(\mathbf{x}, \mathbf{z}) R_{b_0}(\mathbf{z}, \mathbf{y}; \xi).$$

Let us note that  $\phi(\mathbf{x}, \mathbf{z}) + \phi(\mathbf{z}, \mathbf{y}) = \text{fl}(\mathbf{x}, \mathbf{z}, \mathbf{y}) + \phi(\mathbf{x}, \mathbf{y})$ . Then from (1.3) together with (2.9):

$$(H_b \tilde{R}_b(\xi))(\mathbf{x}, \mathbf{y}) = e^{i\delta b \phi(\mathbf{x}, \mathbf{y})} \sum_{\mathbf{z} \in \Lambda} H_{b_0}(\mathbf{x}, \mathbf{z}) R_{b_0}(\mathbf{z}, \mathbf{y}; \xi) + \tilde{T}_b(\mathbf{x}, \mathbf{y}; \xi).$$

Since  $(H_{b_0} R_{b_0}(\xi))(\mathbf{x}, \mathbf{y}) = \delta_{\mathbf{x}, \mathbf{y}} + \xi R_{b_0}(\mathbf{x}, \mathbf{y})$  with  $\delta_{\mathbf{x}, \mathbf{y}} := \delta_{x_1, y_1} \delta_{x_2, y_2}$  (the Kronecker symbol), then:

$$\forall (\mathbf{x}, \mathbf{y}) \in \Lambda^2, \quad ((H_b - \xi) \tilde{R}_b(\xi))(\mathbf{x}, \mathbf{y}) = e^{i\delta b \phi(\mathbf{x}, \mathbf{y})} \delta_{\mathbf{x}, \mathbf{y}} + \tilde{T}_b(\mathbf{x}, \mathbf{y}; \xi).$$

This implies in the bounded operators sense on  $l^2(\Lambda)$ :

$$(H_b - \xi) \tilde{R}_b(\xi) = \mathbb{I} + \tilde{T}_b(\xi). \quad (2.13)$$

Next use that there exists a constant  $\varsigma_K > 0$  s.t.  $K \subset \rho(H_b)$  for all  $0 \leq |b - b_0| \leq \varsigma_K$ , see [11]. Thus for such  $b$ 's, we can invert the operator  $(H_b - \xi)$  with  $\xi \in K$  as above. The proof is over. ■

**Proof of Proposition 2.3.** Let  $b_0 \in \mathbb{R}$  and  $K \subset \rho(H_{b_0})$  be a compact subset. Pick a  $\xi \in K$ . The starting point is the identity (2.11) twice iterated. Written in the kernels sense, one has  $\forall b \in \mathbb{R}$  satisfying  $|b - b_0| \leq \varsigma_K$  and  $\forall (\mathbf{x}, \mathbf{y}) \in \Lambda^2$ :

$$R_b(\mathbf{x}, \mathbf{y}; \xi) = \tilde{R}_b(\mathbf{x}, \mathbf{y}; \xi) - (\tilde{R}_b(\xi) \tilde{T}_b(\xi))(\mathbf{x}, \mathbf{y}) + (\tilde{R}_b(\xi) \tilde{T}_b^2(\xi))(\mathbf{x}, \mathbf{y}) - (R_b(\xi) \tilde{T}_b^3(\xi))(\mathbf{x}, \mathbf{y}). \quad (2.14)$$

For each kernel in the r.h.s. of (2.14), we expand the exponential phase factor appearing in the kernels of  $\tilde{R}_b(\xi)$  and  $\tilde{T}_b(\xi)$  in Taylor series up to the second order in  $\delta b$ . Then we have:

$$\tilde{R}_b(\mathbf{x}, \mathbf{y}; \xi) = \sum_{k=0}^2 (\delta b)^k \frac{(i\phi(\mathbf{x}, \mathbf{y}))^k}{k!} R_{b_0}(\mathbf{x}, \mathbf{y}; \xi) + \mathcal{R}_b^{(0)}(\mathbf{x}, \mathbf{y}; \xi),$$

$$\begin{aligned} (\tilde{R}_b(\xi) \tilde{T}_b(\xi))(\mathbf{x}, \mathbf{y}) &= \sum_{\mathbf{z}_1, \mathbf{z}_2 \in \Lambda} R_{b_0}(\mathbf{x}, \mathbf{z}_1; \xi) \left\{ i\delta b - (\delta b)^2 \left[ (\phi(\mathbf{x}, \mathbf{z}_1) + \phi(\mathbf{z}_1, \mathbf{y})) + \frac{1}{2} \text{fl}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{y}) \right] \right\} \times \\ &\quad \times \text{fl}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{y}) H_{b_0}(\mathbf{z}_1, \mathbf{z}_2) R_{b_0}(\mathbf{z}_2, \mathbf{y}; \xi) + \mathcal{R}_b^{(1)}(\mathbf{x}, \mathbf{y}; \xi), \end{aligned}$$

$$\begin{aligned} (\tilde{R}_b(\xi) \tilde{T}_b^2(\xi))(\mathbf{x}, \mathbf{y}) &= -(\delta b)^2 \sum_{\mathbf{z}_1, \dots, \mathbf{z}_4 \in \Lambda} R_{b_0}(\mathbf{x}, \mathbf{z}_1; \xi) \text{fl}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3) H_{b_0}(\mathbf{z}_1, \mathbf{z}_2) R_{b_0}(\mathbf{z}_2, \mathbf{z}_3; \xi) \times \\ &\quad \times \text{fl}(\mathbf{z}_3, \mathbf{z}_4, \mathbf{y}) H_{b_0}(\mathbf{z}_3, \mathbf{z}_4) R_{b_0}(\mathbf{z}_4, \mathbf{y}; \xi) + \mathcal{R}_b^{(2)}(\mathbf{x}, \mathbf{y}; \xi), \end{aligned}$$

where the remainder terms  $\mathcal{R}_b^{(l)}(\mathbf{x}, \mathbf{y}; \xi)$ ,  $l = 0, 1, 2$  satisfy the property that their first two derivatives at  $b_0 = 0$  are identically zero. Note that we can disregard the last term in the r.h.s. of (2.14) since by virtue of (2.10), its  $l^2$ -norm behaves like  $\mathcal{O}(|\delta b|^3)$  when  $|\delta b| \rightarrow 0$  uniformly in  $\xi \in K$ . Next from the above expansions, for  $b \in [b_0 - \varsigma_K, b_0 + \varsigma_K]$  sufficiently close to  $b_0$ , it holds on  $\Lambda^2$ :

$$\begin{aligned} R_b(\mathbf{x}, \mathbf{y}; \xi) - R_{b_0}(\mathbf{x}, \mathbf{y}; \xi) &= \delta b \{i\phi(\mathbf{x}, \mathbf{y})R_{b_0}(\mathbf{x}, \mathbf{y}; \xi) + \\ &\quad - i \sum_{\mathbf{z}_1, \mathbf{z}_2 \in \Lambda} R_{b_0}(\mathbf{x}, \mathbf{z}_1; \xi) \text{fl}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{y}) H_{b_0}(\mathbf{z}_1, \mathbf{z}_2) R_{b_0}(\mathbf{z}_2, \mathbf{y}; \xi)\} + o(\delta b). \end{aligned}$$

Perform the limit  $b \rightarrow b_0$ , and we get that the map  $b \mapsto R_b(\mathbf{x}, \mathbf{y}; \xi)$  is differentiable at  $b_0$  with:

$$(\partial_b R)_{b_0}(\mathbf{x}, \mathbf{y}; \xi) := i\phi(\mathbf{x}, \mathbf{y})R_{b_0}(\mathbf{x}, \mathbf{y}; \xi) - i \sum_{\mathbf{z}_1, \mathbf{z}_2 \in \Lambda} R_{b_0}(\mathbf{x}, \mathbf{z}_1; \xi) \text{fl}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{y}) H_{b_0}(\mathbf{z}_1, \mathbf{z}_2) R_{b_0}(\mathbf{z}_2, \mathbf{y}; \xi). \quad (2.15)$$

This result can be extended on the whole of  $(b_0 - \varsigma_K, b_0 + \varsigma_K)$ . The Proposition follows by induction. Let us note that from (2.11) iterated  $n$ -times and written in the kernels sense, our method allows to get the expansion of the kernel of the resolvent in power series of  $\delta b$  up to the  $n$ -th order. By an iterating procedure, one therefore can prove that  $b \mapsto R_b(\mathbf{x}, \mathbf{y}; \xi)$  is a  $C^\infty$ -function near  $b_0$ ,  $\forall \xi \in K$  and  $\forall (\mathbf{x}, \mathbf{y}) \in \Lambda^2$ . Though the formula obtained for the  $n$ -th derivative  $(\partial_b^n R)_{b_0}(\mathbf{x}, \mathbf{y}; \xi)$  is quite complicated, its diagonal part only involves the flux  $\text{fl}$  with the kernels  $H_0(\cdot, \cdot)$  and  $R_0(\cdot, \cdot; \xi)$ . ■

### 3 Proof of Theorem 1.2.

#### 3.1 Proof of (i).

We start by writing down a general formula for the bulk orbital susceptibility under the grand-canonical conditions. From (1.9) together with the results of Section 2.3, one directly obtains:

$$\mathcal{X}(\beta, z, b) = \left(\frac{e}{c}\right)^2 \frac{1}{\beta} \frac{1}{\text{Card}(\Theta)} \frac{i}{2\pi} \int_{\Gamma} d\xi \text{f}(\beta, z; \xi) \sum_{\underline{\mathbf{x}} \in \Theta} (\partial_b^2 R)_b(\underline{\mathbf{x}}, \underline{\mathbf{x}}; \xi) \quad \beta > 0, z > 0, b \geq 0, \quad (3.1)$$

where  $(\partial_b^2 R)_b(\cdot, \cdot; \xi)$  is defined in (2.12) (only the two last terms survive for the diagonal part).

From now on, the density of particles  $\rho_0 > 0$  becomes our fixed external parameter. Under the conditions of Lemma 1.1, the relation between the fugacity and the bulk density in (1.8) can be inverted. This is made possible since for any  $\beta > 0$  and  $b \geq 0$ , the map  $z \mapsto \rho(\beta, z, b)$  is strictly increasing on  $(0, \infty)$ , and eventually defines a  $C^\infty$ -diffeomorphism of  $(0, \infty)$  into itself. Then there exists a unique  $z^{(0)}(\beta, \rho_0, b) \in (0, \infty)$ , and therefore a unique  $\mu^{(0)}(\beta, \rho_0, b) \in \mathbb{R}$  satisfying:

$$\rho(\beta, e^{\beta\mu^{(0)}(\beta, \rho_0, b)}, b) = \rho_0 > 0.$$

In this way, the bulk orbital susceptibility at fixed positive temperature and density is defined as:

$$\mathcal{X}(\beta, \rho_0, b) := \mathcal{X}(\beta, \mu^{(0)}(\beta, \rho_0, b), b) = \mathcal{X}(\beta, e^{\beta\mu^{(0)}(\beta, \rho_0, b)}, b) \quad \beta > 0, b \geq 0. \quad (3.2)$$

Under the zero magnetic field assumption, let us now prepare the formula (3.2) (knowing (3.1)) for the Bloch-Floquet decomposition. To do that, introduce the 'position operator' in  $l^2(\Lambda)$  as:

$$\forall \mathbf{x} = (x_1, x_2) \in \Lambda, \quad (\mathbf{X} \cdot \mathbf{e}_l \psi)(\mathbf{x}) = x_l \psi(\mathbf{x}) \quad \psi \in l_c^2(\Lambda), l = 1, 2, \quad (3.3)$$

where  $l_c^2(\Lambda)$  denotes the space of  $l^2$ -functions with compact support.

For any  $\xi \in \rho(H_0)$  and any  $q, r, s, t \in \{1, 2\}$ , define on  $l^2(\Lambda)$  the following families of operators:

$$\mathcal{S}_{q,r,s,t}(\xi) := R_0(\xi)[\mathbf{X} \cdot \mathbf{e}_q, [\mathbf{X} \cdot \mathbf{e}_r, H_0]][\mathbf{X} \cdot \mathbf{e}_s, [\mathbf{X} \cdot \mathbf{e}_t, R_0(\xi)]], \quad (3.4)$$

$$\mathcal{T}_{q,r,s,t}(\xi) := R_0(\xi)[\mathbf{X} \cdot \mathbf{e}_q, H_0][\mathbf{X} \cdot \mathbf{e}_r, R_0(\xi)][\mathbf{X} \cdot \mathbf{e}_s, H_0][\mathbf{X} \cdot \mathbf{e}_t, R_0(\xi)]. \quad (3.5)$$



Here  $[\cdot, \cdot]$  denotes the commutator on  $l^2(\Lambda)$ . Since the kernels of  $H_0$  and  $R_0(\xi)$  are *EAD*, then all the above commutators are well-defined on  $l^2(\Lambda)$ . As we will see hereafter, the families of operators defined in (3.4) and (3.5) have the feature to commute with the translations of the  $\Upsilon$ -lattice.

Involving (3.4) and (3.5), the bulk zero-field orbital susceptibility at fixed positive temperature and density can be written as:

**Lemma 3.1.** *Let  $\beta > 0$  and  $\rho_0 > 0$  be fixed. Let  $\mu^{(0)} = \mu^{(0)}(\beta, \mu, b = 0) \in \mathbb{R}$  be the unique solution of the equation  $\rho(\beta, e^{\beta\mu}, b = 0) = \rho_0$ . Then one has:*

$$\begin{aligned} \mathcal{X}(\beta, \rho_0, b = 0) = & \frac{1}{4} \left( \frac{e}{c} \right)^2 \frac{1}{\beta} \frac{1}{\text{Card}(\Theta)} \frac{i}{2\pi} \int_{\Gamma} d\xi \mathfrak{f}(\beta, \mu^{(0)}; \xi) \sum_{\underline{\mathbf{x}} \in \Theta} \left\{ \frac{1}{2} \mathcal{S}_{2,2,1,1}(\xi) + \frac{1}{2} \mathcal{S}_{1,1,2,2}(\xi) + \right. \\ & \left. - \mathcal{S}_{1,2,1,2}(\xi) - \mathcal{T}_{2,1,2,1}(\xi) + \mathcal{T}_{2,1,1,2}(\xi) + \mathcal{T}_{1,2,2,1}(\xi) - \mathcal{T}_{1,2,1,2}(\xi) \right\}(\underline{\mathbf{x}}, \underline{\mathbf{x}}). \end{aligned} \quad (3.6)$$

**Proof.** By inserting (2.12) into (3.1), then by setting  $b = 0$  we get:

$$\begin{aligned} \mathcal{X}(\beta, \rho_0, b = 0) = & \frac{1}{2} \left( \frac{e}{c} \right)^2 \frac{1}{\beta} \frac{1}{\text{Card}(\Theta)} \frac{i}{2\pi} \int_{\Gamma} d\xi \mathfrak{f}(\beta, \mu^{(0)}; \xi) \times \\ & \times \sum_{\underline{\mathbf{x}} \in \Theta} \left\{ \sum_{\mathbf{z}_1, \mathbf{z}_2 \in \Lambda} R_0(\underline{\mathbf{x}}, \mathbf{z}_1; \xi) \{\text{fl}(\mathbf{z}_1, \mathbf{z}_2, \underline{\mathbf{x}})\}^2 H_0(\mathbf{z}_1, \mathbf{z}_2) R_0(\mathbf{z}_2, \underline{\mathbf{x}}; \xi) + \right. \\ & \left. - 2 \sum_{\mathbf{z}_1, \dots, \mathbf{z}_4 \in \Lambda} R_0(\underline{\mathbf{x}}, \mathbf{z}_1; \xi) \text{fl}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3) H_0(\mathbf{z}_1, \mathbf{z}_2) R_0(\mathbf{z}_2, \mathbf{z}_3; \xi) \text{fl}(\mathbf{z}_3, \mathbf{z}_4, \underline{\mathbf{x}}) H_0(\mathbf{z}_3, \mathbf{z}_4) R_0(\mathbf{z}_4, \underline{\mathbf{x}}; \xi) \right\}. \end{aligned}$$

From the above expression, it remains to give the both ingredients which lead to (3.6). The first one consists in rewriting the magnetic flux  $\text{fl}(\mathbf{x}, \mathbf{y}, \mathbf{z})$ , for arbitrary  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \Lambda^3$ , as the following:

$$\text{fl}(\mathbf{x}, \mathbf{y}, \mathbf{z}) := \phi(\mathbf{x}, \mathbf{y}) + \phi(\mathbf{y}, \mathbf{z}) + \phi(\mathbf{z}, \mathbf{x}) = \frac{1}{2} [(y_1 - z_1)(x_2 - y_2) - (y_2 - z_2)(x_1 - y_1)],$$

where we used that  $\phi(\mathbf{u}, \mathbf{v}) = \frac{1}{2}(u_2 v_1 - u_1 v_2) = \frac{1}{2} [(u_2 - v_2)v_1 - (u_1 - v_1)v_2]$ . The second one consists in using the following identity. Let  $A(\mathbf{x}, \mathbf{y})$  be the kernel of a bounded operator  $A$ . Then:

$$\forall (\mathbf{x}, \mathbf{y}) \in \Lambda^2, \quad (x_l - y_l)A(\mathbf{x}, \mathbf{y}) = [\mathbf{X} \cdot \mathbf{e}_l, A](\mathbf{x}, \mathbf{y}) \quad l = 1, 2,$$

where  $\mathbf{X} \cdot \mathbf{e}_l$  is the position operator defined in (3.3). Its kernel reads on  $\Lambda^2$  as  $(\mathbf{X} \cdot \mathbf{e}_l)(\mathbf{x}, \mathbf{y}) = x_l \delta_{\mathbf{x}, \mathbf{y}}$ ,  $l = 1, 2$ . The remainder of the proof only is rearranging of terms. Note that  $\mathcal{S}_{1,2,1,2}(\xi)$  in (3.6) can be replaced with  $\mathcal{S}_{1,2,2,1}(\xi)$  or  $\mathcal{S}_{2,1,1,2}(\xi)$  or  $\mathcal{S}_{2,1,2,1}(\xi)$  due to the above arguments.  $\blacksquare$

It turns out that (3.6) eventually is suitable for the Bloch-Floquet decomposition which is at the root of the formula (1.11). Let us recall the framework of it, see e.g. [35, Sect. XIII.16]. Let  $\Omega^*$  be the Brillouin zone of the dual lattice  $\Upsilon^*$ . Introduce the 'Bloch isometry' as:

$$\begin{aligned} \mathcal{U} : l_c^2(\Lambda) & \mapsto \mathfrak{h} := \int_{\Omega^*}^{\oplus} d\mathbf{k} l^2(\Theta) \\ (\mathcal{U}\psi)(\underline{\mathbf{x}}; \mathbf{k}) & = \frac{1}{|\Omega^*|^{\frac{1}{2}}} \sum_{\mathbf{v} \in \Upsilon} e^{-i\mathbf{k} \cdot (\underline{\mathbf{x}} + \mathbf{v})} \psi(\underline{\mathbf{x}} + \mathbf{v}) \quad \mathbf{k} \in \Omega^*, \underline{\mathbf{x}} \in \Theta, \psi \in l_c^2(\Lambda), \end{aligned}$$

which can be extended by continuity in an unitary operator on  $l^2(\Lambda)$ . The adjoint of  $\mathcal{U}$  reads as:

$$(\mathcal{U}^* \phi)(\underline{\mathbf{x}} + \mathbf{v}) = \frac{1}{|\Omega^*|^{\frac{1}{2}}} \int_{\Omega^*} d\mathbf{k} e^{i\mathbf{k} \cdot (\underline{\mathbf{x}} + \mathbf{v})} \phi(\underline{\mathbf{x}}; \mathbf{k}) \quad \underline{\mathbf{x}} \in \Theta, \mathbf{v} \in \Upsilon, \phi(\cdot; \mathbf{k}) \in l^2(\Theta).$$

Since  $H_0$  commutes with the translations of  $\Upsilon$ , then the unitary transformation of  $H_0$  is decomposable into a direct integral  $\mathcal{U}H_0\mathcal{U}^* = \int_{\Omega^*}^{\oplus} d\mathbf{k} H(\mathbf{k})$ . For each  $\mathbf{k} \in \Omega^*$ , the fiber Hamiltonian  $H(\mathbf{k})$  acts on  $l^2(\Theta)$  and it is defined via its kernel which reads as, see e.g. [6, Eq. (2.4)]:

$$\forall(\underline{\mathbf{x}}, \underline{\mathbf{y}}) \in \Theta^2, \quad H(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) := \sum_{\mathbf{v} \in \Upsilon} e^{-i\mathbf{k} \cdot (\underline{\mathbf{x}} + \mathbf{v} - \underline{\mathbf{y}})} H_0(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}}) \quad \mathbf{k} \in \Omega^*.$$

Let us note that for each  $\mathbf{k} \in \Omega^*$ , the fiber  $H(\mathbf{k})$  is a self-adjoint operator on  $l^2(\Theta)$  due to the identity  $H_0(\mathbf{x}, \mathbf{y}) = \overline{H_0(\mathbf{y}, \mathbf{x})}$  (see assumption (P1)) which implies the one  $H(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) = \overline{H(\underline{\mathbf{y}}, \underline{\mathbf{x}}; \mathbf{k})}$ . Hereafter we denote by  $R(\mathbf{k}; \xi) := (H(\mathbf{k}) - \xi)^{-1} \forall \xi \in \rho(H_0)$  the fiber of the resolvent on  $l^2(\Theta)$ .

We now can start the actual proof of Theorem 1.2 (i). In view of (3.6), we need an expression for the fiber of the operator  $\mathcal{V}_{q,r,s,t}(\xi)$ ,  $\mathcal{V} = \mathcal{S}$  or  $\mathcal{T}$ . Below we use the shorthand notation  $Q_0$  to denote equally  $H_0$  or  $R_0(\xi)$ ,  $\xi \in \rho(H_0)$ . Due to the assumption (P3), then  $[\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0]$  with  $\alpha = 1, 2$ , also commute with the lattice translations. By induction, ditto for  $[\mathbf{X} \cdot \mathbf{e}_\gamma, [\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0]]$  with  $\alpha, \gamma = 1, 2$ . Therefore the unitary transformation of  $\mathcal{V}_{q,r,s,t}(\xi)$  is decomposable into a direct integral  $\mathcal{U}\mathcal{V}_{q,r,s,t}(\xi)\mathcal{U}^* = \int_{\Omega^*}^{\oplus} d\mathbf{k} \mathcal{V}_{q,r,s,t}(\mathbf{k}; \xi)$ . For each  $\mathbf{k} \in \Omega^*$  the fiber  $\mathcal{V}_{q,r,s,t}(\mathbf{k}; \xi)$ ,  $\mathcal{V} = \mathcal{S}$  or  $\mathcal{T}$  lives on  $l^2(\Theta)$  and it is defined as the following:

$$\mathcal{S}_{q,r,s,t}(\mathbf{k}; \xi) := R(\mathbf{k}; \xi) ([\mathbf{X} \cdot \mathbf{e}_q, [\mathbf{X} \cdot \mathbf{e}_r, H_0]])(\mathbf{k}) ([\mathbf{X} \cdot \mathbf{e}_s, [\mathbf{X} \cdot \mathbf{e}_t, R_0(\xi)]](\mathbf{k}),$$

$$\mathcal{T}_{q,r,s,t}(\mathbf{k}; \xi) := R(\mathbf{k}; \xi) ([\mathbf{X} \cdot \mathbf{e}_q, H_0])(\mathbf{k}) ([\mathbf{X} \cdot \mathbf{e}_r, R_0(\xi)])(\mathbf{k}) ([\mathbf{X} \cdot \mathbf{e}_s, H_0])(\mathbf{k}) ([\mathbf{X} \cdot \mathbf{e}_t, R_0(\xi)])(\mathbf{k}).$$

Here the fibers  $([\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0])(\mathbf{k})$  and  $([\mathbf{X} \cdot \mathbf{e}_\gamma, [\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0]])(\mathbf{k})$  with  $Q_0 = H_0$  or  $R_0(\xi)$  and  $\alpha, \gamma = 1, 2$  are defined through their kernel which read for each  $\mathbf{k} \in \Omega^*$  and any  $(\underline{\mathbf{x}}, \underline{\mathbf{y}}) \in \Theta^2$  respectively as:

$$\begin{aligned} ([\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0])(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) &:= \sum_{\mathbf{v} \in \Upsilon} e^{-i\mathbf{k} \cdot (\underline{\mathbf{x}} + \mathbf{v} - \underline{\mathbf{y}})} ([\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0])(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}}), \\ ([\mathbf{X} \cdot \mathbf{e}_\gamma, [\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0]])(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) &:= \sum_{\mathbf{v} \in \Upsilon} e^{-i\mathbf{k} \cdot (\underline{\mathbf{x}} + \mathbf{v} - \underline{\mathbf{y}})} ([\mathbf{X} \cdot \mathbf{e}_\gamma, [\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0]])(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}}). \end{aligned}$$

Let  $Q_0(\cdot, \cdot)$  be the kernel of  $Q_0$ . Since  $([\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0])(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}}) = (\underline{x}_\alpha + v_\alpha - \underline{y}_\alpha) Q_0(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}})$  and  $([\mathbf{X} \cdot \mathbf{e}_\gamma, [\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0]])(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}}) = (\underline{x}_\gamma + v_\gamma - \underline{y}_\gamma) (\underline{x}_\alpha + v_\alpha - \underline{y}_\alpha) Q_0(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}})$ , then one has on  $\Theta^2$ :

$$([\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0])(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) = i \frac{\partial}{\partial k_\alpha} Q(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}), \quad ([\mathbf{X} \cdot \mathbf{e}_\gamma, [\mathbf{X} \cdot \mathbf{e}_\alpha, Q_0]])(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) = (i \frac{\partial}{\partial k_\gamma}) i \frac{\partial}{\partial k_\alpha} Q(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}),$$

$$\text{where: } \forall(\underline{\mathbf{x}}, \underline{\mathbf{y}}) \in \Theta^2, \quad Q(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) := \sum_{\mathbf{v} \in \Upsilon} e^{-i\mathbf{k} \cdot (\underline{\mathbf{x}} + \mathbf{v} - \underline{\mathbf{y}})} Q_0(\underline{\mathbf{x}} + \mathbf{v}, \underline{\mathbf{y}}) \quad \mathbf{k} \in \Omega^*.$$

Here these identifications are made possible since the kernels of  $H_0$  and  $R_0(\xi)$  are *EAD*. Denoting by  $(\partial_{k_\alpha} Q)(\mathbf{k})$  and  $(\partial_{k_\gamma} \partial_{k_\alpha} Q)(\mathbf{k})$  the operators on  $l^2(\Theta)$  generated respectively by the kernels:

$$(\partial_{k_\alpha} Q)(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) := \frac{\partial}{\partial k_\alpha} Q(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}), \quad (\partial_{k_\gamma} \partial_{k_\alpha} Q)(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) := \frac{\partial^2}{\partial k_\gamma \partial k_\alpha} Q(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) = \frac{\partial^2}{\partial k_\alpha \partial k_\gamma} Q(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}),$$

then for each  $\mathbf{k} \in \Omega^*$  the fiber  $\mathcal{V}_{q,r,s,t}(\mathbf{k}; \xi)$ ,  $\mathcal{V} = \mathcal{S}$  or  $\mathcal{T}$  can be rewritten as:

$$\mathcal{S}_{q,r,s,t}(\mathbf{k}; \xi) = R(\mathbf{k}; \xi) (\partial_{k_q} \partial_{k_r} H)(\mathbf{k}) (\partial_{k_s} \partial_{k_t} R)(\mathbf{k}; \xi), \quad (3.7)$$

$$\mathcal{T}_{q,r,s,t}(\mathbf{k}; \xi) = R(\mathbf{k}; \xi) (\partial_{k_q} H)(\mathbf{k}) (\partial_{k_r} R)(\mathbf{k}; \xi) (\partial_{k_s} H)(\mathbf{k}) (\partial_{k_t} R)(\mathbf{k}; \xi). \quad (3.8)$$

Since the kernel of  $\mathcal{V}_{q,r,s,t}(\xi)$  is related to the one of the fiber  $\mathcal{V}_{q,r,s,t}(\mathbf{k}; \xi)$  by:

$$\forall(\mathbf{x}, \mathbf{y}) \in \Lambda^2, \quad (\mathcal{V}_{q,r,s,t}(\xi))(\mathbf{x}, \mathbf{y}) = \frac{1}{|\Omega^*|} \int_{\Omega^*} d\mathbf{k} e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})} (\mathcal{V}_{q,r,s,t}(\mathbf{k}, \xi))(\mathbf{x}, \mathbf{y}) \quad \mathcal{V} = \mathcal{S} \text{ or } \mathcal{T},$$

where we used [14, Eq. (4.1)], then the expression in (3.6) can be rewritten as:

$$\begin{aligned} \mathcal{X}(\beta, \rho_0, 0) &= \frac{1}{4} \left( \frac{\epsilon}{c} \right)^2 \frac{1}{\beta \text{Card}(\Theta) |\Omega^*|} \frac{i}{2\pi} \int_{\Gamma} d\xi f(\beta, \mu^{(0)}; \xi) \int_{\Omega^*} d\mathbf{k} \sum_{\underline{\mathbf{x}} \in \Theta} \left\{ \frac{1}{2} \mathcal{S}_{2,2,1,1}(\mathbf{k}; \xi) + \right. \\ &+ \left. \frac{1}{2} \mathcal{S}_{1,1,2,2}(\mathbf{k}; \xi) - \mathcal{S}_{1,2,1,2}(\mathbf{k}; \xi) - \mathcal{T}_{2,1,2,1}(\mathbf{k}; \xi) + \mathcal{T}_{2,1,1,2}(\mathbf{k}; \xi) + \mathcal{T}_{1,2,2,1}(\mathbf{k}; \xi) - \mathcal{T}_{1,2,1,2}(\mathbf{k}; \xi) \right\} (\underline{\mathbf{x}}, \underline{\mathbf{x}}). \end{aligned}$$

Next it remains to insert in (3.7) and (3.8) these both identities:

$$\begin{aligned} (\partial_{k_\alpha} R)(\mathbf{k}; \xi) &= -R(\mathbf{k}; \xi) (\partial_{k_\alpha} H)(\mathbf{k}) R(\mathbf{k}; \xi) \quad \mathbf{k} \in \Omega^*, \alpha = 1, 2, \\ (\partial_{k_\gamma} \partial_{k_\alpha} R)(\mathbf{k}; \xi) &= R(\mathbf{k}; \xi) (\partial_{k_\gamma} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_\alpha} H)(\mathbf{k}) R(\mathbf{k}; \xi) - R(\mathbf{k}; \xi) (\partial_{k_\gamma} \partial_{k_\alpha} H)(\mathbf{k}) R(\mathbf{k}; \xi) + \\ &+ R(\mathbf{k}; \xi) (\partial_{k_\alpha} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_\gamma} H)(\mathbf{k}) R(\mathbf{k}; \xi) \quad \mathbf{k} \in \Omega^*, \alpha, \gamma = 1, 2. \end{aligned}$$

Thus (1.11) follows by gathering together the terms by decreasing number of fiber of resolvent.

### 3.2 Proof of (ii) – (iii).

The starting point is the expression of the bulk zero-field orbital susceptibility at fixed positive temperature and density obtained in (1.11). In view of (1.15) we need to subject this formula to some transformations again so as to perform the integration w.r.t. the  $\xi$ -variable. Only after we will perform the zero-temperature limit. Let us remark that the presence of fiber resolvents in (1.11) indicates the possibility to bring into play, through their kernel, the eigenvalues and associated eigenfunctions of the fiber Hamiltonian. As we will see below, this will turn out to be essential to perform the integral w.r.t. the  $\xi$ -variable by the residue theorem.

Before beginning, let us fix notation. For each  $\mathbf{k} \in \Omega^*$ , denote by  $\{E_j(\mathbf{k})\}_{j=1}^{\text{Card}(\Theta)}$  the set of eigenvalues of the fiber Hamiltonian  $H(\mathbf{k})$  counting multiplicities and in increasing order. Due to this choice of indexation, the  $E_j$ 's are  $\Upsilon^*$ -periodic and continuous functions, but they are not differentiable at (possible) crossing-points. Denote by  $\{u_j(\cdot; \mathbf{k})\}_{j=1}^{\text{Card}(\Theta)}$  the set of associated eigenfunctions. They form a complete orthonormal system in  $l^2(\Theta)$ . Unlike the  $E_j$ 's, the eigenfunctions  $\mathbf{k} \mapsto u_j(\cdot; \mathbf{k})$  may be not continuous at (possible) crossing-points since they are defined up to a  $\mathbf{k}$ -dependent phase factor which cannot be always chosen to be continuous at crossing points.

For any integer  $l, m = 1, \dots, \text{Card}(\Theta)$  and  $\alpha, \gamma = 1, 2$ , let us define the quantities:

$$\hat{\pi}_{l,m}(\alpha; \mathbf{k}) := \sum_{\underline{\mathbf{x}} \in \Theta} \overline{u_l(\underline{\mathbf{x}}; \mathbf{k})} [(\partial_{k_\alpha} H)(\mathbf{k}) u_m(\underline{\mathbf{x}}; \mathbf{k})] \quad \mathbf{k} \in \Omega^*, \quad (3.9)$$

$$\hat{\sigma}_{l,m}(\alpha, \gamma; \mathbf{k}) := \sum_{\underline{\mathbf{x}} \in \Theta} \overline{u_l(\underline{\mathbf{x}}; \mathbf{k})} [(\partial_{k_\alpha} \partial_{k_\gamma} H)(\mathbf{k}) u_m(\underline{\mathbf{x}}; \mathbf{k})] \quad \mathbf{k} \in \Omega^*. \quad (3.10)$$

Due to the phase factors presence in the  $u_j$ 's, the both above quantities may not be continuous at (possible) crossing points. Besides from (3.9), one has  $\hat{\pi}_{l,m}(\alpha; \mathbf{k}) = \overline{\hat{\pi}_{m,l}(\alpha; \mathbf{k})}$  since:

$$\sum_{\underline{\mathbf{x}} \in \Theta} \overline{u_l(\underline{\mathbf{x}}; \mathbf{k})} \sum_{\underline{\mathbf{y}} \in \Theta} \frac{\partial}{\partial k_\alpha} H(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) u_m(\underline{\mathbf{y}}; \mathbf{k}) = \sum_{\underline{\mathbf{y}} \in \Theta} u_m(\underline{\mathbf{y}}; \mathbf{k}) \overline{\sum_{\underline{\mathbf{x}} \in \Theta} \frac{\partial}{\partial k_\alpha} H(\underline{\mathbf{y}}, \underline{\mathbf{x}}; \mathbf{k}) u_l(\underline{\mathbf{x}}; \mathbf{k})},$$

where we used that  $H(\underline{\mathbf{x}}, \underline{\mathbf{y}}; \mathbf{k}) = \overline{H(\underline{\mathbf{y}}, \underline{\mathbf{x}}; \mathbf{k})} \forall (\underline{\mathbf{x}}, \underline{\mathbf{y}}) \in \Theta^2$ . The same property holds for  $\hat{\sigma}_{l,m}(\alpha, \gamma; \mathbf{k})$ .

The sequel of this paragraph is divided into two parts. In the first one, we treat the generical situation in which the  $E_j(\cdot)$ 's are non-degenerate outside a subset of  $\Omega^*$  with Lebesgue-measure zero. In the second one, we restrict the number of sites in  $\Theta$  to two sites and we consider the situation in which both  $E_1(\cdot)$  and  $E_2(\cdot)$  are non-degenerate everywhere on  $\Omega^*$ .

### 3.2.1 Proof of (ii).

In what follows we tacitly suppose the assumption (A1), see page 5. Here is the last rewriting of the zero-field susceptibility before the zero-temperature limit:

**Proposition 3.2.** *Let  $\beta > 0$  and  $\rho_0 > 0$  be fixed. Let  $\mu^{(0)} = \mu^{(0)}(\beta, \mu, 0) \in \mathbb{R}$  be the unique solution of the equation  $\rho(\beta, e^{\beta\mu}, 0) = \rho_0$ . Then for each integer  $j = 1, \dots, \text{Card}(\Theta)$  there exists four families of functions  $\mathfrak{d}_{j,l}(\cdot)$  with  $l = 0, 1, 2, 3$  defined on  $\Omega^*$  outside a set of Lebesgue-measure zero, s.t. the integrand below can be extended by continuity to the whole of  $\Omega^*$ , and:*

$$\mathcal{X}(\beta, \rho_0, 0) = -\frac{1}{4} \left( \frac{e}{c} \right)^2 \frac{1}{\text{Card}(\Theta) |\Omega^*| \beta} \frac{1}{\sum_{j=1}^{\text{Card}(\Theta)}} \int_{\Omega^*} d\mathbf{k} \sum_{l=0}^3 \frac{\partial^l \mathfrak{f}}{\partial \xi^l}(\beta, \mu^{(0)}; E_j(\mathbf{k})) \mathfrak{d}_{j,l}(\mathbf{k}), \quad (3.11)$$

with the convention  $(\partial_\xi^0 \mathfrak{f})(\beta, \mu^{(0)}; E_j(\mathbf{k})) := \mathfrak{f}(\beta, \mu^{(0)}; E_j(\mathbf{k})) = \ln(1 + e^{\beta(\mu^{(0)} - E_j(\mathbf{k}))})$ .

The special feature of this formula lies in the fact that each function  $\mathfrak{d}_{j,l}(\cdot)$  can be only expressed in terms of eigenvalues and associated eigenfunctions of the Bloch Hamiltonian  $H(\mathbf{k})$ , together with the derivatives (up to the second order) w.r.t. the  $k_\alpha$ -variables ( $\alpha = 1, 2$ ) of the  $\text{Card}(\Theta) \times \text{Card}(\Theta)$ -matrix elements of  $H(\mathbf{k})$ . By way of example, the functions  $\mathfrak{d}_{j,3}(\cdot)$ ,  $j = 1, \dots, \text{Card}(\Theta)$  are identified in (3.22). Note that each and every one functions  $\mathfrak{d}_{j,l}(\cdot)$ ,  $l = 0, 1, 2$  can also be written down, but their explicit expression is not necessary for the proof of Theorem 1.2 (ii). Further, the expansion obtained in (3.11) makes the choice of phase for the  $u_j(\cdot; \mathbf{k})$ 's irrelevant.

Now we give the proof of Proposition 3.2; it follows the outline of the proof of [5, Thm 3.1]. For reader's convenience we collect in the appendix of this section the proofs of intermediary results. For the sake of simplicity, let us rewrite the zero-field orbital susceptibility formula in (1.11) as:

$$\mathcal{X}(\beta, \rho_0, 0) = \frac{1}{4} \left( \frac{e}{c} \right)^2 \frac{1}{\text{Card}(\Theta) |\Omega^*| \beta} \sum_{l=3}^5 \mathcal{W}_l(\beta, \mu^{(0)}, 0), \quad (3.12)$$

$$\text{where: } \mathcal{W}_l(\beta, \mu^{(0)}, 0) := \frac{i}{2\pi} \int_{\Gamma} d\xi \mathfrak{f}(\beta, \mu^{(0)}; \xi) \int_{\Omega^*} d\mathbf{k} \text{Tr}_{l^2(\Theta)}(\mathcal{P}_l(\mathbf{k}; \xi)) \quad l = 3, 4, 5.$$

The first step consists in expressing the kernel of each fiber resolvent appearing in  $\mathcal{P}_l(\mathbf{k}; \xi)$ ,  $l = 3, 4, 5$  (see formulas (1.14)-(1.12)) in terms of eigenvalues and eigenfunctions of the fiber Hamiltonian (via the spectral theorem). Thus the quantities  $\mathcal{W}_l(\beta, \mu^{(0)}, 0)$  can be rewritten as:

**Lemma 3.3.** *Let  $\beta > 0$  and  $\rho_0 > 0$  be fixed. Then one has:*

$$\begin{aligned} \mathcal{W}_5(\beta, \mu^{(0)}, 0) = & - \sum_{j_1, \dots, j_4=1}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \mathcal{C}_{j_1, j_2, j_3, j_4}(\mathbf{k}) \times \\ & \times \frac{1}{2i\pi} \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi) (E_{j_3}(\mathbf{k}) - \xi) (E_{j_4}(\mathbf{k}) - \xi)}, \end{aligned} \quad (3.13)$$

$$\begin{aligned} \mathcal{W}_4(\beta, \mu^{(0)}, 0) = & - \sum_{j_1, \dots, j_3=1}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \mathcal{C}_{j_1, j_2, j_3}(\mathbf{k}) \times \\ & \times \frac{1}{2i\pi} \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi) (E_{j_3}(\mathbf{k}) - \xi)}, \end{aligned} \quad (3.14)$$

$$\mathcal{W}_3(\beta, \mu^{(0)}, 0) = - \sum_{j_1, j_2=1}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \mathcal{C}_{j_1, j_2}(\mathbf{k}) \frac{1}{2i\pi} \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi)}, \quad (3.15)$$

where  $\forall \mathbf{k} \in \Omega^*$ , the functions  $\mathcal{C}_{j_1, j_2, j_3, j_4}(\cdot)$ ,  $\mathcal{C}_{j_1, j_2, j_3}(\cdot)$  and  $\mathcal{C}_{j_1, j_2}(\cdot)$  are respectively defined by:

$$\begin{aligned} \mathcal{C}_{j_1, j_2, j_3, j_4}(\mathbf{k}) := & \{\hat{\pi}_{j_1, j_2}(1; \mathbf{k})\hat{\pi}_{j_2, j_3}(2; \mathbf{k}) - \hat{\pi}_{j_1, j_2}(2; \mathbf{k})\hat{\pi}_{j_2, j_3}(1; \mathbf{k})\} \times \\ & \times \{\hat{\pi}_{j_3, j_4}(2; \mathbf{k})\hat{\pi}_{j_4, j_1}(1; \mathbf{k}) - \hat{\pi}_{j_3, j_4}(1; \mathbf{k})\hat{\pi}_{j_4, j_1}(2; \mathbf{k})\}, \end{aligned} \quad (3.16)$$

$$\begin{aligned} \mathcal{C}_{j_1, j_2, j_3}(\mathbf{k}) := & \hat{\sigma}_{j_1, j_2}(1, 1; \mathbf{k})\hat{\pi}_{j_2, j_3}(2; \mathbf{k})\hat{\pi}_{j_3, j_1}(2; \mathbf{k}) + \hat{\sigma}_{j_1, j_2}(2, 2; \mathbf{k})\hat{\pi}_{j_2, j_3}(1; \mathbf{k})\hat{\pi}_{j_3, j_1}(1; \mathbf{k}) + \\ & - \hat{\sigma}_{j_1, j_2}(1, 2; \mathbf{k})\hat{\pi}_{j_2, j_3}(1; \mathbf{k})\hat{\pi}_{j_3, j_1}(2; \mathbf{k}) - \hat{\sigma}_{j_1, j_2}(1, 2; \mathbf{k})\hat{\pi}_{j_2, j_3}(2; \mathbf{k})\hat{\pi}_{j_3, j_1}(1; \mathbf{k}), \end{aligned} \quad (3.17)$$

$$\mathcal{C}_{j_1, j_2}(\mathbf{k}) := -\Re\{\hat{\sigma}_{j_1, j_2}(1, 1; \mathbf{k})\hat{\sigma}_{j_2, j_1}(2, 2; \mathbf{k})\} + \hat{\sigma}_{j_1, j_2}(1, 2; \mathbf{k})\hat{\sigma}_{j_2, j_1}(1, 2; \mathbf{k}). \quad (3.18)$$

The quantities  $\hat{\pi}_{l, m}(\alpha; \mathbf{k})$  and  $\hat{\sigma}_{l, m}(\alpha, \gamma; \mathbf{k})$ ,  $\alpha, \gamma = 1, 2$  are respectively defined in (3.9) and (3.10).

The second step consists in applying the residue theorem in (3.13)-(3.15). It directly provides:

**Lemma 3.4.** For each integer  $j_1 = 1, \dots, \text{Card}(\Theta)$ , there exists four families of functions  $\mathbf{a}_{j_1, l}(\cdot)$  with  $l = 0, 1, 2, 3$  defined on  $\Omega^*$  outside a set of Lebesgue-measure zero s.t.:

$$\mathcal{W}_5(\beta, \mu^{(0)}, 0) = - \sum_{j_1=1}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \sum_{l=0}^3 \frac{\partial^l \mathbf{f}}{\partial \xi^l}(\beta, \mu^{(0)}; E_{j_1}(\mathbf{k})) \mathbf{a}_{j_1, l}(\mathbf{k}). \quad (3.19)$$

In particular, for Lebesgue-almost all  $\mathbf{k} \in \Omega^*$  the functions  $\mathbf{a}_{j_1, 3}(\cdot)$  is defined by:

$$\mathbf{a}_{j_1, 3}(\mathbf{k}) := \frac{1}{3!} \sum_{\substack{j_2=1 \\ j_2 \neq j_1}}^{\text{Card}(\Theta)} \frac{\mathcal{C}_{j_1, j_1, j_2, j_1}(\mathbf{k})}{E_{j_2}(\mathbf{k}) - E_{j_1}(\mathbf{k})}.$$

**Lemma 3.5.** For each integer  $j_1 = 1, \dots, \text{Card}(\Theta)$ , there exists four families of functions  $\mathbf{b}_{j_1, l}(\cdot)$  with  $l = 0, 1, 2, 3$  defined on  $\Omega^*$  outside a set of Lebesgue-measure zero s.t.:

$$\mathcal{W}_4(\beta, \mu^{(0)}, 0) = - \sum_{j_1=1}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \sum_{l=0}^3 \frac{\partial^l \mathbf{f}}{\partial \xi^l}(\beta, \mu^{(0)}; E_{j_1}(\mathbf{k})) \mathbf{b}_{j_1, l}(\mathbf{k}). \quad (3.20)$$

In particular, for (Lebesgue-almost) all  $\mathbf{k} \in \Omega^*$  the function  $\mathbf{b}_{j_1, 3}(\cdot)$  is defined by:

$$\mathbf{b}_{j_1, 3}(\mathbf{k}) := \frac{1}{3!} \mathcal{C}_{j_1, j_1, j_1}(\mathbf{k}).$$

**Lemma 3.6.** For each integer  $j_1 = 1, \dots, \text{Card}(\Theta)$ , there exists three families of functions  $\mathbf{c}_{j_1, l}(\cdot)$  with  $l = 0, 1, 2$  defined on  $\Omega^*$  outside a set of Lebesgue-measure zero s.t.:

$$\mathcal{W}_3(\beta, \mu^{(0)}, 0) = - \sum_{j_1=1}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \sum_{l=0}^2 \frac{\partial^l \mathbf{f}}{\partial \xi^l}(\beta, \mu^{(0)}; E_{j_1}(\mathbf{k})) \mathbf{c}_{j_1, l}(\mathbf{k}). \quad (3.21)$$

We stress the point that each and every one functions appearing in the expansions (3.19)-(3.21) can be written down since a such result is based only on identities provided by the residue theorem. This will be done in the second part of this paragraph when considering the case of  $\text{Card}(\Theta) = 2$ .

Thus Lemmas 3.4, 3.5 and 3.6 provide an expansion of the type announced in (3.11), where the coefficients read for  $j_1 = 1, \dots, \text{Card}(\Theta)$  and almost everywhere on  $\Omega^*$  as:

$$\mathfrak{d}_{j_1, 3}(\mathbf{k}) := \frac{1}{3!} \left\{ \sum_{\substack{j_2=1 \\ j_2 \neq j_1}}^{\text{Card}(\Theta)} \frac{\mathcal{C}_{j_1, j_1, j_2, j_1}(\mathbf{k})}{E_{j_2}(\mathbf{k}) - E_{j_1}(\mathbf{k})} + \mathcal{C}_{j_1, j_1, j_1}(\mathbf{k}) \right\}, \quad (3.22)$$

$$\mathfrak{d}_{j_1, l}(\mathbf{k}) := \mathbf{a}_{j_1, l}(\mathbf{k}) + \mathbf{b}_{j_1, l}(\mathbf{k}) + \mathbf{c}_{j_1, l}(\mathbf{k}) \quad l = 0, 1, 2. \quad (3.23)$$

To conclude the proof of Proposition 3.2, we need the following result:

**Lemma 3.7.** *The functions  $\mathfrak{a}_{j_1,l}(\cdot)$ ,  $\mathfrak{b}_{j_1,l}(\cdot)$  with  $l = 0, 1, 2, 3$  and  $\mathfrak{c}_{j_1,l}(\cdot)$ ,  $l = 0, 1, 2$  are bounded and continuous on any compact subset of  $\Omega^*$  where  $E_{j_1}$  is isolated from the rest of the spectrum.*

We do not give a proof for Lemma 3.7 since all details can be found in the proof of [5, Lem. 3.7]. Thereby the functions  $\mathfrak{d}_{j_1,l}(\cdot)$  in (3.22)-(3.23) might be singular on a set of Lebesgue-measure zero where  $E_{j_1}$  can touch the neighboring bands. However the whole integrand in (3.11) comes from the complex integrals (3.13)-(3.15) which have no local singularities in  $\mathbf{k}$  (they eventually are bounded uniformly in  $\mathbf{k}$ , see the proof of Lemma 3.3). The proof of Proposition 3.2 is now over.

We now can start the actual proof of Theorem 1.2 (ii); it follows the outline of [5, Sect. 4.1]. Let  $\rho_0 > 0$  be fixed. Let  $\mu^{(0)} = \mu^{(0)}(\beta, \rho_0, 0) \in \mathbb{R}$  be the unique solution of  $\rho(\beta, e^{\beta\mu}, 0) = \rho_0$ . Let  $\mathfrak{f}_{FD}(\beta, \mu^{(0)}; \xi) := (e^{\beta(\xi - \mu^{(0)})} + 1)^{-1} = -\beta^{-1}(\partial_\xi \mathfrak{f})(\beta, \mu^{(0)}; \xi)$  be the Fermi-Dirac (F-D) distribution. By involving the F-D distribution and its derivatives, the expansion (3.11) can be rewritten as:

$$\mathcal{X}(\beta, \rho_0, 0) = \frac{1}{4} \left( \frac{e}{c} \right)^2 \frac{1}{\text{Card}(\Theta)|\Omega^*|} \sum_{j=1}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \left\{ \sum_{l=0}^2 \frac{\partial^l \mathfrak{f}_{FD}}{\partial \xi^l}(\beta, \mu^{(0)}; E_j(\mathbf{k})) \mathfrak{d}_{j,l+1}(\mathbf{k}) + \right. \\ \left. - \frac{1}{\beta} \mathfrak{f}(\beta, \mu^{(0)}; E_j(\mathbf{k})) \mathfrak{d}_{j,0}(\mathbf{k}) \right\}. \quad (3.24)$$

Consider the semiconducting situation, i.e. assume that the Fermi energy lies in the middle of a non-trivial gap. Then there exists  $M \in \{1, \dots, \text{Card}(\Theta) - 1\}$  such that  $\mathcal{E}_M < \mathcal{E}_{M+1}$ , and moreover,  $\lim_{\beta \rightarrow \infty} \mu^{(0)}(\beta, \rho_0) = (\max \mathcal{E}_M + \min \mathcal{E}_{M+1})/2$ . From the pointwise convergences [5, Eqs. (4.2)-(4.3)] together with the estimates [5, Eq. (4.4)], then all terms in (3.24) containing derivatives of the Fermi-Dirac distribution will converge to zero in the limit  $\beta \rightarrow \infty$ . This leads to (1.15).

### 3.2.2 Proof of (iii).

In this paragraph, we implicitly suppose the assumption (A2), see page 5. Supposing (A2) instead of (A1) does not bring anything to the expansion obtained in (3.11), with the only difference that the functions  $\mathfrak{d}_{j,l}(\cdot)$ ,  $j = 1, \dots, \text{Card}(\Theta)$  and  $l = 0, 1, 2, 3$  (which are the same) are in this instance smooth w.r.t. the  $\mathbf{k}$ -variable. That is ensured by the analytic perturbation theory, see e.g. [35, Sect. XII] and also [29, Thm 3.5]. Nevertheless, considering the particular case of  $\text{Card}(\Theta) = 2$  allows us to write down concise and explicit formulas for the functions appearing in an expansion of type (3.11). Here is the counterpart of Proposition 3.2 when  $\text{Card}(\Theta) = 2$ :

**Lemma 3.8.** *Let  $\beta > 0$  and  $\rho_0 > 0$  be fixed. Let  $\mu^{(0)} = \mu^{(0)}(\beta, \mu, 0) \in \mathbb{R}$  be the unique solution of the equation  $\rho(\beta, e^{\beta\mu}, 0) = \rho_0$ . Then:*

$$\mathcal{X}(\beta, \rho_0, 0) = -\frac{1}{4} \left( \frac{e}{c} \right)^2 \frac{1}{2|\Omega^*|} \frac{1}{\beta} \sum_{j=1}^2 \int_{\Omega^*} d\mathbf{k} \sum_{l=0}^3 \frac{\partial^l \mathfrak{f}}{\partial \xi^l}(\beta, \mu^{(0)}; E_j(\mathbf{k})) \hat{\mathfrak{d}}_{j,l}(\mathbf{k}), \quad (3.25)$$

where  $\forall \mathbf{k} \in \Omega^*$ , the functions  $\hat{\mathfrak{d}}_{j,l}(\cdot)$ ,  $j = 1, 2$ ,  $l = 0, 1, 2, 3$  are given by (below  $m = 1, 2$ ,  $m \neq j$ ):

$$\hat{\mathfrak{d}}_{j,3}(\mathbf{k}) := \frac{1}{3!} \left\{ \frac{\mathcal{C}_{j,j,m,j}(\mathbf{k})}{E_m(\mathbf{k}) - E_j(\mathbf{k})} + \mathcal{C}_{j,j,j}(\mathbf{k}) \right\}, \quad (3.26)$$

$$\hat{\mathfrak{d}}_{j,2}(\mathbf{k}) := -\frac{1}{2} \left\{ \frac{\mathcal{C}_{j,j,m,m}(\mathbf{k}) + \mathcal{C}_{j,m,j,m}(\mathbf{k}) + \mathcal{C}_{j,m,m,j}(\mathbf{k})}{(E_m(\mathbf{k}) - E_j(\mathbf{k}))^2} + \frac{\mathcal{C}_{j,m,j}(\mathbf{k}) + \mathcal{C}_{j,j,m}(\mathbf{k})}{E_m(\mathbf{k}) - E_j(\mathbf{k})} + \mathcal{C}_{j,j}(\mathbf{k}) \right\}, \quad (3.27)$$

$$\hat{\mathfrak{d}}_{j,1}(\mathbf{k}) := \sum_{n=0}^2 \frac{u_{j,n}(\mathbf{k})}{(E_m(\mathbf{k}) - E_j(\mathbf{k}))^{n+1}}, \quad \hat{\mathfrak{d}}_{j,0}(\mathbf{k}) := \sum_{n=0}^2 \frac{v_{j,n}(\mathbf{k})}{(E_m(\mathbf{k}) - E_j(\mathbf{k}))^{n+2}}; \quad (3.28)$$

and the functions  $\mathbf{u}_{j,n}(\cdot)$  and  $\mathbf{v}_{j,n}(\cdot)$ ,  $j = 1, 2$ ,  $n = 0, 1, 2$  are given by (below  $m = 1, 2$ ,  $m \neq j$ ):

$$\mathbf{u}_{j,0}(\mathbf{k}) := \mathcal{C}_{j,m}(\mathbf{k}), \quad (3.29)$$

$$\mathbf{u}_{j,1}(\mathbf{k}) := \mathcal{C}_{j,m,m}(\mathbf{k}) + \mathcal{C}_{m,j,j}(\mathbf{k}) - \mathcal{C}_{j,m,j}(\mathbf{k}) - \mathcal{C}_{j,j,m}(\mathbf{k}), \quad (3.30)$$

$$\mathbf{u}_{j,2}(\mathbf{k}) := \mathcal{C}_{j,m,m,m}(\mathbf{k}) - \mathcal{C}_{m,j,j,j}(\mathbf{k}) - \mathcal{C}_{j,j,m,m}(\mathbf{k}) - \mathcal{C}_{j,m,j,m}(\mathbf{k}) - \mathcal{C}_{j,m,m,j}(\mathbf{k}), \quad (3.31)$$

$$\mathbf{v}_{j,0}(\mathbf{k}) := 0, \quad (3.32)$$

$$\mathbf{v}_{j,1}(\mathbf{k}) := 2\mathcal{C}_{j,m,m}(\mathbf{k}) + 2\mathcal{C}_{m,j,j}(\mathbf{k}) - \mathcal{C}_{j,m,j}(\mathbf{k}) - \mathcal{C}_{j,j,m}(\mathbf{k}) - \mathcal{C}_{m,j,m}(\mathbf{k}) - \mathcal{C}_{m,m,j}(\mathbf{k}), \quad (3.33)$$

$$\mathbf{v}_{j,1}(\mathbf{k}) := 2\mathcal{C}_{j,m,m,m}(\mathbf{k}) - 2\mathcal{C}_{m,j,j,j}(\mathbf{k}). \quad (3.34)$$

The functions  $\mathcal{C}_{j_1,j_2,j_3,j_4}(\cdot)$ ,  $\mathcal{C}_{j_1,j_2,j_3}(\cdot)$  and  $\mathcal{C}_{j_1,j_2}(\cdot)$ ,  $j_1, j_2, j_3, j_4 = 1, 2$  are defined in Lemma 3.3.

The above lemma is obtained from Lemma 3.3 by setting  $\text{Card}(\Theta) = 2$ , then by mimicking the proof of Lemmas 3.4-3.6. Since it is simply calculations based on identities provided by the residue theorem, we do not give further details. Let us note that the above functions  $\hat{\mathfrak{d}}_{j,l}(\cdot)$ , with  $j = 1, 2$ ,  $l = 0, 1, 2, 3$  have the same special feature than the one mentioned for the functions  $\mathfrak{d}_{j,l}(\cdot)$  appearing in the expansion (3.11), see below Proposition 3.2.

The non-degeneracy assumption for the  $E_j$ 's,  $j = 1, 2$  allows us to use the regular perturbation theory, which from the Hellman-Feynman formula (see e.g. [18, Thm 4.1]), provides the identities:

$$\frac{\partial E_j(\mathbf{k})}{\partial k_\alpha} = \hat{\pi}_{j,j}(\alpha; \mathbf{k}) \quad \alpha = 1, 2, \quad (3.35)$$

$$\frac{\partial^2 E_j(\mathbf{k})}{\partial k_\alpha^2} = \hat{\sigma}_{j,j}(\alpha, \alpha; \mathbf{k}) + 2 \sum_{m \neq j} \frac{|\hat{\pi}_{m,j}(\alpha; \mathbf{k})|^2}{E_j(\mathbf{k}) - E_m(\mathbf{k})} \quad \alpha = 1, 2, \quad (3.36)$$

$$\frac{\partial^2 E_j(\mathbf{k})}{\partial k_\alpha \partial k_\gamma} = \hat{\sigma}_{j,j}(\alpha, \gamma; \mathbf{k}) + 2 \sum_{m \neq j} \frac{\Re\{\hat{\pi}_{m,j}(\alpha; \mathbf{k})\hat{\pi}_{j,m}(\gamma; \mathbf{k})\}}{E_j(\mathbf{k}) - E_m(\mathbf{k})} = \frac{\partial^2 E_j(\mathbf{k})}{\partial k_\gamma \partial k_\alpha} \quad \alpha, \gamma = 1, 2. \quad (3.37)$$

The derivatives of higher order can also be identified, but we will not use them in the following.

The use of the regular perturbation theory plays a crucial role in the next result: it yields the identification of the interband contributions coming from the expansion (3.25):

**Proposition 3.9.** *Let  $\beta > 0$  and  $\rho_0 > 0$  be fixed. Let  $\mu^{(0)} = \mu^{(0)}(\beta, \mu, 0) \in \mathbb{R}$  be the unique solution of the equation  $\rho(\beta, e^{\beta\mu}, 0) = \rho_0$ . Then (3.25) can be split into two contributions:*

$$\mathcal{X}(\beta, \rho_0, 0) = \mathcal{X}_P(\beta, \rho_0, 0) + \mathcal{X}_{\text{Ib}}(\beta, \rho_0, 0), \quad (3.38)$$

with, by using the shorthand notations  $\frac{\partial^l \mathfrak{f}}{\partial \xi^l}(\beta, \mu^{(0)}; \cdot) = \frac{\partial^l \mathfrak{f}}{\partial \xi^l}(\cdot)$ ,  $l = 0, 1, 2$ :

$$\begin{aligned} \mathcal{X}_P(\beta, \rho_0, 0) &:= -\frac{1}{48} \left(\frac{e}{c}\right)^2 \frac{1}{|\Omega^*|} \frac{1}{\beta} \sum_{j=1}^2 \int_{\Omega^*} d\mathbf{k} \frac{\partial^2 \mathfrak{f}}{\partial \xi^2}(E_j(\mathbf{k})) \left\{ \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1^2} \frac{\partial^2 E_j(\mathbf{k})}{\partial k_2^2} - \left(\frac{\partial^2 E_j(\mathbf{k})}{\partial k_1 \partial k_2}\right)^2 \right\}, \\ \mathcal{X}_{\text{Ib}}(\beta, \rho_0, 0) &:= -\frac{1}{8} \left(\frac{e}{c}\right)^2 \frac{1}{|\Omega^*|} \frac{1}{\beta} \sum_{j=1}^2 \int_{\Omega^*} d\mathbf{k} \left\{ \frac{\partial^2 \mathfrak{f}}{\partial \xi^2}(E_j(\mathbf{k})) \mathfrak{J}_j(\mathbf{k}) + \sum_{l=0}^1 \frac{\partial^l \mathfrak{f}}{\partial \xi^l}(E_j(\mathbf{k})) \hat{\mathfrak{d}}_{j,l}(\mathbf{k}) \right\}, \end{aligned} \quad (3.39)$$

where  $\forall \mathbf{k} \in \Omega^*$ , the function  $\mathfrak{J}_j(\cdot)$ ,  $j = 1, 2$  is given by (below  $m = 1, 2$  with  $m \neq j$ ):

$$\mathfrak{J}_j(\mathbf{k}) := \frac{1}{2} \Re \left\{ -\frac{\mathcal{C}_{j,j,m,m}(\mathbf{k}) + \mathcal{C}_{j,m,j,m}(\mathbf{k}) + \mathcal{C}_{j,m,m,j}(\mathbf{k})}{(E_j(\mathbf{k}) - E_m(\mathbf{k}))^2} + i \frac{\Im\{\mathcal{C}_{j,m,j}(\mathbf{k})\}}{E_j(\mathbf{k}) - E_m(\mathbf{k})} \right\}, \quad (3.40)$$

and the functions  $\hat{\mathfrak{d}}_{j,l}(\cdot)$ , with  $j = 1, 2$  and  $l = 0, 1$  are defined in Lemma 3.8.

The proof of the above result can be found in the appendix of this section. The term  $\mathcal{X}_P(\beta, \rho_0, 0)$  is nothing but the so-called Peierls contribution, see e.g. [33, Eq. (62c)] and [5, Eq. (4.10)]. As for the term  $\mathcal{X}_{\text{Ib}}(\beta, \rho_0, 0)$ , it stands for the interband (or band-to-band) contributions. As we will see below, only the functions multiplying  $(\partial_{\xi}^l \mathfrak{f})(\beta, \mu^{(0)}; \cdot)$  with  $l = 0, 1$  (and therefore, only a great part of these interband contributions) will give rise to (1.16).

We now can start the actual proof of Theorem 1.2 (iii). By using the F-D distribution together with its partial derivatives, (3.25) reads as:

$$\mathcal{X}(\beta, \rho_0, 0) = \frac{1}{8} \left( \frac{e}{c} \right)^2 \frac{1}{|\Omega^*|} \sum_{j=1}^2 \int_{\Omega^*} d\mathbf{k} \left\{ \sum_{l=0}^2 \frac{\partial^l \mathfrak{f}_{FD}}{\partial \xi^l}(\beta, \mu^{(0)}; E_j(\mathbf{k})) \hat{\mathfrak{d}}_{j,l+1}(\mathbf{k}) + \right. \\ \left. - \frac{1}{\beta} \mathfrak{f}(\beta, \mu^{(0)}; E_j(\mathbf{k})) \hat{\mathfrak{d}}_{j,0}(\mathbf{k}) \right\}.$$

Assume that the Fermi energy is in the middle of the gap separating the band  $\mathcal{E}_1$  from  $\mathcal{E}_2$ . From [5, Eqs. (4.2)-(4.3)] together with [5, Eq. (4.4)] again, then all terms in (3.24) containing derivatives of the Fermi-Dirac distribution will converge to zero in the limit  $\beta \rightarrow \infty$ . Hence, we get (1.16). Let us note that the expansion in (3.38) also leads to (1.16) in the semiconducting situation.

### 3.3 Appendix: proofs of the intermediate results.

**Proof of Lemma 3.3.** We start with the quantity  $\mathcal{W}_5(\beta, \mu^{(0)}, 0)$ . Take a generical term of it:

$$w_5(\beta, \mu^{(0)}, 0) := \frac{i}{2\pi} \int_{\Gamma} d\xi \mathfrak{f}(\beta, \mu^{(0)}; \xi) \int_{\Omega^*} d\mathbf{k} \text{Tr}_{l^2(\Theta)}(p_5(\mathbf{k}; \xi)), \quad \text{with:} \\ p_5(\mathbf{k}; \xi) := R(\mathbf{k}; \xi) (\partial_{k_\alpha} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_\gamma} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_\gamma} H)(\mathbf{k}) R(\mathbf{k}; \xi) (\partial_{k_\alpha} H)(\mathbf{k}) R(\mathbf{k}; \xi),$$

where  $\alpha, \gamma = 1, 2$ . Without loss of generality, let us take for the moment  $\gamma = \alpha$  (needed to simplify the notations). By the mean of kernels, the trace of  $p_5(\mathbf{k}; \xi)$  can be written:

$$\sum_{\mathbf{x} \in \Theta} \sum_{\mathbf{z}_1 \in \Theta} R(\mathbf{x}, \mathbf{z}_1; \mathbf{k}, \xi) \left( \prod_{m=1}^3 \sum_{\mathbf{z}_m, \mathbf{z}_{m+1} \in \Theta} (\partial_{k_\alpha} H)(\mathbf{z}_{2m-1}, \mathbf{z}_{2m}; \mathbf{k}) R(\mathbf{z}_{2m}, \mathbf{z}_{2m+1}; \mathbf{k}, \xi) \right) R(\mathbf{z}_8, \mathbf{x}; \mathbf{k}, \xi). \quad (3.41)$$

Next use that the kernel of  $R(\mathbf{k}; \xi)$  express in terms of eigenfunctions of the fiber  $H(\mathbf{k})$  as:

$$\forall (\mathbf{x}, \mathbf{y}) \in \Theta^2, \quad R(\mathbf{x}, \mathbf{y}; \mathbf{k}, \xi) = \sum_{j=1}^{\text{Card}(\Theta)} \frac{u_j(\mathbf{x}; \mathbf{k}) \overline{u_j(\mathbf{y}; \mathbf{k})}}{E_j(\mathbf{k}) - \xi} \quad \mathbf{k} \in \Omega^*.$$

The replacement of each kernel of the resolvent in (3.41) with the above expression leads to:

$$w_5(\beta, \mu^{(0)}, 0) = \sum_{j_1, \dots, j_5=1}^{\text{Card}(\Theta)} \frac{i}{2\pi} \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi) \cdots (E_{j_5}(\mathbf{k}) - \xi)} \int_{\Omega^*} d\mathbf{k} \overbrace{\sum_{\mathbf{x} \in \Theta} \overline{u_{j_5}(\mathbf{x}; \mathbf{k})} u_{j_1}(\mathbf{x}; \mathbf{k})}^{=\delta_{j_1, j_5}} \times \\ \times \prod_{m=1}^4 \sum_{\mathbf{z}_{2m-1} \in \Theta} \overline{u_{j_m}(\mathbf{z}_{2m-1}; \mathbf{k})} \sum_{\mathbf{z}_{2m} \in \Theta} (\partial_{k_\alpha} H)(\mathbf{z}_{2m-1}, \mathbf{z}_{2m}; \mathbf{k}) u_{j_{m+1}}(\mathbf{z}_{2m}; \mathbf{k}).$$

From the definition (3.9), one therefore obtains:

$$w_5(\beta, \mu^{(0)}, 0) = \sum_{j_1, \dots, j_4=1}^{\text{Card}(\Theta)} \frac{i}{2\pi} \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi) \cdots (E_{j_4}(\mathbf{k}) - \xi)} \int_{\Omega^*} d\mathbf{k} c_{j_1, j_2, j_3, j_4}(\mathbf{k}), \quad (3.42)$$



where  $c_{j_1, j_2, j_3, j_4}(\mathbf{k}) := \hat{\pi}_{j_1, j_2}(\alpha; \mathbf{k}) \hat{\pi}_{j_2, j_3}(\gamma; \mathbf{k}) \hat{\pi}_{j_3, j_4}(\gamma; \mathbf{k}) \hat{\pi}_{j_4, j_1}(\alpha; \mathbf{k})$ ,  $\mathbf{k} \in \Omega^*$ . Let us now prove:

$$\int_{\Gamma} |d\xi| \left| c_{j_1, j_2, j_3, j_4}(\mathbf{k}) \frac{f(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi) \cdots (E_{j_4}(\mathbf{k}) - \xi)} \right| \leq C, \quad (3.43)$$

for some  $\mathbf{k}$ -independent constants  $C > 0$ , what allows to change the order of integration in (3.42). We only give the two main ingredients leading to (3.43). Let  $\xi_0 > 0$  so that  $-\xi_0 < \inf \sigma(H_0)$  and large enough. On the one hand, from [14, Eq. (4.8)] there exists a  $C = C_{\xi_0}(\beta, \mu^{(0)}) > 0$  s.t.:

$$\int_{\Gamma} |d\xi| \frac{|f(\beta, \mu^{(0)}; \xi)|}{|E_{j_1}(\mathbf{k}) - \xi|^2 |E_{j_2}(\mathbf{k}) - \xi| \cdots |E_{j_4}(\mathbf{k}) - \xi|} \leq C (E_{j_1}(\mathbf{k}) + \xi_0)^{-2}.$$

On the other hand, one has the following rough estimate. There exists another  $C = C_{\xi_0} > 0$  s.t.:

$$|\hat{\pi}_{l, m}(\alpha; \mathbf{k})| \leq C (E_l(\mathbf{k}) + \xi_0) \quad l, m = 1, \dots, \text{Card}(\Theta), \alpha = 1, 2.$$

This follows from [14, Eq. (4.6)] together with the fact that  $(\partial_{k_\alpha} H)(\mathbf{k})$  is bounded from  $l^2(\Theta)$  to  $l^\infty(\Theta)$ . To see that, use that  $\|(\partial_{k_\alpha} H)(\mathbf{k})\|_{2, \infty}^2 = \text{ess sup}_{\mathbf{x} \in \Theta} \sum_{\mathbf{y} \in \Theta} |(\partial_{k_\alpha} H)(\mathbf{x}, \mathbf{y}; \mathbf{k})|^2$  combined with (1.10) and the estimate (1.2). It remains to involve the continuity of the  $E_j(\cdot)$ 's to get (3.43). To conclude the proof for the quantity  $\mathcal{W}_5(\beta, \mu^{(0)}, 0)$ , it is sufficient to use that it is made of a linear combination of  $w_5(\beta, \mu^{(0)}, 0)$ -like terms. The quantities  $\mathcal{W}_l(\beta, \mu^{(0)})$ , with  $l = 3, 4$  can be treated by following a similar method.  $\blacksquare$

**Proof of Lemma 3.4.** Denote the integrand appearing in (3.13) by:

$$\mathfrak{g}_{j_1, j_2, j_3, j_4}(\beta, \mu^{(0)}; \xi) := \frac{f(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi) (E_{j_3}(\mathbf{k}) - \xi) (E_{j_4}(\mathbf{k}) - \xi)} \quad j_1, j_2, j_3, j_4 \in \mathbb{N}^*.$$

At first sight,  $\mathfrak{g}_{j_1, j_2, j_3, j_4}(\beta, \mu^{(0)}; \cdot)$  can have poles from the first order up to at most fifth order (in the case when  $j_1 = j_2 = j_3 = j_4$ ). Therefore we expect the integral w.r.t.  $\xi$  of  $\mathfrak{g}_{j_1, j_2, j_3, j_4}(\beta, \mu^{(0)}; \cdot)$  to make appear partial derivatives of  $f(\beta, \mu^{(0)}; \cdot)$  with order at most equal to four. However the factor multiplying  $(\partial_\xi^4 f)(\beta, \mu^{(0)}; \cdot)$  is identically zero. Indeed, in view of the function  $\mathcal{C}_{j_1, j_2, j_3, j_4}(\cdot)$  defined in (3.16), it is identically zero for the combinations of subscripts:

$$j_1 = j_2 = j_3 = j_4; \quad j_1 = j_2 = j_3 \neq j_4; \quad j_1 = j_3 = j_4 \neq j_2. \quad (3.44)$$

Ergo the expansion of (3.13) consists of partial derivatives of  $f(\beta, \mu^{(0)}; \cdot)$  of order at most equal to three. Then by virtue of (3.44), the quadruple summation in the r.h.s. of (3.13) is reduced to:

$$\begin{aligned} \mathcal{W}_5(\beta, \mu^{(0)}, 0) = & - \sum_{j_1=1}^{\text{Card}(\Theta)} \sum_{\substack{j_3=1 \\ j_3 \neq j_1}}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \mathcal{C}_{j_1, j_1, j_3, j_1}(\mathbf{k}) \frac{1}{2i\pi} \int_{\Gamma} d\xi \frac{f(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^4 (E_{j_3}(\mathbf{k}) - \xi)} + \\ & - \underbrace{\sum_{\substack{j_1, \dots, j_4=1 \\ \text{at most 3 equal} \\ \text{subscripts}}}^{\text{Card}(\Theta)}} \int_{\Omega^*} d\mathbf{k} \mathcal{C}_{j_1, j_2, j_3, j_4}(\mathbf{k}) \frac{1}{2i\pi} \int_{\Gamma} d\xi \frac{f(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi) (E_{j_3}(\mathbf{k}) - \xi) (E_{j_4}(\mathbf{k}) - \xi)}. \end{aligned}$$

By applying the residue theorem in the first term of the above r.h.s., one has:

$$\begin{aligned} & \sum_{\substack{j_3=1 \\ j_3 \neq j_1}}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \mathcal{C}_{j_1, j_1, j_3, j_1}(\mathbf{k}) \frac{1}{2i\pi} \int_{\Gamma} d\xi \frac{f(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^4 (E_{j_3}(\mathbf{k}) - \xi)} = \sum_{\substack{j_3=1 \\ j_3 \neq j_1}}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \mathcal{C}_{j_1, j_1, j_3, j_1}(\mathbf{k}) \times \\ & \times \left\{ \frac{1}{3!} \frac{1}{E_{j_3}(\mathbf{k}) - E_{j_1}(\mathbf{k})} \frac{\partial^3 f}{\partial \xi^3}(\beta, \mu^{(0)}; E_{j_1}(\mathbf{k})) + \text{others terms involving } \frac{\partial^l f}{\partial \xi^l}(\beta, \mu^{(0)}; \cdot) \text{ with } l \leq 2 \right\}. \end{aligned}$$

The function  $\mathcal{C}_{j_1, j_1, j_3, j_1}(\cdot)$  appearing in front of  $(\partial_\xi^3 \mathfrak{f})(\beta, \mu^{(0)}; E_{j_1}(\mathbf{k}))$  corresponds to  $3! \mathfrak{a}_{j_1, 3}(\cdot)$ . The remainder of the proof only is plain computation using the residue theorem.  $\blacksquare$

**Proof of Lemma 3.5.** Denote the integrand appearing in (3.14) by:

$$\mathfrak{h}_{j_1, j_2, j_3}(\beta, \mu^{(0)}; \xi) := \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi) (E_{j_3}(\mathbf{k}) - \xi)} \quad j_1, j_2, j_3 \in \mathbb{N}^*.$$

Let us note that  $\mathfrak{h}_{j_1, j_2, j_3}(\beta, \mu^{(0)}; \cdot)$  can have poles from the first order up to at most fourth order (in the case when  $j_1 = j_2 = j_3$ ). Therefore we expect the integral w.r.t.  $\xi$  of  $\mathfrak{h}_{j_1, j_2, j_3}(\beta, \mu^{(0)}; \cdot)$  to make appear partial derivatives of  $\mathfrak{f}(\beta, \mu^{(0)}; \cdot)$  with order at most equal to three.

The triple summation in the r.h.s. of (3.14) can be rewritten:

$$\begin{aligned} \mathcal{W}_4(\beta, \mu^{(0)}, 0) = & - \sum_{j_1=1}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \mathcal{C}_{j_1, j_1, j_1}(\mathbf{k}) \frac{1}{2i\pi} \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^4} + \\ & - \sum_{\substack{j_1, j_2, j_3=1 \\ \text{at most 2 equal} \\ \text{subscripts}}}^{\text{Card}(\Theta)} \int_{\Omega^*} d\mathbf{k} \mathcal{C}_{j_1, j_2, j_3}(\mathbf{k}) \frac{1}{2i\pi} \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi) (E_{j_3}(\mathbf{k}) - \xi)}. \end{aligned}$$

The application of the residue theorem in the first term of the above r.h.s. yields:

$$\frac{1}{2i\pi} \int_{\Gamma} d\xi \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^4} = \frac{1}{3!} \frac{\partial^3 \mathfrak{f}}{\partial \xi^3}(\beta, \mu^{(0)}; E_{j_1}(\mathbf{k})).$$

This is only the one term which provides a third-order partial derivative of  $\mathfrak{f}(\beta, \mu^{(0)}; \cdot)$ .  $\blacksquare$

**Proof of Lemma 3.6.** Denote the integrand appearing in (3.15) by:

$$\mathfrak{i}_{j_1, j_2}(\beta, \mu^{(0)}; \xi) := \frac{\mathfrak{f}(\beta, \mu^{(0)}; \xi)}{(E_{j_1}(\mathbf{k}) - \xi)^2 (E_{j_2}(\mathbf{k}) - \xi)} \quad j_1, j_2 \in \mathbb{N}^*.$$

Let us note that  $\mathfrak{i}_{j_1, j_2, j_3}(\beta, \mu^{(0)}; \cdot)$  has poles from the first order up to at most third order (in the case when  $j_1 = j_2$ ). Therefore the integral w.r.t.  $\xi$  of  $\mathfrak{i}_{j_1, j_2}(\beta, \mu^{(0)}; \cdot)$  will make appear partial derivatives of  $\mathfrak{f}(\beta, \mu^{(0)}; \cdot)$  with order at most equal to two. We do not give further details.  $\blacksquare$

**Proof of Proposition 3.9.** Consider the quantity  $\hat{\mathfrak{d}}_{j,3}(\mathbf{k})$ ,  $j = 1, 2$  defined in (3.26). Let  $m = 1, 2$  with  $m \neq j$ . From (3.16) and (3.17) together with the identities (3.35)-(3.37) we have:

$$\begin{aligned} \frac{\mathcal{C}_{j,j,m,j}(\mathbf{k})}{E_m(\mathbf{k}) - E_j(\mathbf{k})} = & \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \right)^2 \frac{1}{2} \left[ \hat{\sigma}_{j,j}(2, 2; \mathbf{k}) - \frac{\partial^2 E_j(\mathbf{k})}{\partial k_2^2} \right] + \\ & + \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \right)^2 \frac{1}{2} \left[ \hat{\sigma}_{j,j}(1, 1; \mathbf{k}) - \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1^2} \right] - \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \right) \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \right) \left[ \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) - \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1 \partial k_2} \right], \\ \mathcal{C}_{j,j,j}(\mathbf{k}) = & \hat{\sigma}_{j,j}(1, 1; \mathbf{k}) \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \right)^2 + \hat{\sigma}_{j,j}(2, 2; \mathbf{k}) \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \right)^2 - 2 \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \right) \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \right). \end{aligned}$$

Thus the quantity  $\hat{\mathfrak{d}}_{j,3}(\mathbf{k})$  can be rewritten  $\hat{\mathfrak{d}}_{j,3}(\mathbf{k}) = \mathfrak{m}_{j,3}(\mathbf{k}) + \mathfrak{n}_{j,3}(\mathbf{k})$ ,  $\mathbf{k} \in \Omega^*$  where respectively:

$$\begin{aligned} \mathfrak{m}_{j,3}(\mathbf{k}) := & \frac{-1}{12} \left\{ \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \right)^2 \frac{\partial^2 E_j(\mathbf{k})}{\partial k_2^2} + \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \right)^2 \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1^2} - 2 \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \right) \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \right) \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1 \partial k_2} \right\}, \\ \mathfrak{n}_{j,3}(\mathbf{k}) := & \frac{1}{4} \left\{ \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \right)^2 \hat{\sigma}_{j,j}(2, 2; \mathbf{k}) + \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \right)^2 \hat{\sigma}_{j,j}(1, 1; \mathbf{k}) - 2 \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \right) \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \right) \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) \right\}. \end{aligned}$$

Hereafter we use the shorthand notations  $f(\cdot) = f(\beta, \mu^{(0)}; \cdot)$ . Let us now prove the following:

$$\int_{\Omega^*} d\mathbf{k} \frac{\partial^3 f}{\partial \xi^3}(E_j(\mathbf{k})) \mathbf{m}_{j,3}(\mathbf{k}) = \frac{1}{6} \int_{\Omega^*} d\mathbf{k} \frac{\partial^2 f}{\partial \xi^2}(E_j(\mathbf{k})) \left\{ \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1^2} \frac{\partial^2 E_j(\mathbf{k})}{\partial k_2^2} - \left( \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1 \partial k_2} \right)^2 \right\}. \quad (3.45)$$

From [5, Eq. (4.15)] and after rearranging of terms, one has:

$$\begin{aligned} \int_{\Omega^*} d\mathbf{k} \frac{\partial^3 f}{\partial \xi^3}(E_j(\mathbf{k})) \mathbf{m}_{j,3}(\mathbf{k}) &= \frac{1}{6} \int_{\Omega^*} d\mathbf{k} \frac{\partial^3 f}{\partial \xi^3}(E_j(\mathbf{k})) \left\{ \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \right) \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \right) \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1 \partial k_2} \right\} + \\ &+ \frac{1}{12} \int_{\Omega^*} d\mathbf{k} \frac{\partial^2 f}{\partial \xi^2}(E_j(\mathbf{k})) \left\{ 2 \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1^2} \frac{\partial^2 E_j(\mathbf{k})}{\partial k_2^2} + \frac{\partial E_j(\mathbf{k})}{\partial k_1} \frac{\partial}{\partial k_2} \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1 \partial k_2} + \frac{\partial E_j(\mathbf{k})}{\partial k_2} \frac{\partial}{\partial k_1} \frac{\partial^2 E_j(\mathbf{k})}{\partial k_2 \partial k_1} \right\}. \end{aligned}$$

The above identity corresponds to [5, Eq. (4.16)] added with [5, Eq. (4.13)] by setting  $N = j$ . By following the same method as the one concluding the proof of [5, Prop. 4.2], one obtains (3.45). Next, let us consider (3.27). In view of (3.40) what we still have to do is prove:

$$\begin{aligned} \int_{\Omega^*} d\mathbf{k} \frac{\partial^3 f}{\partial \xi^3}(E_j(\mathbf{k})) \mathbf{n}_{j,3}(\mathbf{k}) - \frac{1}{2} \int_{\Omega^*} d\mathbf{k} \frac{\partial^2 f}{\partial \xi^2}(E_j(\mathbf{k})) \left\{ \frac{\mathcal{C}_{j,m,j}(\mathbf{k}) + \mathcal{C}_{j,j,m}(\mathbf{k})}{E_m(\mathbf{k}) - E_j(\mathbf{k})} + \mathcal{C}_{j,j}(\mathbf{k}) \right\} = \\ \frac{i}{2} \int_{\Omega^*} d\mathbf{k} \frac{\partial^2 f}{\partial \xi^2}(E_j(\mathbf{k})) \frac{\Im\{\mathcal{C}_{j,m,j}(\mathbf{k})\}}{E_j(\mathbf{k}) - E_m(\mathbf{k})}. \quad (3.46) \end{aligned}$$

On the one hand, from the definition (3.17) together with the identities (3.35)-(3.37):

$$\begin{aligned} \frac{\mathcal{C}_{j,j,m}(\mathbf{k})}{E_m(\mathbf{k}) - E_j(\mathbf{k})} &= \hat{\sigma}_{j,j}(1, 1; \mathbf{k}) \left[ \frac{1}{2} \hat{\sigma}_{j,j}(2, 2; \mathbf{k}) - \frac{1}{2} \frac{\partial^2 E_j(\mathbf{k})}{\partial k_2^2} \right] + \\ &+ \hat{\sigma}_{j,j}(2, 2; \mathbf{k}) \left[ \frac{1}{2} \hat{\sigma}_{j,j}(1, 1; \mathbf{k}) - \frac{1}{2} \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1^2} \right] - \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) \left[ \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) - \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1 \partial k_2} \right], \end{aligned}$$

Then from the definition (3.18) we get after rearranging of terms:

$$\begin{aligned} -\frac{1}{2} \int_{\Omega^*} d\mathbf{k} \frac{\partial^2 f}{\partial \xi^2}(E_j(\mathbf{k})) \left\{ \frac{\mathcal{C}_{j,m,j}(\mathbf{k}) + \mathcal{C}_{j,j,m}(\mathbf{k})}{E_m(\mathbf{k}) - E_j(\mathbf{k})} + \mathcal{C}_{j,j}(\mathbf{k}) \right\} &= \frac{1}{4} \int_{\Omega^*} d\mathbf{k} \frac{\partial^2 f}{\partial \xi^2}(E_j(\mathbf{k})) \times \\ \times \left\{ \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1^2} \hat{\sigma}_{j,j}(2, 2; \mathbf{k}) + \frac{\partial^2 E_j(\mathbf{k})}{\partial k_2^2} \hat{\sigma}_{j,j}(1, 1; \mathbf{k}) - 2 \frac{\partial^2 E_j(\mathbf{k})}{\partial k_1 \partial k_2} \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) - 2 \frac{\mathcal{C}_{j,m,j}(\mathbf{k})}{E_m(\mathbf{k}) - E_j(\mathbf{k})} \right\}. \quad (3.47) \end{aligned}$$

On the other hand, the definition of  $\mathbf{n}_{j,3}(\mathbf{k})$  together with the use of [5, Eq. (4.15)] lead to:

$$\begin{aligned} \int_{\Omega^*} d\mathbf{k} \frac{\partial^3 f}{\partial \xi^3}(E_j(\mathbf{k})) \mathbf{n}_{j,3}(\mathbf{k}) &= -\frac{1}{4} \int_{\Omega^*} d\mathbf{k} \frac{\partial^2 f}{\partial \xi^2}(E_j(\mathbf{k})) \left\{ \frac{\partial}{\partial k_1} \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \hat{\sigma}_{j,j}(2, 2; \mathbf{k}) \right) + \right. \\ &+ \left. \frac{\partial}{\partial k_2} \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \hat{\sigma}_{j,j}(1, 1; \mathbf{k}) \right) - \frac{\partial}{\partial k_1} \left( \frac{\partial E_j(\mathbf{k})}{\partial k_2} \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) \right) - \frac{\partial}{\partial k_2} \left( \frac{\partial E_j(\mathbf{k})}{\partial k_1} \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) \right) \right\}. \quad (3.48) \end{aligned}$$

By adding (3.47) with (3.48) and using the definition of  $\mathcal{C}_{j,m,j}(\mathbf{k})$ , the l.h.s. of (3.46) is equal to:

$$\begin{aligned} \frac{1}{2} \int_{\Omega^*} d\mathbf{k} \frac{\partial^2 f}{\partial \xi^2}(E_j(\mathbf{k})) \left\{ \frac{\partial E_j(\mathbf{k})}{\partial k_1} \left( \frac{\hat{\sigma}_{j,m}(2, 2; \mathbf{k}) \hat{\pi}_{m,j}(1; \mathbf{k})}{E_j(\mathbf{k}) - E_m(\mathbf{k})} - \frac{1}{2} \frac{\partial}{\partial k_1} \hat{\sigma}_{j,j}(2, 2; \mathbf{k}) + \right. \right. \\ \left. \left. - \frac{\hat{\sigma}_{j,m}(1, 2; \mathbf{k}) \hat{\pi}_{m,j}(2; \mathbf{k})}{E_j(\mathbf{k}) - E_m(\mathbf{k})} + \frac{1}{2} \frac{\partial}{\partial k_2} \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) \right) + \frac{\partial E_j(\mathbf{k})}{\partial k_2} \left( \frac{\hat{\sigma}_{j,m}(1, 1; \mathbf{k}) \hat{\pi}_{m,j}(2; \mathbf{k})}{E_j(\mathbf{k}) - E_m(\mathbf{k})} + \right. \right. \\ \left. \left. - \frac{1}{2} \frac{\partial}{\partial k_2} \hat{\sigma}_{j,j}(1, 1; \mathbf{k}) - \frac{\hat{\sigma}_{j,m}(1, 2; \mathbf{k}) \hat{\pi}_{m,j}(1; \mathbf{k})}{E_j(\mathbf{k}) - E_m(\mathbf{k})} + \frac{1}{2} \frac{\partial}{\partial k_1} \hat{\sigma}_{j,j}(1, 2; \mathbf{k}) \right) \right\}. \end{aligned}$$

To get (3.46), it remains to use this identity (obtained via the regular perturbation theory):

$$\frac{\partial}{\partial k_\delta} \hat{\sigma}_{j,j}(\alpha, \gamma; \mathbf{k}) = \hat{\tau}_{j,j}(\delta, \alpha, \gamma; \mathbf{k}) + \frac{2\Re\{\hat{\sigma}_{j,m}(\alpha, \gamma; \mathbf{k})\hat{\pi}_{m,j}(\delta; \mathbf{k})\}}{E_j(\mathbf{k}) - E_m(\mathbf{k})} \quad \mathbf{k} \in \Omega^*, \alpha, \gamma, \delta = 1, 2,$$

with  $\hat{\tau}_{j,j}(\alpha, \alpha, \gamma; \mathbf{k}) := \langle u_j(\cdot; \mathbf{k}), (\partial_{k_\alpha} \partial_{k_\alpha} \partial_{k_\gamma} H)(\mathbf{k}) u_j(\cdot; \mathbf{k}) \rangle = \hat{\tau}_{j,j}(\alpha, \gamma, \alpha; \mathbf{k}) = \hat{\tau}_{j,j}(\gamma, \alpha, \alpha; \mathbf{k})$ . ■

## 4 Acknowledgments.

B.S. warmly thanks Horia D. Cornean for many fruitful discussions. A part of this work was done while the author was visiting the Department of Mathematical Sciences of Aalborg University. B.S. is thankful for the hospitality and financial support from the Department. As well, B.S. is indebted to Mathieu Beau for making many references available to the author.

## References

- [1] Angelescu N., Bundaru M., Nenciu G., *On the Landau diamagnetism*, Commun. Math. Phys. **42**(1), 9–28 (1975)
- [2] Adams E.N., *Magnetic susceptibility of a diamagnetic electron gas - The role of small effective electron mass*, Phys. Rev. **89**(3), 633–647 (1953)
- [3] Briet P., Cornean H.D., Louis D., *Diamagnetic expansions for perfect quantum gases*, J. Math. Phys. **47**(8) 083511, (2006)
- [4] Briet P., Cornean H.D., Savoie B., *Diamagnetism of quantum gases with singular potentials*, J. Phys. A: Math. Theor. **43** (2010), 474008
- [5] Briet P., Cornean H.D., Savoie B., *A rigorous proof of the Landau-Peierls formula and much more*, Ann. Henri Poincaré **13**, 1–40 (2012)
- [6] Briet P., Cornean H.D., Zagrebnov V., *Do bosons condense in a homogeneous magnetic field ?*, J. Stat. Phys. **116**(5-6), 1545–1578 (2004)
- [7] Briet P., Savoie B., *A rigorous approach to the magnetic response in disordered systems*, preprint arXiv:1112.6177v1 (2011)
- [8] Brynildsen M.H., Cornean H.D., *On the Verdet constant and Faraday rotation for graphene-like materials*, preprint arXiv:1112.2613v1 (2011)
- [9] Castro Neto A.H., Guinea F., Peres N.M.R., Novoselov K.S., Geim A.K., *The electronic properties of graphene*, Rev. Mod. Phys. **81**(1), 109–162 (2009)
- [10] Cornean H.D., *On the magnetization of a charged Bose gas in the canonical ensemble*, Commun. Math. Phys. **212** (2000), 1–27
- [11] Cornean H.D., *On the Lipschitz continuity of spectral bands of Harper-like and magnetic Schrödinger operators*, Ann. Henri Poincaré **11**, 973–990 (2010)
- [12] Cornean H.D., Nenciu G., Pedersen, T.G., *The Faraday effect revisited: General theory*, J. Math. Phys. **47**, 013511 (2006)
- [13] Cornean H.D., Nenciu G., *The Faraday effect revisited: Thermodynamic limit*, J. Funct. Anal. **257**(7), 2024–2066 (2009)
- [14] Cornean H.D., Nenciu G., *Faraday effect revisited: sum rules and convergence issues*, J. Phys. A: Math. Theor. **43**, 474012 (2010)
- [15] Dunford N., Schwartz J.T., *Linear Operators Part 1: General Theory*, Interscience Publishers, New York, 1958.
- [16] Fukuyama H., *Theory of orbital magnetism of Bloch electrons: Coulomb interactions*, Prog. Theor. Phys. **45**(3), 704–729 (1971)
- [17] Helffer B., Sjöstrand J., *On diamagnetism and the de Haas-Van Alphen effect*, Ann. Inst. H. Poincaré Phys. Théor. **52**(4), 303–375 (1990)

- [18] Hislop P.D., *Lectures on random Schrödinger operators*, Contemporary Mathematics, **476**, 41–131 (2008)
- [19] Hove J.E., *Theory of the magnetic susceptibility of graphite*, Phys. Rev. **100**(2), 645–649 (1955)
- [20] Koshino M., Ando T., *Diamagnetism in disordered graphene*, Phys. Rev. B **75**(23), 235333 (2007)
- [21] Koshino M., Ando T., *Orbital diamagnetism in multilayer graphenes: Systematic study with the effective mass approximation*, Phys. Rev. B **76**(8), 085425 (2007)
- [22] Kuchment P., *Floquet Theory for Partial Differential Equations*, Birkhäuser Verlag, Basel, 1993
- [23] Levintovich Ya., Kotosonov A.S., *Diamagnetism of narrow-gap semiconductors*, Sov. Phys. J. **30**(2), 153–156 (1987)
- [24] McClure J.W., *Diamagnetism of graphite*, Phys. Rev. **104**(3), 666–671 (1956)
- [25] McClure J.W., *Theory of diamagnetism of graphite*, Phys. Rev. **119**(2), 606–613 (1960)
- [26] Misra P.K., Kleinman L., *Theory of the diamagnetic susceptibility of Bloch electrons*, Phys. Rev. B **5**(11), 4581–4597 (1972)
- [27] Nakamura M., *Orbital magnetism and transport phenomena in two-dimensional Dirac fermions in a weak magnetic field*, Phys. Rev. B **76**, 113301 (2007). *Erratum*, Phys. Rev. B **77**, 129903(E) (2008)
- [28] Nenciu G., *On asymptotic perturbation theory for quantum mechanics: Almost invariant subspaces and gauge invariant magnetic perturbation theory*, J. Math. Phys. **43**(3), 1273–1298 (2002)
- [29] Nenciu G., *Dynamics of band electrons in electric and magnetic fields: rigorous justification of the effective Hamiltonians*, Rev. Mod. Phys. **63**(1), 91–127 (1991)
- [30] Novoselov K.S., Geim A.K., Morozov S.V., Jiang D., Zhang Y., Dubonos S.V., Gregorieva I.V., Firsov A.A., *Electric field effect in atomically thin carbon films*, Science **306**, 666–669 (2004)
- [31] Pedersen J.G., Pedersen T.G., *Tight-binding study of the magneto-optical properties of gapped Graphene*, Phys. Rev. B **84**, 115424 (2011)
- [32] Pedersen J.G., Pedersen T.G., *Band gaps in graphene via periodic electrostatic gating*, preprint arXiv:1203.3629v1
- [33] Peierls R., *Zur Theorie des Diamagnetismus von Leitungselektronen. Zeitschrift für Physik A. Hadrons and Nuclei*, **80**, 2–19 (1933)
- [34] Robinson D.W., *The Thermodynamic Pressure in Quantum Statistical Mechanics*, Lecture Notes in Physics 9, Springer-Verlag, Berlin-Heidelberg, 1971
- [35] Reed M., Simon B., *Methods of Modern Mathematical Physics, IV : Analysis of Operators*, Academic Press, San Diego, 1978
- [36] Safran S.A., DiSalvo F.J., *Theory of magnetic susceptibility of graphite intercalation compounds*, Phys. Rev. B **20**(12), 4889–4895 (1979)
- [37] Safran S.A., *Stage dependence of magnetic susceptibility of intercalated graphite*, Phys. Rev. B **30**(1), 421–423 (1984)
- [38] Sharma M.P., Johnson L.G., McClure J.W., *Diamagnetism of graphite*, Phys. Rev. B **9**(6), 2467–2475 (1974)
- [39] Smoluchowski R., *Magnetic susceptibility of bromine-graphite*, Rev. Mod. Phys. **25**(1), 178–180 (1953)
- [40] Wallace P.R., *The band theory of graphite*, Phys. Rev. **71**(9), 622–634 (1947). *Erratum: The band theory of graphite*, Phys. Rev. **72**(3), 258–258 (1947)
- [41] Wannier G.H., Upadhyaya U.N., *Zero-field susceptibility of Bloch electrons*, Phys. Rev. **136**(3A), 803–810 (1964)