



HAL
open science

Deformation of the Hopf algebra of plane posets

Loïc Foissy

► **To cite this version:**

Loïc Foissy. Deformation of the Hopf algebra of plane posets. European Journal of Combinatorics, 2014. hal-00687491v2

HAL Id: hal-00687491

<https://hal.science/hal-00687491v2>

Submitted on 23 Nov 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Deformation of the Hopf algebra of plane posets

Loïc Foissy

*Laboratoire de Mathématiques, Université de Reims
Moulin de la Housse - BP 1039 - 51687 REIMS Cedex 2, France
e-mail : loic.foissy@univ-reims.fr*

ABSTRACT. We describe and study a four parameters deformation of the two products and the coproduct of the Hopf algebra of plane posets. We obtain a family of braided Hopf algebras, which are generically self-dual. We also prove that in a particular case (when the second parameter goes to zero and the first and third parameters are equal), this deformation is isomorphic, as a self-dual braided Hopf algebra, to a deformation of the Hopf algebra of free quasi-symmetric functions **FQSym**.

KEYWORDS. Plane posets; Deformation; Braided Hopf algebras; Self- duality.

AMS CLASSIFICATION. 06A11, 16W30, 16S80.

Contents

1	Recalls and notations	3
1.1	Double and plane posets	3
1.2	Algebraic structures on plane posets	4
1.3	Pairings	6
1.4	Morphism to free quasi-symmetric functions	6
2	Deformation of the products	7
2.1	Construction	7
2.2	Particular cases	9
2.3	Subalgebras and quotients	11
3	Dual coproducts	12
3.1	Constructions	12
3.2	Particular cases	13
3.3	Compatibilities with the products	15
4	Self-duality results	16
4.1	A first pairing on \mathcal{H}_q	16
4.2	Properties of the pairing	18
4.3	Comparison of pairings with colinear parameters	20
4.4	Non-degeneracy of the pairing	21
5	Morphism to free quasi-symmetric functions	23
5.1	A second Hopf pairing on $\mathcal{H}_{(q_1,0,q_3,q_4)}$	23
5.2	Quantization of the Hopf algebra of free quasi-symmetric functions	26

Introduction

A double poset is a finite set with two partial orders. As explained in [8], the space generated by the double posets inherits two products and one coproduct, here denoted by m , ζ and Δ , making it both a Hopf and an infinitesimal Hopf algebra [6]. Moreover, this Hopf algebra is self dual. A double poset is plane if its two partial orders satisfy a (in)compatibility condition, see definition 1. The subspace $\mathcal{H}_{\mathcal{PP}}$ generated by plane posets is stable under the two products and the coproduct, and is self-dual [2, 3]: in particular, two Hopf pairings are defined on it, using the notion of picture. Moreover, as proved in [3], it is isomorphic to the Hopf algebra of free quasi-symmetric functions **FQSym**, also known as the Malvenuto-Reutenauer Hopf algebra of permutations. An explicit isomorphism Θ is given by the linear extensions of plane posets, see definition 11.

We define in this text a four-parameters deformation of the products and the coproduct of $\mathcal{H}_{\mathcal{PP}}$, together with a deformation of the two pairings and of the morphism from $\mathcal{H}_{\mathcal{PP}}$ to **FQSym**. If $q = (q_1, q_2, q_3, q_4) \in K^4$, the product $m_q(P \otimes Q)$ of two plane posets P and Q is a linear span of plane posets R such that $R = P \sqcup Q$ as a set, P and Q being plane subposets of R . The coefficients are defined with the help of the two partial orders of R , see theorem 14, and are polynomials in q . In particular:

$$\begin{cases} m_{(1,0,0,0)} &= \zeta, \\ m_{(0,1,0,0)} &= \zeta^{op}, \\ m_{(0,0,1,0)} &= m, \\ m_{(0,0,0,1)} &= m^{op}. \end{cases}$$

We also obtain the product dual to the coproduct Δ (considering the basis of double posets as orthonormal) as $m_{(1,0,1,1)}$, and its opposite given by $m_{(0,1,1,1)}$. Dually, we define a family of coassociative coproducts Δ_q . For any plane poset P , $\Delta(P)$ is a linear span of terms $(P \setminus I) \otimes I$, running over the plane subposets I of P , the coefficients being polynomials in q . In the particular cases where at least one of the components of q is zero, only h -ideals, r -ideals or biideals can appear in this sum (definition 7 and proposition 22). We study the compatibility of Δ_q with both products m and ζ on $\mathcal{H}_{\mathcal{PP}}$ (proposition 23). For example, $(\mathcal{H}_{\mathcal{PP}}, m, \Delta_q)$ satisfies the axiom

$$\Delta_q(xy) = \sum \sum q_3^{|x'_q||y''_q|} q_4^{|x''_q||y'_q|} (x'_q y'_q) \otimes (x''_q y''_q).$$

If $q_3 = 1$, it is a braided Hopf algebra, with braiding given by $c_q(P \otimes Q) = q_4^{|P||Q|} Q \otimes P$; in particular, if $q_3 = 1$ and $q_4 = 1$ this is a Hopf algebra, and if $q_3 = 1$ and $q_4 = 0$ it is an infinitesimal Hopf algebra. If $q_4 = 1$, it is the coopposite (or the opposite) of a braided Hopf algebra. Similar results hold if we consider the second product ζ , permuting the roles of (q_3, q_4) and (q_1, q_2) .

We define a symmetric pairing $\langle -, - \rangle_q$ such that:

$$\langle x \otimes y, \Delta_q(z) \rangle_q = \langle xy, z \rangle_q \text{ for all } x, y, z \in \mathcal{H}_{\mathcal{PP}}.$$

If $q = (1, 0, 1, 1)$, we recover the first "classical" pairing of $\mathcal{H}_{\mathcal{PP}}$. We prove that in the case $q_2 = 0$, this pairing is nondegenerate if, and only if, $q_1 \neq 0$ (corollary 36). Consequently, this pairing is generically nondegenerate.

The coproduct of **FQSym** is finally deformed, in such a way that the algebra morphism Θ from $\mathcal{H}_{\mathcal{PP}}$ to **FQSym** becomes compatible with Δ_q , if q has the form $q = (q_1, 0, q_1, q_4)$. Deforming the second pairing $\langle -, - \rangle'$ of $\mathcal{H}_{\mathcal{PP}}$ and the usual Hopf pairing of **FQSym**, the map Θ becomes also an isometry (theorem 40). Consequently, the deformation $\langle -, - \rangle'_q$ is nondegenerate if, and only if, $q_1 q_4 \neq 0$.

- If $i, j \in P_\sigma$, $i \leq_h j$ if $i \leq j$ and $\sigma(i) \leq \sigma(j)$.
- If $i, j \in P_\sigma$, $i \leq_r j$ if $i \leq j$ and $\sigma(i) \geq \sigma(j)$.

Note that the total order on $\{1, \dots, n\}$ induced by this plane poset structure is the usual one. Then for all $n \geq 0$, the following map is a bijection:

$$\Psi_n : \begin{cases} \mathfrak{S}_n & \longrightarrow \mathcal{PP}(n) \\ \sigma & \longrightarrow P_\sigma. \end{cases}$$

Examples.

$$\begin{aligned} \Psi_2((12)) &= \mathbf{1}, & \Psi_2((21)) &= \mathbf{..}, & \Psi_3(123) &= \mathbf{\dot{1}}, & \Psi_3((132)) &= \mathbf{V}, \\ \Psi_3(213) &= \mathbf{\wedge}, & \Psi_3((231)) &= \mathbf{1.}, & \Psi_3(312) &= \mathbf{.1}, & \Psi_3((321)) &= \mathbf{...} \end{aligned}$$

We define several bijections on \mathcal{PP} :

Definition 4 Let $P = (P, \leq_h, \leq_r) \in \mathcal{PP}$. We put:

$$\begin{cases} \iota(P) &= (P, \leq_r, \leq_h), \\ \alpha(P) &= (P, \geq_h, \leq_r), \\ \beta(P) &= (P, \leq_h, \geq_r), \\ \gamma(P) &= (P, \geq_h, \geq_r). \end{cases}$$

Remarks.

1. Graphically:

- A Hasse graph of $\alpha(P)$ is obtained from a Hasse graph of P by a horizontal symmetry.
- A Hasse graph of $\beta(P)$ is obtained from a Hasse graph of P by a vertical symmetry.
- A Hasse graph of $\gamma(P)$ is obtained from a Hasse graph of P by a rotation of angle π .

2. These bijections generate a group of permutations of \mathcal{PP} of cardinality 8. It is described by the following array:

\circ	α	β	γ	ι	$\iota \circ \alpha$	$\iota \circ \beta$	$\iota \circ \gamma$
α	Id	γ	β	$\iota \circ \beta$	$\iota \circ \gamma$	ι	$\iota \circ \alpha$
β	γ	Id	α	$\iota \circ \alpha$	ι	$\iota \circ \gamma$	$\iota \circ \beta$
γ	β	α	Id	$\iota \circ \gamma$	$\iota \circ \beta$	$\iota \circ \alpha$	ι
ι	$\iota \circ \alpha$	$\iota \circ \beta$	$\iota \circ \gamma$	Id	α	β	γ
$\iota \circ \alpha$	ι	$\iota \circ \gamma$	$\iota \circ \beta$	β	γ	Id	α
$\iota \circ \beta$	$\iota \circ \gamma$	ι	$\iota \circ \alpha$	α	Id	γ	β
$\iota \circ \gamma$	$\iota \circ \beta$	$\iota \circ \alpha$	ι	γ	β	α	Id

This is a dihedral group D_4 .

1.2 Algebraic structures on plane posets

Two products are defined on \mathcal{PP} . The first is called *composition* in [8] and denoted by \rightsquigarrow in [2]. We shall shortly denote it by m in this text.

Definition 5 Let $P, Q \in \mathcal{PP}$.

1. The double poset $PQ = m(P \otimes Q)$ is defined as follows:

- $PQ = P \sqcup Q$ as a set, and P, Q are plane subposets of PQ .

- if $x \in P$ and $y \in Q$, then $x \leq_r y$ in PQ .
2. The double poset $P \dot{\sqcup} Q$ is defined as follows:
- $P \dot{\sqcup} Q = P \sqcup Q$ as a set, and P, Q are plane subposets of $P \dot{\sqcup} Q$.
 - if $x \in P$ and $y \in Q$, then $x \leq_h y$ in $P \dot{\sqcup} Q$.

Examples.

1. The Hasse graph of PQ is the concatenation of the Hasse graphs of P and Q .
2. Here are examples for $\dot{\sqcup}$: $\cdot \dot{\sqcup} \mathbf{1} = \mathbf{\dot{1}}$, $\mathbf{1} \dot{\sqcup} \cdot = \mathbf{\dot{1}}$, $\cdot \dot{\sqcup} \cdot = \mathbf{\vee}$, $\cdot \dot{\sqcup} \cdot = \mathbf{\wedge}$.

The vector space generated by \mathcal{PP} is denoted by $\mathcal{H}_{\mathcal{PP}}$. These two products are linearly extended to $\mathcal{H}_{\mathcal{PP}}$; then $(\mathcal{H}_{\mathcal{PP}}, m)$ and $(\mathcal{H}_{\mathcal{PP}}, \dot{\sqcup})$ are two associative, unitary algebras, sharing the same unit 1, which is the empty plane poset. Moreover, they are both graded by the cardinality of plane posets. They are free algebras, as implied that the following theorem, proved in [2]:

- Theorem 6** 1. (a) Let P be a nonempty plane poset. We shall say that P is h -irreducible if for all $Q, R \in \mathcal{PP}$, $P = QR$ implies that $Q = 1$ or $R = 1$.
- (b) Any plane poset P can be uniquely written as $P = P_1 \dots P_k$, where P_1, \dots, P_k are h -irreducible. We shall say that P_1, \dots, P_k are the h -irreducible components of P .
2. (a) Let P be a nonempty plane poset. We shall say that P is r -irreducible if for all $Q, R \in \mathcal{PP}$, $P = Q \dot{\sqcup} R$ implies that $Q = 1$ or $R = 1$.
- (b) Any plane poset P can be uniquely written as $P = P_1 \dot{\sqcup} \dots \dot{\sqcup} P_k$, where P_1, \dots, P_k are r -irreducible. We shall say that P_1, \dots, P_k are the r -irreducible components of P .

Remark. The Hasse graphs of the h -irreducible components of H are the connected components of the Hasse graph of (P, \leq_h) , whereas the Hasse graphs of the r -irreducible components of H are the connected components of the Hasse graph of (P, \leq_r) .

Definition 7 Let $P = (P, \leq_h, \leq_r)$ be a plane poset, and let $I \subseteq P$.

1. We shall say that I is a h -ideal of P , if, for all $x, y \in P$:

$$(x \in I, x \leq_h y) \implies (y \in I).$$

2. We shall say that I is a r -ideal of P , if, for all $x, y \in P$:

$$(x \in I, x \leq_r y) \implies (y \in I).$$

3. We shall say that I is a biideal of P if it both an h -ideal and a r -ideal. Equivalently, I is a biideal of P if, for all $x, y \in P$:

$$(x \in I, x \leq y) \implies (y \in I).$$

The following proposition is proved in [2] (proposition 29):

Proposition 8 $\mathcal{H}_{\mathcal{PP}}$ is given a coassociative, counitary coproduct in the following way: for any plane poset P ,

$$\Delta(P) = \sum_{I \text{ } h\text{-ideal of } P} (P \setminus I) \otimes I.$$

Moreover, $(\mathcal{H}_{\mathcal{PP}}, m, \Delta)$ is a Hopf algebra, and $(\mathcal{H}_{\mathcal{PP}}, \zeta, \Delta)$ is an infinitesimal Hopf algebra [6], both graded by the cardinality of the plane posets. In other words, using Sweedler's notations $\Delta(x) = \sum x^{(1)} \otimes x^{(2)}$, for all $x, y \in \mathcal{H}_{\mathcal{PP}}$:

$$\begin{cases} \Delta(xy) &= \sum x^{(1)}y^{(1)} \otimes x^{(2)}y^{(2)}, \\ \Delta(x\zeta y) &= \sum x\zeta y^{(1)} \otimes y^{(2)} + \sum x^{(1)} \otimes x^{(2)}\zeta y - x \otimes y. \end{cases}$$

Remarks. The following compatibilities are satisfied:

1. For all $P, Q \in \mathcal{PP}$:

$$\begin{cases} \iota(PQ) &= \iota(P)\zeta\iota(Q), & \iota(P\zeta Q) &= \iota(P)\iota(Q), \\ \alpha(PQ) &= \alpha(P)\alpha(Q), & \alpha(P\zeta Q) &= \alpha(Q)\zeta\alpha(P), \\ \beta(PQ) &= \beta(Q)\beta(P), & \beta(P\zeta Q) &= \beta(P)\zeta\beta(Q), \\ \gamma(PQ) &= \gamma(Q)\gamma(P), & \gamma(P\zeta Q) &= \gamma(Q)\zeta\gamma(P). \end{cases}$$

2. Moreover, $\Delta \circ \alpha = (\alpha \otimes \alpha) \circ \Delta^{op}$, $\Delta \circ \beta = (\beta \otimes \beta) \circ \Delta$, and $\Delta \circ \gamma = (\gamma \otimes \gamma) \circ \Delta^{op}$.

1.3 Pairings

We also defined two pairings on $\mathcal{H}_{\mathcal{PP}}$, using the notion of *pictures*:

Definition 9 Let P, Q be two elements of \mathcal{PP} .

1. We denote by $S(P, Q)$ the set of bijections $\sigma : P \rightarrow Q$ such that, for all $i, j \in P$:

- $(i \leq_h j \text{ in } P) \implies (\sigma(i) \leq_r \sigma(j) \text{ in } Q)$.
- $(\sigma(i) \leq_h \sigma(j) \text{ in } Q) \implies (i \leq_r j \text{ in } P)$.

2. We denote by $S'(P, Q)$ the set of bijections $\sigma : P \rightarrow Q$ such that, for all $i, j \in P$:

- $(i \leq_h j \text{ in } P) \implies (\sigma(i) \leq \sigma(j) \text{ in } Q)$.
- $(\sigma(i) \leq_h \sigma(j) \text{ in } Q) \implies (i \leq j \text{ in } P)$.

The following theorem is proved in [2, 3, 8]:

Theorem 10 We define two pairings:

$$\begin{aligned} \langle -, - \rangle &: \begin{cases} \mathcal{H}_{\mathcal{PP}} \otimes \mathcal{H}_{\mathcal{PP}} &\longrightarrow K \\ P \otimes Q &\longrightarrow \langle P, Q \rangle = \text{Card}(S(P, Q)), \end{cases} \\ \langle -, - \rangle' &: \begin{cases} \mathcal{H}_{\mathcal{PP}} \otimes \mathcal{H}_{\mathcal{PP}} &\longrightarrow K \\ P \otimes Q &\longrightarrow \langle P, Q \rangle' = \text{Card}(S'(P, Q)). \end{cases} \end{aligned}$$

They are both homogeneous, symmetric, nondegenerate Hopf pairings on the Hopf algebra $\mathcal{H}_{\mathcal{PP}} = (\mathcal{H}_{\mathcal{PP}}, m, \Delta)$.

1.4 Morphism to free quasi-symmetric functions

We here briefly recall the construction of the Hopf algebra **FQSym** of free quasi-symmetric functions, also called the Malvenuto-Reutenauer Hopf algebra [1, 7]. As a vector space, a basis of **FQSym** is given by the disjoint union of the symmetric groups \mathfrak{S}_n , for all $n \geq 0$. By convention, the unique element of \mathfrak{S}_0 is denoted by 1. The product of **FQSym** is given, for $\sigma \in \mathfrak{S}_k$, $\tau \in \mathfrak{S}_l$, by:

$$\sigma\tau = \sum_{\epsilon \in Sh(k, l)} (\sigma \otimes \tau) \circ \epsilon,$$

where $Sh(k, l)$ is the set of (k, l) -shuffles, that is to say permutations $\epsilon \in \mathfrak{S}_{k+l}$ such that $\epsilon^{-1}(1) < \dots < \epsilon^{-1}(k)$ and $\epsilon^{-1}(k+1) < \dots < \epsilon^{-1}(k+l)$. In other words, the product of σ and τ is given by shifting the letters of the word representing τ by k , and then summing all the possible shufflings of this word and of the word representing σ . For example:

$$(123)(21) = (12354) + (12534) + (15234) + (51234) + (12543) \\ + (15243) + (51243) + (15423) + (51423) + (54123).$$

Let $\sigma \in \Sigma_n$. For all $0 \leq k \leq n$, there exists a unique triple $(\sigma_1^{(k)}, \sigma_2^{(k)}, \zeta_k) \in \mathfrak{S}_k \times \mathfrak{S}_{n-k} \times Sh(k, n-k)$ such that $\sigma = \zeta_k^{-1} \circ (\sigma_1^{(k)} \otimes \sigma_2^{(k)})$. The coproduct of \mathbf{FQSym} is then defined by:

$$\Delta(\sigma) = \sum_{k=0}^n \sigma_1^{(k)} \otimes \sigma_2^{(k)}.$$

For example:

$$\Delta((43125)) = 1 \otimes (43125) + (1) \otimes (3124) + (21) \otimes (123) \\ + (321) \otimes (12) + (4312) \otimes (1) + (43125) \otimes 1.$$

Note that $\sigma_1^{(k)}$ and $\sigma_2^{(k)}$ are obtained by cutting the word representing σ between the k -th and the $(k+1)$ -th letter, and then *standardizing* the two obtained words, that is to say applying to their letters the unique increasing bijection to $\{1, \dots, k\}$ or $\{1, \dots, n-k\}$. Moreover, \mathbf{FQSym} has a nondegenerate, homogeneous, Hopf pairing defined by $\langle \sigma, \tau \rangle = \delta_{\sigma, \tau^{-1}}$ for all permutations σ and τ .

Definition 11 [4, 8]

1. Let $P = (P, \leq_h, \leq_r)$ a plane poset. Let $x_1 < \dots < x_n$ be the elements of P , which is totally ordered. A linear extension of P is a permutation $\sigma \in \mathfrak{S}_n$ such that, for all $i, j \in \{1, \dots, n\}$:

$$(x_i \leq_h x_j) \implies (\sigma^{-1}(i) < \sigma^{-1}(j)).$$

The set of linear extensions of P will be denoted by S_P .

2. The following map is an isomorphism of Hopf algebras:

$$\Theta : \begin{cases} \mathcal{H}_{\mathcal{PP}} & \longrightarrow \mathbf{FQSym} \\ P \in \mathcal{PP} & \longrightarrow \sum_{\sigma \in S_P} \sigma. \end{cases}$$

moreover, for all $x, y \in \mathcal{H}_{\mathcal{PP}}$, $\langle \Theta(x), \Theta(y) \rangle' = \langle x, y \rangle$.

2 Deformation of the products

2.1 Construction

Definition 12 Let $P \in \mathcal{PP}$ and $X, Y \subseteq P$. We put:

1. $h_X^Y = \#\{(x, y) \in X \times Y / x \leq_h y \text{ in } P\}$.
2. $r_X^Y = \#\{(x, y) \in X \times Y / x \leq_r y \text{ in } P\}$.

Lemma 13 Let X and Y be disjoint parts of a plane poset P . Then:

$$h_X^Y + h_Y^X + r_X^Y + r_Y^X = |X||Y|.$$

Proof. Indeed, $h_X^Y + h_Y^X + r_X^Y + r_Y^X = \#\{(x, y) \in X \times Y \mid x < y \text{ or } x > y\} = |X \times Y|$. \square

Remark. If it more generally possible to prove that for any part X and Y of a plane poset:

$$h_X^Y + h_Y^X + r_X^Y + r_Y^X = 3|X \cap Y|^2 + |X||Y|.$$

Theorem 14 Let $q = (q_1, q_2, q_3, q_4) \in K^4$. We consider the following map:

$$m_q : \begin{cases} \mathcal{H}_{\mathcal{PP}} \otimes \mathcal{H}_{\mathcal{PP}} & \longrightarrow & \mathcal{H}_{\mathcal{PP}} \\ P \otimes Q & \longrightarrow & \sum_{\substack{(R,I) \in \mathcal{PP}^2 \\ I \subseteq R, R \setminus I = P, I = Q}} q_1^{h_{R \setminus I}^I} q_2^{h_I^{R \setminus I}} q_3^{r_{R \setminus I}^I} q_4^{r_I^{R \setminus I}} R, \end{cases}$$

where $P, Q \in \mathcal{PP}$. Then $(\mathcal{H}_{\mathcal{PP}}, m_q)$ is an associative algebra, and its unit is the empty plane poset 1 .

Proof. Let $P, Q, R \in \mathcal{PP}$. We put:

$$\begin{cases} P_1 & = & (m_q \otimes Id) \circ m_q(P \otimes Q \otimes R), \\ P_2 & = & (Id \otimes m_q) \circ m_q(P \otimes Q \otimes R). \end{cases}$$

Then:

$$\begin{cases} P_1 & = & \sum_{(R_1, I_1, R_2, I_2) \in E_1} q_1^{h_{R_1 \setminus I_1}^{I_1} + h_{R_2 \setminus I_2}^{I_2}} q_2^{h_{I_1}^{R_1 \setminus I_1} + h_{I_2}^{R_2 \setminus I_2}} q_3^{r_{R_1 \setminus I_1}^{I_1} + r_{R_2 \setminus I_2}^{I_2}} q_4^{r_{I_1}^{R_1 \setminus I_1} + r_{I_2}^{R_2 \setminus I_2}} R_2, \\ P_2 & = & \sum_{(R_1, I_1, R_2, I_2) \in E_2} q_1^{h_{R_1 \setminus I_1}^{I_1} + h_{R_2 \setminus I_2}^{I_2}} q_2^{h_{I_1}^{R_1 \setminus I_1} + h_{I_2}^{R_2 \setminus I_2}} q_3^{r_{R_1 \setminus I_1}^{I_1} + r_{R_2 \setminus I_2}^{I_2}} q_4^{r_{I_1}^{R_1 \setminus I_1} + r_{I_2}^{R_2 \setminus I_2}} R_2, \end{cases}$$

With:

$$\begin{aligned} E_1 &= \{(R_1, I_1, R_2, I_2) \in \mathcal{PP}^4 \mid I_1 \subseteq R_1, I_1 = Q, R_1 \setminus I_1 = P, I_2 \subseteq R_2, I_2 = R, R_2 \setminus I_2 = R_1\}, \\ E_2 &= \{(R_1, I_1, R_2, I_2) \in \mathcal{PP}^4 \mid I_1 \subseteq R_1, I_1 = R, R_1 \setminus I_1 = Q, I_2 \subseteq R_2, I_2 = R_1, R_2 \setminus I_2 = P\}. \end{aligned}$$

We shall also consider:

$$E = \{(R, J_1, J_2, J_3) \in \mathcal{PP}^4 \mid R = J_1 \sqcup J_2 \sqcup J_3, J_1 = P, J_2 = Q, J_3 = R\}.$$

First step. Let us consider the following maps:

$$\begin{aligned} \phi &: \begin{cases} E_1 & \longrightarrow & E \\ (R_1, I_1, R_2, I_2) & \longrightarrow & (R_2, R_1 \setminus I_1, I_1, I_2), \end{cases} \\ \phi' &: \begin{cases} E & \longrightarrow & E_1 \\ (R, J_1, J_2, J_3) & \longrightarrow & (J_1 \sqcup J_2, J_2, R, J_3). \end{cases} \end{aligned}$$

By definition of E and E_1 , these maps are well-defined, and an easy computation shows that $\phi \circ \phi' = Id_E$ and $\phi' \circ \phi = Id_{E_1}$, so ϕ is a bijection. Let $(R_1, I_1, R_2, I_2) \in E_1$. We put $\phi(R_1, I_1, R_2, I_2) = (R, J_1, J_2, J_3)$. Then:

$$h_{R_1 \setminus I_1}^{I_1} + h_{R_2 \setminus I_2}^{I_2} = h_{J_1}^{J_2} + h_{J_1 \sqcup J_2}^{J_3} = h_{J_1}^{J_2} + h_{J_1}^{J_3} + h_{J_2}^{J_3}.$$

Similar computations finally give:

$$P_1 = \sum_{(R, J_1, J_2, J_3) \in E} q_1^{h_{J_1}^{J_2} + h_{J_1}^{J_3} + h_{J_2}^{J_3}} q_2^{h_{J_2}^{J_1} + h_{J_3}^{J_1} + h_{J_3}^{J_2}} q_3^{r_{J_1}^{J_2} + r_{J_1}^{J_3} + r_{J_2}^{J_3}} q_4^{r_{J_2}^{J_1} + r_{J_3}^{J_1} + r_{J_3}^{J_2}} R.$$

Second step. Let us consider the following maps:

$$\begin{aligned} \psi &: \begin{cases} E_2 & \longrightarrow E \\ (R_1, I_1, R_2, I_2) & \longrightarrow (R_2, R_1 \setminus R_1, R_1 \setminus I_1, I_1), \end{cases} \\ \psi' &: \begin{cases} E & \longrightarrow E_2 \\ (R, J_1, J_2, J_3) & \longrightarrow (J_2 \sqcup J_3, J_3, R, J_2 \sqcup J_3). \end{cases} \end{aligned}$$

By definition of E and E_2 , these maps are well-defined, and a simple computation shows that $\psi \circ \psi' = Id_E$ and $\psi' \circ \psi = Id_{E_2}$, so ψ is a bijection. Let $(R_1, I_1, R_2, I_2) \in E_2$. We put $\psi(R_1, I_1, R_2, I_2) = (R, J_1, J_2, J_3)$. Then:

$$h_{R_1 \setminus I_1}^{I_1} + h_{R_2 \setminus I_2}^{I_2} = h_{J_2}^{J_3} + h_{J_1}^{J_2 \sqcup J_3} = h_{J_1}^{J_2} + h_{J_1}^{J_3} + h_{J_2}^{J_3}.$$

Similar computations finally give:

$$P_2 = \sum_{(R, J_1, J_2, J_3) \in E} q_1^{h_{J_1}^{J_2} + h_{J_1}^{J_3} + h_{J_2}^{J_3}} q_2^{h_{J_2}^{J_1} + h_{J_3}^{J_1} + h_{J_3}^{J_2}} q_3^{r_{J_1}^{J_2} + r_{J_1}^{J_3} + r_{J_2}^{J_3}} q_4^{r_{J_2}^{J_1} + r_{J_3}^{J_1} + r_{J_3}^{J_2}} R.$$

So m_q is associative.

Last step. Let $P \in \mathcal{PP}$. Then:

$$P.1 = \sum_{\substack{(R, I) \in \mathcal{PP}^2 \\ I \subseteq R, R \setminus I = P, I = 1}} h_{R \setminus I}^I q_1^{R \setminus I} q_2^{R \setminus I} q_3^{R \setminus I} q_4^{R \setminus I} R = q_1^{h_P^1} q_2^{h_P^1} q_3^{r_P^1} q_4^{r_P^1} P = P.$$

Similarly, for all $Q \in \mathcal{PP}$, $1.Q = Q$. □

Examples.

$$\begin{aligned} m_q(\cdot \otimes \cdot) &= q_3 q_4 \bullet \bullet + q_1 q_2 \mathbf{1}, \\ m_q(\cdot \otimes \mathbf{1}) &= q_3^2 \bullet \mathbf{1} + q_4^2 \mathbf{1} \bullet + q_2(q_3 + q_4) \mathbf{V} + q_1(q_3 + q_4) \mathbf{\Lambda} + (q_1^2 + q_1 q_2 + q_2^2) \mathbf{1} \mathbf{1}, \\ m_q(\cdot \otimes \bullet \bullet) &= q_1^2 \mathbf{V} + q_2^2 \mathbf{\Lambda} + (q_1 + q_2) q_4 \bullet \mathbf{1} + (q_1 + q_2) q_3 \mathbf{1} \bullet + (q_3^2 + q_3 q_4 + q_4^2) \bullet \bullet, \\ m_q(\mathbf{1} \otimes \cdot) &= q_3^2 \bullet \mathbf{1} + q_4^2 \mathbf{1} \bullet + q_1(q_3 + q_4) \mathbf{V} + q_2(q_3 + q_4) \mathbf{\Lambda} + (q_1^2 + q_1 q_2 + q_2^2) \mathbf{1} \mathbf{1}, \\ m_q(\bullet \bullet \otimes \cdot) &= q_2^2 \mathbf{V} + q_1^2 \mathbf{\Lambda} + (q_1 + q_2) q_4 \bullet \mathbf{1} + (q_1 + q_2) q_3 \mathbf{1} \bullet + (q_3^2 + q_3 q_4 + q_4^2) \bullet \bullet. \end{aligned}$$

The following result is immediate:

Proposition 15 *Let $(q_1, q_2, q_3, q_4) \in K^4$. Then:*

$$\left\{ \begin{array}{l} m_{(q_1, q_2, q_3, q_4)}^{op} = m_{(q_2, q_1, q_4, q_3)}, \\ m_{(q_1, q_2, q_3, q_4)} \circ (\iota \otimes \iota) = \iota \circ m_{(q_3, q_4, q_1, q_2)}, \\ m_{(q_1, q_2, q_3, q_4)} \circ (\alpha \otimes \alpha) = \alpha \circ m_{(q_2, q_1, q_3, q_4)}, \\ m_{(q_1, q_2, q_3, q_4)} \circ (\beta \otimes \beta) = \beta \circ m_{(q_1, q_2, q_4, q_3)}, \\ m_{(q_1, q_2, q_3, q_4)} \circ (\gamma \otimes \gamma) = \gamma \circ m_{(q_2, q_1, q_4, q_3)}. \end{array} \right.$$

2.2 Particular cases

Lemma 16 *Let $P \in \mathcal{PP}$ and $X \subseteq P$.*

1. X is a h -ideal of P if, and only if, $h_X^{P \setminus X} = 0$.

2. X is a r -ideal of P if, and only if, $r_X^{P \setminus X} = 0$.
3. The h -irreducible components of P are the h -irreducible components of X and $P \setminus X$ if, and only if, $h_{P \setminus X}^X = h_X^{P \setminus X} = 0$.
4. The r -irreducible components of P are the r -irreducible components of X and $P \setminus X$ if, and only if, $r_{P \setminus X}^X = r_X^{P \setminus X} = 0$.
5. $P = X(P \setminus X)$ if, and only if, $h_{P \setminus X}^X = h_X^{P \setminus X} = r_{P \setminus X}^X = 0$.
6. $P = X \dot{\setminus} (P \setminus X)$ if, and only if, $r_{P \setminus X}^X = r_X^{P \setminus X} = h_{P \setminus X}^X = 0$.

Proof. We give the proofs of points 1, 3 and 5. The others are similar.

1. \Leftarrow . Let $x \in X$, $y \in P$, such that $x \leq_h y$. As $h_X^{P \setminus X} = 0$, $y \notin P \setminus X$, so $y \in X$: X is a h -ideal.

\Rightarrow . Then, for all $x \in X$, $y \in P \setminus X$, $x \leq_h y$ is not possible. So $h_X^{P \setminus X} = 0$.

3. \Rightarrow . Let us put $P = P_1 \dots P_k$, where P_1, \dots, P_k are the h -irreducible components of P . By hypothesis, X is the disjoint union of certain P_i 's, and $P \setminus X$ is the disjoint union of the other P_i 's. So $h_{P \setminus X}^X = h_X^{P \setminus X} = 0$.

\Leftarrow . Let I be a h -irreducible component of P . If $I \cap X$ and $I \cap (P \setminus X)$, then for any $x \in I \cap X$ and any $y \in I \cap (P \setminus X)$, x and y are not comparable for \leq_h : contradiction. So I is included in I or in $P \setminus X$, so is a h -irreducible component of X or $P \setminus X$.

5. \Rightarrow . If $x \in X$ and $y \in P \setminus X$, then $x <_r y$, so $h_{P \setminus X}^X = h_X^{P \setminus X} = r_{P \setminus X}^X = 0$.

\Leftarrow . If $x \in X$ and $y \in P \setminus X$, by hypothesis we do not have $x <_h y$, $x >_h y$ nor $x >_r y$, so $x <_r y$. Hence, $P = X(P \setminus X)$. \square

Proposition 17 1. Let us assume that $q_3 = q_4 = 0$. Let $P = P_1 \dot{\setminus} \dots \dot{\setminus} P_k$ and $P' = P_{k+1} \dot{\setminus} \dots \dot{\setminus} P_{k+l}$ be two plane posets, decomposed into their r -connected components. Then:

$$m_q(P \otimes P') = \sum_{\sigma \in Sh(k,l)} \prod_{1 \leq i \leq k < j \leq k+l} Q_{i,j}(\sigma)^{|P_i||P_j|} P_{\sigma(1)} \dot{\setminus} \dots \dot{\setminus} P_{\sigma(k+l)},$$

where $Q_{i,j}(\sigma) = q_1$ if $\sigma^{-1}(i) < \sigma^{-1}(j)$ and q_2 if $\sigma^{-1}(i) > \sigma^{-1}(j)$.

2. Let us assume that $q_1 = q_2 = 0$. Let $P = P_1 \dots P_k$ and $P' = P_{k+1} \dots P_{k+l}$ be two plane posets, decomposed into their r -connected components. Then:

$$m_q(P \otimes P') = \sum_{\sigma \in Sh(k,l)} \prod_{1 \leq i \leq k < j \leq k+l} Q'_{i,j}(\sigma)^{|P_i||P_j|} P_{\sigma(1)} \dots P_{\sigma(k+l)},$$

where $Q'_{i,j}(\sigma) = q_3$ if $\sigma^{-1}(i) < \sigma^{-1}(j)$ and q_4 if $\sigma^{-1}(i) > \sigma^{-1}(j)$.

Proof. Let us prove the first point; the proof of the second point is similar. Let us consider a plane poset R such that the coefficient of R in $m_q(P \otimes P')$ is not zero. So there exists $I \subseteq R$ such that $R \setminus I = P$ and $I = P'$. Moreover, as $q_3 = q_4 = 0$, $r_{R \setminus I}^I = r_I^{R \setminus I} = 0$. By lemma 16-4, the r -connected components of R are the r -components of $R \setminus I$ and I . As a consequence, there exists a (k, l) -shuffle σ , such that $R = P_{\sigma(1)} \dot{\setminus} \dots \dot{\setminus} P_{\sigma(k+l)}$. Then $h_{R \setminus I}^I$ is the sum of $|P_i||P_j|$, where $1 \leq i \leq k < j \leq k+l$, such that $\sigma^{-1}(i) < \sigma^{-1}(j)$; $h_I^{R \setminus I}$ is the sum of $|P_i||P_j|$, where $1 \leq i \leq k < j \leq k+l$, such that $\sigma^{-1}(i) > \sigma^{-1}(j)$. This immediately implies the announced result. \square

Remarks.

1. The first point implies that:

- $m_{(q_1,0,0,0)}(P \otimes P') = q_1^{|P||P'|} P \not\leq P'$. In particular, $m_{(1,0,0,0)} = \not\leq$.
- $m_{(0,q_2,0,0)}(P \otimes P') = q_2^{|P||P'|} P' \not\leq P$. In particular, $m_{(0,1,0,0)} = \not\leq^{op}$.
- $m_{(0,0,q_3,0)}(P \otimes P') = q_3^{|P||P'|} P P'$. In particular, $m_{(1,0,0,0)} = m$.
- $m_{(0,0,0,q_4)}(P \otimes P') = q_4^{|P||P'|} P' P$. In particular, $m_{(1,0,0,0)} = m^{op}$.

2. It is possible to define m_q on the space of double posets. The same arguments prove that it is still associative. However, $m_{(1,0,0,0)}$ is not equal to $\not\leq$ on $\mathcal{H}_{\mathcal{PP}}$ and $m_{(0,0,1,0)}$ is not equal to m ; for example, if we denote by \wp_2 the double poset with two elements x, y , x and y being not comparable for \leq_h and \leq_r :

$$m_{(1,0,0,0)}(\cdot \otimes \cdot) = \mathbf{1} + 2\wp_2, \quad m_{(0,0,1,0)}(\cdot \otimes \cdot) = \dots + 2\wp_2.$$

2.3 Subalgebras and quotients

These two particular families of plane posets are used in [2, 3]:

Definition 18 *Let $P \in \mathcal{PP}$.*

1. *We shall say that P is a plane forest if it does not contain Λ as a plane subposet. The set of plane forests is denoted by \mathcal{PF} .*
2. *We shall say that P is WN ("without N") if it does not contain \mathbb{N} nor \mathbb{N} . The set of WN posets is denoted by \mathcal{WNP} .*

Examples. A plane poset is a plane forest if, and only if, its Hasse graph is a rooted forest.

$$\begin{aligned} \mathcal{PF}(0) &= \{1\}, \\ \mathcal{PF}(1) &= \{\cdot\}, \\ \mathcal{PF}(2) &= \{\dots, \mathbf{1}\}, \\ \mathcal{PF}(3) &= \{\dots, \cdot\mathbf{1}, \mathbf{1}\cdot, \mathbb{V}, \mathbf{1}\}, \\ \mathcal{PF}(4) &= \left\{ \dots, \dots\mathbf{1}, \cdot\mathbf{1}\cdot, \mathbf{1}\dots, \cdot\mathbb{V}, \mathbb{V}\cdot, \cdot\mathbf{1}\cdot, \mathbf{1}\mathbf{1}, \mathbb{W}, \mathbb{V}, \mathbb{V}, \mathbb{Y}, \mathbf{1}\mathbf{1} \right\}, \\ \mathcal{WNP}(0) &= \{1\}, \\ \mathcal{WNP}(1) &= \{\cdot\}, \\ \mathcal{WNP}(2) &= \{\dots, \mathbf{1}\}, \\ \mathcal{WNP}(3) &= \{\dots, \cdot\mathbf{1}, \mathbf{1}\cdot, \mathbb{V}, \mathbf{1}\}, \\ \mathcal{WNP}(4) &= \left\{ \dots, \dots\mathbf{1}, \cdot\mathbf{1}\cdot, \mathbf{1}\dots, \cdot\mathbb{V}, \mathbb{V}\cdot, \cdot\mathbf{1}\cdot, \mathbf{1}\mathbf{1}, \cdot\mathbb{A}, \mathbb{A}\cdot, \mathbf{1}\mathbf{1}, \mathbb{W}, \mathbb{V}, \mathbb{V}, \mathbb{Y}, \mathbf{1}\mathbf{1}, \mathbb{A}, \mathbf{1}\mathbb{A}, \mathbb{A}\mathbf{1}, \mathbb{A}, \mathbb{N}, \mathbb{D} \right\}. \end{aligned}$$

Definition 19 *We denote by:*

- $\mathcal{H}_{\mathcal{WNP}}$ the subspace of $\mathcal{H}_{\mathcal{PP}}$ generated by WN plane posets.
- $\mathcal{H}_{\mathcal{PF}}$ the subspace of $\mathcal{H}_{\mathcal{PP}}$ generated by plane forests.
- $I_{\mathcal{WNP}}$ the subspace of $\mathcal{H}_{\mathcal{PP}}$ generated by plane posets which are not WN.
- $I_{\mathcal{PF}}$ the subspace of $\mathcal{H}_{\mathcal{PP}}$ generated by plane posets which are not plane forests.

Note that $\mathcal{H}_{\mathcal{WN}\mathcal{P}}$ and $\mathcal{H}_{\mathcal{P}\mathcal{F}}$ are naturally identified with $\mathcal{H}_{\mathcal{P}\mathcal{P}}/I_{\mathcal{WN}\mathcal{P}}$ and $\mathcal{H}_{\mathcal{P}\mathcal{P}}/I_{\mathcal{P}\mathcal{F}}$.

Proposition 20 Let $q = (q_1, q_2, q_3, q_4) \in K^4$.

1. $\mathcal{H}_{\mathcal{WN}\mathcal{P}}$ is a subalgebra of $(\mathcal{H}_{\mathcal{P}\mathcal{P}}, m_q)$ if and only if, $q_1 = q_2 = 0$ or $q_3 = q_4 = 0$.
2. $\mathcal{H}_{\mathcal{P}\mathcal{F}}$ is a subalgebra of $(\mathcal{H}_{\mathcal{P}\mathcal{P}}, m_q)$ if and only if, $q_1 = q_2 = 0$.
3. $I_{\mathcal{WN}\mathcal{P}}$ and $I_{\mathcal{P}\mathcal{F}}$ are ideals of $(\mathcal{H}_{\mathcal{P}\mathcal{P}}, m_q)$.

Proof. 1. \Leftarrow . We use the notations of proposition 17-1. If $q_3 = q_4 = 0$, let us consider two WN posets P and P' . then the P_i 's are also WN, so for any $\sigma \in \mathfrak{S}_{k+l}$, $P_{\sigma^{-1}(1)} \dot{\downarrow} \dots \dot{\downarrow} P_{\sigma^{-1}(k+l)}$ is WN. As a conclusion, $m_q(P \otimes P') \in \mathcal{H}_{\mathcal{WN}\mathcal{P}}$. The proof is similar if $q_1 = q_2 = 0$, using proposition 17-2.

1. \Rightarrow . Let us consider the coefficients of \mathcal{U} and \mathcal{N} in certain products. We obtain:

	\mathcal{U}	\mathcal{N}
$m_q(\mathcal{A} \otimes \cdot)$	$q_1 q_3^2$	$q_1 q_4^2$
$m_q(\cdot \otimes \mathcal{A})$	$q_2 q_4^2$	$q_2 q_3^2$

If $\mathcal{H}_{\mathcal{WN}\mathcal{P}}$ is a subalgebra of $(\mathcal{H}_{\mathcal{P}\mathcal{P}}, m_q)$, then these four coefficients are zero, so, from the first row, $q_1 = 0$ or $q_3 = q_4 = 0$ and from the second row, $q_2 = 0$ or $q_3 = q_4 = 0$. As a conclusion, $q_1 = q_2 = 0$ or $q_3 = q_4 = 0$.

2. \Leftarrow . We use the notations of proposition 17-2. If $q_1 = q_2 = 0$, let us consider two plane forests P and P' . Then the P_i 's are plane trees, so for any $\sigma \in \mathfrak{S}_{k+l}$, $P_{\sigma^{-1}(1)} \dot{\downarrow} \dots \dot{\downarrow} P_{\sigma^{-1}(k+l)}$ is a plane forest. As a conclusion, $m_q(P \otimes P') \in \mathcal{H}_{\mathcal{P}\mathcal{F}}$.

2. \Rightarrow . Let us consider the coefficients of \mathcal{A} in certain products. We obtain:

	\mathcal{A}
$m_q(\cdot \otimes \cdot)$	q_2^2
$m_q(\cdot \otimes \cdot)$	q_1^2

If $\mathcal{H}_{\mathcal{P}\mathcal{F}}$ is a subalgebra of $(\mathcal{H}_{\mathcal{P}\mathcal{P}}, m_q)$, then $q_1 = q_2 = 0$.

3. Let P and P' be two plane posets such that P or P' is not WN. Let us consider a plane poset R such that the coefficient of R in $m_q(P \otimes P')$ is not zero. There exists $I \subseteq R$, such that $R \setminus I = P$ and $I = P'$. As P or P' is not WN, I or $R \setminus I$ contains \mathcal{U} or \mathcal{N} , so R contains \mathcal{U} or \mathcal{N} : R is not WN. So $m_q(P \otimes P') \subseteq I_{\mathcal{WN}\mathcal{P}}$. The proof is similar for $I_{\mathcal{P}\mathcal{F}}$, using \mathcal{A} instead of \mathcal{U} and \mathcal{N} . \square

3 Dual coproducts

3.1 Constructions

Dually, we give $\mathcal{H}_{\mathcal{P}\mathcal{P}}$ a family of coproducts Δ_q , for $q \in K^4$, defined for all $P \in \mathcal{P}\mathcal{P}$ by:

$$\Delta_q(P) = \sum_{I \subseteq P} q_1^{h_{P \setminus I}^I} q_2^{h_I^{P \setminus I}} q_3^{r_{P \setminus I}^I} q_4^{r_I^{P \setminus I}} (P \setminus I) \otimes I.$$

These coproducts are coassociative; their common counit is given by:

$$\varepsilon : \begin{cases} \mathcal{H}_{\mathcal{P}\mathcal{P}} & \longrightarrow K \\ P \in \mathcal{P}\mathcal{P} & \longrightarrow \delta_{1,P}. \end{cases}$$

Examples. We put, for all $P \in \mathcal{PP}$, nonempty, $\tilde{\Delta}_q(P) = \Delta(P) - P \otimes 1 - 1 \otimes P$.

$$\begin{aligned}
\tilde{\Delta}_q(\mathbf{1}) &= (q_1 + q_2) \cdot \otimes \cdot, \\
\tilde{\Delta}_q(\cdot \cdot) &= (q_3 + q_4) \otimes \cdot, \\
\tilde{\Delta}(\mathbf{!}) &= (q_1^2 + q_1 q_2 + q_2^2) \cdot \otimes \mathbf{!} + (q_1^2 + q_1 q_2 + q_2^2) \mathbf{!} \otimes \cdot, \\
\tilde{\Delta}(\mathbf{V}) &= q_2(q_3 + q_4) \cdot \otimes \mathbf{!} + q_1(q_3 + q_4) \mathbf{!} \otimes \cdot + q_1^2 \cdot \otimes \cdot + q_2^2 \cdot \otimes \cdot, \\
\tilde{\Delta}(\mathbf{\wedge}) &= q_1(q_3 + q_4) \cdot \otimes \mathbf{!} + q_2(q_3 + q_4) \mathbf{!} \otimes \cdot + q_2^2 \cdot \otimes \cdot + q_1^2 \cdot \otimes \cdot, \\
\tilde{\Delta}(\mathbf{!} \cdot) &= q_4^2 \cdot \otimes \mathbf{!} + q_3^2 \mathbf{!} \otimes \cdot + (q_1 + q_2) q_3 \cdot \otimes \cdot + (q_1 + q_2) q_4 \cdot \otimes \cdot, \\
\tilde{\Delta}(\cdot \mathbf{!}) &= q_3^2 \cdot \otimes \mathbf{!} + q_4^2 \mathbf{!} \otimes \cdot + (q_1 + q_2) q_4 \cdot \otimes \cdot + (q_1 + q_2) q_3 \cdot \otimes \cdot, \\
\tilde{\Delta}(\cdot \cdot \cdot) &= (q_3^2 + q_3 q_4 + q_4^2) \cdot \otimes \cdot \cdot + (q_3^2 + q_3 q_4 + q_4^2) \cdot \cdot \otimes \cdot.
\end{aligned}$$

Dualizing proposition 15:

Proposition 21 Let $(q_1, q_2, q_3, q_4) \in K^4$. Then:

$$\left\{ \begin{array}{l} \Delta_{(q_1, q_2, q_3, q_4)}^{op} = \Delta_{(q_2, q_1, q_4, q_3)}, \\ (\iota \otimes \iota) \circ \Delta_{(q_1, q_2, q_3, q_4)} = \Delta_{(q_3, q_4, q_1, q_2)} \circ \iota, \\ (\alpha \otimes \alpha) \circ \Delta_{(q_1, q_2, q_3, q_4)} = \Delta_{(q_2, q_1, q_3, q_4)} \circ \alpha, \\ (\beta \otimes \beta) \circ \Delta_{(q_1, q_2, q_3, q_4)} = \Delta_{(q_1, q_2, q_4, q_3)} \circ \beta, \\ (\gamma \otimes \gamma) \circ \Delta_{(q_1, q_2, q_3, q_4)} = \Delta_{(q_2, q_1, q_4, q_3)} \circ \gamma. \end{array} \right.$$

3.2 Particular cases

Proposition 22 Let $P \in \mathcal{PP}$. Then:

1. $\Delta_{(q, q, q, q)}(P) = \sum_{I \subseteq P} q^{|P \setminus I| |I|} (P \setminus I) \otimes I$.

2.

$$\left\{ \begin{array}{l} \Delta_{(0, q_2, q_3, q_4)}(P) = \sum_{I \text{ h-ideal of } P} q_2^{h_{P \setminus I}^I} q_3^{r_{P \setminus I}^I} q_4^{r_{P \setminus I}^I} I \otimes (P \setminus I), \\ \Delta_{(q_1, 0, q_3, q_4)}(P) = \sum_{I \text{ h-ideal of } P} q_1^{h_{P \setminus I}^I} q_4^{r_{P \setminus I}^I} q_3^{r_{P \setminus I}^I} (P \setminus I) \otimes I. \end{array} \right.$$

3.

$$\left\{ \begin{array}{l} \Delta_{(q_1, q_2, 0, q_4)}(P) = \sum_{I \text{ r-ideal of } P} q_1^{h_{P \setminus I}^I} q_2^{h_{P \setminus I}^I} q_4^{r_{P \setminus I}^I} I \otimes (P \setminus I), \\ \Delta_{(q_1, q_2, q_3, 0)}(P) = \sum_{I \text{ r-ideal of } P} q_2^{h_{P \setminus I}^I} q_1^{h_{P \setminus I}^I} q_3^{r_{P \setminus I}^I} (P \setminus I) \otimes I. \end{array} \right.$$

4.

$$\left\{ \begin{array}{l} \Delta_{(0, q_2, 0, q_4)}(P) = \sum_{I \text{ biideal of } P} q_2^{h_{P \setminus I}^I} q_4^{r_{P \setminus I}^I} I \otimes (P \setminus I), \\ \Delta_{(q_1, 0, q_3, 0)}(P) = \sum_{I \text{ biideal of } P} q_1^{h_{P \setminus I}^I} q_3^{r_{P \setminus I}^I} (P \setminus I) \otimes I. \end{array} \right.$$

5. If $P = P_1 \cdots P_k$, where the P_i 's are h-connected,

$$\Delta_{(0, 0, q_3, q_4)}(P) = \sum_{I \subseteq \{1, \dots, k\}} q_3^{\alpha_P(I)} q_4^{\alpha_P(\{1, \dots, k\} \setminus I)} P_I \otimes P_{\{1, \dots, k\} \setminus I},$$

with, for all $J = \{j_1, \dots, j_l\}$, $1 \leq j_1 < \dots < j_l \leq k$, $P_J = P_{j_1} \cdots P_{j_l}$, and $\alpha_P(J) = \sum_{i \in J, j \notin J, i < j} |P_i||P_j|$.

6. If $P = P_1 \dot{\downarrow} \cdots \dot{\downarrow} P_k$, where the P_i 's are r -connected,

$$\Delta_{(q_1, q_2, 0, 0)}(P) = \sum_{I \subseteq \{1, \dots, k\}} q_1^{\beta_P(I)} q_2^{\beta_P(\{1, \dots, k\} \setminus I)} P_I^{\dot{\downarrow}} \otimes P_{\{1, \dots, k\} \setminus I}^{\dot{\downarrow}},$$

with, for all $J = \{j_1, \dots, j_l\}$, $1 \leq j_1 < \dots < j_l \leq k$, $P_J \dot{\downarrow} = P_{j_1} \dot{\downarrow} \cdots \dot{\downarrow} P_{j_l}$, and $\beta_P(J) = \sum_{i \in J, j \notin J, i < j} |P_i||P_j|$.

7.

$$\left\{ \begin{array}{l} \Delta_{(q, 0, 0, 0)}(P) = \sum_{P_1 \dot{\downarrow} P_2 = P} q^{|P_1||P_2|} P_1 \otimes P_2, \\ \Delta_{(0, q, 0, 0)}(P) = \sum_{P_1 \dot{\downarrow} P_2 = P} q^{|P_1||P_2|} P_2 \otimes P_1, \\ \Delta_{(0, 0, q, 0)}(P) = \sum_{P_1 P_2 = P} q^{|P_1||P_2|} P_1 \otimes P_2, \\ \Delta_{(0, 0, 0, q)}(P) = \sum_{P_1 P_2 = P} q^{|P_1||P_2|} P_2 \otimes P_1. \end{array} \right.$$

8. $\Delta_{(0, 0, 0, 0)}(P) = P \otimes 1 + 1 \otimes P$ if $P \neq 1$.

Proof. We only prove points 1, 2, 5, 7 and 8; the others are proved in the same way.

1. Immediate, as $h_{P \setminus I}^I + h_I^{P \setminus I} + r_{P \setminus I}^I + r_I^{P \setminus I} = |I||P \setminus I|$ by lemma 13.

2. By lemma 16:

$$\Delta_{(q_1, 0, q_3, q_4)}(P) = \sum_{\substack{I \subseteq P \\ h_{P \setminus I}^I = 0}} q_1^{h_{P \setminus I}^I} q_3^{r_{P \setminus I}^I} q_4^{r_I^{P \setminus I}} (R \setminus I) \otimes I = \sum_{P \setminus I \text{ } h\text{-ideal of } P} q_1^{h_{P \setminus I}^I} q_3^{r_{P \setminus I}^I} q_4^{r_I^{P \setminus I}} (P \setminus I) \otimes I.$$

As $\Delta_{(0, q_2, q_3, q_4)} = \Delta_{(0, q_2, q_4, q_3)}^{op}$, we obtain also the first assertion.

5. By lemma 16, point 3:

$$\Delta_{(0, 0, q_3, q_4)}(P) = \sum_{\substack{I \subseteq P \\ h_{P \setminus I}^I = h_I^{P \setminus I} = 0}} q_3^{r_{P \setminus I}^I} q_4^{r_I^{P \setminus I}} (R \setminus I) \otimes I = \sum_{I \subseteq \{1, \dots, k\}} q_3^{r_{P \setminus P_I}^{P_I}} q_4^{r_{P_I}^{P \setminus P_I}} P_I^{\otimes} P_{\{1, \dots, k\} \setminus I},$$

and it is immediate that $r_{P \setminus P_I}^{P_I} = \alpha_P(I)$ and $r_{P_I}^{P \setminus P_I} = \alpha_P(\{1, \dots, k\} - I)$.

7. By lemma 16:

$$\Delta_{(0, 0, q, 0)}(P) = \sum_{\substack{I \subseteq P \\ h_{P \setminus I}^I = h_I^{P \setminus I} = r_I^{P \setminus I} = 0}} q^{r_{P \setminus I}^I} (P \setminus I) \otimes I = \sum_{P_1 P_2 = P} q^{r_{P_1}^{P_2}} P_1 \otimes P_2 = \sum_{P_1 P_2 = P} q^{|P_1||P_2|} P_1 \otimes P_2.$$

8. Let $I \subseteq P$, such that $h_{P \setminus I}^I = h_I^{P \setminus I} = r_{P \setminus I}^I = r_I^{P \setminus I} = 0$. Then:

$$|I||P \setminus I| = h_{P \setminus I}^I + h_I^{P \setminus I} + r_{P \setminus I}^I + r_I^{P \setminus I} = 0,$$

so $I = 1$ or $I = P$. □

Remark. In particular, the coproduct defined in section 1 is $\Delta_{(1,0,1,1)}$. The coproduct of deconcatenation, dual of m , is $\Delta_{(0,0,1,0)}$ and the coproduct of deconcatenation, dual of ζ , is $\Delta_{(1,0,0,0)}$.

3.3 Compatibilities with the products

Proposition 23 *Let $x, y \in \mathcal{H}_{\mathcal{PP}}$. We put $\Delta_q(x) = \sum x'_q \otimes x''_q$ and $\Delta_q(y) = \sum y'_q \otimes y''_q$, with the x'_q 's, x''_q 's, y'_q 's, y''_q 's homogeneous. Then:*

1. $\Delta_q(xy) = \sum \sum q_3^{|x'_q||y''_q|} q_4^{|x''_q||y'_q|} (x'_q y'_q) \otimes (x''_q y''_q)$.
2. $\Delta_q(x \zeta y) = \sum \sum q_1^{|x'_q||y''_q|} q_2^{|x''_q||y'_q|} (x'_q \zeta y'_q) \otimes (x''_q \zeta y''_q)$.

Proof. We only prove the first point; the proof of the second point is similar. Let $P, Q \in \mathcal{PP}$. Then:

$$\begin{aligned} \Delta_q(PQ) &= \sum_{I \subseteq P, J \subseteq Q} q_1^{h_{(PQ) \setminus (IJ)}^{IJ}} q_2^{h_{IJ}^{(PQ) \setminus (IJ)}} q_3^{r_{(PQ) \setminus (IJ)}^{IJ}} q_4^{r_{IJ}^{(PQ) \setminus (IJ)}} ((PQ) \setminus (IJ)) \otimes (IJ) \\ &= \sum_{I \subseteq P, J \subseteq Q} q_1^{h_{(PQ) \setminus (IJ)}^{IJ}} q_2^{h_{IJ}^{PQ \setminus IJ}} q_3^{r_{PQ \setminus IJ}^{IJ}} q_4^{r_{IJ}^{PQ \setminus IJ}} ((P \setminus I)(Q \setminus J)) \otimes (IJ). \end{aligned}$$

Moreover, for all $I \subseteq P, J \subseteq Q$:

- For all $x \in P \setminus I, y \in J, x <_r y$ in PQ , so $h_{P \setminus I}^J = 0$. Similarly, $h_{Q \setminus J}^I = 0$. Hence:

$$h_{PQ \setminus IJ}^{IJ} = h_{P \setminus I}^I + h_{Q \setminus J}^J + h_{P \setminus I}^J + h_{Q \setminus J}^I = h_{P \setminus I}^I + h_{Q \setminus J}^J.$$

- In the same way, $h_{IJ}^{PQ \setminus IJ} = h_I^{P \setminus I} + h_J^{Q \setminus J} + h_J^{P \setminus I} + h_I^{Q \setminus J} = h_I^{P \setminus I} + h_J^{Q \setminus J}$.
- For all $x \in P \setminus I, y \in J, x <_r y$ in PQ , so $r_{P \setminus I}^J = |P \setminus I||J|$. For all $x \in Q \setminus J, y \in I, x >_r y$, so $r_{Q \setminus J}^I = 0$. Hence:

$$r_{PQ \setminus IJ}^{IJ} = r_{P \setminus I}^I + r_{Q \setminus J}^J + r_{P \setminus I}^J + r_{Q \setminus J}^I = h_{P \setminus I}^I + h_{Q \setminus J}^J + |P \setminus I||J|.$$

- In the same way, $r_{IJ}^{PQ \setminus IJ} = r_I^{P \setminus I} + r_J^{Q \setminus J} + r_J^{P \setminus I} + r_I^{Q \setminus J} = r_I^{P \setminus I} + r_J^{Q \setminus J} + |I||Q \setminus J|$.

So:

$$\Delta_q(PQ) = \sum \sum q_3^{|P'_q||Q''_q|} q_4^{|P''_q||Q'_q|} (P'_q Q'_q) \otimes (P''_q Q''_q),$$

which is the announced formula. □

Examples.

1. If $q_3 = 1$, then $(\mathcal{H}_{\mathcal{PP}}, m, \Delta_q)$ is a braided Hopf algebra. The braiding is given, for all $P, Q \in \mathcal{PP}$, by:

$$c_{q_4}(P \otimes Q) = q_4^{|P||Q|} Q \otimes P.$$

In particular, if $q_4 = 1$, it is a Hopf algebra; if $q_4 = 0$, it is an infinitesimal Hopf algebra.

2. If $q_4 = 0$, the compatibility becomes the following:

$$\Delta_q(xy) = \sum q_3^{|x||y''_q|} (xy'_q) \otimes y''_q + \sum q_3^{|x'_q||y|} x'_q \otimes (x''_q y) - q_3^{|x||y|} x \otimes y.$$

In particular, if $q_3 = 1$, then it is an infinitesimal Hopf algebra.

3. If $q_2 = q_4 = 0$, then for all $x, y \in \mathcal{HPP}$:

$$\begin{aligned}\Delta_q(xy) &= (xy) \otimes 1 + 1 \otimes (xy) + \varepsilon(x)(\Delta_q(y) - y \otimes 1 - 1 \otimes y) \\ &\quad + \varepsilon(y)(\Delta_q(x) - x \otimes 1 - 1 \otimes x) + \varepsilon(x)\varepsilon(y)1 \otimes 1.\end{aligned}$$

In other terms, for all $x, y \in \mathcal{HPP}$, such that $\varepsilon(x) = \varepsilon(y) = 0$, xy is primitive.

4 Self-duality results

4.1 A first pairing on \mathcal{H}_q

Notations. If $P, Q \in \mathcal{PP}$, we denote by $Bij(P, Q)$ the set of bijections from P to Q .

Definition 24 Let P, Q be two double posets and let $\sigma \in Bij(P, Q)$. We put:

$$\left\{ \begin{array}{l} \phi_1(\sigma) = \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma(x) <_h \sigma(y)\} \\ \quad + \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma(x) >_h \sigma(y)\} \\ \quad + \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma(x) <_r \sigma(y)\} \\ \quad + \#\{(x, y) \in P^2 \mid x <_r y \text{ and } \sigma(x) <_h \sigma(y)\}, \\ \phi_2(\sigma) = \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma(x) <_h \sigma(y)\} \\ \quad + \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma(x) >_h \sigma(y)\} \\ \quad + \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma(x) >_r \sigma(y)\} \\ \quad + \#\{(x, y) \in P^2 \mid x <_r y \text{ and } \sigma(x) >_h \sigma(y)\}, \\ \phi_3(\sigma) = \#\{(x, y) \in P^2 \mid x <_r y \text{ and } \sigma(x) <_r \sigma(y)\}, \\ \phi_4(\sigma) = \#\{(x, y) \in P^2 \mid x <_r y \text{ and } \sigma(x) >_r \sigma(y)\}.\end{array} \right.$$

Lemma 25 Let P, Q be two double posets and let $\sigma \in Bij(P, Q)$. For all $i \in \{1, \dots, 4\}$, $\phi_i(\sigma^{-1}) = \phi_i(\sigma)$.

Proof. Indeed, as σ is a bijection:

$$\begin{aligned}\phi_1(\sigma^{-1}) &= \#\{(x, y) \in Q^2 \mid x <_h y \text{ and } \sigma^{-1}(x) <_h \sigma^{-1}(y)\} \\ &\quad + \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma^{-1}(x) >_h \sigma^{-1}(y)\} \\ &\quad + \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma^{-1}(x) <_r \sigma^{-1}(y)\} \\ &\quad + \#\{(x, y) \in P^2 \mid x <_r y \text{ and } \sigma^{-1}(x) <_h \sigma^{-1}(y)\} \\ &= \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma(x) <_h \sigma(y)\} \\ &\quad + \#\{(x, y) \in P^2 \mid x >_h y \text{ and } \sigma(x) <_h \sigma(y)\} \\ &\quad + \#\{(x, y) \in P^2 \mid x <_h y \text{ and } \sigma(x) <_r \sigma(y)\} \\ &\quad + \#\{(x, y) \in P^2 \mid x <_r y \text{ and } \sigma(x) <_h \sigma(y)\} \\ &= \phi_1(\sigma).\end{aligned}$$

The other equalities are proved similarly. □

Lemma 26 Let P_1, P_2, Q be double posets. There is a bijection:

$$\left\{ \begin{array}{l} Bij(P_1 P_2, Q) \longrightarrow \bigcup_{I \subseteq Q} Bij(P_1, R \setminus I) \times R(P_2, I) \\ \sigma \longrightarrow (\sigma|_{P_1}, \sigma|_{P_2}), \text{ with } I = \sigma(P_2).\end{array} \right.$$

Let $\sigma \in \text{Bij}(P_1 P_2, Q)$ and let (σ_1, σ_2) be its image by this bijection. Then:

$$\begin{cases} \phi_1(\sigma) &= \phi_1(\sigma_1) + \phi_1(\sigma_2) + h_{Q \setminus I}^I, \\ \phi_2(\sigma) &= \phi_2(\sigma_1) + \phi_2(\sigma_2) + h_I^{Q \setminus I}, \\ \phi_3(\sigma) &= \phi_3(\sigma_1) + \phi_3(\sigma_2) + r_{Q \setminus I}^I, \\ \phi_4(\sigma) &= \phi_4(\sigma_1) + \phi_4(\sigma_2) + r_I^{Q \setminus I}. \end{cases}$$

Proof. We put $P = P_1 P_2$. if $x \in P_1$ and $y \in P_2$, then $x <_r y$, so:

$$\begin{aligned} \phi_3(\sigma) &= \#\{(x, y) \in P_1^2 \mid x <_r y \text{ and } \sigma(x) <_r \sigma(y)\} \\ &\quad + \#\{(x, y) \in P_2^2 \mid x <_r y \text{ and } \sigma(x) <_r \sigma(y)\} \\ &\quad + \#\{(x, y) \in P_1 \times P_2 \mid x <_r y \text{ and } \sigma(x) <_r \sigma(y)\} \\ &\quad + \#\{(x, y) \in P_2 \times P_1 \mid x <_r y \text{ and } \sigma(x) <_r \sigma(y)\} \\ &= \phi_3(\sigma) + \phi_3(\sigma_2) + \#\{(x, y) \in P_1 \times P_2 \mid \sigma(x) <_r \sigma(y)\} + 0 \\ &= \phi_3(\sigma) + \phi_3(\sigma_2) + \#\{(x, y) \in (Q \setminus I) \times I \mid x <_r y\} \\ &= \phi_3(\sigma_1) + \phi_3(\sigma_2) + r_{Q \setminus I}^I. \end{aligned}$$

The other equalities are proved similarly. □

Theorem 27 Let $q = (q_1, q_2, q_3, q_4) \in K^4$. We define a pairing on $\mathcal{H}_{\mathcal{P}\mathcal{P}}$ by:

$$\langle P, Q \rangle_q = \sum_{\sigma \in \text{Bij}(P, Q)} q_1^{\phi_1(\sigma)} q_2^{\phi_2(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)}.$$

This pairing is symmetric and for all $x, y, z \in \mathcal{H}_{\mathcal{P}\mathcal{P}}$, $\langle xy, z \rangle_q = \langle x \otimes y, \Delta_q(z) \rangle_q$.

Proof. Let P, Q be two double posets. Then, by lemma 25:

$$\begin{aligned} \langle P, Q \rangle_q &= \sum_{\sigma \in \text{Bij}(P, Q)} q_1^{\phi_1(\sigma)} q_2^{\phi_2(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)} \\ &= \sum_{\sigma \in \text{Bij}(Q, P)} q_1^{\phi_1(\sigma^{-1})} q_2^{\phi_2(\sigma^{-1})} q_3^{\phi_3(\sigma^{-1})} q_4^{\phi_4(\sigma^{-1})} \\ &= \sum_{\sigma \in \text{Bij}(Q, P)} q_1^{\phi_1(\sigma)} q_2^{\phi_2(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)} \\ &= \langle Q, P \rangle_q. \end{aligned}$$

So this pairing is symmetric. Let P_1, P_2, Q be three double posets. By lemma 26:

$$\begin{aligned} \langle P_1 P_2, Q \rangle_q &= \sum_{\sigma \in \text{Bij}(P_1 P_2, Q)} q_1^{\phi_1(\sigma)} q_2^{\phi_2(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)} \\ &= \sum_{I \subseteq Q} \sum_{\sigma_1 \in \text{Bij}(P_1, Q \setminus I)} \sum_{\sigma_2 \in \text{Bij}(P_2, I)} q_1^{\phi_1(\sigma_1) + \phi_1(\sigma_2) + h_{Q \setminus I}^I} \\ &\quad \times q_2^{\phi_2(\sigma_1) + \phi_2(\sigma_2) + h_I^{Q \setminus I}} q_3^{\phi_3(\sigma_1) + \phi_3(\sigma_2) + r_{Q \setminus I}^I} q_4^{\phi_4(\sigma_1) + \phi_4(\sigma_2) + r_I^{Q \setminus I}} \\ &= \sum_{I \subseteq Q} q_1^{h_{R \setminus I}^I} q_2^{h_I^{R \setminus I}} q_3^{r_{R \setminus I}^I} q_4^{r_I^{R \setminus I}} \langle P_1, Q \setminus I \rangle_q \langle P_2, I \rangle_q \\ &= \langle P_1 \otimes P_2, \Delta_q(Q) \rangle_q. \end{aligned}$$

So $\langle -, - \rangle_q$ is a Hopf pairing. □

Remark. More generally, adapting this proof, it is possible to show that for all $x, y, z \in \mathcal{H}_{PP}$:

$$\langle m_{(0,0,q,0)}(x \otimes y), \Delta(z) \rangle_{(q_1, q_2, q_3, q_4)} = \langle x \otimes y, \Delta_{(qq_1, qq_2, qq_3, qq_4)}(z) \rangle_{(q_1, q_2, q_3, q_4)}.$$

Examples. $\langle \cdot, \cdot \rangle_q = 1$. The pairing of plane posets of degree 2 is given by the following array:

	1	..
1	$2q_1q_2$	$q_1 + q_2$
..	$q_1 + q_2$	$q_3 + q_4$

Proposition 28 For all double posets P, Q :

$$\langle P, Q \rangle_{(q_1, 0, q_3, q_4)} = \sum_{\sigma \in S(P, Q)} q_1^{\phi_1(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)}.$$

Consequently, $\langle -, - \rangle_{(1,0,1,1)}$ is the pairing $\langle -, - \rangle$ described in section 1.

Proof. Let $\sigma \in \text{Bij}(P, Q)$, such that the contribution of σ in the sum defining $\langle P, Q \rangle_{(q_1, 0, q_3, q_4)}$ is non zero. As $q_2 = 0$, necessarily $\phi_2(\sigma) = 0$. Hence, if $x <_h y$ in P , then $\sigma(x) <_r \sigma(y)$ in Q ; if $\sigma(x) <_h \sigma(y)$ in Q , then $x <_r y$ in P . So $\sigma \in S(P, Q)$. \square

4.2 Properties of the pairing

Lemma 29 Let P_1, P_2, Q be double posets. There is a bijection:

$$\begin{cases} \text{Bij}(P_1 \dot{\cup} P_2, Q) & \longrightarrow \bigcup_{I \subseteq Q} \text{Bij}(P_1, R \setminus I) \times R(P_2, I) \\ \sigma & \longrightarrow (\sigma|_{P_1}, \sigma|_{P_2}), \text{ with } I = \sigma(P_2). \end{cases}$$

Let $\sigma \in \text{Bij}(P_1 P_2, Q)$ and let (σ_1, σ_2) be its image by this bijection. Then:

$$\begin{cases} \phi_1(\sigma) & = \phi_1(\sigma_1) + \phi_1(\sigma_2) + h_{Q \setminus I}^I + h_I^{Q \setminus I} + r_{Q \setminus I}^I, \\ \phi_2(\sigma) & = \phi_2(\sigma_1) + \phi_2(\sigma_2) + h_{Q \setminus I}^I + h_I^{Q \setminus I} + r_I^{Q \setminus I}, \\ \psi_2(\sigma) & = \psi_2(\sigma_1) + \psi_2(\sigma_2) + h_{Q \setminus I}^I + h_I^{Q \setminus I}, \\ \phi_3(\sigma) & = \phi_3(\sigma_1) + \phi_3(\sigma_2), \\ \phi_4(\sigma) & = \phi_4(\sigma_1) + \phi_4(\sigma_2). \end{cases}$$

Proof. We put $P = P_1 \dot{\cup} P_2$. Let $x, y \in P$. If $x <_h y$, then $x, y \in P_1$, or $x, y \in P_2$, or $x \in P_1, y \in P_2$. If $x <_r y$, then $x, y \in P_1$, or $x, y \in P_2$. Moreover, if $x \in P_1$ and $x \in P_2$, then $x <_h y$. So:

$$\begin{aligned} \phi_1(\sigma) & = \phi_1(\sigma_1) + \phi_1(\sigma_2) \\ & \quad + \#\{(x, y) \in P_1 \times P_2 \mid (x <_h y) \text{ and } \sigma(x) <_h \sigma(y)\} \\ & \quad + \#\{(x, y) \in P_1 \times P_2 \mid (x <_h y) \text{ and } \sigma(x) >_h \sigma(y)\} \\ & \quad + \#\{(x, y) \in P_1 \times P_2 \mid (x <_h y) \text{ and } \sigma(x) <_r \sigma(y)\} \\ & = \phi_1(\sigma_1) + \phi_1(\sigma_2) \\ & \quad + \#\{(x, y) \in P_1 \times P_2 \mid \sigma(x) <_h \sigma(y)\} \\ & \quad + \#\{(x, y) \in P_1 \times P_2 \mid \sigma(x) >_h \sigma(y)\} \\ & \quad + \#\{(x, y) \in P_1 \times P_2 \mid \sigma(x) <_r \sigma(y)\} \\ & = \phi_1(\sigma_1) + \phi_1(\sigma_2) + h_{Q \setminus I}^I + h_I^{Q \setminus I} + r_{Q \setminus I}^I. \end{aligned}$$

The other equalities are proved similarly. \square

Proposition 30 For all $x, y, z \in \mathcal{HPP}$, $\langle x \downarrow y, z \rangle_q = \langle x \otimes y, \Delta_{(q_1 q_2, q_1 q_2, q_1, q_2)}(z) \rangle_q$.

Proof. Let P_1, P_2, Q be three double posets. By lemma 29:

$$\begin{aligned}
\langle P_1 \downarrow P_2, Q \rangle_q &= \sum_{\sigma \in \text{Bij}(P_1 \downarrow P_2, Q)} q_1^{\phi_1(\sigma)} q_2^{\phi_2(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)} \\
&= \sum_{I \subseteq Q} \sum_{\sigma_1 \in \text{Bij}(P_1, Q \setminus I)} \sum_{\sigma_2 \in \text{Bij}(P_2, I)} q_1^{\phi_1(\sigma_1) + \phi_1(\sigma_2) + h_{Q \setminus I}^I + h_I^{Q \setminus I} + r_{Q \setminus I}^I} \\
&\quad \times q_2^{\phi_2(\sigma_1) + \phi_2(\sigma_2) + h_{Q \setminus I}^I + h_I^{Q \setminus I} + r_I^{Q \setminus I}} q_3^{\phi_3(\sigma_1) + \phi_3(\sigma_2)} q_4^{\phi_4(\sigma_1) + \phi_4(\sigma_2)} \\
&= \sum_{I \subseteq Q} (q_1 q_2)^{h_{R \setminus I}^I} (q_1 q_2)^{h_I^{R \setminus I}} q_1^{r_{R \setminus I}^I} q_2^{r_I^{R \setminus I}} \langle P_1, Q \setminus I \rangle_q \langle P_2, I \rangle_q \\
&= \langle P_1 \otimes P_2, \Delta_{(q_1 q_2, q_1 q_2, q_1, q_2)}(Q) \rangle_q,
\end{aligned}$$

which is the announced formula. \square

Remarks.

1. In particular, if $q = (1, 0, 1, 1)$, for all $P, Q, R \in \mathcal{PP}$:

$$\langle P \downarrow Q, R \rangle = \langle P \otimes Q, \Delta_{(0,0,1,0)}(R) \rangle = \sum_{R=R_1 R_2} \langle P \otimes Q, R_1 \otimes R_2 \rangle.$$

this formula is already proved in [2].

2. More generally, it is possible to show that for all $x, y, z \in \mathcal{HPP}$:

$$\langle m_{(q,0,0,0)}(x \otimes y), \Delta(z) \rangle_{(q_1, q_2, q_3, q_4)} = \langle x \otimes y, \Delta_{(qq_1 q_2, qq_1 q_2, qq_1, qq_2)}(z) \rangle_{(q_1, q_2, q_3, q_4)}.$$

Proposition 31 For all $x, y \in \mathcal{HPP}$:

$$\begin{aligned}
\langle x, \beta(y) \rangle_{(q_1, q_2, q_3, q_4)} &= \langle x, y \rangle_{(q_1, q_2, q_4, q_3)}, & \langle x, \gamma(y) \rangle_{(q_1, q_2, q_3, q_4)} &= \langle x, y \rangle_{(q_2, q_1, q_4, q_3)}, \\
\langle \alpha(x), \alpha(y) \rangle_{(q_1, q_2, q_3, q_4)} &= \langle x, y \rangle_{(q_2, q_1, q_3, q_4)}, & \langle \alpha(x), \beta(y) \rangle_{(q_1, q_2, q_3, q_4)} &= \langle x, y \rangle_{(q_1, q_2, q_4, q_3)}, \\
\langle \alpha(x), \gamma(y) \rangle_{(q_1, q_2, q_3, q_4)} &= \langle x, y \rangle_{(q_2, q_1, q_4, q_3)}, & \langle \beta(x), \beta(y) \rangle_{(q_1, q_2, q_3, q_4)} &= \langle x, y \rangle_{(q_2, q_1, q_3, q_4)}, \\
\langle \gamma(x), \gamma(y) \rangle_{(q_1, q_2, q_3, q_4)} &= \langle x, y \rangle_{(q_1, q_2, q_3, q_4)}.
\end{aligned}$$

Proof. Let P, Q be two double posets. We put $P' = \alpha(P)$ and $Q' = \alpha(Q)$. Note that $\text{Bij}(P, Q) = \text{Bij}(P', Q')$. Let $\sigma \in \text{Bij}(P, Q)$. We denote it by σ' is we consider it as an element of $\text{Bij}(P', Q')$. By definition of P' and Q' , it is clear that $\phi_1(\sigma') = \phi_2(\sigma)$, $\phi_2(\sigma') = \phi_2(\sigma)$, $\phi_3(\sigma') = \phi_3(\sigma)$, and $\phi_4(\sigma') = \phi_4(\sigma)$. Hence:

$$\begin{aligned}
\langle P', Q' \rangle_{(q_1, q_2, q_3, q_4)} &= \sum_{\sigma' \in \text{Bij}(P', Q')} q_1^{\phi_1(\sigma')} q_2^{\phi_2(\sigma')} q_3^{\phi_3(\sigma')} q_4^{\phi_4(\sigma')} \\
&= \sum_{\sigma \in \text{Bij}(P, Q)} q_1^{\phi_2(\sigma)} q_2^{\phi_1(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)} \\
&= \langle P, Q \rangle_{(q_2, q_1, q_3, q_4)}.
\end{aligned}$$

The other assertions are proved in the same way. \square

Remark. In general, the pairing defined by $x \otimes y \rightarrow \langle x, \alpha(y) \rangle_{(q_1, q_2, q_3, q_4)}$ is not a pairing $\langle -, - \rangle_q$. However, when $q_1 = q_2$, it is possible to prove that for all $x, y \in \mathcal{HPP}$, $\langle x, \alpha(y) \rangle_{(q_1, q_1, q_3, q_4)} = \langle x, y \rangle_{(q_1, q_1, q_3, q_4)}$. Similarly, $\langle \beta(x), \gamma(y) \rangle_{(q_1, q_1, q_3, q_4)} = \langle x, y \rangle_{(q_1, q_1, q_4, q_3)}$.

4.3 Comparison of pairings with colinear parameters

Proposition 32 *Let $q \in K$. We define the following map:*

$$v_q : \begin{cases} \mathcal{H}_{\mathcal{PP}} & \longrightarrow \mathcal{H}_{\mathcal{PP}} \\ P \in \mathcal{PP} & \longrightarrow q^{h(P)}P, \end{cases}$$

where $h(P) = \#\{(x, y) \in P \mid x <_h y\}$. Then v_q is an algebra and coalgebra morphism from $(\mathcal{H}_{\mathcal{PP}}, m, \Delta_{(qq_1, qq_2, q_3, q_4)})$ to $(\mathcal{H}_{\mathcal{PP}}, m, \Delta_{(q_1, q_2, q_3, q_4)})$. Moreover, for all $x, y \in \mathcal{H}_{\mathcal{PP}}$:

$$\langle v_q(x), v_q(y) \rangle_{(q_1, q_2, q_3, q_4)} = \langle x, y \rangle_{(qq_1, qq_2, q_3, q_4)}.$$

Proof. Let $P_1, P_2 \in \mathcal{PP}$. Then $h(P_1 P_2) = h(P_1) + h(P_2)$, so $v_q(P_1 P_2) = v_q(P_1) v_q(P_2)$, and v_q is an algebra morphism. Let $P \in \mathcal{PP}$. For any $I \subseteq P$, as $P^2 = I^2 \sqcup (P \setminus I)^2 \sqcup (I \times (P \setminus I)) \sqcup ((P \setminus I) \times I)$:

$$h(P) = h(I) + h(P \setminus I) + h_{P \setminus I}^I + h_I^{P \setminus I}.$$

Consequently:

$$\begin{aligned} \Delta_{(q_1, q_2, q_3, q_4)} \circ v_q(P) &= \sum_{I \subseteq P} (qq_1)^{h_{P \setminus I}^I} (qq_2)^{h_I^{P \setminus I}} q_3^{r_{P \setminus I}^I} q_4^{r_I^{P \setminus I}} q^{h(P \setminus I)} (P \setminus I) \otimes q^{h(I)} I \\ &= (v_q \otimes v_q) \circ \Delta_{(qq_1, qq_2, q_3, q_4)}(P). \end{aligned}$$

So v_q is a coalgebra morphism. Let $P, Q \in \mathcal{PP}$ and let $\sigma \in \text{Bij}(P, Q)$. We define:

$$\begin{aligned} a_1 &= \#\{(x, y) \in P^2 \mid x <_h y, \sigma(x) <_h \sigma(y)\} \\ a_2 &= \#\{(x, y) \in P^2 \mid x <_h y, \sigma(x) >_h \sigma(y)\} \\ a_3 &= \#\{(x, y) \in P^2 \mid x <_h y, \sigma(x) <_r \sigma(y)\} \\ a_4 &= \#\{(x, y) \in P^2 \mid x <_h y, \sigma(x) >_r \sigma(y)\} \\ a_5 &= \#\{(x, y) \in P^2 \mid x <_r y, \sigma(x) <_h \sigma(y)\} \\ a_6 &= \#\{(x, y) \in P^2 \mid x <_r y, \sigma(x) >_h \sigma(y)\} \\ a_7 &= \#\{(x, y) \in P^2 \mid x <_r y, \sigma(x) <_r \sigma(y)\} \\ a_8 &= \#\{(x, y) \in P^2 \mid x <_r y, \sigma(x) >_r \sigma(y)\}. \end{aligned}$$

In order to sum up the notations, we give a following array:

$\#\{(x, y) \in P^2 \mid \dots\}$	$x <_h y, \dots$	$x >_h y, \dots$	$x <_r y, \dots$	$x >_r y, \dots$
$\sigma(x) <_h \sigma(y)$	a_1	a_2	a_5	a_6
$\sigma(x) >_h \sigma(y)$	a_2	a_1	a_6	a_5
$\sigma(x) <_r \sigma(y)$	a_3	a_4	a_7	a_8
$\sigma(x) >_r \sigma(y)$	a_4	a_3	a_8	a_7

So $\phi_1(\sigma) = a_1 + a_2 + a_3 + a_5$, $\phi_2(\sigma) = a_1 + a_2 + a_4 + a_6$, $\phi_3(\sigma) = a_7$ and $\phi_4(\sigma) = a_8$; $h(P) = a_1 + a_2 + a_3 + a_4$ and $h(Q) = a_1 + a_2 + a_5 + a_6$. Then:

$$\begin{aligned} q^{h(P)} q^{h(Q)} q_1^{\phi_1(\sigma)} q_2^{\phi_2(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)} &= (q^2 q_1 q_2)^{a_1} (q^2 q_1 q_2)^{a_2} (qq_1)^{a_3} (qq_2)^{a_4} (qq_1)^{a_5} (qq_2)^{a_6} q_3^{a_7} q_4^{a_8} \\ &= (qq_1)^{\phi_1(\sigma)} (qq_2)^{\phi_2(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)}. \end{aligned}$$

Finally:

$$\begin{aligned} \langle v_q(P), v_q(Q) \rangle_{(q_1, q_2, q_3, q_4)} &= \sum_{\sigma \in \text{Bij}(P, Q)} q^{h(P)} q^{h(Q)} q_1^{\phi_1(\sigma)} q_2^{\phi_2(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)} \\ &= \sum_{\sigma \in \text{Bij}(P, Q)} (qq_1)^{\phi_1(\sigma)} (qq_2)^{\phi_2(\sigma)} q_3^{\phi_3(\sigma)} q_4^{\phi_4(\sigma)} \\ &= \langle P, Q \rangle_{(qq_1, qq_2, q_3, q_4)}. \end{aligned}$$

So v_q is an isometry. □

Remark. It is easy to prove that for $P, Q \in \mathcal{PP}$, $v_q(P \dot{\downarrow} Q) = v_q(P) \dot{\downarrow} v_q(Q)$.

Similarly, one can prove:

Proposition 33 *Let $q \in K$. We define the following map:*

$$v'_q : \begin{cases} \mathcal{H}_{\mathcal{PP}} & \longrightarrow \mathcal{H}_{\mathcal{PP}} \\ P \in \mathcal{PP} & \longrightarrow q^{r(P)}P, \end{cases}$$

where $r(P) = \#\{(x, y) \in P \mid x <_r y\}$. Then v'_q is an algebra and coalgebra morphism from $(\mathcal{H}_{\mathcal{PP}}, \dot{\downarrow}, \Delta_{(q_1, q_2, qq_3, qq_4)})$ to $(\mathcal{H}_{\mathcal{PP}}, \dot{\downarrow}, \Delta_{(q_1, q_2, q_3, q_4)})$. Moreover, for all $x, y \in \mathcal{H}_{\mathcal{PP}}$:

$$\langle v'_q(x), v'_q(y) \rangle_{(q_1, q_2, q_3, q_4)} = \langle x, y \rangle_{(q_1, q_2, qq_3, qq_4)}.$$

4.4 Non-degeneracy of the pairing

Lemma 34 *For all $P \in \mathcal{PP}(n)$, $S(P, \iota(P))$ is reduced to a single element and:*

$$\langle P, \iota(P) \rangle_{(q_1, 0, q_3, q_4)} = q_1^{\frac{n(n-1)}{2}}.$$

Proof. Clearly, $Id_P : P \longrightarrow P$ belongs to $S(P, \iota(P))$. Let $\sigma : P \longrightarrow P$ be a bijection. Then $\sigma \in S(P, \iota(P))$ if, and only if, for all $i, j \in P$:

- $(i \leq_h j \text{ in } P) \implies (\sigma(i) \leq_h \sigma(j) \text{ in } P)$.
- $(\sigma(i) \leq_r \sigma(j)) \text{ in } P \implies (i \leq_r j \text{ in } P)$.

It is clear $S(P, \iota(P))$ is a submonoid of \mathfrak{S}_P . As the group \mathfrak{S}_P is finite, $S(P, \iota(P))$ is a subgroup of \mathfrak{S}_P . So if $\sigma \in S(P, \iota(P))$, then $\sigma^{-1} \in S(P, \iota(P))$. So, if $\sigma \in S(P, \iota(P))$:

- $(i \leq_h j \text{ in } P) \iff (\sigma(i) \leq_h \sigma(j) \text{ in } P)$.
- $(i \leq_r j \text{ in } P) \iff (\sigma(i) \leq_r \sigma(j) \text{ in } P)$.

So σ is the unique increasing bijection from P to P , that is to say Id_P . Moreover:

$$\begin{aligned} \phi_1(Id_P) &= \#\{(x, y) \in P^2 \mid x <_h y \text{ and } x <_r y\} + \#\{(x, y) \in P^2 \mid x <_h y \text{ and } x >_r y\} \\ &\quad + \#\{(x, y) \in P^2 \mid x <_h y \text{ and } x <_h y\} + \#\{(x, y) \in P^2 \mid x <_r y \text{ and } x <_r y\} \\ &= 0 + 0 + \#\{(x, y) \in P^2 \mid x <_h y\} + \#\{(x, y) \in P^2 \mid x <_r y\} \\ &= \#\{(x, y) \in P^2 \mid x < y\} \\ &= \frac{n(n-1)}{2}, \\ \phi_3(Id_P) &= \#\{(x, y) \in P^2 \mid (x <_r y) \text{ and } (x <_h y)\} \\ &= 0, \\ \phi_4(Id_P) &= \#\{(x, y) \in P^2 \mid (x >_r y) \text{ and } (x >_h y)\} \\ &= 0. \end{aligned}$$

So $\langle P, \iota(P) \rangle_{(q_1, 0, q_2, q_3)} = q_1^{\frac{n(n-1)}{2}}$. □

For all $n \in \mathbb{N}$, we give \mathfrak{S}_n the lexicographic order. For example, if $n = 3$:

$$(123) \leq (132) \leq (213) \leq (231) \leq (312) \leq (321).$$

We then define a total order \ll on $\mathcal{PP}(n)$ by $P \ll Q$ if, and only if, $\Psi_n(P) \leq \Psi_n(Q)$ in \mathfrak{S}_n . For example, if $n = 3$:

$$\mathfrak{I} \ll \mathfrak{V} \ll \mathfrak{A} \ll \mathfrak{I} \ll \mathfrak{I} \ll \dots$$

For all $P \in \mathcal{PP}(n)$, we put:

$$m(P) = \min_{\ll} \{Q \in \mathcal{PP}(n) / S(P, Q) \neq \emptyset\}.$$

Lemma 35 *For all $P \in \mathcal{PP}$, $m(P) = \iota(P)$.*

Proof. By lemma 34, $m(P) \ll \iota(P)$. Let $Q \in \mathcal{PP}$, such that $S(P, Q) \neq \emptyset$. Let us prove that $\iota(P) \ll Q$. We denote $\sigma = \Psi_n(P)$ and $\tau = \Psi_n(Q)$; we can suppose that $P = \Phi_n(\sigma)$ and $Q = \Phi_n(\tau)$. Moreover, it is not difficult to prove that $\Psi_n(\iota(P)) = (n \cdots 1) \circ \sigma$.

First step. Let us prove that $\tau(1) \geq n - \sigma(1) + 1$. In $P = \Psi_n(\sigma)$, $1 \leq_h j$ if, and only if, $\sigma(1) \leq \sigma(j)$. So there are exactly $n - \sigma(1) + 1$ elements of P which satisfy $1 \leq_h j$ in P . Let $\alpha \in S(P, Q)$ (which is non-empty by hypothesis). Then if $1 \leq_h j$, $\alpha(1) \leq_r \alpha(j)$: there are at least $n - \sigma(1) + 1$ elements of Q which satisfy $\alpha(1) \leq_r j$. Let us put $\alpha^{-1}(1) = i$. Then $\alpha(i) = 1 \leq \alpha(1)$ in Q , that is to say $1 \leq_h \alpha(1)$ or $1 \leq_r \alpha(1)$ in Q . If $1 = \alpha(i) \leq_h \alpha(1)$ in Q , then $i \leq_r 1$ in P , so $i = 1$: in both cases, $1 \leq_r \alpha(1)$ in P . So there are at least $n - \sigma(1) + 1$ elements of Q which satisfy $1 \leq_r j$. In Q , $1 \leq_r j$ if, and only if, $\tau(j) \leq \tau(1)$, so there are exactly $\tau(1)$ such elements. As a consequence, $\tau(1) \geq n - \sigma(1) + 1$. Moreover, if there is equality, necessarily $1 = \alpha(1)$ for all $\alpha \in S(P, Q)$.

Second step. We consider the assertion H_i : if $\tau(1) = n - \sigma(1) + 1, \dots, \tau(i-1) = n - \sigma(i-1) + 1$, then $\tau(i) \geq n - \sigma(i) + 1$ and, if there is equality, then $\alpha(1) = 1, \dots, \alpha(i) = i$ for all $\alpha \in S(P, Q)$. We proved H_0 in the first step. Let us prove H_i by induction on i . Let us assume H_{i-1} , $1 \leq i \leq n$, and let us prove H_i . Let $\alpha \in S(P, Q)$ (non empty by hypothesis). By H_{i-1} , as the equality is satisfied, $\alpha(1) = 1, \dots, \alpha(i-1) = i-1$.

In P , the number N_i of elements j such that $i \leq_h j$ is the cardinality of $\sigma^{-1}(\{\sigma(i), \dots, n\}) \cap \{i, \dots, n\}$, so $N_i = n - \sigma(i) + 1 - |\{k \leq i / \sigma(k) > \sigma(i)\}|$. Using α , there exists at least N_i elements j of Q such that $\alpha(i) \leq_r j$ in Q .

Let us put $\alpha^{-1}(i) = j$. As $\alpha(1) = 1, \dots, \alpha(i-1) = i-1$, $j \geq i$ and $i \leq \alpha(i)$. If $i \leq_h \alpha(i)$ in Q , then $j \leq_r i$ in P , so $j = i$. So we always have $i \leq_r \alpha(i)$ in Q , so at least N_i elements k of Q satisfy $i \leq_r j$ in Q . Hence:

$$\begin{aligned} N_i &\leq |\tau^{-1}(\{1, \dots, \tau(i)\}) \cap \{i, \dots, n\}| \\ n - \sigma(i) + 1 - |\{k \leq i / \sigma(k) > \sigma(i)\}| &\leq \tau(i) - |\{k \leq i / \tau(k) < \tau(i)\}| \\ &\leq \tau(i) - |\{k \leq i / n - \sigma(k) + 1 < n - \sigma(i) + 1\}| \\ &\leq \tau(i) - |\{k \leq i / \sigma(k) > \sigma(i)\}|, \end{aligned}$$

so $\tau(i) \geq n - \sigma(i) + 1$. If there is equality, then necessarily $i = \alpha(i)$ for all $\alpha \in S(P, Q)$.

Conclusion. The hypothesis H_i is true for all $0 \leq i \leq n$. So $\tau \geq (n \cdots 1) \circ \sigma$ in \mathfrak{S}_n , so $\iota(P) \ll Q$ in $\mathcal{PP}(n)$. As a conclusion, $\iota(P) \ll m(P)$. \square

Corollary 36 *Let us assume that $q_2 = 0$. Then the pairing $\langle -, - \rangle_q$ is non-degenerate if, and only if, $q_1 \neq 0$.*

Proof. Let us fix an integer $n \in \mathbb{N}$. We consider the basis $\mathcal{PP}(n)$ of $\mathcal{H}(n)$, totally ordered by \ll . In this basis, the matrix of $\langle -, - \rangle_q$ restricted to $\mathcal{H}(n)$ has the following form, coming from

lemmas 34 and 35:

$$\begin{pmatrix} 0 & \cdots & \cdots & 0 & q_1^{\frac{n(n-1)}{2}} \\ \vdots & & & \frac{n(n-1)}{2} & * \\ \vdots & & \ddots & q_1 & \vdots \\ 0 & q_1^{\frac{n(n-1)}{2}} & \ddots & \ddots & \vdots \\ q_1^{\frac{n(n-1)}{2}} & * & \cdots & \cdots & * \end{pmatrix},$$

so its determinant is $\pm q_1^{\frac{n!n(n-1)}{2}}$. Hence, $\langle -, - \rangle_q$ is non-degenerate if, and only if, $q_1 \neq 0$. \square

Remark. With the help of the isometry α (proposition 31), it is possible to prove that if $q_1 = 0$, $\langle -, - \rangle_q$ is non-degenerate if, and only if, $q_2 \neq 0$.

Corollary 37 *If q_1, q_2, q_3, q_4 are algebraically independent over \mathbb{Z} , then the pairing $\langle -, - \rangle_q$ is non-degenerate.*

Proof. For all n , let us consider the matrix of the pairing restricted to $\mathcal{H}_{\mathcal{PP}}(n)$ in the basis formed by the plane posets of degree n . Its determinant D_n is clearly an element of $\mathbb{Z}[q_1, q_2, q_3, q_4]$. Moreover, $D_n(1, 0, q_3, q_4) \neq 0$ by corollary 36, so D_n is a non-zero polynomial. As a consequence, if q_1, q_2, q_3, q_4 are algebraically independent over \mathbb{Z} , $D_n(q_1, q_2, q_3, q_4) \neq 0$. \square

5 Morphism to free quasi-symmetric functions

5.1 A second Hopf pairing on $\mathcal{H}_{(q_1, 0, q_1, q_4)}$

We here assume that $q_2 = 0$ and $q_1 = q_3$. For all double poset P ,

$$\Delta_{(q_1, 0, q_1, q_4)}(P) = \sum_{I \text{ } h\text{-ideal of } P} q_1^{h_{R \setminus I}^I + r_{R \setminus I}^I} q_4^{r_I^{P \setminus I}} (P \setminus I) \otimes I.$$

For any h -ideal I of P , $h_I^{P \setminus I} = 0$, so:

$$\begin{aligned} h_{R \setminus I}^I + r_{R \setminus I}^I &= \#\{(x, y) \in (P \setminus I) \times I \mid x < y\}, \\ r_I^{P \setminus I} &= r_I^{P \setminus I} + h_I^{P \setminus I} \\ &= \#\{(x, y) \in (P \setminus I) \times I \mid x > y\}. \end{aligned}$$

Hence:

$$\Delta_{(q_1, 0, q_1, q_4)}(P) = \sum_{I \text{ } h\text{-ideal of } P} q_1^{\#\{(x, y) \in (P \setminus I) \times I \mid x < y\}} q_4^{\#\{(x, y) \in (P \setminus I) \times I \mid x > y\}} (P \setminus I) \otimes I.$$

Notations. Let $P, Q \in \mathcal{P}(n)$ and let $\sigma \in \text{Bij}(P, Q)$. As totally ordered sets, $P = Q = \{1, \dots, n\}$, so σ can be seen as an element of the symmetric group \mathfrak{S}_n . Its *length* $\ell(\sigma)$ is then its length in the Coxeter group \mathfrak{S}_n [5].

Theorem 38 *We define a pairing $\langle -, - \rangle'_q : \mathcal{H}_q \otimes \mathcal{H}_q \longrightarrow K$ by:*

$$\langle P, Q \rangle'_q = \sum_{\sigma \in S'(P, Q)} q_1^{\frac{n(n-1)}{2} - \ell(\sigma)} q_4^{\ell(\sigma)},$$

for all $P, Q \in \mathcal{PP}$, with $n = \text{deg}(P)$. Then $\langle -, - \rangle'_q$ is an homogeneous symmetric Hopf pairing on the braided Hopf algebra $\mathcal{H}_q = (\mathcal{H}, m, \Delta_q)$.

Proof. This pairing is clearly homogeneous. For any double posets P and Q , the map from $S'(P, Q)$ to $S'(Q, P)$ sending σ to its inverse is a bijection and conserves the length: this implies that the $\langle P, Q \rangle'_q = \langle Q, P \rangle'_q$.

Let P, Q, R be three double posets. There is a bijection:

$$\Theta : \begin{cases} \text{Bij}(PQ, R) & \longrightarrow \bigcup_{I \subseteq R} \text{Bij}(P, R \setminus I) \times \text{Bij}(Q, I) \\ \sigma & \longrightarrow (\sigma|_P, \sigma|_Q), \end{cases}$$

with $I = \sigma(Q)$. Let us first prove that $\sigma \in S'(PQ, R)$ if, and only if, I is a h -ideal of R and $(\sigma|_P, \sigma|_Q) \in S'(P, R \setminus I) \times S'(Q, I)$.

\implies . Let $x' \in I$ and let $y' \in R$ such that $x' \leq_h y'$. We put $x' = \sigma(x)$ and $y' = \sigma(y)$. As $\sigma \in S'(PQ, R)$, $x \leq y$. As $x \in Q$, $y \in Q$, so $y' \in I$: I is a h -ideal of R . By restriction, $\sigma|_P \in S'(P, R \setminus I)$ and $\sigma|_Q \in S'(Q, I)$.

\impliedby . Let us assume that $x \leq_h y$ in PQ . Then $x, y \in P$ or $x, y \in Q$. As $\sigma|_P \in S'(P, R \setminus I)$ and $\sigma|_Q \in S'(Q, I)$, $\sigma(x) \leq \sigma(y)$ in R . Let us assume that $\sigma(x) \leq_h \sigma(y)$ in R . As I is a h -ideal, there are two possibilities:

- $\sigma(x), \sigma(y) \in R \setminus I$ or $\sigma(x), \sigma(y) \in I$. Hence, $x, y \in P$ or $x, y \in Q$. As $\sigma|_P \in S'(P, R \setminus I)$ and $\sigma|_Q \in S'(Q, I)$, $x \leq y$ in PQ .
- $\sigma(x) \in R \setminus I$ and $\sigma(y) \in I$. Hence, $x \in P$ and $y \in Q$, so $x \leq y$ in PQ .

Finally, we obtain a bijection:

$$\Theta : \begin{cases} S'(PQ, R) & \longrightarrow \bigcup_{I \text{ } h\text{-ideal of } R} S'(P, R \setminus I) \times S'(Q, I) \\ \sigma & \longrightarrow (\sigma|_P, \sigma|_Q), \end{cases}$$

Let $\sigma \in S'(PQ, R)$ and we put $\Theta(\sigma) = (\sigma_1, \sigma_2)$. Let $R \setminus I = \{i_1, \dots, i_k\}$ and $I = \{j_1, \dots, j_l\}$. Let $\zeta \in \mathfrak{S}_{k+l}$ defined by $\zeta^{-1}(1) = i_1, \dots, \zeta^{-1}(k) = i_k, \zeta^{-1}(k+1) = j_1, \dots, \zeta^{-1}(k+l) = j_l$. Then ζ is a (k, l) -shuffle and $\sigma = \zeta^{-1} \circ (\sigma_1 \otimes \sigma_2)$, so $\ell(\sigma) = \ell(\zeta) + \ell(\sigma_1) + \ell(\sigma_2)$. Moreover:

$$\ell(\zeta) = \#\{(p, q) \mid j_q < i_p\} = \#\{(i, j) \in (R \setminus I) \times I \mid j < i\} = r_I^{R \setminus I} + h_I^{R \setminus I} = r_I^{R \setminus I},$$

as I is a h -ideal of R . Finally, $\ell(\sigma) = \ell(\sigma_1) + \ell(\sigma_2) + r_I^{R \setminus I}$. Moreover, if $|R| = n$ and $|I| = k$, by lemma 13:

$$h_{R \setminus I}^I + r_{R \setminus I}^I + r_I^{R \setminus I} = h_{R \setminus I}^I + r_{R \setminus I}^I + h_I^{R \setminus I} + r_I^{R \setminus I} = k(n - k),$$

so:

$$\begin{aligned} & h_{R \setminus I}^I + r_{R \setminus I}^I + \frac{k(k-1)}{2} + \frac{(n-k)(n-k-1)}{2} \\ &= k(n-k) + \frac{k(k-1)}{2} + \frac{(n-k)(n-k-1)}{2} - r_I^{R \setminus I} \\ &= \frac{n(n-1)}{2} - r_I^{R \setminus I}. \end{aligned}$$

Finally:

$$\begin{aligned}
\langle P, Q \rangle'_q &= \sum_{\sigma \in S'(PQ, R)} q_1^{\frac{n(n-1)}{2} - \ell(\sigma)} q_4^{\ell(\sigma)} \\
&= \sum_{I \text{ h-ideal of } R} \sum_{\sigma_1 \in S'(P, R \setminus I), \sigma_2 \in S'(Q, I)} q_1^{\frac{n(n-1)}{2} - \ell(\sigma_1) - \ell(\sigma_2) - r_I^{R \setminus I}} q_4^{\ell(\sigma_1) + \ell(\sigma_2) + r_I^{R \setminus I}} \\
&= \sum_{I \text{ h-ideal of } R} \sum_{\sigma_1 \in S'(P, R \setminus I), \sigma_2 \in S'(Q, I)} q_1^{h_{R \setminus I}^I + r_{R \setminus I}^I + \frac{k(k-1)}{2} + \frac{(n-k)(n-k-1)}{2}} q_4^{\ell(\sigma_1) + \ell(\sigma_2) + r_I^{R \setminus I}} \\
&= \sum_{I \text{ h-ideal of } R} q_1^{h_{R \setminus I}^I + r_{R \setminus I}^I} q_4^{r_I^{R \setminus I}} \langle P, R \setminus I \rangle'_q \langle Q, I \rangle'_q \\
&= \langle P \otimes Q, \Delta_q(R) \rangle'_q.
\end{aligned}$$

So this pairing is a Hopf pairing on \mathcal{H}_q . □

Remark. In particular, $\langle -, - \rangle'_{(1,0,1,1)}$ is the pairing $\langle -, - \rangle'$ of the section 1.

Examples. Here are the matrices of the pairing $\langle -, - \rangle'_q$ restricted to $\mathcal{H}_q(n)$, for $n = 1, 2, 3$.

$$\begin{array}{c|c} \cdot & \cdot \\ \hline \cdot & 1 \end{array} \quad \begin{array}{c|c|c} & \mathbf{1} & \dots \\ \hline \mathbf{1} & q_1 & q_1 \\ \hline \dots & q_1 & q_1 + q_4 \end{array}$$

	$\mathbf{1}$	\vee	\wedge	$\mathbf{1.}$	$\mathbf{.1}$	\dots
$\mathbf{1}$	q_1^3	q_1^3	q_1^3	q_1^3	q_1^3	q_1^3
\vee	q_1^3	$q_1^2(q_1 + q_4)$	q_1^3	$q_1^2(q_1 + q_4)$	q_1^3	$q_1^2(q_1 + q_4)$
\wedge	q_1^3	q_1^3	$q_1^2(q_1 + q_4)$	q_1^3	$q_1^2(q_1 + q_4)$	$q_1^2(q_1 + q_4)$
$\mathbf{1.}$	q_1^3	$q_1^2(q_1 + q_4)$	q_1^3	$q_1^2(q_1 + q_4)$	$q_1(q_1^2 + q_1^4)$	$q_1(q_1^2 + q_1q_4 + q_4^2)$
$\mathbf{.1}$	q_1^3	q_1^3	$q_1^2(q_1 + q_4)$	$q_1(q_1^2 + q_1^4)$	$q_1^2(q_1 + q_4)$	$q_1(q_1^2 + q_1q_4 + q_4^2)$
\dots	q_1^3	$q_1^2(q_1 + q_4)$	$q_1^2(q_1 + q_4)$	$q_1(q_1^2 + q_1q_4 + q_4^2)$	$q_1(q_1^2 + q_1q_4 + q_4^2)$	$(q_1 + q_4)(q_1^2 + q_1q_4 + q_4^2)$

Proposition 39 For all $x, y, z \in \mathcal{H}_q$, $\langle x \dot{\smile} y, z \rangle'_q = \langle x \otimes y, \Delta_{(q_1, 0, q_1, 0)}(z) \rangle'_q$.

Proof. Let $P, Q, R \in \mathcal{PP}$. We consider the following map:

$$\Theta : \begin{cases} \text{Bij}(P \dot{\smile} Q, R) & \longrightarrow \bigcup_{I \subseteq R} \text{Bij}(P, R \setminus I) \times \text{Bij}(Q, I) \\ \sigma & \longrightarrow (\sigma|_P, \sigma|_Q), \end{cases}$$

with $I = \sigma(Q)$. Let us first prove that $\sigma \in S'(PQ, R)$ if, and only if, I is a biideal of R and $(\sigma_1, \sigma_2) \in S'(P, R \setminus I) \times S'(Q, I)$.

\implies . Obviously, $(\sigma_1, \sigma_2) \in S'(P, R \setminus I) \times S'(Q, I)$. Let $x' \in I$, $y' \in R$, such that $x' \leq y'$. We put $x' = \sigma(x)$ and $y' = \sigma(y)$. If $y \notin Q$, then $y \in P$ and $x \in Q$, so $y <_h x$. As $\sigma \in S'(P \dot{\smile} Q, R)$, $y' < x'$: contradiction. So $y \in Q$ and $y' \in I$.

\impliedby . Let $x, y \in P \dot{\smile} Q$, such that $x \leq_h y$. Two cases are possible.

- If $x, y \in P$ or $x, y \in Q$, as $(\sigma_1, \sigma_2) \in S'(P, R \setminus I) \times S'(Q, I)$, $\sigma(x) \leq \sigma(y)$ in R .
- If $x \in P$ and $y \in Q$, as $I = \sigma(Q)$ is a biideal, it is not possible to have $y \leq x$, so $x \leq y$.

Let $x, y \in P \not\downarrow Q$, such that $\sigma(x) \leq_h \sigma(y)$ in R . As $I = \sigma(Q)$ is a biideal, two cases are possible.

- If $x, y \in P$ or $x, y \in Q$, as $(\sigma_1, \sigma_2) \in S'(P, R \setminus I) \times S'(Q, I)$, $x \leq y$ in $P \not\downarrow Q$.
- If $x \in P$ and $y \in Q$, then $x \leq y$ in $P \not\downarrow Q$.

Moreover, if $I = \sigma(Q)$ is a biideal of R , then $I = \{k+1, \dots, k+l\} \subseteq R$, where $k = |P|$ and $l = |Q|$. Then $\sigma = \sigma_1 \otimes \sigma_2$, so $\ell(\sigma) = \ell(\sigma_1) + \ell(\sigma_2)$. So, with $n = |R|$:

$$\begin{aligned}
\langle P \not\downarrow Q, R \rangle'_q &= \sum_{I \text{ biideal of } R} \sum_{\sigma_1 \in S'(P, R \setminus I), \sigma_2 \in S'(Q, I)} q_1^{\frac{n(n-1)}{2} - \ell(\sigma_1) - \ell(\sigma_2)} q_4^{\ell(\sigma_1) + \ell(\sigma_2)} \\
&= \sum_{I \text{ biideal of } R} q_1^{\frac{n(n-1)}{2} - \frac{k(k-1)}{2} - \frac{(n-k)(n-k-1)}{2}} \langle P \otimes Q, (R \setminus I) \otimes I \rangle'_q \\
&= \sum_{I \text{ biideal of } R} q_1^{(n-k)k} \langle P \otimes Q, (R \setminus I) \otimes I \rangle'_q \\
&= \sum_{I \text{ biideal of } R} q_1^{|R \setminus I| \cdot |I|} \langle P \otimes Q, (R \setminus I) \otimes I \rangle'_q \\
&= \sum_{I \text{ biideal of } R} q_1^{h_{R \setminus I}^I + r_{R \setminus I}^I} \langle P \otimes Q, (R \setminus I) \otimes I \rangle'_q \\
&= \langle P \otimes Q, \Delta_{(q_1, 0, q_1, 0)}(R) \rangle'_q,
\end{aligned}$$

with the observation that, as I is a biideal, $h_I^{R \setminus I} = r_I^{R \setminus I} = 0$, so $|R \setminus I| \cdot |I| = h_{R \setminus I}^I + r_{R \setminus I}^I$. \square

Remark. In particular, if $q = (1, 0, 1, 1)$, for all $P, Q, R \in \mathcal{PP}$:

$$\langle P \not\downarrow Q, R \rangle' = \langle P \otimes Q, \Delta_{(1, 0, 1, 0)}(R) \rangle' = \sum_{I \text{ biideal of } R} \langle P \otimes Q, (R \setminus I) \otimes I \rangle'.$$

This formula was already proved in [3].

5.2 Quantization of the Hopf algebra of free quasi-symmetric functions

Theorem 40 1. We define a coproduct on \mathbf{FQSym} in the following way: for all $\sigma \in \mathfrak{S}_n$,

$$\Delta_q(\sigma) = \sum_{k=0}^n q_1^{k(n-k) - \ell(\sigma) + \ell(\sigma_k^{(1)}) + \ell(\sigma_k^{(2)})} q_4^{\ell(\sigma) - \ell(\sigma_k^{(1)}) - \ell(\sigma_k^{(2)})} \sigma_k^{(1)} \otimes \sigma_k^{(2)}.$$

For all $q \in K$, $\mathbf{FQSym}_q = (\mathbf{FQSym}, m, \Delta_q)$ is a graded braided Hopf algebra.

2. One defines a Hopf pairing on \mathbf{FQSym}_q by $\langle \sigma, \tau \rangle' = q_1^{\frac{n(n-1)}{2} - \ell(\sigma)} q_4^{\ell(\sigma)} \delta_{\sigma, \tau^{-1}}$.

3. The following map is an isomorphism of braided Hopf algebras:

$$\Theta : \begin{cases} \mathcal{H}_q & \longrightarrow \mathbf{FQSym}_q \\ P \in \mathcal{PP} & \longrightarrow \sum_{\sigma \in S_P} \sigma. \end{cases}$$

Moreover, for all $x, y \in \mathcal{H}_q$, $\langle \Theta(x), \Theta(y) \rangle'_q = \langle x, y \rangle'_q$.

Remarks.

1. Let $\sigma \in \mathfrak{S}_n$. If $0 \leq k \leq n$, we put $\sigma(\{1, \dots, k\}) = \{i_1, \dots, i_k\}$ and $\sigma(\{k+1, \dots, n\}) = \{j_1, \dots, j_l\}$. Let ζ be the (k, l) -shuffle defined by $\zeta^{-1}(1) = i_1, \dots, \zeta^{-1}(k) = i_k, \zeta^{-1}(k+1) = j_1, \dots, \zeta^{-1}(k+l) = j_l$. Then $\sigma = \zeta^{-1} \circ (\sigma_k^{(1)} \otimes \sigma_k^{(2)})$, and $\ell(\sigma) = \ell(\zeta) + \ell(\sigma_k^{(1)}) + \ell(\sigma_k^{(2)})$. Moreover, $0 \leq \ell(\zeta) \leq k(n-k)$, so:

$$0 \leq \ell(\sigma) - \ell(\sigma_k^{(1)}) - \ell(\sigma_k^{(2)}) \leq k(n-k).$$

As a consequence, the coproduct on \mathbf{FQSym}_q is well-defined, even if $q_1 = 0$ or $q_4 = 0$.

2. It is clear that the pairing $\langle -, - \rangle'_q$ defined on \mathbf{FQSym}_q is non degenerate if, and only if, $q_1 \neq 0$ and $q_2 \neq 0$.

Proof. We already know that Θ is an algebra isomorphism. Moreover, it is also proved there that $S'(P, Q) = S(P) \cap S(Q)^{-1}$ for any $P, Q \in PP$, so:

$$\begin{aligned} \langle \Theta(P), \Theta(Q) \rangle'_q &= \sum_{\sigma \in S(P), \tau \in S(Q)} \langle \sigma, \tau \rangle'_q \\ &= \sum_{\sigma \in S(P), \tau \in S(Q)} q_1^{\frac{n(n-1)}{2} - \ell(\sigma)} q_4^{\ell(\sigma)} \delta_{\sigma, \tau^{-1}} \\ &= \sum_{\sigma \in S(P) \cap S(Q)^{-1}} q_1^{\frac{n(n-1)}{2} - \ell(\sigma)} q_4^{\ell(\sigma)} \\ &= \sum_{\sigma \in S'(P, Q)} q_1^{\frac{n(n-1)}{2} - \ell(\sigma)} q_4^{\ell(\sigma)} \\ &= \langle P, Q \rangle'_q. \end{aligned}$$

So Θ is an isometry.

As a consequence, for all $x \in \mathcal{H}_q, y, z \in \mathbf{FQSym}_q$:

$$\begin{aligned} \langle \Delta \circ \Theta(x), y \otimes z \rangle'_q &= \langle \Theta(x), yz \rangle'_q \\ &= \langle x, \Theta^{-1}(yz) \rangle'_q \\ &= \langle x, \Theta^{-1}(y)\Theta^{-1}(z) \rangle'_q \\ &= \langle \Delta_q(x), \Theta^{-1}(y) \otimes \Theta^{-1}(z) \rangle'_q \\ &= \langle (\Theta \otimes \Theta) \circ \Delta_q(x), y \otimes z \rangle'_q. \end{aligned}$$

Let us now assume that $q_1, q_4 \neq 0$. Then the pairing $\langle -, - \rangle'_q$ on \mathbf{FQSym}_q is non-degenerate, so $\Delta \circ \Theta = (\Theta \otimes \Theta) \circ \Delta$. Hence, Θ is an isometric isomorphism of braided Hopf algebras. This proves the three points immediately.

Up to an extension, we can assume that K is infinite. It is not difficult to show that the set A of elements $(q_1, q_4) \in K^2$ such that the three points are satisfied is given by polynomial equations with coefficients in \mathbb{Z} . As it contains $(K - \{0\})^2$ from the preceding observations, which is dense in K^2 , it is equal to K^2 , so the result also holds for any (q_1, q_4) . \square

Corollary 41 *The pairing $\langle -, - \rangle'_q$ is non-degenerate if, and only if, $q_1 \neq 0$ and $q_4 \neq 0$.*

Proof. As the isometric pairing $\langle -, - \rangle'_q$ on \mathbf{FQSym}_q is non-degenerate if, and only if, $q_1 \neq 0$ and $q_4 \neq 0$. \square

References

- [1] Gérard Duchamp, Florent Hivert, and Jean-Yves Thibon, *Noncommutative symmetric functions. VI. Free quasi-symmetric functions and related algebras*, *Internat. J. Algebra Comput.* **12** (2002), no. 5, 671–717, arXiv:math/0105065.
- [2] Loïc Foissy, *Algebraic structures on double and plane posets*, to be published in *Journal of Algebraic Combinatorics*, arXiv:1101.5231, 2011.
- [3] ———, *Plane posets, special posets, and permutations*, arXiv:1109.1101, 2011.
- [4] Loïc Foissy and Jeremie Unterberger, *Ordered forests, permutations and iterated integrals*, *Int. Math. Res. Notices* (2012), doi: 10.1093/imrn/rnr273 F, arXiv:1004.5208.
- [5] James E. Humphreys, *Reflection groups and Coxeter groups*, *Cambridge Studies in Advanced Mathematics*, vol. 29, Cambridge University Press, Cambridge, 1990.
- [6] Jean-Louis Loday and María Ronco, *On the structure of cofree Hopf algebras*, *J. Reine Angew. Math.* **592** (2006), 123–155, arXiv:math/0405330.
- [7] Claudia Malvenuto and Christophe Reutenauer, *Duality between quasi-symmetric functions and the Solomon descent algebra*, *J. Algebra* **177** (1995), no. 3, 967–982.
- [8] Claudia Malvenuto and Christophe Reutenauer, *A self-dual Hopf algebra on double partially ordered sets*, arXiv:0905.3508.