



HAL
open science

Homologies of Algebraic Structures via Braidings and Quantum Shuffles

Victoria Lebed

► **To cite this version:**

Victoria Lebed. Homologies of Algebraic Structures via Braidings and Quantum Shuffles. 2012. hal-00687866v2

HAL Id: hal-00687866

<https://hal.science/hal-00687866v2>

Preprint submitted on 27 Oct 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Homologies of Algebraic Structures via Braidings and Quantum Shuffles

Victoria Lebed
lebed@math.jussieu.fr

October 27, 2012

Abstract

In this paper we construct “structural” pre-braidings characterizing different algebraic structures: a rack, an associative algebra, a Leibniz algebra and their representations. Some of these pre-braidings seem original. On the other hand, we propose a general homology theory for pre-braided vector spaces and braided modules, based on the quantum co-shuffle comultiplication. Applied to the structural pre-braidings above, it gives a generalization and a unification of many known homology theories. All the constructions are categorified, resulting in particular in their super- and co-versions. Loday’s hyper-boundaries, as well as certain homology operations are efficiently treated using the “shuffle” tools.

Keywords: pre-braiding; braided (co)algebra; braided homology; character; braided module; quantum shuffle algebra; Koszul complex; rack homology; Hochschild homology; Leibniz algebra; pre-braided object.

Mathematics Subject Classification 2010: 18D10, 20F36, 18G60, 55N35, 16E40, 17A32, 17D99, 18G30, 05E99.

Contents

1	Introduction	2
2	Braided world: a short reminder	5
3	Homologies of braided vector spaces	9
3.1	Pre-braiding + character \mapsto homology	9
3.2	Comultiplication \mapsto degeneracies	11
3.3	Arrow and concatenation operations	13
3.4	Loday’s hyper-boundaries	15
4	Basic examples: familiar complexes recovered	16
4.1	Vector spaces	17
4.2	Self-distributive structures	17
4.3	Associative algebras	21
4.4	Leibniz algebras	24
5	An upper world: categories	25
5.1	Categorifying braided differentials	26
5.2	Basic examples revisited	28
5.3	The super trick	30
5.4	Co-world, or the world upside down	31
5.5	Right-left duality	34
6	Braided modules and homologies with coefficients	34

1 Introduction

The aim of this paper is to develop a unifying framework for (co)homologies of algebraic structures. Our starting point is the following fundamental procedure:

$$\boxed{\text{algebraic structure} \rightsquigarrow \text{chain complex.}}$$

Figure 1: Homology of algebraic structures

The step \rightsquigarrow is far from being canonical, and can be dictated by motivations of very different nature: structure deformations and obstructions, classification questions, topological applications etc. Here we propose to regard it from a purely combinatorial viewpoint, in the spirit of operad theory. The complexes one associates in practice to basic algebraic structures on a vector space V usually have the same flavor: they are all **signed** sums $d_n : V^{\otimes n} \rightarrow V^{\otimes(n-1)}$ of **terms of the same nature** $d_{n;i}$, one for each component $1, 2, \dots, n$ of $V^{\otimes n}$.

The examples we have in mind are

1. *Koszul complex* for vector spaces,
2. *bar and Hochschild complexes* for associative algebras,
3. *Chevalley-Eilenberg complex* for Lie algebras,
4. *rack and shelf (or one-term distributive) complexes* for self-distributive (= SD) structures.

Verifying that one has indeed a differential, i.e. $d_{n-1} \circ d_n = 0$, can be reduced to checking some **local** algebraic identities (which mysteriously coincide with the defining properties for our algebraic structure!) coupled with a sign manipulation, no less mysterious.

For many algebraic structures, their chain complexes can be refined by introducing a (weakly) (pre)(bi)**simplicial structure** on $T(V)$ (see section 3.2 or J.-L.Loday’s book [23] for the simplicial vocabulary). Moreover, the degree -1 differentials can be generalized to **Loday’s hyperboundaries** of arbitrary degree (see the definitions from section 3.4, or exercise E.2.2.7 in [23], from which this notion takes inspiration). Some **homology operations**, similar for different algebraic structures, are also to be mentioned here. Such common features are presented, for the example of associative and SD structures, in J.Przytycki’s paper [31].

In this work, we propose to interpret and partially explain these parallels (typed in bold letters above) and mysteries by adding a new step to the scheme in figure 1:

$$\boxed{\text{algebraic structure} \xrightarrow{\text{case by case}} \text{pre-braiding} \xrightarrow{\text{theorem 2}} \text{chain complex.}}$$

Figure 2: Homology of algebraic structures via pre-braidings

After a short reminder on braided structures in section 2, we proceed to describing in detail the right part of this new scheme. More precisely, given a vector space endowed with a *pre-braiding* $\sigma : V \otimes V \rightarrow V \otimes V$ satisfying the *Yang-Baxter equation* (=YBE)

$$(\sigma \otimes \text{Id}_V) \circ (\text{Id}_V \otimes \sigma) \circ (\sigma \otimes \text{Id}_V) = (\text{Id}_V \otimes \sigma) \circ (\sigma \otimes \text{Id}_V) \circ (\text{Id}_V \otimes \sigma) \in \text{End}(V \otimes V \otimes V),$$

we associate in theorem 2 a bidifferential on $T(V)$ to any couple of *braided characters* (= elements of V^* “respecting” σ) ϵ and ζ , using *quantum co-shuffle comultiplication* techniques (cf. M.Rosso’s pioneer papers [34],[35]). We call such (bi)differentials *braided*.

In theorem 3 we refine these braided bidifferential structures: we show that they come from a pre-bisimplicial structure on $T(V)$, completed to a weakly bisimplicial one if V is moreover endowed with a coassociative σ -cocommutative *comultiplication* Δ compatible with the pre-braiding σ . This is done using the graphical calculus (in the spirit of J.C.Baez ([1]), S.Majid ([29]) and other authors), appearing naturally due to our use of “braided” techniques. Here are for example the components of the weakly bisimplicial structure from the theorem (all diagrams are to be read from bottom to top here):




Figure 3: Simplicial structure for braided homology

See table 2 for a comparison of the quantum co-shuffle and the graphical approaches to braided differentials. Braided differentials are generalized to hyper-boundaries in section 3.4, and some homology operations for them are studied in section 3.3.

Note that we never demand σ to be invertible, emphasizing it in the term pre-braiding. Rare in literature, this elementary generalization of the notion of braiding allows interesting examples.

Armed with this general homology theory for pre-braided vector spaces, we are now interested in the left part of figure 2. Unfortunately we do not know any systematic way of associating a pre-braiding to an algebraic structure. So we do it by hand in section 4 for each of the four structures in the list above. In each case, the pre-braiding σ we propose encodes surprisingly well the structure in question, in the sense that

- YBE for σ is equivalent to the defining relation for the structure (e.g. the associativity for an algebra), under some mild assumptions concerning units;
- the invertibility condition for σ , when this makes sense, translates important algebraic properties (e.g. the rack condition);
- braided morphisms (= those preserving σ) are precisely algebra morphisms (= those preserving the underlying structure);
- braided characters for σ include the usual characters for the structure (e.g. algebra characters);
- the comultiplication necessary for constructing the degeneracies turns out to be quite characteristic of the structures.

Thus the left part of the scheme in figure 2 can be informally stated in a stronger way:

$$\boxed{\text{“algebraic structure} = \text{pre-braiding”}.$$

Besides its unifying character, our braided homology theory has the following advantages:

1. We produce two compatible differentials – a left and a right one – for each braided character, these differentials often being compatible even for different characters. Their combinations thus give a family of homology theories for the same algebraic structure. For instance, for SD structures we recover
 - (a) usual shelf ([32], [31]), rack ([13]) and quandle ([5]) homology theories;
 - (b) the partial derivatives from [30];
 - (c) and the twisted rack homology from [3].
2. The technical sign manipulation (especially heavy for the Chevalley-Eilenberg complex) is controlled either by using the negative pre-braiding $-\sigma$ in the quantum co-shuffle comultiplication, or by counting the number of intersections in the graphical interpretation.

3. The identities $d_{n-1} \circ d_n = 0$, which are of “global” nature, are replaced with the YBE for the corresponding pre-braiding, which is “local” and thus easier to verify.
4. The decomposition $d_n = \sum_{i=1}^n (-1)^{i-1} d_{n;i}$ becomes natural when one reasons in terms of braids and strands.
5. So do some homology operations – for instance, the generalizations of the homology operations for shelves, defined by M.Niebrzydowski and J.Przytycki in [30].
6. Subscript chasing (in the relations defining simplicial structures for instance) is substituted with the more transparent “strand chasing”.
7. J.-L.Loday’s hyper-boundaries of degree $-i$ arise naturally in the co-shuffle interpretation: one simply replaces the $V^{\otimes n} \rightarrow V^{\otimes 1} \otimes V^{\otimes (n-1)}$ component of the quantum co-shuffle co-product with the $V^{\otimes n} \rightarrow V^{\otimes i} \otimes V^{\otimes (n-i)}$ component.

We thus recover all the common features of different algebraic homology theories mentioned (in bold letters) in the beginning of this introduction. Moreover, we obtain a simplified and conceptual way of proving that $d^2 = 0$, as well as of “guessing” the right boundary map. As an illustration, note that the “braided” considerations naturally lead one to lifting the Chevalley-Eilenberg complex from the external to the tensor algebra of a Lie algebra, reinterpreting the work of J.-L.Loday on *Leibniz algebras* ([23]) from a more conceptual viewpoint. We also recover for Leibniz algebras the braiding studied in the Lie case by A.Crans ([8], [2]).

In section 6 we introduce the notion of a *braided module* over a pre-braided space (V, σ) , unifying, quite unexpectedly, the familiar notions of modules over different structures. Concretely, it is a space M equipped with a linear map $\rho : M \otimes V \rightarrow M$ satisfying

$$\rho \circ (\rho \otimes \text{Id}_V) = \rho \circ (\rho \otimes \text{Id}_V) \circ (\text{Id}_M \otimes \sigma) : M \otimes V \otimes V \rightarrow M.$$

These braided modules are natural candidates for coefficients in the braided complexes, leading us to *braided homologies with coefficients*, recovering for example the Hochschild and Chevalley-Eilenberg homologies.

We now give a table of braided structures encoding the algebraic structures mentioned above, and of the familiar complexes recovered as particular cases of our braided complexes.

structure	pre-braiding	invertibility	Δ	complexes
vector space V	flip $\tau : v \otimes w \mapsto w \otimes v$	$\tau^{-1} = \tau$	–	Koszul
unital associative algebra $(V, \cdot, \mathbf{1})$	$\sigma_\mu : v \otimes w \mapsto \mathbf{1} \otimes vw$	no inverse in general	$\Delta(v) = \mathbf{1} \otimes v$	bar, Hochschild
unital Leibniz algebra $(V, [,], \mathbf{1})$	$\sigma_{[,] : v \otimes w \mapsto w \otimes v + \mathbf{1} \otimes [v, w]$	$\exists \sigma_{[,]^{-1}}$	$\Delta(v) = v \otimes \mathbf{1} + \mathbf{1} \otimes v,$ $\Delta(\mathbf{1}) = \mathbf{1} \otimes \mathbf{1}$	Leibniz, Chevalley-Eilenberg
shelf $(S, \triangleleft), V := \mathbb{k}S$	$\sigma_{\triangleleft} : (a, b) \mapsto (b, a \triangleleft b)$	$\exists \sigma_{\triangleleft}^{-1}$ iff S is a rack	$\Delta(a) = (a, a)$	shelf, quandle, (twisted) rack

structure	braided characters	braided modules	homology operations
vector space	any $\epsilon \in V^*$	space endowed with commutative operators	multiplication by scalars
unital associative algebra	algebra character: $\epsilon(vw) = \epsilon(v)\epsilon(w),$ $\epsilon(\mathbf{1}) = 1$	algebra module: $m \cdot vw = (m \cdot v) \cdot w$	peripheral: $\epsilon \pi_w(v_1 \dots v_{n-1} v_n) = v_1 \dots v_{n-1}(v_n w)$
unital Leibniz algebra	Lie character: $\epsilon([v, w]) = 0,$ $\epsilon(\mathbf{1}) = 1$	Leibniz module ([23]): $m \cdot [v, w] = (m \cdot v) \cdot w - (m \cdot w) \cdot v$	adjoint: $\epsilon \pi_w(v_1 \dots v_n) = \sum_{i=1}^n v_1 \dots [v_i, w] \dots v_n$
shelf	shelf character: $\epsilon(a \triangleleft b) = \epsilon(a)$	shelf module ([7], [17]): $(m \cdot a) \cdot b = (m \cdot b) \cdot (a \triangleleft b)$	diagonal: $\epsilon \pi_b(a_1, \dots, a_n) = (a_1 \triangleleft b, \dots, a_n \triangleleft b)$

Table 1: Braided homology ingredients in concrete algebraic settings

Pre-braidings for vector spaces and self-distributive structures are classical; that for Leibniz algebras was used in the Lie case by A.Crans in [8] (cf. also [2]), but does not seem to be widely known; the author has never met the pre-braiding for associative algebras elsewhere.

Our braided homology theory, as well as the pre-braidings for associative and Leibniz algebras, are raised to the *categorical level* in section 5. The *categorification of SD structures* and of the corresponding pre-braiding is less straightforward and is done in [21] (see also [2] for an alternative construction). Several typical applications of the categorical approach are presented, obtained by changing the underlying category (e.g. the Leibniz superalgebra homology) or using different types of categorical dualities (e.g. cobar and Cartier complexes for coalgebras).

An important feature of our categorified pre-braiding, besides relaxing the invertibility condition, is its “*local*” character: instead of demanding the whole category to be pre-braided, we need a pre-braiding for a single object only, omitting in particular the naturality condition.

This paper contains the results from the first part of the author’s thesis [20], where more details and proofs can be found.

Notations and conventions.

We systematically use notation R for a commutative unital ring, and \mathbb{k} for a field. The word “linear” means R - (or \mathbb{k} -) linear, and all tensor products are over R (or \mathbb{k}), unless we work in the settings of a general monoidal category. The category of R -modules (resp. \mathbb{k} -vector spaces) and linear maps is denoted by \mathbf{Mod}_R (resp. $\mathbf{Vect}_{\mathbb{k}}$).

Notation $T(V) := \bigoplus_{n \geq 0} V^{\otimes n}$ is used for the tensor algebra of an R -module V , with $V^{\otimes 0} := R$. A simplified notation is used for its elements:

$$\bar{v} = v_1 v_2 \dots v_n := v_1 \otimes v_2 \otimes \dots \otimes v_n \in V^{\otimes n},$$

leaving the tensor product sign for

$$v_1 v_2 \dots v_n \otimes w_1 w_2 \dots w_m \in V^{\otimes n} \otimes W^{\otimes m}.$$

We often call the R -module $T(V)$ the *tensor module* of V , emphasizing that it can be endowed with a multiplication different from the usual concatenation. We talk about the *tensor space* of V in the \mathbb{k} -linear setting.

The notation $V^{\otimes n}$ is sometimes reduced to V^n , and $\text{Id}_{V^{\otimes n}}$ to Id_n .

Given an R -module V and a linear map $\varphi : V^{\otimes l} \rightarrow V^{\otimes r}$, the following notations are repeatedly used:

$$\varphi_i := \text{Id}_{i-1} \otimes \varphi \otimes \text{Id}_{k-i+1} : V^{k+l} \rightarrow V^{k+r}, \quad (1)$$

$$\varphi^n := (\varphi_1)^{\circ n} = \varphi_1 \circ \dots \circ \varphi_1 : V^k \rightarrow V^{k+n(r-l)}, \quad (2)$$

where φ_1 is composed with itself n times. Similar notations are used for tensor products of different modules.

The above notations are also used (when they make sense) in the context of a strict monoidal category.

By a *differential* on a graded R -module (for example $T(V)$) we mean a square zero endomorphism of degree $+1$ or -1 , while a *bidifferential* is a pair of anticommuting differentials. The word *complex* always means a differential (co)chain complex here, i.e. a graded R -module endowed with a differential. Similarly, a *bicomplex* is a graded R -module endowed with a bidifferential.

2 Braided world: a short reminder

We recall here various facts about *braided vector spaces* necessary for subsequent sections. For a more systematic treatment of braid groups, [18] is an excellent reference. A particular focus is made here on *quantum (co-)shuffles*, introduced and studied by M.Rosso in [35] and [36]. These structures will provide an important tool for constructing braided space homologies in the next section.

All the notions defined here for vector spaces are directly generalized for R -modules. We prefer the language of vector spaces for its familiarity.

Definition 2.1. \rightarrow A *pre-braiding* on a \mathbb{k} -vector space V is a linear map $\sigma : V \otimes V \rightarrow V \otimes V$ satisfying the *Yang-Baxter equation* (abbreviated as YBE)

$$\sigma_1 \circ \sigma_2 \circ \sigma_1 = \sigma_2 \circ \sigma_1 \circ \sigma_2 : V \otimes V \otimes V \longrightarrow V \otimes V \otimes V, \quad (\text{YB})$$

where σ_i is the braiding σ applied to components i and $i + 1$ of $V^{\otimes 3}$ (cf. notation (1)).

- \rightarrow A *braiding* is an invertible pre-braiding.
- \rightarrow A braiding is called *symmetric* if $\sigma^2 = \text{Id}_{V \otimes V}$.
- \rightarrow A vector space endowed with a (pre-)braiding is called *(pre-)braided*.
- \rightarrow A *braided morphism* between pre-braided spaces (V, σ_V) and (W, σ_W) is a \mathbb{k} -linear map $f : V \rightarrow W$ respecting the pre-braidings:

$$(f \otimes f) \circ \sigma_V = \sigma_W \circ (f \otimes f) : V \otimes V \rightarrow W \otimes W.$$

Note that unlike most authors **we work with pre-braidings**, allowing interesting highly non-invertible examples.

Remark 2.2. A (pre-)braiding on a set is defined similarly: tensor products \otimes are simply replaced by Cartesian products \times . These two settings are particular cases of a more abstract one: they both come from **(pre-)braided categories**, studied in detail in section 5.

Example 2.3. The most familiar braidings are the **flip**, the **signed flip** and their generalization for graded vector spaces, the **Koszul flip**:

$$\begin{aligned} \tau : v \otimes w &\longmapsto w \otimes v, \\ -\tau : v \otimes w &\longmapsto -w \otimes v, \\ \tau_{\text{Koszul}} : v \otimes w &\longmapsto (-1)^{\deg v \deg w} w \otimes v \end{aligned} \quad (3)$$

for homogeneous v and w . The last braiding explains the *Koszul sign convention* in many settings.

Remark 2.4. In general for a (pre-)braiding σ , its opposite $-\sigma : v \otimes w \mapsto -\sigma(v \otimes w)$ is also a (pre-)braiding.

A pre-braiding gives an action of the positive braid monoid B_n^+ on $V^{\otimes n}$, i.e. a monoid morphism

$$\begin{aligned} \rho : B_n^+ &\longrightarrow \text{End}_{\mathbb{k}}(V^{\otimes n}), \\ b &\longmapsto b^\sigma \end{aligned} \quad (4)$$

defined on the generators σ_i of B_n^+ by

$$\sigma_i \longmapsto \text{Id}_{i-1} \otimes \sigma \otimes \text{Id}_{n-i-1}. \quad (5)$$

This action is best depicted in the graphical form





Figure 4: B_n^+ acts via pre-braidings

All diagrams in this work are to be read **from bottom to top**, as indicated by the arrow on the diagram above. One could have presented the crossing as , which is often done in literature. It is just a matter of convention, and the one used here comes from rack theory (section 4.2).

For braidings, the action above is in fact an action of the braid group B_n , and for symmetric braidings it is an action of the symmetric group S_n .

The graphical translation of the YBE for pre-braidings is the third Reidemeister move, which is at the heart of knot theory:




Figure 5: Yang-Baxter equation = Reidemeister move III

Numerous constructions become natural in the graphical settings. For instance,

Remark 2.5. A (pre-)braiding σ on V naturally extends to a (pre-)braiding σ on its tensor space $T(V)$ via

$$\sigma(\bar{v} \otimes \bar{w}) = (\sigma_k \cdots \sigma_1) \cdots (\sigma_{n+k-2} \cdots \sigma_{n-1}) (\sigma_{n+k-1} \cdots \sigma_n) (\bar{v}\bar{w}) \in V^{\otimes k} \otimes V^{\otimes n}$$

for pure tensors $\bar{v} \in V^{\otimes n}$, $\bar{w} \in V^{\otimes k}$ ($\bar{v}\bar{w}$ being simply their concatenation), or, graphically,




Figure 6: Pre-braiding extended to $T(V)$

Recall further the famous set inclusion (not preserving the group structure)

$$\begin{aligned} S_n &\hookrightarrow B_n \\ s = \tau_{i_1} \tau_{i_2} \cdots \tau_{i_k} &\mapsto T_s := \sigma_{i_1} \sigma_{i_2} \cdots \sigma_{i_k} \end{aligned} \quad (6)$$

where

- $\tau_i \in S_n$ is the transposition of the neighboring elements i and $i + 1$,
- σ_i is the corresponding generator of B_n ,
- $\tau_{i_1} \tau_{i_2} \cdots \tau_{i_k}$ is one of the shortest words representing $s \in S_n$.

This inclusion factorizes through

$$S_n \hookrightarrow B_n^+ \hookrightarrow B_n.$$

The following subsets of symmetric groups deserve particular attention:

Definition 2.6. *Shuffle sets* are the permutation sets

$$Sh_{p,q} := \left\{ s \in S_{p+q} \text{ s.t. } \left. \begin{array}{l} s(1) < s(2) < \dots < s(p), \\ s(p+1) < s(p+2) < \dots < s(p+q) \end{array} \right\}.$$

In other words, one permutes $p + q$ elements preserving the order within the first p and the last q elements, just like when shuffling cards, which explains the name. Shuffles and their diverse modifications appear, sometimes quite unexpectedly, in various areas of mathematics.

Everything is now ready for defining quantum shuffle algebras.

Definition 2.7. The *quantum shuffle multiplication* on the tensor space $T(V)$ of a pre-braided vector space (V, σ) is the \mathbb{k} -linear extension of the map

$$\begin{aligned} \Delta_\sigma &= \Delta_\sigma^{p,q} : V^{\otimes p} \otimes V^{\otimes q} \longrightarrow V^{\otimes(p+q)}, \\ \bar{v} \otimes \bar{w} &\longmapsto \bar{v} \Delta_\sigma \bar{w} := \sum_{s \in Sh_{p,q}} T_s^\sigma(\bar{v}\bar{w}). \end{aligned} \quad (7)$$

Notation T_s^σ stands for the lift $T_s \in B_n^+$ (cf. (6)) acting on $V^{\otimes n}$ via the pre-braiding σ (cf. (4)).

The algebra $Sh_\sigma(V) := (T(V), \Delta_\sigma)$ is the *quantum shuffle algebra* of (V, σ) .

By a *(pre-)braided Hopf algebra* (in the sense of S.Majid, cf. Definition 2.2 in [28] for example) we mean an additional structure on a (pre-)braided vector space satisfying all the axioms of a Hopf algebra except for the compatibility between the multiplication and the comultiplication, which is replaced by the braided compatibility (this last notion is recalled in the following theorem).

The quantum shuffle multiplication can be upgraded to an interesting pre-braided Hopf algebra structure:

Theorem 1. *Let (V, σ) be a pre-braided vector space.*

1. *The multiplication Δ_σ of $Sh_\sigma(V)$ is associative.*
2. *If $\sigma^2 = \text{Id}$, then the multiplication Δ_σ is σ -commutative, i.e.*

$$\Delta_\sigma(\bar{v} \otimes \bar{w}) = \Delta_\sigma(\sigma(\bar{v} \otimes \bar{w}))$$

(with the extension σ of σ to $T(V)$ from remark 2.5).

3. *The element $1 \in R$ is a unit for $Sh_\sigma(V)$.*
4. *The **deconcatenation** and the **augmentation** maps*

$$\begin{aligned} \Delta : v_1 v_2 \dots v_n &\longmapsto \sum_{p=0}^n v_1 v_2 \dots v_p \otimes v_{p+1} \dots v_n, & \varepsilon : v_1 v_2 \dots v_n &\longmapsto 0, \\ 1 &\longmapsto 1 \otimes 1, & 1 &\longmapsto 1 \end{aligned}$$

(where an empty product means 1) define, after linearizing, a counital coalgebra structure on $T(V)$.

5. *These algebra and coalgebra structures are σ -compatible, in the sense that*

$$\Delta \circ \Delta_\sigma = (\Delta_\sigma \otimes \Delta_\sigma) \circ \sigma_2 \circ (\Delta \otimes \Delta).$$

6. *An antipode can be given on $Sh_\sigma(V)$ by linearizing the map*

$$\begin{aligned} s : \bar{v} &\longmapsto (-1)^n T_{\Delta_n}^\sigma(\bar{v}), & \bar{v} &\in V^{\otimes n}, \\ 1 &\longmapsto 1, \end{aligned}$$

$$\text{where } \Delta_n := \begin{pmatrix} 1 & 2 & \dots & n \\ n & n-1 & \dots & 1 \end{pmatrix} \in S_n. \quad (8)$$

The pre-braided vector space $(T(V), \sigma)$ becomes thus a pre-braided Hopf algebra.

This result is well-known for invertible braidings ([36]); we point out that it still holds when the pre-braiding admits no inverse. See [20] for a proof.

Remark 2.8. Dually (in the sense to be specified in subsection 5.4), the tensor space of a pre-braided vector space (V, σ) can be endowed with the **quantum co-shuffle comultiplication**:

$$\begin{aligned} \Theta_\sigma|_{V^{\otimes n}} &:= \sum_{p+q=n; p,q \geq 0} \Theta_\sigma^{p,q}, \\ \Theta_\sigma^{p,q} &:= \sum_{s \in Sh_{p,q}} T_{s^{-1}}^\sigma : V^{\otimes n} \longrightarrow V^{\otimes p} \otimes V^{\otimes q}, \end{aligned} \quad (9)$$

which can be upgraded to a pre-braided Hopf algebra structure “dual” to that described in theorem 1, denoted by $\overline{Sh}_\sigma(V)$.

3 Homologies of braided vector spaces

In this section we introduce a homology theory of braided vector spaces. We propose two different viewpoints on our “braided” differentials (all the notions and properties are explained in this section):

approach	subsection	advantages
quantum co-shuffle comultiplication and square zero coelements	3.1	<ul style="list-style-type: none"> →the sign manipulation is hidden in the choice of the negative braiding $-\sigma$, →a subscript-free approach, →compact formulas;
graphical calculus: diagrams, braids	3.2	<ul style="list-style-type: none"> →a tool easy to manipulate, →a finer structure of a pre-bisimplicial complex, →an intuitive definition of degenerate and normalized complexes via braided coalgebras.

Table 2: Two approaches to “braided” differentials

We also study certain homology operations for “braided” homologies, generalizing those introduced by M.Niebrzydowski and J.Przytycki in the context homologies of self-distributive structures ([30], [31]), and we obtain a new interpretation and a generalization of J.-L.Loday’s hyperboundaries, automatically calculating all their compositions (cf. [23], exercise E.2.2.7).

Everything described here can be translated verbatim to the setting of R -modules. We prefer working with vector spaces for the sake of clarity. A categorical version of all the results is given in section 5.

Fix a pre-braided \mathbb{k} -vector space (V, σ) .

3.1 Pre-braiding + character \mapsto homology

We start with distinguishing elements of V^* which behave with respect to the pre-braiding σ as if it were just a flip τ :

Definition 3.1. Two co-elements $f, g \in V^*$ are called σ -compatible if

$$(f \otimes g) \circ \sigma = g \otimes f, \quad \text{and} \quad (g \otimes f) \circ \sigma = f \otimes g. \quad (10)$$

A *braided character* is an $\epsilon \in V^*$ σ -compatible with itself, i.e.

$$(\epsilon \otimes \epsilon) \circ \sigma = \epsilon \otimes \epsilon, \quad (11)$$

or, in the co-shuffle form,

$$(\epsilon \otimes \epsilon) \circ \Theta_{-\sigma}^{1,1} = 0 : V \otimes V \rightarrow \mathbb{k}. \quad (12)$$

We omit the part *braided* of the last term when it does not lead to confusion.

The definition of braided character takes a simple graphical form:




Figure 7: Braided character

The labels ϵ are often omitted when clear from the context.

In section 4, we recover familiar notions of characters for algebraic structures (such as associative algebras) as examples of braided characters for the corresponding “structural” braidings, which justifies our term. Counits often turn out to be braided characters, hence the notation ϵ .

Remark 3.2. Braided characters can also be regarded as *braided morphisms* $\epsilon : V \rightarrow \mathbb{k}$, where \mathbb{k} is endowed with the *trivial braiding* $\mathbb{k} \otimes \mathbb{k} \simeq \mathbb{k} \xrightarrow{\text{Id}_{\mathbb{k}}} \mathbb{k} \simeq \mathbb{k} \otimes \mathbb{k}$. This is consistent with the interpretation of usual characters for algebraic structures as homomorphisms to trivial structures.

A pre-braiding and a braided character are sufficient for constructing homologies:

Theorem 2. *Let (V, σ) be a pre-braided vector space.*

1. *For a braided character ϵ , the maps*

$$\begin{aligned} V^{\otimes n} &\longrightarrow V^{\otimes(n-1)}, \\ \epsilon d : \bar{v} &\longmapsto \epsilon_1 \circ \Theta_{-\sigma}^{1,n-1}(\bar{v}), \\ d^\epsilon : \bar{v} &\longmapsto (-1)^{n-1} \epsilon_n \circ \Theta_{-\sigma}^{n-1,1}(\bar{v}) \end{aligned}$$

(cf. notation (1)) *define a bidifferential on $T(V)$.*

2. *For two braided characters ϵ and ζ , one gets a differential bicomplex $(T(V), \epsilon d, d^\zeta)$. If the braided characters are moreover σ -compatible, then one gets differential bicomplexes $(T(V), \epsilon d, \zeta d)$ and $(T(V), d^\epsilon, d^\zeta)$.*

Proof. The verifications use the coassociativity of $\Theta_{-\sigma}$ and the defining property (12) of braided characters. For example,

$$\begin{aligned} \epsilon d \circ \epsilon d(\bar{v}) &= \epsilon_1 \circ (\epsilon \otimes \Theta_{-\sigma}^{1,n-2}) \circ \Theta_{-\sigma}^{1,n-1}(\bar{v}) \\ &= ((\epsilon \otimes \epsilon) \circ \Theta_{-\sigma}^{1,1})_1 \circ \Theta_{-\sigma}^{2,n-2}(\bar{v}) \\ &= 0 \circ \Theta_{-\sigma}^{2,n-2}(\bar{v}) = 0, \end{aligned}$$

and, for compatible braided characters,

$$\begin{aligned} (\zeta d \circ \epsilon d + \epsilon d \circ \zeta d)(\bar{v}) &= \zeta_1 \circ (\epsilon \otimes \Theta_{-\sigma}^{1,n-2}) \circ \Theta_{-\sigma}^{1,n-1}(\bar{v}) + \epsilon_1 \circ (\zeta \otimes \Theta_{-\sigma}^{1,n-2}) \circ \Theta_{-\sigma}^{1,n-1}(\bar{v}) \\ &= ((\zeta \otimes \epsilon + \epsilon \otimes \zeta) \circ \Theta_{-\sigma}^{1,1})_1 \circ \Theta_{-\sigma}^{2,n-2}(\bar{v}), \end{aligned}$$

which, due to (10), equals

$$((\zeta \otimes \epsilon - \epsilon \otimes \zeta) + (\epsilon \otimes \zeta - \zeta \otimes \epsilon))_1 \circ \Theta_{-\sigma}^{2,n-2}(\bar{v}) = 0. \quad \square$$

This proof can be understood as follows: the multiplication by a square zero element in the pre-braided Hopf algebra dual to $\overline{\text{Sh}}_{-\sigma}$ is a square zero operator.

The theorem gives for two braidings two compatible differentials on $T(V)$. Their linear combinations are then also differentials; $\epsilon d - d^\epsilon$ is a recurrent example in practice. All such (bi)differentials, corresponding (bi)complexes and homologies are called **braided** in what follows.

We now give a survey of existing “braided” homologies, comparing them to our theory.

1. In [4], J.S.Carter, M.Elhamdadi and M.Saito develop a homology theory for solutions (S, σ) of the set-theoretic YBE using combinatorial and geometric methods completely different from ours. They also provide applications to virtual knot invariants. Their differential on $(\mathbb{Z}S)^{\otimes n}$ is our braided differential $\epsilon d - d^\epsilon$, where ϵ is the linearization of the map $\epsilon : S \rightarrow \mathbb{Z}$ assigning 1 to any $a \in S$.
2. Our applications ϵd , where $\epsilon \in V^*$ are not necessarily braided characters, also recover the *braided-differential calculus* of S.Majid ([27]). He defines an addition law (related to the quantum co-shuffle comultiplication) on the quantum plane associated to a braiding, and defines a differentiation as an infinitesimal translation. In particular, taking

- one-variable polynomials $T(V) = \mathbb{k}[x]$ (i.e. $V = \mathbb{k}x$),
- the opposite of the q -flip $x \otimes x \mapsto qx \otimes x$ (with $q \in \mathbb{k}^*$) as a braiding,
- and the linearization of the map $\epsilon(x) = 1$,

one gets the famous q -differentials

$$\varepsilon d(x^{\otimes n}) = (n)_q x^{\otimes(n-1)}, \quad (n)_q := \frac{q^n - 1}{q - 1} = q^{n-1} + \dots + q + 1.$$

3. The last approach to “braided” cohomologies to be mentioned here is M.Eisermann’s *Yang-Baxter cochain complex*, cf. [12]. Motivated by the study of deformations of Yang-Baxter operators, he defines a degree 1 differential on $\text{Hom}_{\mathbb{k}}(V^{\otimes n}, V^{\otimes n})$. His second cohomology groups classify infinitesimal Yang-Baxter deformations. We do not know precisely how his construction is related to ours, but the parallels between the graphical versions of the two are very suggestive.

3.2 Comultiplication \mapsto degeneracies

In a more detailed study of the structure of braided (bi)complexes from theorem 2, the simplicial approach proves to be particularly helpful.

First, recall the notion of simplicial vector spaces (cf. [23] for details and [31] for weak simplicial notions; note that our definition is a shifted version of theirs, and that our definition of bisimplicial vector spaces is different from the one in [23]):

Definition 3.3. Consider a collection of \mathbb{k} -vector spaces V_n , $n \geq 0$, equipped with linear maps $d_{n;i} : V_n \rightarrow V_{n-1}$ (and $d'_{n;i} : V_n \rightarrow V_{n-1}$ and/or $s_{n;i} : V_n \rightarrow V_{n+1}$ when necessary) with $1 \leq i \leq n$, denoted simply by d_i, d'_i, s_i when the subscript n is clear from the context. This datum is called

\rightarrow a *presimplicial vector space* if

$$d_i d_j = d_{j-1} d_i \quad \forall 1 \leq i < j \leq n; \quad (13)$$

\rightarrow a *very weakly simplicial vector space* if moreover

$$s_i s_j = s_{j+1} s_i \quad \forall 1 \leq i \leq j \leq n, \quad (14)$$

$$d_i s_j = s_{j-1} d_i \quad \forall 1 \leq i < j \leq n, \quad (15)$$

$$d_i s_j = s_j d_{i-1} \quad \forall 1 \leq j + 1 < i \leq n; \quad (16)$$

\rightarrow a *weakly simplicial vector space* if moreover

$$d_i s_i = d_{i+1} s_i \quad \forall 1 \leq i \leq n; \quad (17)$$

\rightarrow a *simplicial vector space* if moreover

$$d_i s_i = \text{Id}_{V_n} \quad \forall 1 \leq i \leq n; \quad (18)$$

\rightarrow a *pre-bisimplicial vector space* if (13) holds for the d_i ’s, the d'_i ’s and their mixture:

$$d_i d'_j = d'_{j-1} d_i, \quad d'_i d_j = d_{j-1} d'_i \quad \forall 1 \leq i < j \leq n; \quad (19)$$

\rightarrow a *(weakly / very weakly) bisimplicial vector space* if it is pre-bisimplicial, with (weakly / very weakly) simplicial structures $(V_n, d_{n;i}, s_{n;i})$ and $(V_n, d'_{n;i}, s_{n;i})$.

The omitted subscripts $n, n \pm 1$ are those which guarantee that the source of all the above mentioned morphisms is V_n . The d_i ’s and the s_i ’s are called *face* (resp. *degeneracy*) maps.

Simplicial vector spaces are interesting because of the following properties (see [23] for most proofs):

Proposition 3.4. 1. For any presimplicial vector space $(V_n, d_{n;i})$, the map $\partial_n := \sum_{i=1}^n (-1)^{i-1} d_{n;i}$ is a differential (called the **total differential**) for the graded vector space $\tilde{V} := \bigoplus_{n \geq 0} V_n$.

2. For any pre-bisimplicial vector space $(V_n, d_{n;i}, d'_{n;i})$, there is a bidifferential structure on \tilde{V} given by ∂_n and $\partial'_n := \sum_{i=1}^n (-1)^{i-1} d'_{n;i}$.

3. For any weakly simplicial vector space $(V_n, d_{n;i}, s_{n;i})$, the complex (V_n, ∂_n) contains a sub-complex (called **the degenerate subcomplex**) $D_n := \sum_{i=1}^{n-1} s_{n-1;i}(V_{n-1})$.
4. If our vector space turns out to be simplicial, then the degenerate subcomplex is acyclic, hence V_* is quasi-isomorphic to the **normalized complex** $N_* := V_*/D_*$.
5. In the weakly bisimplicial case, D_* is a sub-bicomplex of V_* , acyclic in the bisimplicial setting.

We will soon show that the bicomplexes from theorem 2 come from pre-bisimplicial structures. As for degeneracies, they arise from the following structure:

Definition 3.5. A pre-braided vector space (V, σ) endowed with a comultiplication $\Delta : V \rightarrow V \otimes V$ is called a *pre-braided coalgebra* if

→ Δ is *co-associative*:

$$(\Delta \otimes \text{Id}_V) \circ \Delta = (\text{Id}_V \otimes \Delta) \circ \Delta : V \rightarrow V \otimes V \otimes V, \quad (20)$$

→ and Δ is *compatible* with the pre-braiding:

$$\Delta_2 \circ \sigma = \sigma_1 \circ \sigma_2 \circ \Delta_1 : V^{\otimes 2} \rightarrow V^{\otimes 3}, \quad (21)$$

$$\Delta_1 \circ \sigma = \sigma_2 \circ \sigma_1 \circ \Delta_2 : V^{\otimes 2} \rightarrow V^{\otimes 3}. \quad (22)$$

One talks about *semi-pre-braided coalgebras* if only (21) holds.

A (semi-)pre-braided coalgebra is called *σ -cocommutative* if

$$\sigma \circ \Delta = \Delta : V \rightarrow V \otimes V. \quad (23)$$

The properties from the definition are graphically depicted as





Figure 8: Coassociativity and σ -cocommutativity



Figure 9: Braided coalgebra

Here the comultiplication Δ is represented as .

Theorem 2 admits now the following simplicial interpretation:

Theorem 3. Let (V, σ) be a pre-braided vector space.

1. For braided characters ϵ and ζ , the maps

$$d_{n;i}(\bar{v}) := \epsilon_1 \circ T_{p_{i,n}}^\sigma(\bar{v}), \quad (24)$$

$$d'_{n;i}(\bar{v}) := \zeta_n \circ T_{p'_{i,n}}^\sigma(\bar{v}), \quad (25)$$

define a pre-bisimplicial structure on the tensor vector space $T(V)$. Here $p_{i,n} \in S_n$ (resp. $p'_{i,n} \in S_n$) is the permutation moving the i th element to the leftmost (resp. rightmost) position, and the notation T_s^σ from (6) and (4) is used.

2. The total differentials ∂ and ∂' coincide with the “shuffle” differentials ϵd and, respectively, d^ζ from theorem 2.
3. If the braided characters are moreover σ -compatible, then the d_i 's for ϵ and the d_i 's for ζ define a pre-bisimplicial structure on $T(V)$.

4. If for a comultiplication Δ the triple (V, σ, Δ) is a pre-braided coalgebra, then the maps

$$s_{n;i} := \Delta_i \tag{26}$$

complete the preceding structures into very weakly bisimplicial ones.

5. If (V, σ, Δ) is a semi-pre-braided coalgebra, then the data $(V^{\otimes n}, d_{n;i}, s_{n;i})$ described above give a very weakly simplicial vector space only.

6. If Δ is moreover σ -cocommutative, then the above structures are weakly (bi)simplicial.

The face and degeneracy maps from the theorem are graphically depicted as




Figure 10: Simplicial structures

Proof. One has to deduce the “simplicial” relations (definition 3.3) from the properties of the structures on V , which were conceived precisely for these relations to hold. This can be done graphically, using the pictorial interpretation of face and degeneracy maps, and the graphical definitions of a (σ -cocommutative) pre-braided coalgebra presented above, as well as the pictorial versions of the YBE (fig. 5) and of the definition of braided characters (fig. 7).

For instance,




Figure 11: Graphical proof of $d_i d_j = d_{j-1} d_i \forall 1 \leq i < j \leq n$.

Here

(1) is a repeated application of YBE;

(2) follows from the definition (11) of a braided character (cf. fig. 7). \square

Remark 3.6. When checking the axioms of different types of simplicial structures in the proof, one can get rid of the tiresome index chasing by reasoning in terms of strands. For example, pulling a strand to the left commutes with applying the branching Δ to any other strand if a strand can pass over a branching.

3.3 Arrow and concatenation operations

The last face map $d_{n+1;n+1}$ on $V^{\otimes(n+1)}$ is of particular interest. It inspires the definition of a useful operation on $T(V)$:

Definition 3.7. Take a braided character ϵ on (V, σ) . For a $w \in V$, we call an *arrow operation* on $T(V)$ the map defined by

$$\epsilon_{\pi_w}(\bar{v}) := d_{n+1;n+1}(\bar{v}w) = \epsilon_1 \circ T_{p_{n+1}, n+1}^\sigma(\bar{v}w), \quad \forall \bar{v} \in V^{\otimes n}.$$

The notation and the name come from the graphical presentation:




Figure 12: Arrow operation

This map will be interpreted in terms of modules over pre-braided vector spaces and adjoint maps in proposition 6.4. Here we study its properties, and get a generalization of some homology operations from [30] and [31]. Our constructions are deeply inspired by those papers.

Start with some technical definitions:

Definition 3.8. Take a pre-braided vector space (V, σ) .

→ A *normalized pair* is an element $w \in V$ and a co-element $\psi \in V^*$ satisfying $\psi(w) = 1$.

→ A $w \in V$ and a $\psi \in V^*$ are called *right σ -compatible* if

$$(\text{Id}_V \otimes \psi) \circ \sigma \circ (v \otimes w) = \psi(v)w \quad \forall v \in V. \quad (27)$$

→ The pre-braiding σ is called *natural* with respect to a $w \in V$ if

$$\sigma \circ (w \otimes v) = v \otimes w \quad \forall v \in V, \quad (28)$$

$$\sigma \circ (v \otimes w) = w \otimes v \quad \forall v \in V, \quad (29)$$

and *semi-natural* (or *demi-natural*) if only (28) (resp. (29)) holds.

The properties from the definition graphically mean




Figure 13: Right σ -compatibility and naturality with respect to an element

The naturality can be interpreted as follows: the element w can “pass through a crossing” to the left / to the right.

Note that condition (29) implies (27) for any ψ .

The compatibility of arrow operations ϵ_{π_w} with the braided differential ϵd from theorem 2 is a consequence of theorem 3, where we interpret ϵd as a total differential for which $\epsilon \pi$ is the last face map. This is an inspiring remark for the following analysis of the behavior of our braided differentials with respect to arrow operations and **concatenation operations** $\bar{v} \mapsto \bar{v}w$ on $T(V)$, for a fixed $w \in V$.

Proposition 3.9. Let (V, σ) be a pre-braided vector space with braided characters ϵ, ξ, ζ , the first two being σ -compatible, and the last one being right σ -compatible with an element $w \in V$.

1. The map ϵ_{π_w} is a bicomplex map for $(T(V), \xi d, d^\zeta)$, i.e.

$$\xi d \circ \epsilon_{\pi_w}(\bar{v}) = \epsilon_{\pi_w} \circ \xi d(\bar{v}),$$

$$d^\zeta \circ \epsilon_{\pi_w}(\bar{v}) = \epsilon_{\pi_w} \circ d^\zeta(\bar{v}).$$

2. The following relations hold between the concatenation operations and the braided differentials:

$$\epsilon d(\bar{v}w) = \epsilon d(\bar{v})w + (-1)^n \epsilon_{\pi_w}(\bar{v}),$$

$$d^\zeta(\bar{v}w) = d^\zeta(\bar{v})w + (-1)^n \zeta(w)\bar{v}.$$

3. If σ is demi-natural with respect to w , then

$$\epsilon\pi_w = \epsilon(w) \text{Id}_{T(V)}.$$

Here the notation \bar{v} stays for any pure tensor in $V^{\otimes n}$.

This result admits an evident “left” version (with respect to w).

Proof. Point 1 follows, in the same way as the proof of theorem 3, from the YBE for σ and from the σ -compatibilities (use for instance the graphical calculus).

Point 2 can be checked using the pre-braided Hopf algebra structure on $\overline{\text{Sh}}_{-\sigma}(V)$. For instance, for the left differentials one has

$$\begin{aligned} \epsilon d(\bar{v}w) &= (\epsilon \otimes \text{Id}_n) \circ \Theta_{-\sigma}^{1,n}(\bar{v}w) \\ &\stackrel{(*)}{=} (\epsilon \otimes \text{Id}_n)(\Theta_{-\sigma}^{1,n-1}(\bar{v})w + T_{p_{n+1},n+1}^{-\sigma}(\bar{v}w)) \\ &= (\epsilon \otimes \text{Id}_n)(\Theta_{-\sigma}^{1,n-1}(\bar{v})w) + (-1)^n(\epsilon \otimes \text{Id}_n) \circ T_{p_{n+1},n+1}^{\sigma}(\bar{v}w) \\ &= \epsilon d(\bar{v})w + (-1)^n \epsilon\pi_w(\bar{v}). \end{aligned}$$

Equality (*) is the compatibility between the multiplication and the comultiplication in $\overline{\text{Sh}}_{-\sigma}(V)$.

Point 3 is straightforward. \square

Corollary 3.10. 1. In the settings of the previous proposition, the arrow operation $\epsilon\pi_w$ is homotopic to zero on the complex $(T(V), \epsilon d)$, and to $\zeta(w) \text{Id}_{T(V)}$ on $(T(V), \epsilon d - d^\zeta)$.

2. The complex $(T(V), \epsilon d)$ is acyclic if σ is demi-natural with respect to w and the pair (w, ϵ) is normalized.

3. The complex $(T(V), d^\zeta)$ is acyclic if the pair (w, ζ) is normalized.

Proof. The concatenation map $\bar{v} \mapsto \bar{v}w$ gives the demanded homotopies. \square

3.4 Loday’s hyper-boundaries

Our quantum shuffle setting provides a natural interpretation for J.-L.Loday’s hyper-boundaries (see [23], exercise E.2.2.7), which we redefine as generalizations of the “shuffle” differentials from theorem 2.

Definition 3.11. Let (V, σ) be a pre-braided vector space with a braided character ϵ . The maps

$$\begin{aligned} V^{\otimes n} &\longrightarrow V^{\otimes(n-k)}, \\ \epsilon^{(k)}d : \bar{v} &\longmapsto (\epsilon^{\otimes k} \otimes \text{Id}_{n-k}) \circ \Theta_{-\sigma}^{k,n-k}(\bar{v}), \\ d^{\epsilon^{(k)}} : \bar{v} &\longmapsto (-1)^{kn - \frac{k(k+1)}{2}} (\text{Id}_{n-k} \otimes \epsilon^{\otimes k}) \circ \Theta_{-\sigma}^{n-k,k}(\bar{v}) \end{aligned}$$

are called *hyper-boundaries* on $T(V)$.

The last sign should be understood as $(-1)^{n-1}(-1)^{n-2} \dots (-1)^{n-k}$.

For $k = 1$ one recovers the braided differentials ϵd and d^ϵ .

The next step is to understand compositions of hyper-boundaries, generalizing

$$d^{(1)} \circ d^{(1)} = 0 = {}^{(1)}d \circ {}^{(1)}d.$$

We start with a kind of a special case. This result seems to be well-known, but a written proof is difficult to find in literature. See for example [20].

Lemma 3.12. Consider a vector space W and an element $w \in W$. One has

$$w^{\otimes m} \underset{-\tau}{\Delta} w^{\otimes k} = \binom{m+k}{k}_{-1} w^{\otimes(m+k)}, \quad \binom{m+k}{k}_{-1} := \begin{cases} 0 & \text{if } mk \text{ is odd,} \\ \binom{\lfloor (m+k)/2 \rfloor}{\lfloor k/2 \rfloor} & \text{otherwise.} \end{cases}$$

The brackets $[\cdot]$ stand here for the lower integral part of a number.

This lemma is crucial in the calculations giving

Theorem 4. Let (V, σ) be a pre-braided vector space with a braided character ϵ . One has

$$\begin{aligned} \epsilon, (m)d \circ \epsilon, (k)d &= \binom{m+k}{k}_{-1} \epsilon, (m+k)d, \\ d^{\epsilon, (m)} \circ d^{\epsilon, (k)} &= \binom{m+k}{k}_{-1} d^{\epsilon, (m+k)}. \end{aligned}$$

Proof. We prove the first formula only. By definition,

$$\epsilon, (m)d \circ \epsilon, (k)d(\bar{v}) = (\epsilon \otimes \cdots \otimes \epsilon \otimes \text{Id}_{n-k-m}) \circ (\epsilon \otimes \cdots \otimes \epsilon \otimes \underset{-\sigma}{\Theta}^{m, n-k-m}) \circ \underset{-\sigma}{\Theta}^{k, n-k}(\bar{v}).$$

By the coassociativity of the quantum co-shuffle comultiplication, it equals

$$(\epsilon \otimes \cdots \otimes \epsilon \otimes \text{Id}_{n-k-m}) \circ (\underset{-\sigma}{\Theta}^{k, m} \otimes \text{Id}_{n-k-m}) \circ \underset{-\sigma}{\Theta}^{m+k, n-m-k}(\bar{v}).$$

Now ϵ is a braided character, so

$$(\epsilon \otimes \epsilon) \circ \sigma = \epsilon \otimes \epsilon = (\epsilon \otimes \epsilon) \circ \tau,$$

thus

$$\epsilon, (m)d \circ \epsilon, (k)d(\bar{v}) = (\epsilon \otimes \cdots \otimes \epsilon \otimes \text{Id}_{n-k-m}) \circ (\underset{-\tau}{\Theta}^{k, m} \otimes \text{Id}_{n-k-m}) \circ \underset{-\sigma}{\Theta}^{m+k, n-m-k}(\bar{v}).$$

The dual version of the previous lemma calculates

$$(\epsilon \otimes \cdots \otimes \epsilon) \circ \underset{-\tau}{\Theta}^{k, m} = \binom{m+k}{k}_{-1} \epsilon \otimes \cdots \otimes \epsilon,$$

and the previous expression becomes $\binom{m+k}{k}_{-1} \epsilon, (m+k)d$. \square

The relations from J.-L.Loday's exercise, which are particular cases of the above theorem for several values of m and k , are thus easily proved and generalized thanks to our quantum co-shuffle interpretation.

4 Basic examples: familiar complexes recovered

Now we consider a \mathbb{k} -vector space (or an R -module) V with some algebraic structure, and we look for a **pre-braiding σ encoding the properties of this structure**. Such pre-braidings are informally called *structural*. Certain algebraic properties of the initial structure are coded by the **invertibility** condition for the corresponding pre-braiding. In each case, **braided characters** are determined, always up to scalar multiples, recovering the usual algebraic notions of characters. Theorem 2 then gives numerous bicomplex structures on $T(V)$. We calculate explicitly some of the differentials obtained this way, recognizing many familiar complexes. **Arrow operations** are also considered, showing the triviality of some of the appearing homologies (cf. corollary 3.10). In some cases, V is endowed with a (semi-)pre-braided coalgebra structure, giving, according to theorem 3, a (very) weakly bisimplicial structure on $T(V)$. The **comultiplications** Δ we use always arise naturally from the original algebraic structure.

A typical subsection of this section contains thus several lemmas, one for each question emphasized above, followed by propositions explicitly describing the bidifferential or simplicial structures obtained for "interesting" characters. Graphical calculus is extensively used.

4.1 Vector spaces

Following a nice mathematical tradition, the first example we consider is the trivial one: that of an “empty” structure. Take any vector space V and the flip

$$\tau : v \otimes w \longmapsto w \otimes v$$

as its braiding. Each $\epsilon \in V^*$ is automatically a braided character. In particular,

$$\epsilon d = d^\epsilon : v_1 \dots v_n \longmapsto \sum_{i=1}^n (-1)^{i-1} \epsilon(v_i) v_1 \dots \widehat{v}_i \dots v_n$$

gives the well-known **Koszul differential**, in its simplest form.

Further, a (τ -cocommutative) braided coalgebra structure on (V, τ) is precisely a (cocommutative) comultiplication in the usual sense. The corresponding very weakly simplicial structure on $T(V)$ is simplicial if and only if ϵ is the *counit* for the comultiplication Δ . In the last case the cocommutativity is not necessary for the structure to be simplicial. Thus, according to theorem 3, one can quotient the Koszul complex by the images of $s_{n;i} := \Delta_i$ without changing the homology.

4.2 Self-distributive structures

The simplest non-trivial example of a pre-braiding naturally coming from an algebraic structure is the following. Take a set S with a binary operation $\triangleleft : S \times S \rightarrow S$. Define an application

$$\begin{aligned} \sigma = \sigma_{\triangleleft} : S \times S &\longrightarrow S \times S, \\ (a, b) &\longmapsto (b, a \triangleleft b). \end{aligned} \tag{30}$$

It is very familiar to topologists, since it can be interpreted in terms of the fundamental group of the complement of a knot. See for instance the seminal paper [15], or [16] for a very readable introduction. Graphically σ_{\triangleleft} becomes

$$\begin{array}{ccc} b & a \triangleleft b & \\ & \times & \\ a & b & . \end{array}$$

Figure 14: Pre-braiding for shelves

All the “braided” notions are to be understood in the **set-theoretic sense** in this subsection (cf. remark 2.2).

The structure for which σ_{\triangleleft} is a pre-braiding is well-known (see for instance [8] or [20]):

Lemma 4.1. The map σ_{\triangleleft} is a pre-braiding if and only if \triangleleft is **self-distributive**:

$$(a \triangleleft b) \triangleleft c = (a \triangleleft c) \triangleleft (b \triangleleft c) \quad \forall a, b, c \in S. \tag{SD}$$

Definition 4.2. A pair (S, \triangleleft) satisfying (SD) is called a *shelf*, or a *self-distributive system*.

Lemma 4.3. A map f between two shelves (S_1, \triangleleft_1) and (S_2, \triangleleft_2) is a shelf morphism (i.e. $f(a \triangleleft_1 b) = f(a) \triangleleft_2 f(b)$) if and only if it is a braided morphism from $(S_1, \sigma_{\triangleleft_1})$ to $(S_2, \sigma_{\triangleleft_2})$.

These “if and only if” lemmas show that the pre-braiding σ_{\triangleleft} **encodes** the defining property of a shelf, just as we wanted.

Fix a shelf (S, \triangleleft) until the end of this section.

Lemma 4.4. The pre-braiding σ_{\triangleleft} is a braiding if and only if the application $a \mapsto a \triangleleft b$ is a bijection on S for every $b \in S$, that is if there exists an application $\widetilde{\triangleleft} : S \times S \rightarrow S$ such that

$$(a \triangleleft b) \widetilde{\triangleleft} b = (a \widetilde{\triangleleft} b) \triangleleft b = a \quad \forall a, b \in S. \tag{R}$$

Definition 4.5. A triple $(S, \triangleleft, \widetilde{\triangleleft})$ satisfying (SD) and (R) is called a *rack*.

Now *linearize* a shelf (S, \triangleleft) : put $V := \mathbb{k}S$, where \mathbb{k} is a field, and extend the braiding σ_{\triangleleft} to V linearly. One gets a pre-braided vector space $(V, \sigma_{\triangleleft})$.

Lemma 4.6. Characters $\epsilon \in V^*$ are characterized by $\epsilon(a \triangleleft b) = \epsilon(a)$ for all $a, b \in S$ with $\epsilon(b) \neq 0$.

In the R -linear setting, this condition is sufficient but not necessary in general. Here are some examples of braided characters:

Example 4.7. 1. The linearization of

$$\epsilon : a \mapsto 1 \quad \forall a \in S \quad (31)$$

is always a character.

2. The linearization of a “*Dirac map*”

$$\varphi_a(b) := \delta_{a,b} = \begin{cases} 1 & \text{if } b = a, \\ 0 & \text{for other } b \in S \end{cases} \quad (32)$$

(here $\delta_{a,b}$ is the Kronecker delta) is a character if and only if a is idempotent and satisfies $b \triangleleft a \neq a$ for $b \neq a$. In particular, if S is a *quandle*, i.e. a rack with idempotent elements:

$$a \triangleleft a = a \quad \forall a \in S, \quad (\text{Q})$$

then all the φ_a ’s are characters.

We finish with a more conceptual construction of a class of braided characters. Recall that a character for an algebraic structure is usually defined as a morphism to the trivial structure. It is natural (having in mind the conjugation quandle) to define the *trivial shelf structure* on a set X by $x \triangleleft y = x$ for all $x, y \in X$.

Definition 4.8. A *shelf character* for a shelf (S, \triangleleft) is a shelf morphism $\epsilon : S \rightarrow \mathbb{k}$, where \mathbb{k} is endowed with the trivial shelf structure. In other words, ϵ satisfies

$$\epsilon(a \triangleleft b) = \epsilon(a) \quad \forall a, b \in S.$$

Lemma 4.6 then implies

Lemma 4.9. The linearization of a shelf character is always a braided character for the pre-braiding σ_{\triangleleft} . Moreover, two braided characters coming from shelf characters are automatically σ_{\triangleleft} -compatible.

The last ingredient we need is a comultiplication. The one proposed here is quite classical in the self-distributive world:

Lemma 4.10. Let $\Delta_D : V \rightarrow V \otimes V$ be the linearization of the *diagonal map*

$$\begin{aligned} D : S &\longrightarrow S \times S, \\ a &\longmapsto (a, a) \quad \forall a \in S. \end{aligned}$$

1. $(V, \sigma_{\triangleleft}, \Delta_D)$ is a semi-pre-braided coalgebra.
2. This coalgebra is pre-braided if and only if

$$a \triangleleft b = (a \triangleleft b) \triangleleft b \quad \forall a, b \in S.$$

3. The σ_{\triangleleft} -cocommutativity for Δ_D is equivalent to (Q).

Definition 4.11. A shelf satisfying (Q) is called a *spindle*.

Remark 4.12. The image of the map $s_{n,i} := (\Delta_D)_i$ is the linear span of the elements $(a_1, \dots, a_{n+1}) \in S^{\times(n+1)}$ with $a_i = a_{i+1}$.

It is now time to put together all the ingredients and to make some concrete calculations of bidifferentials. Start with the character ε defined by (31).

Proposition 4.13. *Take a shelf (S, \triangleleft) .*

1. A bicomplex structure can be defined on $T(\mathbb{k}S)$ by

$$\begin{aligned}\varepsilon d(a_1, \dots, a_n) &= \sum_{i=1}^n (-1)^{i-1} ((a_1 \triangleleft a_i), \dots, (a_{i-1} \triangleleft a_i), a_{i+1}, \dots, a_n), \\ d^\varepsilon(a_1, \dots, a_n) &= \sum_{i=1}^n (-1)^{i-1} (a_1, \dots, \widehat{a}_i, \dots, a_n).\end{aligned}$$

2. This bidifferential comes from a pre-bisimplicial structure given by

$$\begin{aligned}d_{n;i}(a_1, \dots, a_n) &= ((a_1 \triangleleft a_i), \dots, (a_{i-1} \triangleleft a_i), a_{i+1}, \dots, a_n), \\ d'_{n;i}(a_1, \dots, a_n) &= (a_1, \dots, \widehat{a}_i, \dots, a_n).\end{aligned}$$

3. If our shelf is moreover a spindle, then $(T(\mathbb{k}S), d_{n;i}, d'_{n;i}, s_{n;i} := (\Delta_D)_i)$ is a weakly bisimplicial vector space. As a consequence, the linear span $C_*^D(S)$ of the elements $(a_1, \dots, a_n) \in S^{\times n}$ with $a_i = a_{i+1}$ for one of the subscripts i forms a sub-bicomplex of $(T(\mathbb{k}S), \varepsilon d, d^\varepsilon)$.

Proof. Points 1 and 2 are direct applications of theorems 2 and 3 respectively to the pre-braiding σ_\triangleleft from (30) and the character ε , combined with the lemmas from this section.

As for point 3, theorem 3 gives only a half of this assertion: $(T(\mathbb{k}S), d_{n;i}, s_{n;i})$ is a weakly simplicial vector space, hence $C_*^D(S)$ is a subcomplex of $(T(\mathbb{k}S), \partial = \varepsilon d)$. Since $(\mathbb{k}S, \sigma_\triangleleft, \Delta_D)$ is only a semi-pre-braided coalgebra in general, the compatibilities (15) between the $d'_{n;i}$'s and the $s_{n;i}$'s should be verified by hand, which is an easy exercise. Finally, the explicit description of the degenerate sub-bicomplex follows from remark 4.12. \square

Let us point out familiar complexes recovered in this proposition. We freely replace the field \mathbb{k} by \mathbb{Z} since, as noted above, all the constructions here work in the R -linear setting.

Example 4.14. 1. The *rack homology* ([13]) appears as

$$C_*^R(S) := (T(\mathbb{Z}S), \varepsilon d - d^\varepsilon).$$

2. The *shelf, or one-term distributive, homology* ([31], [32]) appears as

$$C_*^\triangleleft(S) := (T(\mathbb{Z}S), \varepsilon d).$$

3. The *quandle homology* ([5]) is the quotient

$$C_*^Q(S) := C_*^R(S) / C_*^D(S).$$

The map $\varepsilon\pi$ takes the familiar *diagonal* form here:

$$\varepsilon\pi_b(a_1, \dots, a_n) = (a_1 \triangleleft b, \dots, a_n \triangleleft b).$$

Moreover, ε is right σ_\triangleleft -compatible and forms a normalized pair with any $b \in S$. Proposition 3.9 and corollary 3.10 are then applicable, recovering some results on homology operations from [30], [31] and [32] and their consequences:

Proposition 4.15. 1. *The complex $(T(\mathbb{k}S), d^\varepsilon)$ is acyclic.*

2. *If there exists an element $b \in S$ such that the application $a \mapsto a \triangleleft b$ is a bijection on S , then the complex $(T(\mathbb{k}S), \varepsilon d)$ is acyclic.*

3. If there exists an $a \in S$ stable by all the inner shelf morphisms, i.e.

$$a \triangleleft b = a \quad \forall b \in S, \quad (33)$$

then the complex $(T(\mathbb{k}S), \varepsilon d)$ is acyclic.

Proof. Point 1 follows from corollary 3.10 (point 3).

Point 2 follows from corollary 3.10 (point 1), since the arrow operation $\varepsilon\pi_b$, shown there to be homotopic to zero, is now invertible, the inverse given by $(a_1, \dots, a_n) \mapsto (a_1 \widetilde{\triangleleft} b, \dots, a_n \widetilde{\triangleleft} b)$, where $a \mapsto a \widetilde{\triangleleft} b$ denotes the map inverse to $a \mapsto a \triangleleft b$.

Point 3 follows from the “right” version of corollary 3.10 (point 3): condition (33) means precisely that a is left σ_{\triangleleft} -compatible with ε . \square

Thus, the complex $(T(\mathbb{k}S), \varepsilon d)$ is acyclic for a rack. However, it can be highly non-trivial for shelves (cf. ([31], [32])).

Further, let us turn to the characters given by Dirac maps (32). Theorem 3 applied to the pre-braiding σ_{\triangleleft} and the character φ_a gives

Proposition 4.16. *Take a quandle $(S, \triangleleft, \widetilde{\triangleleft})$ with a fixed element a . A pre-bisimplicial structure can be given on $T(\mathbb{k}S)$ by*

$$\begin{aligned} d_{n;i}(a_1, \dots, a_n) &= \delta_{a,a_i}((a_1 \triangleleft a_i), \dots, (a_{i-1} \triangleleft a_i), a_{i+1}, \dots, a_n), \\ d'_{n;i}(a_1, \dots, a_n) &= \delta_{a,(\dots(a_i \triangleleft a_{i+1}) \triangleleft \dots) \triangleleft a_n}(a_1, \dots, \widehat{a}_i, \dots, a_n). \end{aligned}$$

Moreover, the face maps $d_{n;i}$ combined with degeneracies $s_{n;i} := \Delta_i$ give a weakly simplicial structure.

The total differentials

$$\varphi_a d(a_1, \dots, a_n) = \sum_{i=1}^n (-1)^{i-1} \delta_{a,a_i}((a_1 \triangleleft a_i), \dots, (a_{i-1} \triangleleft a_i), a_{i+1}, \dots, a_n)$$

are called **partial derivatives** and are denoted by $\frac{\partial^1}{\partial a}$ in [30]. Our general setting thus contains some results of [30].

Remark 4.17. One can not talk about weakly bisimplicial structure here, since the coalgebra $(\mathbb{k}S, \sigma_{\triangleleft}, \Delta_D)$ is only semi-braided, and the compatibilities (15) between the $d'_{n;i}$'s and the $s_{n;i}$'s, which are automatical for pre-braided coalgebras and happen to hold in proposition 4.13, are no longer true for here. However, $C_*^D(S)$ is still a sub-bicomplex of $(T(RS), \varphi_a d, d^{\varphi_a})$: indeed, $d'_{n+1;i} \circ s_{n;j}(a_1, \dots, a_n)$ is proportional to $s_{n-1;j-1}(a_1, \dots, \widehat{a}_i, \dots, a_n)$ and is thus still in the image of $s_{n-1;j-1}$ for all $1 \leq i < j \leq n$.

We finish with an example where different characters are used on the right and on the left. This construction is inspired by [3].

Proposition 4.18. *Take a shelf (S, \triangleleft) and work with its linearization ΛS , $\Lambda = \mathbb{Z}[T^{\pm 1}]$. The pre-braiding σ_{\triangleleft} , combined with the shelf characters ε and*

$$\begin{aligned} \varepsilon_T : \Lambda S &\longrightarrow \Lambda, \\ a &\longmapsto T \quad \forall a \in S \end{aligned}$$

define, via theorem 2, a bicomplex structure on $T(\Lambda S)$ by

$$\begin{aligned} \varepsilon d(a_1, \dots, a_n) &= \sum_{i=1}^n (-1)^{i-1} ((a_1 \triangleleft a_i), \dots, (a_{i-1} \triangleleft a_i), a_{i+1}, \dots, a_n), \\ d^{\varepsilon T}(a_1, \dots, a_n) &= \sum_{i=1}^n (-1)^{i-1} T(a_1, \dots, \widehat{a}_i, \dots, a_n), \end{aligned}$$

refined, as usual, to a pre-bisimplicial structure.

The differential $\varepsilon d - d^{\varepsilon T}$ defines the **twisted rack homology** from [3].

4.3 Associative algebras

Now take a \mathbb{k} -vector space V endowed with a bilinear operation $\mu : V \otimes V \rightarrow V$ and a distinguished element $\mathbf{1} \in V$, sometimes regarded as a linear map

$$\begin{aligned} \nu : \mathbb{k} &\longrightarrow V, \\ \alpha &\longmapsto \alpha \mathbf{1}. \end{aligned}$$

In this subsection we construct quite an exotic non-invertible pre-braiding on V which encodes the associativity of μ , and complete it with an exotic comultiplication.

Consider the bilinear application

$$\begin{aligned} \sigma = \sigma_\mu : V \otimes V &\longrightarrow V \otimes V, \\ v \otimes w &\longmapsto \mathbf{1} \otimes \mu(v \otimes w) \end{aligned} \quad (34)$$

or, graphically,

$$\begin{array}{c} \mathbf{1} \quad \mu(v \otimes w) \\ \nu \quad \mu \\ v \quad w \end{array} .$$

Figure 15: Pre-braiding for associative algebras

Lemma 4.19. Suppose that $\mathbf{1}$ is a right unit for μ , i.e. $\mu(v \otimes \mathbf{1}) = v \forall v \in V$. Then the map σ_μ is a pre-braiding if and only if μ is associative on V , i.e.

$$\mu(\mu(v \otimes w) \otimes u) = \mu(v \otimes \mu(w \otimes u)) \quad \forall v, w, u \in V, \quad (\text{Ass})$$

$$\begin{array}{c} \mathbf{1} \\ \mu \\ v \quad w \end{array} = \begin{array}{c} \mathbf{1} \\ \mu \\ v \quad \mu(w \otimes u) \end{array} .$$

Figure 16: Associativity

Proof. Graphically, YBE for σ_μ means

$$\begin{array}{c} \mathbf{1} \quad \mathbf{1} \quad \mu(\mu(v \otimes w) \otimes u) \\ \mu \\ \mathbf{1} \quad \mu(v \otimes w) \\ v \quad w \quad u \end{array} = \begin{array}{c} \mathbf{1} \quad \mathbf{1} \quad \mu(v \otimes \mu(w \otimes u)) \\ \mu \\ \mathbf{1} \quad \mu(w \otimes u) \\ v \quad w \quad u \end{array} .$$

Figure 17: Pictorial proof of lemma 4.19

This is equivalent to the associativity condition (Ass) for μ . □

One thus gets, like in the case of shelves, a pre-braiding subtly encoding the algebraic structure “associative algebra”.

Remark 4.20. The braiding σ_μ is **highly non-invertible**. More precisely, $\sigma_\mu^2 = \sigma_\mu$ if $\mathbf{1}$ is moreover a left unit.

Fix an associative \mathbb{k} -algebra (V, μ) with a right unit $\mathbf{1}$ until the end of this section. Such algebras are called **right-unital** here.

Definition 4.21. An *algebra character* is a unital algebra morphism $\epsilon : V \rightarrow \mathbb{k}$ to the trivial algebra \mathbb{k} , i.e.

$$\begin{aligned} \epsilon(\mu(v \otimes w)) &= \epsilon(v)\epsilon(w) \quad \forall v, w \in V, \\ \epsilon(\mathbf{1}) &= 1, \end{aligned} \quad (35)$$

Figure 18: Algebra character

A *non-unital algebra character* satisfies the first condition only.

Lemma 4.22. Braided characters are precisely maps $\epsilon \in V^*$ satisfying

$$\epsilon(\mathbf{1})\epsilon(\mu(v \otimes w)) = \epsilon(v)\epsilon(w) \quad \forall v, w \in V. \quad (36)$$

In particular, every algebra character is a braided character. On the other hand, any non-zero solution of (36) is a scalar multiple of an algebra character.

Working over a ring R instead of a field \mathbb{k} , one has to drop the last statement.

Lemma 4.23. The linear map

$$\begin{aligned} \Delta_{\mathbf{1}} : V &\longrightarrow V \otimes V, \\ v &\longmapsto \mathbf{1} \otimes v \end{aligned}$$

endows (V, σ_μ) with a pre-braided coalgebra structure, σ_μ -cocommutative if and only if $\mathbf{1}$ is also a left unit.

The last remarks concern arrow operations and the special role of the right unit $\mathbf{1}$ in our braided story. Recall definition 3.8 and corollary 3.10.

Lemma 4.24. 1. The arrow operations give *peripheral* actions:

$${}^\epsilon \pi_w(v_1 \dots v_{n-1} v_n) = \epsilon(\mathbf{1})v_1 \dots v_{n-1}\mu(v_n \otimes w) \quad \forall v_i, w \in V.$$

2. In particular, the right unit $\mathbf{1}$ acts by identity if ϵ is an algebra character:

$${}^\epsilon \pi_{\mathbf{1}} = \text{Id}_{T(V)}.$$

3. The pre-braiding σ_μ is demi-natural with respect to $\mathbf{1}$. Consequently, $\mathbf{1}$ is right σ_μ -compatible with any $f \in V^*$.

We now turn to concrete computations:

Proposition 4.25. Take a right-unital associative algebra $(V, \mu, \mathbf{1})$ and two algebra characters ϵ and ζ .

1. A bicomplex structure can be defined on $T(V)$ by

$$\begin{aligned} d(v_1 \dots v_n) &= \epsilon(v_1)v_2 \dots v_n + \sum_{i=1}^{n-1} (-1)^i v_1 \dots v_{i-1} \mu(v_i \otimes v_{i+1}) v_{i+2} \dots v_n, \\ d^\zeta(v_1 \dots v_n) &= (-1)^{n-1} \zeta(v_n) v_1 \dots v_{n-1} \\ &\quad + \sum_{i=0}^{n-2} (-1)^i \zeta(v_{i+1}) \dots \zeta(v_n) v_1 \dots v_i \mathbf{1} \dots \mathbf{1}. \end{aligned}$$

2. This bidifferential comes from the pre-bisimplicial structure

$$\begin{aligned} d_{n;1}(v_1 \dots v_n) &= \epsilon(v_1)v_2 \dots v_n, \\ d_{n;i+1}(v_1 \dots v_n) &= v_1 \dots v_{i-1} \mu(v_i \otimes v_{i+1}) v_{i+2} \dots v_n, & 1 \leq i \leq n-1, \\ d'_{n;i}(v_1 \dots v_n) &= \zeta(v_i) \dots \zeta(v_n) v_1 \dots v_{i-1} \mathbf{1} \dots \mathbf{1}, & 1 \leq i \leq n-1, \\ d'_{n;n}(v_1 \dots v_n) &= \zeta(v_n) v_1 \dots v_{n-1}. \end{aligned}$$

3. The complex $(T(V), \epsilon d)$ is acyclic.

4. If $\mathbf{1}$ is a two-sided unit, then the above structure becomes weakly bisimplicial, with

$$s_{n;i}(v_1 \dots v_n) = v_1 \dots v_{i-1} \mathbf{1} v_i \dots v_n, \quad 1 \leq i \leq n.$$

5. In this case the structure $(T(V), d_{n;i}, s_{n;i})$ is even simplicial.

6. In the normalized bicomplex, $d'_{n;i} = 0$ for $i < n - 1$.

7. Still supposing the unit $\mathbf{1}$ two-sided, the differential $\epsilon d - d^\zeta$ descends to $T(V')$, where $V' := V/\mathbb{k}\mathbf{1}$, giving the differential

$$\begin{aligned} \epsilon d^\zeta(v_1 \dots v_n) &:= \epsilon(v_1)v_2 \dots v_n \\ &+ \sum_{i=1}^{n-1} (-1)^i v_1 \dots v_{i-1} \mu(v_i \otimes v_{i+1}) v_{i+2} \dots v_n, \\ &+ (-1)^n \zeta(v_n) v_1 \dots v_{n-1}. \end{aligned}$$

Proof. Most of the assertions follow from theorems 2 and 3 applied to the pre-braiding σ_μ from (34) and the algebra, hence braided, characters ϵ and ζ , combined with preceding lemmas.

Point 3 is the corollary 3.10 applied to the element $\mathbf{1}$, possessing the “nice” properties described in lemma 4.24. Point 5 also follows from the properties of $\mathbf{1}$.

More work is needed for proving point 7. Point 4 ensures that tensors $v_1 \dots v_{i-1} \mathbf{1} v_i \dots v_n$ with $1 \leq i \leq n$ generate a sub-bicomplex of $T(V)$, hence a subcomplex of $(T(V), \epsilon d - d^\zeta)$. Further, proposition 3.9 implies that the concatenation map $\bar{v} \mapsto \bar{v}\mathbf{1}$ is a differential complex endomorphism of $(T(V), \epsilon d - d^\zeta)$, thus its image $T(V) \otimes \mathbf{1}$ is a subcomplex. Forming the quotient by these two subcomplexes, one gets the desired differential on $T(V')$. \square

Differential ϵd^ζ defines a (generalization of a) homology sometimes called the **group homology**, which can also be regarded as the *Hochschild homology with trivial coefficients*.

We finish with a “non-unital” remark:

Remark 4.26. In the *non-unital case*, i.e. when V is endowed with a bilinear operation μ only, one enriches V with a formal two-sided unit $\tilde{V} := V \oplus \mathbb{k}\mathbf{1}$, extending μ by

$$\mu(\mathbf{1} \otimes v) = \mu(v \otimes \mathbf{1}) = v \quad \forall v \in \tilde{V}.$$

Due to the equivalence of the associativity of μ on V and on \tilde{V} , lemma 4.19 asserts that σ_μ is a pre-braiding on \tilde{V} if and only if μ is associative on V . Take the character $\epsilon(V) \equiv 0, \epsilon(\mathbf{1}) = 1$ on \tilde{V} . The differential ϵd^ϵ descends to $T((\tilde{V})') \simeq T(V)$, as explained in the previous proposition. One recovers the well-known **bar (or standard) differential**:

$$d_{bar}(v_1 \dots v_n) = \sum_{i=1}^{n-1} (-1)^i v_1 \dots v_{i-1} \mu(v_i \otimes v_{i+1}) v_{i+2} \dots v_n.$$

Moreover, a non-unital algebra character $\epsilon \in V^*$ extends to an algebra character on \tilde{V} by imposing $\epsilon(\mathbf{1}) = 1$. Two such non-unital algebra characters then define a differential ϵd^ζ on $T((\tilde{V})') \simeq T(V)$.

This trick of adding formal elements will often be handy in what follows.

Remark 4.27. One can also obtain the bar differential without doing this formal unit gymnastics. It suffices to replace the total differential with a “cut version” $\partial_{cut} := \sum_{i=2}^n d_{n;i}$ for the pre-simplicial structure from point 2 of proposition 4.25.

4.4 Leibniz algebras

Leibniz algebras are “non-commutative” versions of Lie algebras. They were discovered by A. Bloh in 1965, but it was J.-L. Loday who woke the general interest in this structure around 1989 by, firstly, lifting the classical Chevalley-Eilenberg boundary map from the exterior to the tensor algebra, which yields a new interesting chain complex, and, secondly, by observing that the antisymmetry condition could be omitted (cf. [23],[24],[25],[9]). Here we recover Loday’s complex guided by our “braided” considerations. Our interpretation explains the somewhat mysterious element ordering and signs in the formula given by Loday.

We give here short statements of the main results only, since the proofs are analogous to the associative algebra case; see [20] for details.

Lemma 4.28. Take a \mathbb{k} -vector space V equipped with a bilinear operation $[\cdot, \cdot] : V \otimes V \rightarrow V$ and a *Lie unit*, i.e. a central element, $\mathbf{1}$ in V :

$$[\mathbf{1}, v] = [v, \mathbf{1}] = 0 \quad \forall v \in V. \quad (37)$$

Then the bilinear application

$$\begin{aligned} \sigma = \sigma_{[\cdot, \cdot]} : V \otimes V &\longrightarrow V \otimes V, \\ v \otimes w &\longmapsto w \otimes v + \mathbf{1} \otimes [v, w] \end{aligned} \quad (38)$$

is a pre-braiding if and only if

$$[v, [w, u]] = [[v, w], u] - [[v, u], w] \quad \forall v, w, u \in V, \quad (\text{Lei})$$

Figure 19: Leibniz condition

Note that for the “only if” part of the statement, it is essential to work over a field \mathbb{k} , or to demand another technical condition (cf. lemma 5.9).

Definition 4.29. A pair $(V, [\cdot, \cdot])$ satisfying (Lei) is a *Leibniz algebra*, called *unital* if endowed with a Lie unit $\mathbf{1}$.

One gets the notion of *Lie algebra* when adding the antisymmetry condition.

Lemma 4.30. The pre-braiding $\sigma_{[\cdot, \cdot]}$ is invertible, the inverse given by

$$\sigma_{[\cdot, \cdot]}^{-1} : v \otimes w \longmapsto w \otimes v - [w, v] \otimes \mathbf{1}.$$

Definition 4.31. A *Lie (or Leibniz) character* is a unital Leibniz algebra morphism $\epsilon : V \rightarrow \mathbb{k}$ to the trivial (with the zero bracket) unital (with $1 \in \mathbb{k}$ as a Lie unit) Lie algebra \mathbb{k} , i.e.

$$\begin{aligned} \epsilon([v, w]) &= 0 \quad \forall v, w \in V, \\ \epsilon(\mathbf{1}) &= 1. \end{aligned} \quad (39)$$

A *non-unital Lie character* satisfies the first condition only.

Lemma 4.32. A Lie character is automatically a braided character for the braiding $\sigma_{[\cdot, \cdot]}$.

The comultiplication we choose for Leibniz algebras is what one expects:

Definition 4.33. We call a unital Leibniz algebra V *split* if there is a Leibniz sub-algebra V' of V and a Leibniz algebra decomposition $V \simeq V' \oplus \mathbb{k}\mathbf{1}$.

Lemma 4.34. Take a split unital Leibniz algebra $(V, \sigma_{[\cdot, \cdot]}, \mathbf{1})$. The linear map

$$\begin{aligned} \Delta_{pr} : V &\longrightarrow V \otimes V, \\ v &\longmapsto \mathbf{1} \otimes v + v \otimes \mathbf{1} \quad \forall v \in V', \\ \mathbf{1} &\longmapsto \mathbf{1} \otimes \mathbf{1} \end{aligned}$$

endows it with a semi-braided $\sigma_{[\cdot, \cdot]}$ -cocommutative coalgebra structure.

This comultiplication turns all the elements of V' into primitive ones.

Like in the associative algebra case, the Lie unit $\mathbf{1}$ enjoys important properties with respect to arrow operations:

Lemma 4.35. 1. The arrow operations give *adjoint actions*:

$$\epsilon_{\pi_w}(v_1 \dots v_n) = \epsilon(\mathbf{1}) \sum_{i=1}^n v_1 \dots [v_i, w] \dots v_n + \epsilon(w)v_1 \dots v_n \quad \forall v_i, w \in V.$$

2. In particular, the Lie unit $\mathbf{1}$ acts by scalars:

$$\epsilon_{\pi_{\mathbf{1}}} = \varepsilon(\mathbf{1}) \text{Id}_{T(V)},$$

which is simply $\text{Id}_{T(V)}$ if ε is a Lie character.

3. The pre-braiding $\sigma_{[\cdot]}$ is natural with respect to the Lie unit $\mathbf{1}$. Thus $\mathbf{1}$ is right $\sigma_{[\cdot]}$ -compatible with any $f \in V^*$.

Everything is now ready for explicit calculations of differentials. Only the left ones give something interesting:

Proposition 4.36. *Take a unital Leibniz algebra $(V, [\cdot], \mathbf{1})$ over \mathbb{k} . The braiding $\sigma_{[\cdot]}$ from (38) and a braided character ϵ (for instance, a Lie character) define the following differential on $T(V)$:*

$$\begin{aligned} \epsilon^d(v_1 \dots v_n) &= \epsilon(\mathbf{1}) \sum_{1 \leq i < j \leq n} (-1)^{j-1} v_1 \dots v_{i-1} [v_i, v_j] v_{i+1} \dots \widehat{v}_j \dots v_n + \\ &+ \sum_{1 \leq j \leq n} (-1)^{j-1} \epsilon(v_j) v_1 \dots \widehat{v}_j \dots v_n. \end{aligned}$$

It comes, as usual, from the obvious pre-simplicial structure, completed into a weakly simplicial one by putting

$$s_{n,i}(v_1 \dots v_n) = \begin{cases} v_1 \dots v_{i-1} \mathbf{1} v_i \dots v_n + v_1 \dots v_i \mathbf{1} v_{i+1} \dots v_n & \text{if } v_i \in V', \\ v_1 \dots v_{i-1} \mathbf{1} \mathbf{1} v_{i+1} \dots v_n & \text{if } v_i = \mathbf{1}. \end{cases}$$

The complex $(T(V), \epsilon^d)$ is acyclic if $\epsilon(\mathbf{1}) = 1$.

Remark 4.37. Like for associative algebras, in the *non-unital case* one enriches V with a formal Lie unit $\widetilde{V} := V \oplus \mathbb{k}\mathbf{1}$. The map $\sigma_{[\cdot]}$ is a braiding on \widetilde{V} if and only if $[\cdot]$ is Leibniz on V . Taking the Lie character $\varepsilon(V) \equiv 0, \varepsilon(\mathbf{1}) = 1$ on \widetilde{V} and restricting ϵ^d to the subcomplex $T(V) \subset T(\widetilde{V})$, one recovers the familiar **Leibniz differential**:

$$\epsilon^d(v_1 \dots v_n) = \sum_{1 \leq i < j \leq n} (-1)^{j-1} v_1 \dots v_{i-1} [v_i, v_j] v_{i+1} \dots \widehat{v}_j \dots v_n.$$

5 An upper world: categories

We now present a categorical version of our braided homology theory and of the pre-braidings and other “braided” ingredients for associative and Leibniz algebras. We work in the settings of a preadditive strict monoidal category, symmetric for Leibniz algebras. The world *strict* is omitted but always implied here. This categorification is rather straightforward. A (more subtle and technical) categorical version of shelves and racks and of the corresponding braided differentials can be found in [21].

Only the basic tools of category theory are used here; the famous books [26] and [37] are excellent references for the general and, respectively, “braided” aspects of category theory. We also recommend the preprint [39], where most of the categorical notions used here are nicely presented and illustrated. See also [20].

Notations (1) and (2) are frequently used in this section.

5.1 Categorifying braided differentials

Start with a “local” categorical notion of pre-braiding:

Definition 5.1. \rightarrow An *object* V in a monoidal category $(\mathcal{C}, \otimes, \mathbf{I})$ is called *pre-braided* if it is endowed with a “local” *pre-braiding*, i.e. a morphism $\sigma_V : V \otimes V \rightarrow V \otimes V$ satisfying (a categorical version of) the Yang-Baxter equation (YB):

$$(\sigma_V \otimes \text{Id}_V) \circ (\text{Id}_V \otimes \sigma_V) \circ (\sigma_V \otimes \text{Id}_V) = (\text{Id}_V \otimes \sigma_V) \circ (\sigma_V \otimes \text{Id}_V) \circ (\text{Id}_V \otimes \sigma_V).$$

\rightarrow One talks about a *braided* object if σ_V is an isomorphism.

\rightarrow A *braided morphism* between pre-braided objects (V, σ_V) and (W, σ_W) in \mathcal{C} is a morphism $f : V \rightarrow W$ respecting the pre-braidings:

$$(f \otimes f) \circ \sigma_V = \sigma_W \circ (f \otimes f) : V \otimes V \rightarrow W \otimes W. \quad (40)$$

\rightarrow A *braided character* for a pre-braided object (V, σ_V) is a braided morphism to $(\mathbf{I}, \text{Id}_{\mathbf{I}})$, i.e.

$$(\epsilon \otimes \epsilon) \circ \sigma_V = \epsilon \otimes \epsilon : V \otimes V \rightarrow \mathbf{I} \otimes \mathbf{I} = \mathbf{I}.$$

Every object in a (pre-)braided category (\mathcal{C}, c) is (pre-)braided, with $\sigma_V = c_{V,V}$, since the YBE is automatic in \mathcal{C} . However, the most interesting situation is that of a pre-braiding proper to an object. The idea of working with “*local*” pre-braidings on V instead of demanding the whole category \mathcal{C} to be “*globally*” braided is similar to what is done in [14], where self-invertible YB operators are considered in order to define YB-Lie algebras in an additive monoidal category \mathcal{C} . Note that, contrary to their operator, our pre-braiding is **not necessarily invertible**.

Any pre-braided object (V, σ) in a monoidal category comes with an action of the monoid B_n^+ on $V^{\otimes n}$ for each $n \geq 1$, defined by formula (5). If the category is moreover preadditive, one can mimic the construction of the *quantum (co)shuffle (co)multiplication* to get morphisms

$$\begin{aligned} \Delta_{\sigma}^{p,q} : V^{\otimes n} &= V^{\otimes p} \otimes V^{\otimes q} \rightarrow V^{\otimes n}, \\ \Theta_{\sigma}^{p,q} : V^{\otimes n} &\rightarrow V^{\otimes p} \otimes V^{\otimes q} = V^{\otimes n}. \end{aligned}$$

Here $n = p + q$. Still in the preadditive context, $-\sigma$ is well defined and gives a new pre-braiding for V .

The definition 3.5 of (semi-)pre-braided coalgebra is directly transported to the setting of a monoidal category. A categorical version of definition 3.8 is also straightforward:

Definition 5.2. Take an object V and two morphisms $\eta : \mathbf{I} \rightarrow V$ and $\psi : V \rightarrow \mathbf{I}$ in a monoidal category.

\rightarrow η and ψ are said to form a *normalized pair* if $\psi \circ \eta = \text{Id}_{\mathbf{I}}$.

\rightarrow η and ψ are called *right σ -compatible* for a pre-braiding σ on V if

$$(\text{Id}_V \otimes \psi) \circ \sigma \circ (\text{Id}_V \otimes \eta) = \eta \circ \psi : V \rightarrow V.$$

\rightarrow A pre-braiding σ on V is called *natural* with respect to η if

$$\sigma \circ (\eta \otimes \text{Id}_V) = \text{Id}_V \otimes \eta, \quad (41)$$

$$\sigma \circ (\text{Id}_V \otimes \eta) = \eta \otimes \text{Id}_V, \quad (42)$$

and *semi-natural* (or *demi-natural*) if only (41) (resp. (42)) holds.

Further, recall the categorical notions of (bi)differentials:

Definition 5.3. Take a preadditive category \mathcal{C} .

→ A degree -1 differential for a family of objects $\{V_n\}_{n \geq 0}$ in \mathcal{C} is a family of morphisms $\{d_n : V_n \rightarrow V_{n-1}\}_{n > 0}$, satisfying $d_{n-1} \circ d_n = 0 \ \forall n > 1$.

→ A bidegree -1 bidifferential is a pair of morphism families $\{d_n, d'_n : V_n \rightarrow V_{n-1}\}_{n > 0}$ s.t.

$$d_{n-1} \circ d_n = d'_{n-1} \circ d'_n = d'_{n-1} \circ d_n + d_{n-1} \circ d'_n = 0 \quad \forall n > 1. \quad (43)$$

→ Given a degree -1 differential $\{d_n\}_{n > 0}$ for a family of objects $\{V_n\}_{n \geq 0}$, one defines a *contracting homotopy* as a family of morphisms $\{h_n : V_n \rightarrow V_{n+1}\}_{n \geq 0}$, satisfying

$$h_{n-1} \circ d_n + d_{n+1} \circ h_n = \text{Id}_{V_n} \quad \forall n > 0.$$

→ Different types of *simplicial objects* in \mathcal{C} are defined by replacing the words “vector space” by “object in \mathcal{C} ” in the definition 3.3.

→ If \mathcal{C} is moreover monoidal, a degree -1 tensor (bi)differential for a $V \in \text{Ob}(\mathcal{C})$ is a degree -1 (bi)differential for the family of objects $\{V^{\otimes n}\}_{n \geq 0}$.

Points 1 - 3 of proposition 3.4 remain valid for simplicial objects in a preadditive category \mathcal{C} .

The presence of a contracting homotopy means, in the category $\mathcal{C} = \mathbf{Mod}_R$, that the complex (V_n, d_n) is acyclic.

With all the preparatory work above, theorems 2 and 3 and proposition 3.9 (with their proofs!) are generalized as follows (we freely use the notations from those theorems here):

Theorem 5. *In a preadditive monoidal category $(\mathcal{C}, \otimes, \mathbf{I})$, take a pre-braided object (V, σ) endowed with braided characters ϵ and ζ .*

1. A bidegree -1 tensor bidifferential for V can be defined by the morphism families

$$\begin{aligned} (\epsilon^d)_n &:= \epsilon_1 \circ \Theta_{-\sigma}^{1, n-1}, \\ (d^\zeta)_n &:= (-1)^{n-1} \zeta_n \circ \Theta_{-\sigma}^{n-1, 1}. \end{aligned}$$

2. This bidifferential can be refined into a pre-bisimplicial structure on $(V^{\otimes n})_{n \geq 0}$ given by

$$\begin{aligned} d_{n;i} &:= \epsilon_1 \circ T_{p_{i,n}}^\sigma, \\ d'_{n;i} &:= \zeta_n \circ T_{p'_{i,n}}^\sigma. \end{aligned}$$

3. If a comultiplication Δ endows (V, σ) with a pre-braided coalgebra structure, then the morphisms $s_{n;i} := \Delta_i$ complete the preceding structure into a very weakly bisimplicial one.

4. If (V, σ, Δ) is a semi-pre-braided coalgebra, then one obtains a very weakly simplicial object $(V^{\otimes n}, d_{n;i}, s_{n;i})$ only.

5. If Δ is moreover σ -cocommutative, then the above structures on $(V^{\otimes n})_{n \geq 0}$ are weakly (bi)simplicial.

6. Take a morphism $\eta : \mathbf{I} \rightarrow V$. The family $h_n := (-1)^n \text{Id}_n \otimes \eta : V^{\otimes n} \rightarrow V^{\otimes(n+1)}$

→ is a contracting homotopy for $(V^{\otimes n}, (\epsilon^d)_n)$ if the pair (η, ϵ) is normalized, and σ is demi-natural with respect to η .

→ is a contracting homotopy for $(V^{\otimes n}, (d^\zeta)_n)$ if the pair (η, ζ) is normalized and right σ -compatible.

5.2 Basic examples revisited

Recall the categorical definitions of associative and Leibniz algebras:

Definition 5.4. \rightarrow A *(unital) associative algebra* in a monoidal category \mathcal{C} , abbreviated as (U)AA, is an object V together with morphisms $\mu : V \otimes V \rightarrow V$ (and $\nu : \mathbf{I} \rightarrow V$), satisfying the associativity (and the unit) conditions:

$$\begin{aligned}\mu \circ (\text{Id}_V \otimes \mu) &= \mu \circ (\mu \otimes \text{Id}_V) : V^{\otimes 3} \rightarrow V, \\ \mu \circ (\text{Id}_V \otimes \nu) &= \mu \circ (\nu \otimes \text{Id}_V) = \text{Id}_V.\end{aligned}$$

For a *right-unital associative algebra* we demand only the $\mu \circ (\text{Id}_V \otimes \nu) = \text{Id}_V$ part of the last condition.

- \rightarrow The subcategory of UAAs and unital algebra morphisms (i.e. morphisms respecting μ and ν) in \mathcal{C} is denoted by $\mathbf{UAlg}(\mathcal{C})$, or $\mathbf{Alg}(\mathcal{C})$ in the non-unital case.
- \rightarrow An *algebra character* for $V \in \mathbf{UAlg}(\mathcal{C})$ is a unital algebra morphism $\epsilon \in \text{Hom}_{\mathbf{UAlg}(\mathcal{C})}(V, \mathbf{I})$, with the *default UAA structure on \mathbf{I}* ($\mu = \nu = \text{Id}_{\mathbf{I}}$). A *non-unital algebra character* is a morphism in $\mathbf{Alg}(\mathcal{C})$ only.

Definition 5.5. \rightarrow A *(unital) Leibniz algebra* in a symmetric preadditive category \mathcal{C} , abbreviated as (U)LA, is an object V together with morphisms $[\cdot, \cdot] : V \otimes V \rightarrow V$ (and $\nu : \mathbf{I} \rightarrow V$) satisfying the generalized Leibniz (and the Lie unit) conditions:

$$\begin{aligned}[\cdot, \cdot] \circ (\text{Id}_V \otimes [\cdot, \cdot]) &= [\cdot, \cdot] \circ ([\cdot, \cdot] \otimes \text{Id}_V) - [\cdot, \cdot] \circ ([\cdot, \cdot] \otimes \text{Id}_V) \circ (\text{Id}_V \otimes c_{V,V}) : V^{\otimes 3} \rightarrow V, \\ [\cdot, \cdot] \circ (\text{Id}_V \otimes \nu) &= [\cdot, \cdot] \circ (\nu \otimes \text{Id}_V) = 0 : V \rightarrow V.\end{aligned}$$

- \rightarrow The subcategory of ULAs and unital Leibniz algebra morphisms (i.e. morphisms respecting $[\cdot, \cdot]$ and ν) in \mathcal{C} is denoted by $\mathbf{ULei}(\mathcal{C})$, or $\mathbf{Lei}(\mathcal{C})$ in the non-unital case.
- \rightarrow A *Lie character* for $V \in \mathbf{ULei}(\mathcal{C})$ is an $\epsilon \in \text{Hom}_{\mathbf{ULei}(\mathcal{C})}(V, \mathbf{I})$, with the *default ULA structure on \mathbf{I}* (zero bracket and $\nu = \text{Id}_{\mathbf{I}}$). A *non-unital Lie character* is an $\epsilon \in \text{Hom}_{\mathbf{Lei}(\mathcal{C})}(V, \mathbf{I})$.

See for instance [1] and [28] for the definition of algebras in a monoidal category, and [14] for a survey on categorical Lie algebras. Note that $\mathbf{UAlg}(\mathbf{Vect}_{\mathbb{k}})$ and $\mathbf{ULei}(\mathbf{Vect}_{\mathbb{k}})$ are the familiar categories of \mathbb{k} -linear unital associative and Leibniz algebras respectively.

Definition 5.6. A *normalized morphism* $\varphi : V \rightarrow W$ for $V, W \in \mathbf{UAlg}(\mathcal{C})$ or $\in \mathbf{ULei}(\mathcal{C})$ is a morphism in \mathcal{C} respecting the units, i.e. $\varphi \circ \nu_V = \nu_W$.

For $W = \mathbf{I}$ this means that (ν_V, φ) is a normalized pair.

Now, the content of subsections 4.3 and 4.4 can be categorified as follows:

Theorem 6. 1. Take a right-unital associative algebra (V, μ, ν) in a monoidal category $(\mathcal{C}, \otimes, \mathbf{I})$.

(a) V can be endowed with a pre-braiding

$$\sigma_{Ass} := \nu \otimes \mu : V \otimes V = \mathbf{I} \otimes V \otimes V \rightarrow V \otimes V. \quad (44)$$

(b) If ν is moreover a two-sided unit, then the comultiplication

$$\Delta_{Ass} := \nu \otimes \text{Id}_V : V = \mathbf{I} \otimes V \rightarrow V \otimes V$$

completes σ_{Ass} into a σ_{Ass} -cocommutative pre-braided coalgebra structure.

(c) Any algebra character $\epsilon \in \text{Hom}_{\mathbf{UAlg}(\mathcal{C})}(V, \mathbf{I})$ is a braided character for (V, σ_{Ass}) .

(d) The pre-braiding σ_{Ass} is demi-natural with respect to the unit ν . Moreover, for any algebra character ϵ , the pair (ν, ϵ) is normalized.

2. Take a unital Leibniz algebra $(V, [\cdot, \cdot], \nu)$ in a symmetric preadditive category $(\mathcal{C}, \otimes, \mathbf{I}, c)$.

(a) V can be endowed with an invertible braiding

$$\sigma_{Lei} := c_{V,V} + \nu \otimes [\cdot, \cdot]. \quad (45)$$

(b) If \mathcal{C} is additive and if one has a Leibniz algebra decomposition $V \simeq V' \oplus \mathbf{I}$, then

$$\begin{aligned} \Delta_{Lei}|_{V'} &:= \nu \otimes \text{Id}_{V'} + \text{Id}_{V'} \otimes \nu : V' \rightarrow V \otimes V, \\ \Delta_{Lei}|_{\mathbf{I}} &:= \nu \otimes \nu : \mathbf{I} \rightarrow V \otimes V \end{aligned}$$

completes σ_{Lei} into a σ_{Lei} -cocommutative semi-braided coalgebra structure.

(c) Any Lie character $\epsilon \in \text{Hom}_{\mathbf{ULei}(\mathcal{C})}(V, \mathbf{I})$ is a braided character for (V, σ_{Lei}) .

(d) The braiding σ_{Lei} is natural with respect to the unit ν . Moreover, for any Lie character ϵ , the pair (ν, ϵ) is normalized.

Observe that in the Leibniz algebra setting, the naturality (with respect to morphisms ν and $[\cdot, \cdot]$ in particular) and the symmetry of the braiding c are essential in proving that σ_{Lei} is indeed a braiding, while the naturality of c with respect to ϵ shows that ϵ is a braided character for $(V, c_{V,V})$ (which implies that it is a braided character for (V, σ_V) if it preserves the Leibniz structure).

Remark 5.7. According to the theorem, a ULA V is a “doubly braided” object: σ_V and $c_{V,V}$ are indeed two distinct braidings on V . One can say more: they endow the tensor powers of V with an action of the *virtual braid group* ([19], [38]). The close connections between (pre-)braided objects and virtual braid groups are studied in detail in [21].

Theorems 5 and 6 put together give categorical versions of propositions 4.25 and 4.36, as well as of the “non-unital” remarks 4.26 and 4.37:

Corollary 5.8. 1. Any algebra character $\epsilon : V \rightarrow \mathbf{I}$ for a UAA (V, μ, ν) in a preadditive monoidal category \mathcal{C} produces a degree -1 tensor differential

$$({}^\epsilon d)_n := \epsilon_1 + \sum_{i=1}^{n-1} (-1)^i \mu_i,$$

for V , with a contracting homotopy $h_n = (-1)^n \nu_{n+1}$. Any non-unital algebra characters ϵ, ζ for an associative algebra (V, μ) in \mathcal{C} produce a degree -1 tensor differential

$$({}^\epsilon d^\zeta)_n := \epsilon_1 + \sum_{i=1}^{n-1} (-1)^i \mu_i + (-1)^n \zeta_n.$$

2. Any Lie character $\epsilon : V \rightarrow \mathbf{I}$ for a ULA $(V, [\cdot, \cdot], \nu)$ in a symmetric preadditive category \mathcal{C} produces a degree -1 tensor differential

$$({}^\epsilon d)_n := \epsilon_1 \circ \underset{-c}{\Theta}^{1, n-1} + \sum_{1 \leq i < j \leq n} (-1)^{j-1} [\cdot, \cdot]_i \circ (\text{Id}_i \otimes c_{V^{j-i-1}, V} \otimes \text{Id}_{n-j}),$$

for V , with a contracting homotopy $h_n = (-1)^n \nu_{n+1}$. Any non-unital Lie character ϵ for a Leibniz algebra $(V, [\cdot, \cdot])$ in a symmetric additive \mathcal{C} produces a degree -1 tensor differential

$$({}^\epsilon d)_n := \sum_{1 \leq i < j \leq n} (-1)^{j-1} [\cdot, \cdot]_i \circ (\text{Id}_i \otimes c_{V^{j-i-1}, V} \otimes \text{Id}_{n-j}).$$

Working in the category \mathbf{Vect}_k in section 4, we noticed that the pre-braidings obtained for associative and Leibniz algebras **encode** the underlying algebraic structures (cf. lemmas 4.19 and 4.28). It is still true, with some additional technical assumptions, in the categorical setting:

Lemma 5.9. 1. Take an object V in a monoidal category $(\mathcal{C}, \otimes, \mathbf{I})$ endowed with two morphisms $\mu : V \otimes V \rightarrow V$ and $\nu : \mathbf{I} \rightarrow V$, with ν being a two-sided unit for μ . The morphism σ_{Ass} defined by (44) is a pre-braiding if and only if μ is associative.

2. Take an object V in a symmetric preadditive category $(\mathcal{C}, \otimes, \mathbf{I}, c)$ endowed with two morphisms $[\cdot, \cdot] : V \otimes V \rightarrow V$ and $\nu : \mathbf{I} \rightarrow V$, with ν being a Lie unit for $[\cdot, \cdot]$. Additionally suppose the existence of a normalized morphism $\gamma : V \rightarrow \mathbf{I}$. The morphism σ_{Lei} defined by (45) is a braiding if and only if $[\cdot, \cdot]$ satisfies the Leibniz condition.

Proof. One repeats the proofs of lemmas 4.19 and 4.28. The only non-trivial step is to show that

$$\nu \otimes \nu \otimes f = \nu \otimes \nu \otimes g : V^{\otimes 3} \rightarrow V^{\otimes 3}$$

implies

$$f = g : V^{\otimes 3} \rightarrow V.$$

When ν is a left unit for μ , this is done by applying $\mu \circ (\text{Id}_V \otimes \mu)$ to both sides of the first identity. In the Leibniz case, apply $\gamma \otimes \gamma \otimes \text{Id}_V$. \square

Similar considerations lead to an “if and only if” type result for morphisms:

Lemma 5.10. \rightarrow In the settings of theorem 6, any morphism $f : V \rightarrow W$ in $\mathbf{UAlg}(\mathcal{C})$ (resp. $\mathbf{ULei}(\mathcal{C})$) is braided, with V and W endowed with the pre-braidings σ_{Ass} (resp. σ_{Lei}).

- \rightarrow Suppose additionally, for the algebra case, that ν is a two-sided unit, and, for the Leibniz case, the existence of a normalized morphism $\gamma : V \rightarrow \mathbf{I}$. Then any braided normalized (cf. definition 5.6) morphism $f : V \rightarrow W$ in \mathcal{C} necessarily respects the multiplications, i.e. remains a morphism in $\mathbf{UAlg}(\mathcal{C})$ (resp. $\mathbf{ULei}(\mathcal{C})$).

Thus an associative/Leibniz algebra morphism is the same thing (modulo some technical conditions) as a normalized braided morphism for the corresponding “structural” pre-braiding, illustrating the precision with which our pre-braidings capture the algebraic properties of the corresponding structure.

This result can be compactly restated as follows. Denote by $\mathbf{Br}^\nu(\mathcal{C})$ the subcategory of a monoidal category \mathcal{C} with

- \rightarrow as objects, pre-braided objects V endowed with a “unit” $\nu : \mathbf{I} \rightarrow V$;
- \rightarrow as morphisms, braided normalized morphisms in \mathcal{C} .

Proposition 5.11. 1. Given a monoidal category \mathcal{C} , one can see $\mathbf{UAlg}(\mathcal{C})$ as a full subcategory of $\mathbf{Br}^\nu(\mathcal{C})$ via the functor

$$\begin{aligned} \mathbf{UAlg}(\mathcal{C}) &\hookrightarrow \mathbf{Br}^\nu(\mathcal{C}), \\ (V, \mu, \nu) &\mapsto (V, \sigma_{Ass}, \nu), \\ (f : V \rightarrow W) &\mapsto (f : V \rightarrow W). \end{aligned}$$

2. Given a symmetric preadditive category \mathcal{C} , one can see $\mathbf{ULei}^*(\mathcal{C})$ (the category of ULAs V in \mathcal{C} endowed with a normalized morphism $\gamma : V \rightarrow \mathbf{I}$) as a full subcategory of $\mathbf{Br}^\nu(\mathcal{C})$ via

$$\begin{aligned} \mathbf{ULei}^*(\mathcal{C}) &\hookrightarrow \mathbf{Br}^\nu(\mathcal{C}), \\ (V, [\cdot, \cdot], \nu) &\mapsto (V, \sigma_{Lei}, \nu), \\ (f : V \rightarrow W) &\mapsto (f : V \rightarrow W). \end{aligned}$$

5.3 The super trick

The first bonus one generally gains when passing to abstract symmetric categories is the possibility to derive graded and super versions of algebraic results for free, thanks to the Koszul flip τ_{Koszul} from (3). One clearly sees where to put signs, which is otherwise quite difficult to guess. Here is a typical example.

Take a *graded unital Leibniz algebra* $(V, [\cdot, \cdot], \nu)$, i.e. an object of $\mathbf{ULei}(\mathbb{Z}\mathbf{Vect}_k)$, where $\mathbb{Z}\mathbf{Vect}_k$ is the usual additive monoidal category of graded k -vector spaces, endowed with the symmetric braiding τ_{Koszul} . Leibniz condition in this setting is

$$[v, [w, u]] = [[v, w], u] - (-1)^{\deg v \deg w} [[v, u], w]$$

for any homogeneous elements $v, w, u \in V$. Cf. fig. 19 illustrating (Lei), with the crossing on the right corresponding to the “internal” braiding $c_{V,V} = \tau_{Koszul}$.

Theorem 6 gives a braiding for V :

$$\sigma_V : v \otimes w \longmapsto (-1)^{\deg v \deg w} w \otimes v + \mathbf{1} \otimes [v, w],$$

which, together with a Lie character $\epsilon : V_0 \rightarrow \mathbb{k}$, can be fed into the machinery from theorem 5:

Proposition 5.12. 1. A \mathbb{k} -linear graded unital Leibniz algebra $(V, [,], \nu)$ with a Lie character ϵ can be endowed with the degree -1 tensor differential

$$\begin{aligned} \epsilon d(v_1 \dots v_n) &= \sum_{1 \leq i < j \leq n} (-1)^{j-1+\alpha_{i,j}} v_1 \dots v_{i-1} [v_i, v_j] v_{i+1} \dots \widehat{v}_j \dots v_n + \\ &+ \sum_{1 \leq j \leq n} (-1)^{j-1+\alpha_{0,j}} \epsilon(v_j) v_1 \dots \widehat{v}_j \dots v_n, \end{aligned}$$

where $\alpha_{i,j} := \deg(v_j) \sum_{i < k < j} \deg(v_k)$.

2. A \mathbb{k} -linear graded Leibniz algebra $(V, [,]) with a non-unital Lie character ϵ can be endowed with the degree -1 tensor differential$

$$\epsilon d(v_1 \dots v_n) = \sum_{1 \leq i < j \leq n} (-1)^{j-1+\alpha_{i,j}} v_1 \dots v_{i-1} [v_i, v_j] v_{i+1} \dots \widehat{v}_j \dots v_n.$$

All the v_i 's are taken homogeneous here.

Observe that the $(-1)^{\alpha_{i,j}}$ part of the sign comes from the Koszul braiding, while $(-1)^{j-1}$ appears because we take the opposite braiding when defining $(\epsilon d)_n := \epsilon_1 \circ \Theta_{-\sigma}^{1,n-1}$.

Leibniz superalgebras are treated similarly: one has just to work in the category of super vector spaces over \mathbb{k} (cf. [22]). One thus recovers the **Leibniz superalgebra homology**, which is a lift of the Lie superalgebra homology.

Similarly, one gets for free the **color Leibniz algebra homology** (cf. [11], or [33] for a Lie version). Concretely, take a finite abelian group Γ endowed with an antisymmetric bicharacter χ . The category ${}_{\Gamma}\mathbf{Vect}_{\mathbb{k}}$ of \mathbb{k} -vector spaces graded over Γ is symmetric additive, with the usual Γ -graded tensor product, the zero-graded \mathbb{k} as its identity object and, as a braiding, the *color flip*

$$\tau_{color} : v \otimes w \longmapsto \chi(f, g) w \otimes v$$

for homogeneous v and w graded over f and $g \in \Gamma$ respectively. Then color Leibniz algebras are precisely Leibniz algebras in ${}_{\Gamma}\mathbf{Vect}_{\mathbb{k}}$

See also [39] for an excellent survey of different types of braided Lie algebras.

5.4 Co-world, or the world upside down

One more nice feature of the categorical approach is an automatic treatment of **dualities**. The most common notion of duality, the “upside-down” one, is described here, with the cobar complex for coalgebras (cf. [6], [10]) providing an example. In the monoidal context, one has two more dualities, the “right-left” and the combined ones, treated in the next subsection.

Definition 5.13. Given a category \mathcal{C} , its *dual (or opposite) category* \mathcal{C}^{op} is constructed by keeping the objects of \mathcal{C} and reversing all the arrows. One writes $f^{\text{op}} \in \text{Hom}_{\mathcal{C}^{\text{op}}}(W, V)$ for the morphism in \mathcal{C}^{op} corresponding to an $f \in \text{Hom}_{\mathcal{C}}(V, W)$.

We sometimes call \mathcal{C}^{op} a *co-category* in order to avoid confusion with other notions of duality. Observe that this construction is involutive: $(\mathcal{C}^{\text{op}})^{\text{op}} = \mathcal{C}$.

The **duality principle** (cf. [26], section II.2) tells that a “categorical” theorem for \mathcal{C} implies a dual theorem for \mathcal{C}^{op} by reversing all the arrows and the order of arrows in every composition. Our aim here is to apply this principle to theorems 5 and 6.

To get a notion of duality for categorical structures, it suffices to place them into the co-category. For example, a *counital co-Leibniz coalgebra* (= co-ULA) in a symmetric preadditive category \mathcal{C} is an object V together with morphisms $\partial : V \rightarrow V \otimes V$ and $\varepsilon : V \rightarrow \mathbf{I}$, such that $(V, \partial^{\text{op}}, \varepsilon^{\text{op}})$ is a ULA in \mathcal{C}^{op} . In other words, ∂ and ε satisfy

$$\begin{aligned} (\text{Id}_V \otimes \partial) \circ \partial &= (\partial \otimes \text{Id}_V) \circ \partial - (\text{Id} \otimes c_{V,V}) \circ (\partial \otimes \text{Id}) \circ \partial, \\ (\text{Id}_V \otimes \varepsilon) \circ \partial &= (\varepsilon \otimes \text{Id}_V) \circ \partial = 0. \end{aligned} \quad (46)$$

A convenient way to handle the “upside-down” duality is the graphical one: changing from \mathcal{C} to \mathcal{C}^{op} consists simply in turning all the diagrams upside down, i.e. taking a *horizontal mirror image*. Here is an example for the co-Leibniz condition (46) (cf. fig. 19):

Figure 20: Co-Leibniz condition

We now make a list of dualities for the categorical structures relevant for this paper:

unital associative algebra (V, μ, ν)	co-UAA $(V, \mu^{\text{op}}, \nu^{\text{op}})$
unital Leibniz algebra $(V, [,], \nu)$	co-ULA $(V, [,]^{\text{op}}, \nu^{\text{op}})$
algebra character φ for (V, μ, ν)	coalgebra co-character φ^{op} for $(V, \mu^{\text{op}}, \nu^{\text{op}})$
Lie character φ for $(V, [,], \nu)$	co-Lie co-character φ^{op} for $(V, [,]^{\text{op}}, \nu^{\text{op}})$
pre-braiding σ for V	pre-braiding σ^{op} for V
braided character ϵ for (V, σ)	braided co-character ϵ^{op} for (V, σ^{op})
(bi)degree -1 tensor (bi)bidifferential for V	(bi)degree 1 tensor (bi)differential for V

Table 3: Categorical duality for structures

The subcategory of co-UAA's and co-ULA's in \mathcal{C} are denoted by $\mathbf{coUAlg}(\mathcal{C})$ and $\mathbf{coULei}(\mathcal{C})$ respectively.

Remark 5.14. For a pre-braided object (V, σ) and the action (5) of B_n^+ on $V^{\otimes n}$, one has

$$(T_s^\sigma)^{\text{op}} = T_{s^{-1}}^{(\sigma^{\text{op}})} \in \text{End}_{\mathcal{C}^{\text{op}}}(V^{\otimes n}) \quad \forall s \in S_n.$$

Thus, assuming the category preadditive, the definition (9) of *quantum co-shuffle comultiplication* is translated as $\Theta_\sigma^{p,q} = (\Delta_{\sigma^{\text{op}}}^{p,q})^{\text{op}}$, automatically giving all the properties of this structure.

Everything is now ready for dualizing theorems 5 and 6. We present only short versions of these results here, leaving the dualization of the points concerning simplicial and pre-braided coalgebra structures to the reader.

Theorem 5^{co}. *Let \mathcal{C} be a preadditive monoidal category. For any pre-braided object (V, σ) with braided co-characters e and c , a bidegree 1 tensor bidifferential for V can be defined by*

$$\begin{aligned} (d)^\sigma &:= \Delta_{-\sigma}^{1,n} \circ (e \otimes \text{Id}_n), \\ (d_c)^\sigma &:= (-1)^n \Delta_{-\sigma}^{n,1} \circ (\text{Id}_n \otimes c). \end{aligned}$$

Theorem 6^{co}.

1. Take a counital coassociative coalgebra (V, Δ, ε) in a monoidal category \mathcal{C} .

- (a) V can be endowed with a pre-braiding $\sigma_{\text{coAss}} := \varepsilon \otimes \Delta$.
- (b) Any coalgebra co-character $e \in \text{Hom}_{\mathbf{coUAlg}(\mathcal{C})}(\mathbf{I}, V)$ is a braided co-character for $(V, \sigma_{\text{coAss}})$.

2. Take a counital co-Leibniz coalgebra $(V, \partial, \varepsilon)$ in a symmetric preadditive category $(\mathcal{C}, \otimes, \mathbf{I}, c)$.

(a) V can be endowed with a braiding $\sigma_{coLei} := c_{V,V} + \varepsilon \otimes \partial$.

(b) Any co-Lie co-character $e \in \text{Hom}_{\mathbf{coULei}(\mathcal{C})}(\mathbf{I}, V)$ is a braided co-character for (V, σ_{coLei}) .

A graphical depiction of, for instance, σ_{coAss} is by construction the horizontal mirror image of the diagram one had for UAAs:



Figure 21: $\sigma_{coAss} = \text{HorMirror}(\sigma_{Ass})$

A co-version of corollary 5.8 is then formulated in the evident way, with dual explicit formulas. “If and only if” lemmas 5.9 and 5.10 are also dualized directly. In particular, the pre-braidings from the previous theorem encode the co-associativity (resp. co-Leibniz) condition.

We finish this section with some remarks proper to our favorite category $\mathbf{Vect}_{\mathbb{k}}$ (everything remaining valid, as usual, in \mathbf{Mod}_R).

Lemma 5.15. In $\mathbf{Vect}_{\mathbb{k}}$, a map $e : \mathbb{k} \rightarrow V, \alpha \mapsto \alpha \mathbf{e}$ for a co-UAA (V, Δ, ϵ) is a non-unital coalgebra co-character if and only if $\mathbf{e} \in V$ is *group-like*, i.e. $\Delta(\mathbf{e}) = \mathbf{e} \otimes \mathbf{e}$. A non-unital Lie co-character for a co-ULA (V, ∂, ϵ) corresponds to an $\mathbf{e} \in \text{Ker}(\partial)$.

Further, “non-unital” remarks 4.26 and 4.37 admit co-versions. To create a counit for a coassociative or co-Leibniz coalgebra (V, δ) (resp. (V, ∂)), one extends it by adding a formal element $\tilde{V} := V \oplus \mathbb{k}\mathbf{1}$, modifying the comultiplication:

$$\begin{aligned} \Delta(v) &= \delta(v) + \mathbf{1} \otimes v + v \otimes \mathbf{1} \quad \forall v \in V, \\ \Delta(\mathbf{1}) &= \mathbf{1} \otimes \mathbf{1} \end{aligned}$$

in the coassociative coalgebra case, and

$$\partial(\mathbf{1}) = 0,$$

keeping the original ∂ on V , in the co-Leibniz case. The pre-braiding σ_{coAss} (resp. σ_{coLei}) on \tilde{V} now characterizes the coassociativity (resp. co-Leibniz) condition for V . Further, the application $\varepsilon \in (\tilde{V})^*$ given by $\varepsilon(V) \equiv 0, \varepsilon(\mathbf{1}) = 1$ is a coalgebra (resp. co-Lie) counit for Δ (resp. ∂), and $\mathbf{1}$ is a group-like element (resp. $\mathbf{1} \in \text{Ker}(\partial)$).

The left braided differentials obtained in this setting for coalgebras are described in

Proposition 5.16. Given a \mathbb{k} -linear coalgebra (V, δ) , extend it to a counital one $(\tilde{V}, \Delta, \varepsilon)$ as above. Then the group-like $\mathbf{1}$ gives, via theorem 5^{co}, the following differential on $T(\tilde{V})$:

$$\mathbf{1}d(v_1 \dots v_n) = \mathbf{1}v_1 \dots v_n + \sum_{i=1}^n (-1)^i v_1 \dots v_{i-1} \Delta(v_i) v_{i+1} \dots v_n.$$

The ideal $I_{\mathbf{1}}$ of the tensor algebra $T(\tilde{V})$ generated by the element $\mathbf{1}$ is $\mathbf{1}d$ -stable. The differential induced on $T(\tilde{V})/I_{\mathbf{1}} \simeq T(\tilde{V}/\mathbb{k}\mathbf{1}) \simeq T(V)$ is

$$\tilde{\mathbf{1}}d(v_1 \dots v_n) = \sum_{i=1}^n (-1)^i v_1 \dots v_{i-1} \delta(v_i) v_{i+1} \dots v_n.$$

One eagerly recognizes the *cobar differential* for coalgebras.

5.5 Right-left duality

One more notion of duality is available for a monoidal category $(\mathcal{C}, \otimes, \mathbf{I})$. One can simply change its tensor product to the opposite one: $V \otimes^{\text{op}} W := W \otimes V$ for objects, and similarly for morphisms. We call this new monoidal category **monoidally dual** to \mathcal{C} , denoting it by $\mathcal{C}^{\otimes^{\text{op}}}$ (there seem to be no universally accepted notation, some authors even using \mathcal{C}^{op} here and another notation for co-categories). Graphically, the categories \mathcal{C} and $\mathcal{C}^{\otimes^{\text{op}}}$ differ by the *vertical mirror symmetry* for all diagrams.

Applying monoidal duality to a co-category \mathcal{C}^{op} , one gets $\mathcal{C}^{\text{op}, \otimes^{\text{op}}} := (\mathcal{C}^{\text{op}})^{\otimes^{\text{op}}} \simeq (\mathcal{C}^{\otimes^{\text{op}}})^{\text{op}}$. Graphically, it corresponds to the *central symmetry*.

Similarly to what we have seen for \mathcal{C}^{op} , all “categorical” notions and theorems have monoidally dual versions in $\mathcal{C}^{\otimes^{\text{op}}}$. This gives in particular **right differentials** $(d^e)_n, (d_e)^n$, monoidally dual to the left ones $({}^e d)_n, ({}_e d)^n$. Note that these differentials should be endowed with a sign (cf. theorem 2) if one wants a bidifferential structure.

One also has **right braidings**, monoidally dual to those from theorems 6 and 6^{co}. In particular, a new braiding emerges for UAAs:

$$\sigma_{A_{ss}}^r := \mu \otimes \nu = \text{VertMirror}(\sigma_{A_{ss}}) = \begin{array}{c} \mu \quad \nu \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} .$$

Remark that the Leibniz algebra structure is not right-left symmetric: a Leibniz algebra in $\mathcal{C}^{\otimes^{\text{op}}}$ is in fact a **left Leibniz algebra** in \mathcal{C} (cf. [25]). Thus one automatically obtains braided homology theories for left Leibniz algebras.

6 Braided modules and homologies with coefficients

We introduce here the notion of (bi)modules over a pre-braided object V in a monoidal category. These “braided” modules generalize, in quite an unexpected manner, the usual notions of (bi)modules for algebraic structures. Since at the same time a braided module generalizes a braided character, one naturally arrives to homologies of pre-braided objects with coefficients. As particular cases, we point out Hochschild and Chevalley-Eilenberg complexes.

Fix a monoidal category $(\mathcal{C}, \otimes, \mathbf{I})$.

Definition 6.1. \rightarrow A *right module* over a pre-braided object (V, σ) is an object $M \in \text{Ob}(\mathcal{C})$ equipped with a morphism $\rho : M \otimes V \rightarrow M$ satisfying

$$\rho \circ (\rho \otimes \text{Id}_V) = \rho \circ (\rho \otimes \text{Id}_V) \circ (\text{Id}_M \otimes \sigma) : M \otimes V \otimes V \rightarrow M, \quad (47)$$

$$\begin{array}{c} M \\ \rho \\ \rho \\ \text{---} \\ MVV \end{array} = \begin{array}{c} M \\ \rho \\ \rho \quad \sigma \\ \text{---} \\ MVV \end{array} .$$

Figure 22: Braided module

We talk about *braided V -modules* when the pre-braiding σ is clear from the context.

- \rightarrow A *left module* is a right one in $\mathcal{C}^{\otimes^{\text{op}}}$.
- \rightarrow A right (or left) *comodule* is a right (resp. left) module in \mathcal{C}^{op} .
- \rightarrow A *braided V -module morphism* is a morphism φ between braided V -modules (M, ρ) and (N, π) such that $\varphi \circ \rho = \pi \circ (\varphi \otimes \text{Id}_V) : M \otimes V \rightarrow N$.

Start as usual with a trivial example: in a preadditive category, any object M equipped with the zero map $M \otimes V \rightarrow M$ is a module over any pre-braided object (V, σ) . We further interpret our new notion in more complicated settings from section 4.

Example 6.2. 1. One recovers the notion of (anti)commuting operators on M when $\sigma = \pm\tau$.

2. Take $\mathcal{C} = \mathbf{Set}$, and as a pre-braiding on a set S take σ_{\triangleleft} from (30), coming from a self-distributive operation \triangleleft . Condition (47) becomes

$$(m \triangleleft a) \triangleleft b = (m \triangleleft b) \triangleleft (a \triangleleft b) \quad \forall m \in M, a, b \in S,$$

which defines precisely a **rack module** (= the *rack-set* from [17], or the *shadow* from [7]), having a knot-theoretical motivation.

3. Any UAA (V, μ, ν) in \mathcal{C} comes with the pre-braiding σ_{Ass} from (44). Take a right module (M, ρ) which we suppose **normalized** here, i.e.

$$\rho \circ (\text{Id}_M \otimes \nu) = \text{Id}_M \tag{48}$$

(morally, “the unit acts by identity”). Condition (47) becomes

$$\rho \circ (\rho \otimes \text{Id}_V) = \rho \circ (\text{Id}_M \otimes \mu),$$

giving the familiar notion of **module over an associative algebra**.

4. Take a ULA $(V, [,], \nu)$ in a symmetric preadditive category \mathcal{C} . Endow V with the braiding σ_{Lei} from (45). Take a normalized right module (M, ρ) . Condition (47) becomes

$$\rho \circ (\rho \otimes \text{Id}_V) = \rho \circ (\rho \otimes \text{Id}_V) \circ (\text{Id}_M \otimes c_{V,V}) + \rho \circ (\text{Id}_M \otimes [,]),$$

giving the familiar notion of **module over a Leibniz algebra** (cf. [23]).

Note that, dually, left modules over associative or left Leibniz algebras are particular cases of left modules over pre-braided objects.

Now, returning to the general monoidal category setting, try a special choice of M :

Lemma 6.3. Take a pre-braided object (V, σ) in \mathcal{C} . For a morphism $\epsilon : V = \mathbf{I} \otimes V = V \otimes \mathbf{I} \rightarrow \mathbf{I}$, the following conditions are equivalent:

1. ϵ defines a right braided V -module \mathbf{I} ;
2. ϵ defines a left braided V -module \mathbf{I} ;
3. ϵ is a braided character.

This observation can be generalized to endow each tensor power of V with a braided V -module structure using a braided character ϵ . Recall notations φ_i from (1).

Proposition 6.4. Given a pre-braided object (V, σ) with a braided character ϵ , the map

$$\epsilon_\pi := \epsilon_1 \circ \sigma_{V^{\otimes n}, V} : V^{\otimes n} \otimes V \rightarrow V^{\otimes n}$$

defines a right braided V -module structure on $V^{\otimes n}$. The braiding σ is extended here to arbitrary powers of V as in remark 2.5.

Proof. The definition of ϵ_π and repeated application of the YBE give

$$\begin{aligned} \epsilon_\pi \circ (\epsilon_\pi \otimes \text{Id}_V) \circ (\text{Id}_n \otimes \sigma) &= \\ (\epsilon \otimes \epsilon \otimes \text{Id}_n) \circ \sigma_{V^n, V^2} \circ (\text{Id}_n \otimes \sigma) &= \\ ((\epsilon \otimes \epsilon) \circ \sigma) \otimes \text{Id}_n \circ \sigma_{V^n, V^2} & \end{aligned}$$

which, by the definition of braided character, is the same as

$$(\epsilon \otimes \epsilon \otimes \text{Id}_n) \circ \sigma_{V^n, V^2} = \epsilon_\pi \circ (\epsilon_\pi \otimes \text{Id}_V).$$

The reader is advised to draw some diagrams to better follow the proof. □

Definition 6.5. We call the modules $(V^{\otimes n}, \epsilon_\pi)$ *adjoint*.

One recognizes the arrow operations ϵ_{π_w} from subsection 3.3. See that subsection for diagrams and some properties. In particular, along the lines of proposition 3.9, one proves

Proposition 6.6. *The action ϵ_{π} on $T(V)$ intertwines the left braided differential ξd for σ -compatible braided characters ϵ and ξ . In other words, ξd is a braided V -module morphism, for the adjoint braided V -module structure on $V^{\otimes n}$.*

The motivation for our term comes from examples, where one recognizes familiar actions on $T(V)$ (cf. table 1).

We have seen that a module over a pre-braided object is a generalization of a braided character. Observe that this generalization picks the right property for a generalized version of theorem 5 (where we replace the braided V -module \mathbf{I} by arbitrary braided modules) to hold:

Theorem 5^{coeffs}. *Let $(\mathcal{C}, \otimes, \mathbf{I})$ be a preadditive monoidal category, (V, σ) a pre-braided object in \mathcal{C} , and (M, ρ) and (N, λ) a right and a left braided V -modules respectively. Then*

$$\begin{aligned} ({}^{\rho}d)_n &:= (\rho \otimes \text{Id}_{n-1} \otimes \text{Id}_N) \circ (\text{Id}_M \otimes \Theta_{-\sigma}^{1, n-1} \otimes \text{Id}_N), \\ (d^{\lambda})_n &:= (-1)^{n-1} (\text{Id}_M \otimes \text{Id}_{n-1} \otimes \lambda) \circ (\text{Id}_M \otimes \Theta_{-\sigma}^{n-1, 1} \otimes \text{Id}_N), \end{aligned}$$

define a bidegree -1 tensor bidifferential for V with coefficient in M and N .

The complicated expression a bidegree -1 tensor bidifferential for V with coefficient in M and N hides what one naturally expects: two families of morphisms $d_n, d'_n : M \otimes V^n \otimes N \rightarrow M \otimes V^{n-1} \otimes N$ satisfying (43).

Pictorially, $({}^{\rho}d)_n$ is for example a signed sum of terms of the form




Figure 23: Braided differentials with coefficients

The proof of this result is a direct generalization of that of theorem 5. Moreover, all the remaining points of that theorem can be generalized to “coefficient” settings.

Remark 6.7. Taking as M or N the unit object \mathbf{I} with the zero module structure, one obtains a degree -1 tensor differential for V with coefficient in the left braided V -module N (resp. right braided V -module M) only.

As usual, everything described here can be dualized, in any of the three senses described in subsections 5.4 and 5.5.

Adjoint modules also admit a version with coefficients:

Proposition 6.8. *Given a pre-braided object (V, σ) and a right braided V -module (M, ρ) , the morphisms*

$${}^{\rho}\pi := \rho_1 \circ (\text{Id}_M \otimes \sigma_{V^n, V}) : M \otimes V^{\otimes n} \otimes V \rightarrow M \otimes V^{\otimes n}$$

define a right braided V -module structure on $M \otimes V^{\otimes n}$, intertwining the left differential ${}^{\rho}d$.

In other words, ${}^{\rho}d$ is a braided V -module morphism.

Having the Hochschild homology in mind, one also needs the notion of bimodules.

Definition 6.9. A *bimodule* over a pre-braided object (V, σ) is an object $M \in \text{Ob}(\mathcal{C})$ equipped with two morphisms $\rho : M \otimes V \rightarrow M$ and $\lambda : V \otimes M \rightarrow M$, turning M into a right and left modules respectively and satisfying the following compatibility condition:

$$\rho \circ (\lambda \otimes \text{Id}_V) = \lambda \circ (\text{Id}_V \otimes \rho) : V \otimes M \otimes V \rightarrow M.$$

Another interpretation of bimodules – in terms of modules over appropriate pre-braided systems – is given in [20].

The bidifferential structure from theorem 5^{coeffs} can be nicely adapted to bimodules:

Proposition 6.10. *Let $(\mathcal{C}, \otimes, \mathbf{I}, c)$ be a symmetric preadditive category, (V, σ) a pre-braided object in \mathcal{C} , and (M, ρ, λ) a bimodule over V . Then the families of morphisms*

$$\begin{aligned} ({}^\rho d)_n &:= (\rho \otimes \text{Id}_{n-1}) \circ (\text{Id}_M \otimes \Theta_{-\sigma}^{1, n-1}), \\ (d^\lambda)_n &:= (-1)^{n-1} c_{M, V^{n-1}}^{-1} \circ (\text{Id}_{n-1} \otimes \lambda) \circ (\Theta_{-\sigma}^{n-1, 1} \otimes \text{Id}_M) \circ c_{M, V^n}, \end{aligned}$$

define a bidegree -1 tensor bidifferential for V with coefficients in M on the left.

By definition, $(d^\lambda)_n$ is a signed sum of terms of the form




Figure 24: Braided differentials with bimodule coefficients

Proof. Relations $({}^\rho d)_{n-1} \circ ({}^\rho d)_n = 0$ and

$$\begin{aligned} (d^\lambda)_{n-1} \circ (d^\lambda)_n &= c_{M, V^{n-2}}^{-1} \circ (d'^\lambda)_{n-1} \circ c_{M, V^{n-1}} \circ c_{M, V^{n-1}}^{-1} \circ (d'^\lambda)_n \circ c_{M, V^n} \\ &= c_{M, V^{n-2}}^{-1} \circ (d'^\lambda)_{n-1} \circ (d'^\lambda)_n \circ c_{M, V^n} = 0, \end{aligned}$$

with $(d'^\lambda)_n := (-1)^{n-1} (\text{Id}_{n-1} \otimes \lambda) \circ (\Theta_{-\sigma}^{n-1, 1} \otimes \text{Id}_M)$, follow directly from the corresponding identities in theorem 5^{coeffs}. To prove the compatibility between $({}^\rho d)_n$ and $(d^\lambda)_n$, observe that

$$(d^\lambda)_n = (-1)^{n-1} ((\lambda \circ c_{M, V}) \otimes \text{Id}_{n-1}) \circ (\text{Id}_M \otimes c_{V, V^{n-1}}^{-1}) \circ (\text{Id}_M \otimes \Theta_{-\sigma}^{n-1, 1}),$$

then use the defining property of bimodule, the naturality of c and the YBE for σ . \square

Remark 6.11. We have kept the notation c^{-1} , redundant for a symmetric c , to be able to treat the non symmetric situation. In this case, on the picture showing $(d^\lambda)_n$, the thick line (corresponding to M) should go behind all normal lines, in order to distinguish c from c^{-1} . One should be careful to differentiate the two pre-braidings, c and σ , which is difficult to do pictorially. For the above theorem to be still valid, one should change the compatibility condition defining a bimodule to the following one, different from the old one in general:

$$\lambda \circ (\text{Id}_V \otimes \rho) \circ c_{M \otimes V, V} = \rho \circ (\lambda \otimes \text{Id}_V) \circ c_{M, V} \circ c_{V, V}^{-1} : M \otimes V \otimes V \rightarrow M,$$




Figure 25: Bimodules in the non symmetric case

All the crossings correspond to the braiding c here.

A more elegant solution for the non symmetric case would be welcome.

We finish this section with examples, interpreting several classical homology theories with coefficients as braided ones:

- Example 6.12.** 1. Taking a vector space V with a simple flip as a braiding and, for instance, its symmetric algebra $S(V)$ as a module over V (with the action coming from concatenation, as usual), one obtains more complicated versions of the Koszul complex.
2. In the case of shelves, one recovers the *shelf and rack homologies with coefficients*, hinted at in [7].
3. For Leibniz algebras, our machinery gives the *Leibniz homology with coefficients*, generalizing the Chevalley-Eilenberg homology (cf. [23]).
- In these three cases one generally puts the coefficients only on the left (cf. remark 6.7).
4. Coefficients on both sides turn out to be particularly useful for associative algebras in a symmetric preadditive category. In this setting, proposition 6.10 gives the following differential for an algebra bimodule (M, ρ, λ) :

$$\begin{aligned}
 ({}^\rho d - d^\lambda)_n &:= \rho_1 + \sum_{i=1}^{n-1} (-1)^i \mu_{i+1} + (-1)^n \lambda_1 \circ c_{M \otimes V^{n-1}, V} \\
 &\quad + \text{some terms involving } \nu.
 \end{aligned}$$

For $\mathcal{C} = \mathbf{Vect}_{\mathbb{k}}$, one can get rid of the terms with ν as it was done in the proof of point 7 of proposition 4.25, getting the *Hochschild differential*.

5. The co-version of the previous differential is the *Cartier differential* for coalgebras (cf. [6], where it was first introduced).

Acknowledgements

I would like to thank Marc Rosso for sharing his passion for quantum shuffles and for his patient encouragements, and Józef Przytycki for introducing me to his work on distributive homology. I am also grateful to Paul-André Melliès for his interest in my work, and to Arnaud Mortier for comments on a preliminary version of this paper. My deep gratitude goes to Jean-Louis Loday whom I have never had the chance to meet in person, but whose mathematics I have always admired.

References

- [1] John C. Baez. Hochschild homology in a braided tensor category. *Trans. Amer. Math. Soc.*, 344(2):885–906, 1994.
- [2] J. Scott Carter, Alissa S. Crans, Mohamed Elhamdadi, and Masahico Saito. Cohomology of categorical self-distributivity. *J. Homotopy Relat. Struct.*, 3(1):13–63, 2008.
- [3] J. Scott Carter, Mohamed Elhamdadi, and Masahico Saito. Twisted quandle homology theory and cocycle knot invariants. *Algebr. Geom. Topol.*, 2:95–135 (electronic), 2002.
- [4] J. Scott Carter, Mohamed Elhamdadi, and Masahico Saito. Homology theory for the set-theoretic Yang-Baxter equation and knot invariants from generalizations of quandles. *Fund. Math.*, 184:31–54, 2004.
- [5] J. Scott Carter, Daniel Jelsovsky, Seiichi Kamada, Laurel Langford, and Masahico Saito. Quandle cohomology and state-sum invariants of knotted curves and surfaces. *Trans. Amer. Math. Soc.*, 355(10):3947–3989, 2003.
- [6] Pierre Cartier. Cohomologie des coalgèbres. *Sém. Sophus Lie*, 5, 1955-1956.
- [7] Wesley Chang and Sam Nelson. Rack shadows and their invariants. *J. Knot Theory Ramifications*, 20(9):1259–1269, 2011.

- [8] Alissa Susan Crans. *Lie 2-algebras*. ProQuest LLC, Ann Arbor, MI, 2004. Thesis (Ph.D.)—University of California, Riverside.
- [9] Christian Cuvier. Homologie de Leibniz et homologie de Hochschild. *C. R. Acad. Sci. Paris Sér. I Math.*, 313(9):569–572, 1991.
- [10] Yukio Doi. Homological coalgebra. *J. Math. Soc. Japan*, 33(1):31–50, 1981.
- [11] A. S. Dzhumadil'daev. Cohomologies of colour Leibniz algebras: pre-simplicial approach. In *Lie theory and its applications in physics, III (Clausthal, 1999)*, pages 124–136. World Sci. Publ., River Edge, NJ, 2000.
- [12] Michael Eisermann. Yang-Baxter deformations of quandles and racks. *Algebr. Geom. Topol.*, 5:537–562 (electronic), 2005.
- [13] Roger Fenn, Colin Rourke, and Brian Sanderson. Trunks and classifying spaces. *Appl. Categ. Structures*, 3(4):321–356, 1995.
- [14] I. Goyvaerts and J. Vercauteren. A Note on the categorification of Lie algebras. *ArXiv e-prints*, February 2012.
- [15] David Joyce. A classifying invariant of knots, the knot quandle. *J. Pure Appl. Algebra*, 23(1):37–65, 1982.
- [16] Seiichi Kamada. Knot invariants derived from quandles and racks. In *Invariants of knots and 3-manifolds (Kyoto, 2001)*, volume 4 of *Geom. Topol. Monogr.*, pages 103–117 (electronic). Geom. Topol. Publ., Coventry, 2002.
- [17] Seiichi Kamada. Quandles with good involutions, their homologies and knot invariants. In *Intelligence of low dimensional topology 2006*, volume 40 of *Ser. Knots Everything*, pages 101–108. World Sci. Publ., Hackensack, NJ, 2007.
- [18] Christian Kassel and Vladimir Turaev. *Braid groups*, volume 247 of *Graduate Texts in Mathematics*. Springer, New York, 2008. With the graphical assistance of Olivier Dodane.
- [19] Louis H. Kauffman. Virtual knot theory. *European J. Combin.*, 20(7):663–690, 1999.
- [20] Victoria Lebed. *Braided Objects: Unifying Algebraic Structures and Categorifying Virtual Braids*. 2012. Thesis (Ph.D.)—Université Paris 7.
- [21] Victoria Lebed. Symmetric Categories as a Means of Virtualization for Braid Groups and Racks. *ArXiv e-prints*, June 2012.
- [22] Dong Liu and Naihong Hu. Leibniz superalgebras and central extensions. *J. Algebra Appl.*, 5(6):765–780, 2006.
- [23] Jean-Louis Loday. *Cyclic homology*, volume 301 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1992. Appendix E by María O. Ronco.
- [24] Jean-Louis Loday. Une version non commutative des algèbres de Lie: les algèbres de Leibniz. *Enseign. Math. (2)*, 39(3-4):269–293, 1993.
- [25] Jean-Louis Loday and Teimuraz Pirashvili. Universal enveloping algebras of Leibniz algebras and (co)homology. *Math. Ann.*, 296(1):139–158, 1993.
- [26] Saunders MacLane. *Categories for the working mathematician*. Springer-Verlag, New York, 1971. Graduate Texts in Mathematics, Vol. 5.
- [27] S. Majid. Free braided differential calculus, braided binomial theorem, and the braided exponential map. *J. Math. Phys.*, 34(10):4843–4856, 1993.

- [28] Shahn Majid. Algebras and Hopf algebras in braided categories. In *Advances in Hopf algebras (Chicago, IL, 1992)*, volume 158 of *Lecture Notes in Pure and Appl. Math.*, pages 55–105. Dekker, New York, 1994.
- [29] Shahn Majid. *Foundations of quantum group theory*. Cambridge University Press, Cambridge, 1995.
- [30] M. Niebrzydowski and J. H. Przytycki. Homology operations on homology of quandles. *J. Algebra*, 324(7):1529–1548, 2010.
- [31] Józef H. Przytycki. Distributivity versus associativity in the homology theory of algebraic structures. *Demonstratio Math.*, 44(4):823–869, 2011.
- [32] Józef H. Przytycki and Adam S. Sikora. Distributive products and their homology. *ArXiv e-prints*, May 2011.
- [33] V. Rittenberg and D. Wyler. Generalized superalgebras. *Nuclear Phys. B*, 139(3):189–202, 1978.
- [34] Marc Rosso. Groupes quantiques et algèbres de battage quantiques. *C. R. Acad. Sci. Paris Sér. I Math.*, 320(2):145–148, 1995.
- [35] Marc Rosso. Integrals of vertex operators and quantum shuffles. *Lett. Math. Phys.*, 41(2):161–168, 1997.
- [36] Marc Rosso. Quantum groups and quantum shuffles. *Invent. Math.*, 133(2):399–416, 1998.
- [37] V. G. Turaev. *Quantum invariants of knots and 3-manifolds*, volume 18 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1994.
- [38] Vladimir V. Vershinin. On homology of virtual braids and Burau representation. *J. Knot Theory Ramifications*, 10(5):795–812, 2001. Knots in Hellas '98, Vol. 3 (Delphi).
- [39] Quinton Westrich. Lie algebras in braided monoidal categories. *Preprint at <http://www.scribd.com/QuintonWestrich>*, 2006.