



HAL
open science

Constructing the set of complete intersection numerical semigroups with a given Frobenius number

Abdallah Assi, Pedro A. García-Sánchez

► **To cite this version:**

Abdallah Assi, Pedro A. García-Sánchez. Constructing the set of complete intersection numerical semigroups with a given Frobenius number. 2012. hal-00688929v3

HAL Id: hal-00688929

<https://hal.science/hal-00688929v3>

Preprint submitted on 19 Jan 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONSTRUCTING THE SET OF COMPLETE INTERSECTION NUMERICAL SEMIGROUPS WITH A GIVEN FROBENIUS NUMBER

A. ASSI AND P. A. GARCÍA-SÁNCHEZ

ABSTRACT. Delorme suggested that the set of all complete intersection numerical semigroups can be computed recursively. We have implemented this algorithm, and particularized it to several subfamilies of this class of numerical semigroups: free and telescopic numerical semigroups, and numerical semigroups associated to an irreducible plane curve singularity. The recursive nature of this procedure allows us to give bounds for the embedding dimension and for the minimal generators of a semigroup in any of these families.

1. INTRODUCTION

Let \mathbb{N} denote the set of nonnegative integers. A *numerical semigroup* Γ is a submonoid of \mathbb{N} with finite complement in \mathbb{N} (this condition is equivalent to $\gcd(\Gamma) = 1$). If Γ is a numerical semigroup, the elements in $\mathbb{N} \setminus \Gamma$ are the *gaps* of Γ . The cardinality of $\mathbb{N} \setminus \Gamma$ is the *genus* of Γ , $g(\Gamma)$. The largest integer not in Γ is called the *Frobenius number* of Γ , and will be denoted by $F(\Gamma)$. Clearly, $F(\Gamma) + 1 + \mathbb{N} \subseteq \Gamma$, and this is why $c(\Gamma) = F(\Gamma) + 1$ is known as the *conductor* of Γ .

Since for every $x \in \Gamma$, $F(\Gamma) - x$ cannot be in Γ , we deduce that $g(\Gamma) \geq \frac{c(\Gamma)}{2}$. We say that Γ is *symmetric* when the equality holds, or equivalently, for every integer x , $x \notin \Gamma$ implies $F(\Gamma) - x \in \Gamma$. In this setting, $c(\Gamma)$ is an even integer, and thus $F(\Gamma)$ is odd.

It can be easily proved that any numerical semigroup admits a unique *minimal generating system* (every element is a linear combination of elements in this set with nonnegative integer coefficients and none of its proper subsets fulfills this condition; see for instance [16, Chapter 1]). If $A = \{r_0, \dots, r_h\}$ is the minimal generating set of Γ , then its elements are called *minimal generators*, and its cardinality is the *embedding dimension* of Γ , $e(\Gamma)$. The smallest minimal generator is the smallest positive integer belonging to the semigroup, and it is known as the *multiplicity* of Γ , denoted by $m(\Gamma)$.

The map

$$\mathbb{N}^{e(\Gamma)} \rightarrow \Gamma, \varphi(a_0, \dots, a_h) = a_0 r_0 + \dots + a_h r_h$$

is a monoid epimorphism. Hence Γ is isomorphic to $\mathbb{N}^{e(\Gamma)} / \ker \varphi$, where $\ker \varphi = \{(a, b) \in \mathbb{N}^{e(\Gamma)} \times \mathbb{N}^{e(\Gamma)} \mid \varphi(a) = \varphi(b)\}$ ($\ker \varphi$ is a congruence on $\mathbb{N}^{e(\Gamma)}$).

A *presentation* for Γ is a set of generators of the congruence φ , and a *minimal presentation* is a set of generators minimal with respect to set inclusion (actually, in our setting also with respect to cardinality; see [16, Corollary 8.13]). It can be shown that the cardinality of any minimal presentation is greater than or equal to $e(\Gamma) - 1$, [16, Theorem 9.6]. A numerical semigroup is a *complete intersection* if this equality holds.

Given A a set positive integers, and $A = A_1 \cup A_2$ a non trivial partition of A , we say that A is the *gluing* of A_1 and A_2 if $\text{lcm}(d_1, d_2) \in \langle A_1 \rangle \cap \langle A_2 \rangle$, where $d_i = \gcd(A_i)$ and $\langle A_i \rangle$ denotes the monoid generated by A_i , $i = 1, 2$. If A is the minimal system of generators of Γ , and Γ_i is the numerical semigroup generated by A_i/d_i , $i = 1, 2$, we also say that Γ is the *gluing* of Γ_1 and

The first author is partially supported by the project GDR CNRS 2945.

The second author is supported by the projects MTM2010-15595, FQM-343, FQM-5849, and FEDER funds. This research was performed while the second author visited the Université d'Angers as invited lecturer, and he wants to thank the Département de Mathématiques of this university for its kind hospitality. The authors would like to thank the referees for their suggestions, comments and examples provided.

Γ_2 . It turns out that $d_1 \in \Gamma_2$, $d_2 \in \Gamma_1$, $\gcd(d_1, d_2) = 1$, and neither d_1 is a minimal generator of Γ_2 nor d_2 is a minimal generator of Γ_1 ([16, Section 8.3]). Delorme proved in [5, Proposition 9] that a numerical semigroup is a complete intersection if and only if it is a gluing of two complete intersection numerical semigroups (though with a different notation; the concept of gluing was introduced in [13]). The gluing of symmetric numerical semigroups is symmetric ([5, Proposition 10 (iii)]), and as a consequence of this, complete intersections are symmetric.

In [17] there is a procedure to construct the set of all numerical semigroups with a given Frobenius number. We show in this manuscript how can we use the concept of gluing to compute the set of all complete intersection numerical semigroups with a given Frobenius number (or equivalently with fixed genus). Recently there have been some experimental results that point out to the possibility that the number of numerical semigroups with a fixed genus has a Fibonacci like behaviour ([4]). Indeed, it is known that asymptotically the number of numerical semigroups with given genus grows as the Fibonacci sequence ([18]). However there is not a proof for this for all genus, and we still do not even have a demonstration that there are more numerical semigroups with genus $g + 1$ than numerical semigroups with genus g . This is not the case for complete intersection numerical semigroups, as we see in the last section.

We also show how to calculate the set of all free (in the sense of [1]) numerical semigroups, which is a special subclass of complete intersections, the set of all telescopic numerical semigroups (contained in the set of free numerical semigroups), and that of numerical semigroups associated to an irreducible plane curve singularity (these are a particular case of telescopic numerical semigroups).

The recursive nature of gluing also allows us to give some bounds for the generators and embedding dimension for these families of semigroups when we fix the Frobenius number. The deeper we go in the chain of inclusions given in the preceding paragraph, the smaller are the bounds.

2. THE FROBENIUS NUMBER AND MULTIPLICITY OF A COMPLETE INTERSECTION

Let Γ be a numerical semigroup. We know that Γ is a complete intersection if and only if it is the gluing of two complete intersections. Delorme (though with a different notation) highlighted in [5, Section 11] that this fact can be used to determine if a numerical semigroup is a complete intersection (this idea has already been exploited in [2]; and in [15] one can find a procedure to determine if an affine semigroup is the gluing of two affine semigroups), and also to compute the set of all complete intersections. In order to construct the set of all complete intersection numerical semigroups with given Frobenius number, we can proceed recursively by using the following formula for the Frobenius number of the gluing of two numerical semigroups, which is just a reformulation of Delorme's description of the conductor of a gluing.

Proposition 1. *Assume that Γ is a numerical semigroup minimally generated by $A = A_1 \cup A_2$, and that A is the gluing of A_1 and A_2 . Let $d_1 = \gcd(A_1)$ and $d_2 = \gcd(A_2)$. Define $\Gamma_1 = \langle A_1/d_1 \rangle$ and $\Gamma_2 = \langle A_2/d_2 \rangle$. Then*

$$F(\Gamma) = d_1 F(\Gamma_1) + d_2 F(\Gamma_2) + d_1 d_2.$$

Proof. Observe that $\Gamma = d_1 \Gamma_1 + d_2 \Gamma_2$. By [5, Proposition 10 (i)],

$$(1) \quad c(\Gamma) = d_1 c(\Gamma_1) + d_2 c(\Gamma_2) + (d_1 - 1)(d_2 - 1).$$

Having in mind the relationship between Frobenius number and conductor, the formula follows easily. \square

In Proposition 1, $\Gamma = d_1 \Gamma_1 + d_2 \Gamma_2$ and $d = d_1 d_2 = d_1 \Gamma_1 \cap d_2 \Gamma_2$. The integer d is the element where the gluing takes place. If we repeat the process with $d_1 \Gamma_1$ and $d_2 \Gamma_2$ in this result, we construct a decomposition tree of Γ , whose leaves are copies isomorphic to \mathbb{N} (this was the idea followed in [3]). Assume that $d^{(1)}, \dots, d^{(h)}$ are the elements where the gluings take place in this splitting. The Frobenius number of Γ is precisely $\sum_{i=1}^h d^{(i)} - \sum_{a \in A} a$ (see [5, Section 11] where it is highlighted that this formula is a particular case of a result given in [10]).

Example 2. Let $\Gamma = \langle 10, 14, 15, 21 \rangle$. Then $\Gamma = \langle 10, 15 \rangle + \langle 14, 21 \rangle$ and $35 = 5 \times 7 \in \langle 10, 15 \rangle$. We repeat the process for $\langle 10, 15 \rangle = \langle 10 \rangle + \langle 15 \rangle$ and $\langle 14, 21 \rangle = \langle 14 \rangle + \langle 21 \rangle$. We get $30 \in \langle 10 \rangle \cap \langle 15 \rangle$ and $42 \in \langle 14 \rangle \cap \langle 21 \rangle$. Hence the gluings take place at 35, 30 and 42. Thus $F(\Gamma) = (35 + 30 + 42) - (10 + 14 + 15 + 21) = 47$.

Example 3. We construct a complete intersection numerical semigroup with four generators, by gluing two embedding dimension two numerical semigroups.

```
gap> s:=NumericalSemigroup(10,11);;
gap> t:=NumericalSemigroup(7,9);;
gap> g:=NumericalSemigroup(16*10,16*11,21*7,21*9);;
gap> FrobeniusNumber(g);
2747
gap> 16*FrobeniusNumber(s)+21*FrobeniusNumber(t)+16*21;
2747
```

Remark 4. For $\Gamma = \mathbb{N}$, we have

$$c(\mathbb{N}) = 0, \quad F(\mathbb{N}) = -1, \quad g(\mathbb{N}) = 0, \quad m(\mathbb{N}) = 1, \quad e(\mathbb{N}) = 1.$$

Proposition 5. *If Γ is a complete intersection, then*

$$m(\Gamma) \geq 2^{e(\Gamma)-1}.$$

Proof. Let $h = e(\Gamma) - 1$. We use induction on h . For $h = 1$, the statement follows trivially. As Γ is a complete intersection, if A is its minimal set of generators, we can find a partition of $A = A_1 \cup A_2$ such that A is the gluing of A_1 and A_2 . Set as above $d_i = \gcd(A_i)$, and $\Gamma_i = \langle A_i/d_i \rangle$. Let $h_i = e(\Gamma_i) - 1$. Hence $h = h_1 + h_2 + 1$. By induction hypothesis $m(\Gamma_i) \geq 2^{h_i}$. Recall that $d_1 \in \Gamma_2$ and $d_2 \in \Gamma_1$, and they are not minimal generators. Thus $d_1 \geq 2m(\Gamma_2) \geq 2^{h_2+1}$, and analogously $d_2 \geq 2^{h_1+1}$. For every $a \in A_1$, a/d_1 is a minimal generator of Γ_1 , whence $a/d_1 \geq 2^{h_1}$. Therefore $a \geq 2^{h_1+h_2+1} = 2^h$. The same argument shows that any element in A_2 is greater than or equal to 2^h . \square

Example 6. We construct recursively a family $\{\Gamma^{(n)}\}_{n \in \mathbb{N}}$ of complete intersection numerical semigroups reaching the bound of Proposition 5.

We start with $\Gamma^{(1)} = \langle 2, 3 \rangle$, and the general element in the sequence is defined as $\Gamma^{(n+1)} = 2\Gamma^{(n)} + (2^{n+1} + 1)\mathbb{N}$.

For instance, $\Gamma^{(2)} = 2\langle 2, 3 \rangle + 5\mathbb{N} = \langle 4, 5, 6 \rangle$, $\Gamma^{(3)} = 2\langle 4, 5, 6 \rangle + 9\mathbb{N} = \langle 8, 9, 10, 12 \rangle$, and so on.

It is not hard to prove that

$$\begin{aligned} \Gamma^{(n+1)} &= \langle 2^{n+1}, 2^{n+1} + 1, 2^{n+1} + 2, 2^{n+1} + 2^2, \dots, 2^{n+1} + 2^n \rangle \\ &= 2\langle 2^n, 2^n + 1, \dots, 2^n + 2^{n-1} \rangle + (2^{n+1} + 1)\mathbb{N}. \end{aligned}$$

Notice that $\Gamma^{(n+1)}$ is a gluing of $\Gamma^{(n)}$ and \mathbb{N} , since

- $2 \in \mathbb{N}$ and 2 is not a minimal generator of \mathbb{N} ,
- $2^{n+1} + 1$ is the sum of the two smallest minimal generators of $\Gamma^{(n)}$; thus $2^{n+1} + 1$ belongs to $\Gamma^{(n)}$ and it is not a minimal generator of $\Gamma^{(n)}$,
- $\gcd(2, 2^{n+1} + 1) = 1$.

It follows that $m(\Gamma^{(n)}) = 2^n$ and $e(\Gamma^{(n)}) = n + 1$. Thus the bound in Proposition 5 is attained.

Corollary 7. *If Γ is a complete intersection numerical semigroup other than \mathbb{N} , then*

$$e(\Gamma) \leq \log_2(c(\Gamma)) + 1.$$

Proof. By Proposition 5, $2^{e(\Gamma)-1} \leq m(\Gamma)$. Since $\Gamma \neq \mathbb{N}$, we have that $m(\Gamma) \leq c(\Gamma)$, and the bound follows. \square

Remark 8. Notice that in the proof of Corollary 7 we use $m(\Gamma) \leq c(\Gamma)$. For $\Gamma = \langle 2, 3 \rangle$, we get an equality and also the bound given in this corollary is reached. If $m(\Gamma) = c(\Gamma)$, then $\Gamma = \langle m, m+1, \dots, 2m-1 \rangle$, with $m = m(\Gamma)$. Hence $e(\Gamma) = m$, that is, Γ has maximal embedding dimension (it is easy to see that the embedding dimension of a numerical semigroup is always less than or equal to its multiplicity; see for instance [16, Chapter 1]). It is well known that the cardinality of a minimal presentation of a maximal embedding dimension numerical semigroup with multiplicity m is $\frac{m(m-1)}{2}$ (see for instance [16, Corollary 8.27]). Hence a maximal embedding dimension numerical semigroup with multiplicity m is a complete intersection if and only if $\frac{m(m-1)}{2} = m-1$, or equivalently, either the numerical semigroup is \mathbb{N} or $m = 2$. If in addition we impose that the conductor and the multiplicity agree, then the only two possibilities are \mathbb{N} and $\langle 2, 3 \rangle$.

From the definitions of multiplicity and conductor, it is easy to see that there is no numerical semigroup Γ such that $c(\Gamma) = 1 + m(\Gamma)$.

If $c(\Gamma) = 2 + m(\Gamma)$, then $\Gamma = \langle m, m+2, m+3, \dots, 2m-1, 2m+1 \rangle$, which is a maximal embedding dimension numerical semigroup. So the only complete intersection with $c(\Gamma) = 2 + m(\Gamma)$ is $\langle 2, 5 \rangle$.

The case $c(\Gamma) = 3 + m(\Gamma)$ requires more effort. In this setting $m = m(\Gamma) > 2$. We have two possibilities.

- $\Gamma = \langle m, m+3, m+4, \dots, 2m-1, 2m+1, 2m+2 \rangle$, which has maximal embedding dimension, and so it cannot be a complete intersection numerical semigroup, because $m > 2$.
- $\Gamma = \langle m, m+1, m+3, m+4, \dots, 2m-1 \rangle$. Here $e(\Gamma) = m-1$ and the minimum element in Γ congruent with 2 modulo m is $2m+1 = (m+1) + (m+1)$. Thus in view of [14, Theorem 1(2)], the cardinality of a minimal presentation for Γ is $\frac{(m-1)(m-2)}{2}$. We conclude that Γ is a complete intersection if and only if $\frac{(m-1)(m-2)}{2} = m-2$, and as $m > 2$, this is equivalent to $m = 3$. Hence $\Gamma = \langle 3, 4 \rangle$.

Therefore, if we assume that $\Gamma \notin \{\mathbb{N}, \langle 2, 3 \rangle, \langle 2, 5 \rangle, \langle 3, 4 \rangle\}$, and Γ is a complete intersection numerical semigroup, then we can assert that $c(\Gamma) \geq m(\Gamma) + 4$, and the bound in Corollary 7 can be slightly improved to

$$e(\Gamma) \leq \log_2(c(\Gamma) - 4) + 1.$$

This bound is attained for instance by $\langle 2, 7 \rangle$, $\langle 4, 5, 6 \rangle$ and $\langle 4, 6, 7 \rangle$.

By using [14, Section 1.2], we can determine those complete intersections with $c(\Gamma) = m(\Gamma) + 4$, and thus obtain another small improvement of the above bound.

We can improve this bound by using a different strategy.

Proposition 9. *Let Γ be a complete intersection numerical semigroup. Then*

$$(e(\Gamma) - 1)2^{e(\Gamma)-1} \leq c(\Gamma).$$

Proof. We use induction on the embedding dimension of Γ . If the embedding dimension of Γ is either one or two, then the result holds trivially. So assume that $e(\Gamma) \geq 3$. As Γ is a complete intersection, we know that there exist two complete intersection numerical semigroups Γ_1 and Γ_2 such that Γ is the gluing of Γ_1 and Γ_2 . Thus there exist $d_1 \in \Gamma_2$ and $d_2 \in \Gamma_1$, that are not minimal generators, such that $\Gamma = d_1\Gamma_1 + d_2\Gamma_2$. For sake of simplicity write $c = c(\Gamma)$, $e = e(\Gamma)$, $c_i = c(\Gamma_i)$ and $e_i = e(\Gamma_i)$, $i = 1, 2$. Then from the definition of gluing we already know that $e = e_1 + e_2$.

Since $e \geq 3$, we may assume without loss of generality that $e_1 \geq 2$. As d_1 is not a minimal generator of Γ_2 , $d_1 \geq 2m(\Gamma_2)$, and as $e_1 \geq 2$ and d_2 is not a minimal generator of Γ_1 , $d_2 \geq 2m(\Gamma_1) + 1$. In view of Proposition 5, we deduce $d_1 \geq 2^{e_2}$ and $d_2 \geq 2^{e_1} + 1$.

Now from (1), we have $c = d_1c_1 + d_2c_2 + (d_1 - 1)(d_2 - 1)$. By induction hypothesis and the preceding paragraph, we get $c \geq 2^{e_2}(e_1 - 1)2^{e_1-1} + (2^{e_1} + 1)(e_2 - 1)2^{e_2-1} + (2^{e_2} - 1)2^{e_1} = (e - 2)2^{e-1} + (e_2 - 1)2^{e_2-1} + 2^e - 2^{e_1} \geq (e - 1)2^{e-1} - 2^{e-1} + 2^e - 2^{e_1} = (e - 1)2^{e-1} + 2^{e-1} - 2^{e_1} \geq (e - 1)2^{e-1}$. \square

Example 10. Let $\{\Gamma^{(n)}\}_{n \in \mathbb{N}}$ be the family of numerical semigroups presented in Example 6. By using (1), it is not hard to check inductively that $c(\Gamma^{(n)}) = n2^n$, and thus the bound of Proposition 9 is attained.

If we have a closer look at the proof of Proposition 9, then we easily deduce that for the bound to be attained, the following must hold in all induction steps with $e \geq 3$:

- $(e_2 - 1)2^{e_2 - 1} = 0$ and thus $e_2 = 1$, that is, Γ_2 is \mathbb{N} (we will study these semigroups in the next section);
- from $e_2 = 1$ it follows that $e_1 = e - 1$ and $2^{e-1} - 2^{e_1} = 0$;
- $m(\Gamma_1) = 2^{e_1 - 1}$ and $d_2 = 2m(\Gamma_1) + 1 = 2^{e_1} + 1$, whence $m(\Gamma_1) + 1 \in \Gamma_1$;
- $c_1 = (e_1 - 1)2^{e_1 - 1} = (e - 2)2^{e-2}$;
- $d_1 = 2$.

Also the only embedding dimension two numerical semigroup for which the equality holds is $\langle 2, 3 \rangle$. It follows that the family given in Example 6 contains all possible complete intersection numerical semigroups with the property that the bound in Proposition 9 becomes an equality.

Proposition 11. *Let Γ be a complete intersection numerical semigroup other than \mathbb{N} , minimally generated by $\{r_0, \dots, r_h\}$. If $m(\Gamma) \neq 2$, for all k , $r_k < F(\Gamma)$.*

Proof. Assume without loss of generality that $r_0 = m(\Gamma)$. The numerical semigroup Γ is symmetric and thus for every $i > 0$, $F(\Gamma) + r_0 - r_i \in \Gamma$. If $r_k > F(\Gamma)$, for some $k > 0$, then $F(\Gamma) + r_0 - r_k < r_0$, which forces $F(\Gamma) + r_0 = r_k$.

If $h > 1$, choose $0 < i \neq k$. Then $r_k - r_i = F(\Gamma) + r_0 - r_i \in \Gamma$, contradicting that r_k is a minimal generator. This proves $r_k < F(\Gamma)$, whenever $h > 1$.

For $h = 1$, $F(\Gamma) = (r_0 - 1)(r_1 - 1) - 1$. In this setting, $\Gamma = \langle 2, f + 2 \rangle$ has $F(\Gamma) = f$. For $m(\Gamma) > 2$, we get $F(\Gamma) = (r_0 - 1)(r_1 - 1) - 1 \geq 2(r_1 - 1) - 1 = (r_1 - 1) + (r_1 - 2) \geq r_1$. \square

Remark 12. If we want to compute the set of all complete intersection numerical semigroups with Frobenius number f , then we can use the formula given in Proposition 1. Hence $f = d_1 f_1 + d_2 f_2 + d_1 d_2$, and we recursively construct all possible numerical semigroups with Frobenius number f_1 , and then the set with Frobenius number f_2 . We next give some useful bounds and facts to perform this task. Denote $f + 1$ by c .

- i) $d_1 \neq 1 \neq d_2$. This is because $d_1 \in \Gamma_2$ and it is not a minimal generator of Γ_2 . The only possibility to have $d_1 = 1 \in \Gamma_2$ would be $\Gamma_2 = \mathbb{N} = \langle 1 \rangle$. But then d_1 would be a minimal generator. The same argument is valid for d_2 .
- ii) Since $\gcd(d_1, d_2) = 1$, we can assume without loss of generality that $2 \leq d_2 < d_1$.
- iii) Since $f_1, f_2 \geq -1$, $f \geq -d_1 - d_2 + d_1 d_2 = (d_1 - 1)(d_2 - 1) - 1$. Hence $d_2 \leq \frac{c}{d_1 - 1} + 1$; and consequently, $d_2 \leq \min\{d_1 - 1, \frac{c}{d_1 - 1} + 1\}$.
- iv) $f - d_j f_j \equiv 0 \pmod{d_i}$, $\{i, j\} = \{1, 2\}$. In particular, if $f_j = -1$, then $f + d_j \equiv 0 \pmod{d_i}$.
- v) $d_1 < f$, except in the case $\Gamma = \langle 2 = d_2, f + 2 = d_1 \rangle$.
 - a) If $f_1 = f_2 = -1$, then $\Gamma_1 = \Gamma_2 = \mathbb{N}$, and Γ is $\langle d_2, d_1 \rangle$. If $d_2 \neq 2$, then Proposition 11, asserts that $d_1 < f$.
 - b) If $f_2 > 0$, then $f \geq -d_1 + d_2 + d_1 d_2 = (d_1 + 1)(d_2 - 1) + 1 \geq d_1 + 2$. Hence $d_1 \leq f - 2$.
 - c) If $f_1 > 0$, then $f \geq d_1 - d_2 + d_1 d_2 = (d_1 - 1)(d_2 + 1) + 1 > 3(d_1 - 1) \geq d_1 + 2(d_1 - 1) - 1 > d_1$.
- vi) If $f_1 \neq -1 \neq f_2$, then $f - d_1 d_2 \in \langle d_1, d_2 \rangle$. We are only interested in factorizations $f - d_1 d_2 = a_1 d_1 + a_2 d_2$, $a_1, a_2 \in \mathbb{N}$, with $a_1 \equiv a_2 \equiv 1 \pmod{2}$, since the Frobenius number of a complete intersection is an odd integer.

Example 13. We compute the set of all complete intersection numerical semigroups with Frobenius number 11. First note that $\langle 2, 13 \rangle$ is in this set. The possible d_1 belong to $\{3, \dots, 10\}$.

- $d_1 = 10$. Then $2 \leq d_2 \leq \min\{9, \lfloor \frac{12}{9} \rfloor + 1\} = 2$. Hence d_2 must be 2, but then $\gcd(d_1, d_2) \neq 1$, and we have no complete intersections under these conditions.

- $d_1 = 9$. Then $2 \leq d_2 \leq \min\{8, \lfloor \frac{12}{8} \rfloor + 1\} = 2$. This forces $d_2 = 2$, which in addition is coprime with 9.
 - ★ $11 + 9 \equiv 0 \pmod{2}$, and thus $f_1 = -1$ ($\Gamma_1 = \mathbb{N}$) is a possible choice. In this setting $f_2 = (11 - 18 + 0)/2 = 1$, whence $\Gamma_2 = \langle 2, 3 \rangle$. We obtain a new complete intersection $\Gamma = 9\mathbb{N} + 2\langle 2, 3 \rangle = \langle 4, 6, 9 \rangle$, because $9 \in \langle 2, 3 \rangle$ is not a minimal generator.
 - ★ $11 + 2 \not\equiv 0 \pmod{9}$, so f_2 cannot be -1 .
 - ★ $11 - 18 \notin \langle 2, 9 \rangle$, so we have no more complete intersections with this data.
- For $d_1 = 8$, we have $2 \leq d_2 \leq \min\{7, \lfloor \frac{12}{7} \rfloor + 1\} = 2$. However $\gcd\{d_1, d_2\} \neq 1$.
- If $d_1 = 7$, then $2 \leq d_2 \leq \min\{6, \lfloor \frac{12}{6} \rfloor + 1\} = 3$.
 - ★ $d_2 = 2$.
 - * $11 + 7 \equiv 0 \pmod{2}$, and thus Γ_1 can be \mathbb{N} . But then $f_2 = (11 - 14 + 7)/2 = 2$, which is even. So this case cannot occur.
 - * $11 + 2 \not\equiv 0 \pmod{7}$, and so Γ_2 will not be \mathbb{N} .
 - * Finally, $11 - 14 \notin \langle 2, 7 \rangle$, so no complete intersections can be found with properties.
 - ★ $d_2 = 3$.
 - * $11 + 7 \equiv 0 \pmod{3}$, and thus Γ_1 could be \mathbb{N} . In this setting $f_2 = (11 - 21 + 7)/3 = -1$, and so Γ_2 is also \mathbb{N} . We get a new complete intersection $\Gamma = 7\mathbb{N} + 3\mathbb{N} = \langle 3, 7 \rangle$ with Frobenius number 11.
 - * $11 - 21 \notin \langle 3, 7 \rangle$, so no more complete intersections are obtained for this choice of d_1 and d_2 .
- For $d_1 = 6$, $2 \leq d_2 \leq \min\{5, \lfloor \frac{12}{5} \rfloor + 1\} = 3$, but both 2 and 3 are not coprime with 6.
- $d_1 = 5$. Then $d_2 \in \{2, 3, 4\}$.
 - ★ $d_2 = 2$.
 - * $11 + 5 \equiv 0 \pmod{2}$, and so Γ_1 can possibly be \mathbb{N} . Hence $f_2 = (11 - 10 + 5)/2 = 3$. The only possible complete intersection numerical semigroup with Frobenius number 3 is $\langle 2, 5 \rangle$. But 5 is a minimal generator of this semigroup.
 - * $11 + 2 \not\equiv 0 \pmod{5}$.
 - * $11 - 10 \notin \langle 2, 5 \rangle$.
 - ★ $d_2 = 3$. In this case $11 + 5 \not\equiv 0 \pmod{3}$, $11 + 3 \not\equiv 0 \pmod{5}$, and $11 - 15 \notin \langle 3, 5 \rangle$.
 - ★ $d_2 = 4$.
 - * $11 + 5 \equiv 0 \pmod{4}$, and $f_2 = (11 - 20 + 5)/4 = -1$. So $\Gamma = 5\mathbb{N} + 4\mathbb{N} = \langle 4, 5 \rangle$ is another complete intersection with Frobenius number 11.
 - * $11 - 20 \notin \langle 4, 5 \rangle$.
- $d_1 = 4$, $2 \leq d_2 \leq \min\{3, \lfloor \frac{12}{3} \rfloor + 1\} = 3$, and as $\gcd(2, 4) \neq 1$, we get $d_2 = 3$.
 - ★ $11 + 4 \equiv 0 \pmod{3}$. So Γ_1 could be \mathbb{N} . If this is the case, $f_2 = (11 - 12 + 4)/3 = 1$, which forces Γ_2 to be $\langle 2, 3 \rangle$, and $4 \in \Gamma_2$ is not a minimal generator. So we obtain $\Gamma = 4\mathbb{N} + 3\langle 2, 3 \rangle = \langle 4, 6, 9 \rangle$, which was already computed before.
 - ★ $11 + 3 \not\equiv 0 \pmod{4}$.
 - ★ $11 - 12 \notin \langle 3, 4 \rangle$.
- $d_3 = 3$ and $d_2 = 2$.
 - ★ $11 + 3 \equiv 0 \pmod{2}$, and $\Gamma_1 = \mathbb{N}$ can be a possibility. Then $f_2 = (11 - 6 + 3)/2 = 7$. If we apply this procedure recursively for $f = 7$, we obtain that $\{\langle 2, 9 \rangle, \langle 3, 5 \rangle, \langle 4, 5, 6 \rangle\}$ is the set of all possible complete intersection numerical semigroups with Frobenius number 7. However, $3 \notin \langle 2, 9 \rangle$, 3 is a minimal generator of $\langle 3, 5 \rangle$, and $3 \notin \langle 4, 5, 6 \rangle$.
 - ★ $11 + 2 \not\equiv 0 \pmod{3}$.
 - ★ $11 - 6 = 5 \in \langle 2, 3 \rangle$, and $5 = 1 \cdot 2 + 1 \cdot 3$ is the only factorization. So the only possible choice for f_1 and f_2 is 1. This means that Γ_1 and Γ_2 must be $\langle 2, 3 \rangle$. Again we obtain no new semigroups, since 2 and 3 are minimal generators of $\langle 2, 3 \rangle$.

Thus the set of complete intersection numerical semigroups with Frobenius number 11 is

$$\{\langle 2, 13 \rangle, \langle 4, 6, 9 \rangle, \langle 3, 7 \rangle, \langle 4, 5 \rangle\}.$$

3. FREE NUMERICAL SEMIGROUPS

Throughout this section, let Γ be the numerical semigroup Γ minimally generated by $\{r_0, \dots, r_h\}$. For $k \in \{1, \dots, h+1\}$, set $d_k = \gcd(\{r_0, \dots, r_{k-1}\})$ ($d_1 = r_0$).

Write $\Gamma_k = \left\langle \frac{r_0}{d_{k+1}}, \dots, \frac{r_k}{d_{k+1}} \right\rangle$, and $c_k = c(\Gamma_k)$ for all $k \in \{1, \dots, h\}$. Set $c = c_h = c(\Gamma)$.

We say that Γ is *free* if either $h = 0$ (and thus $r_0 = 1$) or Γ is the gluing of the free numerical semigroup Γ_{h-1} and \mathbb{N} . Free numerical semigroups were introduced in [1]. For other characterizations and properties of free numerical semigroups see [16, Section 8.4].

Example 14. Notice that the order in which the generators are given is crucial. For instance, $S = \langle 8, 10, 9 \rangle$ is free for the arrangement $(8, 10, 9)$ but it is not free for $(8, 9, 10)$. And a numerical semigroup can be free for different arrangements, for example, $S = \langle 4, 6, 9 \rangle$ has this property.

If we take c_0, \dots, c_h pairwise coprime integers greater than one, and $r_i = \prod_{j=0, i \neq j}^h c_j$, $j = 0, \dots, h$, then the numerical semigroup generated by $\{r_0, \dots, r_h\}$ is free for any arrangement of its minimal generating set (see [9]).

According to Proposition 1, with $A_2 = \{r_h\}$, we obtain the following consequence.

Corollary 15. *If Γ is free, then*

$$F(\Gamma) = d_h F(\Gamma_{h-1}) + r_h(d_h - 1).$$

In this way we retrieve Johnson's formula ([11]). Notice also that Γ_{h-1} is again free, so if we expand recursively this formula we obtain the formula given by Bertin and Carbonne for free numerical semigroups (see [1]; these authors named these semigroups in this way).

This equation can be reformulated in terms of the conductor as

$$(2) \quad c(\Gamma) = c_h = d_h c_{h-1} + (d_h - 1)(r_h - 1).$$

Lemma 16. *If Γ is free, then*

- (1) $\gcd(d_h, r_h) = 1$;
- (2) $d_h \mid F(\Gamma) + r_h$ (consequently $d_h \nmid F(\Gamma)$);
- (3) if we define $e_k = \frac{d_k}{d_{k+1}}$, $k = 1, \dots, h$, then $e_k r_k \in \langle r_0, \dots, r_{k-1} \rangle$, for all $k = 1, \dots, h$; in particular, $e_k \geq 2$;
- (4) $d_1 > d_2 > \dots > d_{h+1} = 1$;
- (5) $d_h \leq \frac{c(\Gamma)}{r_h - 1} + 1$;
- (6) for $h \geq 1$, $(d_h - 1)(r_h - 1) \geq 2^h$.

Proof. (1) This follows from the fact that Γ is a numerical semigroup, and thus $\gcd(d_h, r_h) = d_{h+1} = 1$.

(2) $F(\Gamma_{h-1}) = (F(\Gamma) + r_h(1 - d_h))/d_h = (F(\Gamma) + r_h)/d_h - 1$.

(3) As Γ_k is the gluing of Γ_{k-1} and \mathbb{N} , we have that $\frac{r_k}{d_{k+1}} \in \Gamma_{k-1}$. Hence $\frac{d_k}{d_{k+1}} r_k \in \langle r_0, \dots, r_{k-1} \rangle$. If $e_k = 1$, then $r_k \in \langle r_0, \dots, r_{k-1} \rangle$, contradicting that r_k is a minimal generator.

(4) By definition, $d_k \geq d_{k+1}$. As $e_k = \frac{d_k}{d_{k+1}} \geq 2$, we get $d_k > d_{k+1}$.

(5) Notice that $F(\Gamma) \geq (r_h - 1)(d_h - 1) - 1$, since $F(\Gamma_{h-1}) \geq -1$.

(6) If $d_h = 2$, then we show that $r_h > m(\Gamma)$. Assume to the contrary that $r_h = m(\Gamma)$. Then we already proved above that $e_h r_h \in \langle r_0, \dots, r_{h-1} \rangle$. Since $e_h = d_h$ and r_i is a minimal generator of Γ for all i , we deduce that $2r_h = \sum_{i=0}^{h-1} a_i r_i$, with $\sum_{i=0}^{h-1} a_i \geq 2$. As $r_i > r_h$ for every $i = 0, \dots, h-1$, we get $2r_h > r_h \sum_{i=0}^{h-1} a_i$, and thus $\sum_{i=0}^{h-1} a_i < 2$, a contradiction. Thus in view of Proposition 5, we have that $r_h \geq 2^h$, and if $d_h = 2$, then $r_h \geq 2^h + 1$. Hence for $d_h = 2$

the proof follows easily, and for $d_h > 2$ we get $(d_h - 1)(r_h - 1) \geq 2(r_h - 1) \geq 2(2^h - 1) \geq 2^h$ (we are assuming $h \geq 1$).

□

In view of Example 10, the bound proposed in Proposition 9 cannot be improved for free numerical semigroups, since the family introduced in Example 6 consists on free numerical semigroups. However, we can use Proposition 9 to find an upper bound for r_h , as we show next.

For all $h \geq 2$, $c_{h-1} = \frac{c - (d_h - 1)(r_h - 1)}{d_h}$ is an even integer, and $c = c_h \geq h2^h$. In particular,

$$-c_{h-1}d_h \leq -(h-1)2^{h-1}d_h.$$

Hence

$$(r_h - 1)(d_h - 1) \leq c - (h-1)2^{h-1}d_h$$

This gives us the following upper bound for r_h .

$$r_h \leq \frac{c}{d_h - 1} - (h-1)2^{h-1} \frac{d_h}{d_h - 1} + 1.$$

Corollary 17. *For all $h \geq 2$,*

$$2^h + 1 \leq r_h \leq \frac{c}{d_h - 1} - (h-1)2^{h-1} \frac{d_h}{d_h - 1} + 1 \leq c - (h-1)2^{h-1} + 1.$$

Remark 18. In order to compute the set of all free numerical semigroups with a given Frobenius number, we make use of the formula given in Corollary 15, by taking into account the restrictions given in this section for d_h and r_h .

4. TELESCOPIC NUMERICAL SEMIGROUPS

We keep using the same notation as in the preceding section. We say that the numerical semigroup Γ minimally generated by $\{r_0, \dots, r_h\}$ is *telescopic* if it is free for the arrangement of the generators $r_0 < \dots < r_h$ (see for instance [12]). This motivates the notation $\{r_0 < \dots < r_h\}$, that means that the elements in the set $\{r_0, \dots, r_h\}$ fulfill the extra condition $r_0 < \dots < r_h$. We will also write $\Gamma = \langle r_0 < \dots < r_h \rangle$ when $\{r_0, \dots, r_h\}$ is a generating system for Γ and $r_0 < \dots < r_h$.

Notice that in addition to the properties we had for free numerical semigroups, if Γ is telescopic, then

- (1) $d_h < r_h$, because $d_h \mid r_{h-1} < r_h$;
- (2) $F(\Gamma) \geq (r_h - 1)(d_h - 1) - 1 > (d_h - 1)^2 - 1$, whence $d_h \leq \min \left\{ r_h - 1, \frac{c(\Gamma)}{r_h - 1} + 1, \sqrt{c(\Gamma)} + 1 \right\}$.

Proposition 19. *Let Γ be a telescopic numerical semigroup minimally generated by $\{r_0 < \dots < r_h\}$. If $h \geq 2$, then $r_h \geq 2^{h+1} - 1$.*

Proof. Let $h = 2$, and let $\Gamma_1 = \left\langle \frac{r_0}{d_2}, \frac{r_1}{d_2} \right\rangle$. Since $\frac{r_1}{d_2} \geq 3$ and $d_2 \geq 2$, we have $r_1 \geq 6$. Besides, $r_2 > r_1$, whence $r_2 \geq 7$. Note that this bound is attained for $\Gamma_2 = \langle 4, 6, 7 \rangle$.

Assume that $h \geq 3$, and that the formula is true for $h - 1$. We have $r_h \geq r_{h-1} + 1$ and $r_h \in \left\langle \frac{r_0}{d_h}, \dots, \frac{r_{h-1}}{d_h} \right\rangle$. By induction hypothesis, we have $\frac{r_{h-1}}{d_h} \geq 2^h - 1$. Hence $r_h \geq 2(2^h - 1) + 1 = 2^{h+1} - 1$. Note that this bound is reached by $\Gamma_h = \langle 2^h, 3 \cdot 2^{h-1}, 7 \cdot 2^{h-2}, \dots, (2^{k+1} - 1) \cdot 2^{h-k}, \dots, 2^{h+1} - 1 \rangle$. □

As in the free case, we can describe a bound for the embedding dimension of a telescopic numerical semigroup.

Proposition 20. *Let Γ be a telescopic numerical semigroup other than \mathbb{N} . Then*

$$(e(\Gamma) - 2)2^{e(\Gamma)} + 2 \leq c(\Gamma).$$

Proof. Assume that Γ is minimally generated by $\{r_0 < \dots < r_h\}$. Denote as usual $c(\Gamma)$ by c . We use once more induction on h .

The case $h = 1$ is evident.

Suppose that $h \geq 2$, and that our inequality is true for $h - 1$. By (2), we have $c = d_h c_{h-1} + (d_h - 1)(r_h - 1)$. By induction hypothesis, $c_{h-1} \geq (h - 2)2^h + 2$, and as $d_h \geq 2$, and $r_h \geq 2^{h+1} - 1$, we get $c \geq (h - 2)2^{h+1} + 4 + 2^{h+1} - 2 = (h - 1)2^{h+1} + 2$. \square

Note that for all $h \geq 2$, $c_{h-1} = \frac{c - (d_h - 1)(r_h - 1)}{d_h}$ is an even integer, and that $c = c_h \geq (h - 1)2^{h+1} + 2$. In particular

$$-c_{h-1}d_h \leq -((h - 2)2^h + 2)d_h.$$

Hence

$$(r_h - 1)(d_h - 1) \leq c - ((h - 2)2^h + 2)d_h.$$

This gives us the following upper bound for r_h :

$$r_h \leq \frac{c}{d_h - 1} - ((h - 2)2^h + 2)\frac{d_h}{d_h - 1} + 1 \leq c - (h - 2)2^h - 1.$$

Corollary 21. *For all $h \geq 2$, we have*

$$2^{h+1} - 1 \leq r_h \leq \frac{c}{d_h - 1} - ((h - 2)2^h + 2)\frac{d_h}{d_h - 1} + 1 \leq c - (h - 2)2^h - 1.$$

Remark 22. For computing the set of all telescopic numerical semigroups with fixed Frobenius number, we proceed as in the free case, ensuring that r_h is larger than the largest generator of Γ_1 multiplied by d_h . Notice that d_h must now be smaller than r_h .

5. PLANE CURVE SINGULARITIES

Let Γ be the numerical semigroup minimally generated by $\{r_0 < r_1 < \dots < r_h\}$. Let d_k , Γ_k , c_k , and e_k be as in the preceding section. The numerical semigroup Γ is the numerical semigroup associated to an irreducible plane curve singularity if Γ is telescopic and $e_k r_k < r_{k+1}$ for all $k = 1, \dots, h - 1$ (see [19]).

Proposition 23. *Let Γ be the semigroup associated to an irreducible plane curve singularity minimally generated by $\{r_0 < \dots < r_h\}$, with $h \geq 2$. Then $r_h \geq \frac{1}{3}(5 \cdot 2^{2h-1} - 1)$.*

Proof. For $h = 2$, as $\Gamma_1 = \left\langle \frac{r_0}{d_2} < \frac{r_1}{d_2} \right\rangle$, we obtain $\frac{r_1}{2} \geq 3$. Since $d_2 \geq 2$, we deduce that $r_1 \geq 6$. The plane singularity condition implies $r_2 > e_1 r_1 \geq 12$, because we know that $e_1 \geq 2$ (Lemma 16). Hence $r_2 \geq 13$.

Assume that $h \geq 3$, and that the formula is true for $h - 1$. The plane singularity condition for $k = h - 1$ implies that $r_h \geq \frac{r_{h-1}}{d_h} d_{h-1} + 1$. The quotient $\frac{r_{h-1}}{d_h}$ is the largest generator of Γ_{h-1} . The induction hypothesis then asserts that $\frac{r_{h-1}}{d_h} \geq \frac{1}{3}(5 \cdot 2^{2(h-1)-1} - 1)$. By using that $e_k \geq 2$ for all k (Lemma 16), we deduce that $d_{h-1} \geq 4$. By putting all this together, we get $r_h \geq 4\left(\frac{1}{3}(5 \cdot 2^{2(h-1)-1} - 1)\right) + 1 = \frac{1}{3}(5 \cdot 2^{2h-1} - 1)$. \square

Proposition 24. *Let $\Gamma \neq \mathbb{N}$ be the semigroup associated to an irreducible plane curve singularity minimally generated by $\{r_0 < \dots < r_h\}$ and with conductor c . Then*

$$c \geq \frac{5}{3}2^{2h} - 3 \cdot 2^h + \frac{4}{3}.$$

Proof. The case $h = 1$ is evident.

Assume that $h \geq 2$ and that our inequality holds for $h - 1$. We have: $c = d_h c_{h-1} + (d_h - 1)(r_h - 1)$. By induction hypothesis $c_{h-1} \geq \frac{5}{3}2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}$. Notice that $d_h \geq 2$. Thus our assertion follows from Proposition 23. \square

We proceed now as we did in the telescopic case to obtain also an upper bound for r_h . Note that for all $h \geq 2$, $c_{h-1} = \frac{c-(d_h-1)(r_h-1)}{d_h}$ is an even integer, and that $c_{h-1} \geq \frac{5}{3}2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}$. Thus

$$-c_{h-1}d_h \leq -\left(\frac{5}{3}2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}\right)d_h.$$

Hence

$$(r_h - 1)(d_h - 1) \leq c - \left(\frac{5}{3}2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}\right)d_h.$$

This gives us the following upper bound for r_h .

$$r_h \leq \frac{c}{d_h - 1} - \left(\frac{5}{3}2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}\right) \frac{d_h}{d_h - 1} + 1.$$

Corollary 25. *For all $h \geq 2$, we have*

$$\frac{5}{3}2^{2h-1} - \frac{1}{3} \leq r_h \leq \frac{c}{d_h - 1} - \left(\frac{5}{3}2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}\right) \frac{d_h}{d_h - 1} + 1 \leq c - \frac{5}{3}2^{2h-2} - 3 \cdot 2^{h-1} + \frac{7}{3}.$$

A bound for the embedding dimension also follows from the above proposition.

Corollary 26. *If $h \geq 2$, then*

$$h \leq \log_2 \left(\frac{\sqrt{60c+1+9}}{10} \right).$$

Proof. From Proposition 24, $\frac{5}{3}2^{2h} - 3 \cdot 2^h + \frac{4}{3} \leq c$. Write $x = 2^h$, we get $5/3x^2 - 3x + \frac{4}{3} \leq c$. By solving $5/3x^2 - 3x + \frac{4}{3} - c = 0$, we get $x \in \left\{ -\frac{\sqrt{60c+1-9}}{10}, \frac{\sqrt{60c+1+9}}{10} \right\}$. As the minimum of $5/3x^2 - 3x + \frac{4}{3} - c$ is reached in $x = 9/10$, and in our setting $x = 2^h > 1$, we have that the maximum possible $x > 0$ such that $\frac{5}{3}2^{2h} - 3 \cdot 2^h + \frac{4}{3} \leq c$ is $x = \frac{\sqrt{60c+1+9}}{10}$. \square

Remark 27. The set of all numerical semiogroups with fixed Frobenius number associated to an irreducible planar curve singularity is calculated as in the free case, by imposing the condition $e_k r_k < r_{k+1}$.

6. EXPERIMENTAL RESULTS


With the ideas given in the preceding sections, we implemented in **GAP** ([8]), with the help of the **numericalsgps** package ([7]), functions to compute the set of all complete intersection, free and telescopic numerical semigroups, as well as the set of all numerical semigroups associated to irreducible planar curve singularities with fixed Frobenius number (these functions will be included in the next release of this package).

The following table was computed in 6932 milliseconds on a 2.5GHz desktop computer, and it shows, for fixed genus g , the number of complete intersections ($ci(g)$), free ($fr(g)$), telescopic ($tl(g)$), associated to an irreducible planar curve singularity ($pc(g)$) numerical semigroups, respectively. Recall that for a symmetric numerical semigroup its conductor is twice its genus.

Observe that almost all complete intersections in this table are free. This is due to the fact that the embedding dimension of all numerical semigroups appearing there is small, and for embedding dimension three or less, the concepts of free and complete intersections coincide (among the complete intersection numerical semigroups represented in the table 158 of them have embedding dimension 2, 1525 have embedding dimension 3, 1862 have embedding dimension 4, and 205 have embedding dimension 5).

| g | ci(g) | fr(g) | tl(g) | pc(g) | g | ci(g) | fr(g) | tl(g) | pc(g) | g | ci(g) | fr(g) | tl(g) | pc(g) |
|----|-------|-------|-------|-------|----|-------|-------|-------|-------|----|-------|-------|-------|-------|
| 0 | 1 | 1 | 1 | 1 | 19 | 24 | 24 | 12 | 5 | 38 | 61 | 61 | 37 | 12 |
| 1 | 1 | 1 | 1 | 1 | 20 | 16 | 16 | 11 | 6 | 39 | 100 | 100 | 52 | 16 |
| 2 | 1 | 1 | 1 | 1 | 21 | 27 | 27 | 18 | 9 | 40 | 110 | 109 | 54 | 19 |
| 3 | 2 | 2 | 2 | 2 | 22 | 31 | 31 | 19 | 8 | 41 | 80 | 79 | 47 | 12 |
| 4 | 3 | 3 | 2 | 2 | 23 | 21 | 21 | 13 | 6 | 42 | 122 | 120 | 61 | 20 |
| 5 | 2 | 2 | 2 | 1 | 24 | 36 | 35 | 20 | 11 | 43 | 120 | 120 | 60 | 17 |
| 6 | 4 | 4 | 4 | 3 | 25 | 38 | 38 | 22 | 9 | 44 | 94 | 94 | 48 | 15 |
| 7 | 5 | 5 | 3 | 2 | 26 | 27 | 27 | 16 | 8 | 45 | 143 | 142 | 73 | 22 |
| 8 | 3 | 3 | 2 | 2 | 27 | 46 | 46 | 24 | 11 | 46 | 151 | 149 | 72 | 21 |
| 9 | 7 | 7 | 5 | 4 | 28 | 45 | 45 | 25 | 10 | 47 | 108 | 106 | 57 | 15 |
| 10 | 8 | 8 | 6 | 4 | 29 | 34 | 33 | 20 | 7 | 48 | 158 | 157 | 75 | 24 |
| 11 | 5 | 5 | 4 | 2 | 30 | 57 | 57 | 32 | 13 | 49 | 179 | 179 | 84 | 23 |
| 12 | 11 | 11 | 8 | 5 | 31 | 62 | 62 | 31 | 9 | 50 | 128 | 128 | 68 | 20 |
| 13 | 11 | 11 | 8 | 3 | 32 | 43 | 43 | 25 | 10 | 51 | 197 | 194 | 86 | 26 |
| 14 | 9 | 9 | 7 | 4 | 33 | 65 | 65 | 37 | 14 | 52 | 209 | 207 | 89 | 27 |
| 15 | 14 | 14 | 10 | 6 | 34 | 77 | 76 | 39 | 13 | 53 | 142 | 142 | 76 | 20 |
| 16 | 17 | 17 | 9 | 5 | 35 | 53 | 52 | 29 | 11 | 54 | 229 | 227 | 101 | 30 |
| 17 | 12 | 12 | 8 | 3 | 36 | 84 | 83 | 43 | 17 | 55 | 238 | 235 | 104 | 29 |
| 18 | 18 | 18 | 12 | 6 | 37 | 90 | 90 | 47 | 13 | 56 | 172 | 169 | 83 | 24 |

The largest genus, for which the set of numerical semigroups with this genus is known, is 55, and the number of numerical semigroups with genus 55 is 1142140736859 ([6]), while there are just 2496 symmetric numerical semigroup with genus 55 (this last amount can be computed by using the `IrreducibleNumericalSemigroupsWithFrobeniusNumber` command of the `numericalsgps` package). The proportion of complete intersections among symmetric numerical semigroups is small, and tiny compared with the whole set of numerical semigroups.



Also as one of the referees observed, local minimums in the graph are attained when the genus is congruent with 2 modulo 3. We have not a proof for this fact. May be this behavior is inherited from the symmetric case. The sequence

1, 1, 1, 2, 3, 3, 6, 8, 7, 15, 20, 18, 36, 44, 45, 83, 109, 101, 174, 246, 227, 420, 546, 498, 926, 1182, 1121, 2015, 2496, 2436, 4350, 5602, 5317, 8925, 11971, 11276,

represents the number of symmetric numerical semigroups with genus ranging from 0 to 35.

The following table shows that the proportion between complete intersections and free numerical semigroups remains similar even for larger genus. Observe that for genus 310 it takes 70 minutes to compute the set of all complete intersections, while it takes approximately 8 minutes and 30 seconds to determine all free numerical semigroups with this genus. For genus 55, computing the set of all numerical semigroups with this genus might take several months and a few terabytes (this was communicated to us by Manuel Delgado, see [6]).

| g | ci(g) | milliseconds | fr(g) | milliseconds | fr(g)/ci(g) |
|-----|-------|--------------|-------|--------------|-------------|
| 220 | 18018 | 538213 | 17675 | 94134 | 0.98 |
| 230 | 16333 | 660838 | 16026 | 108187 | 0.98 |
| 240 | 24862 | 924409 | 24359 | 153069 | 0.98 |
| 250 | 28934 | 1167901 | 28355 | 158706 | 0.98 |
| 260 | 25721 | 1389167 | 25186 | 177691 | 0.98 |
| 310 | 66335 | 4206374 | 64959 | 509691 | 0.98 |

REFERENCES

- [1] J. Bertin, P. Carbonne, Semi-groupes d'entiers et application aux branches, *J. Algebra* **49** (1977), 81-95.
- [2] I. Bermejo, I. García-Marco, J. J. Salazar-González, An algorithm for checking whether the toric ideal of an affine monomial curve is a complete intersection, *J. Symbolic Comput.* **42** (2007), 971-991.
- [3] I. Bermejo, P. Gimenez, E. Reyes, R. H. Villarreal, Complete intersections in affine monomial curves, *Bol. Soc. Mat. Mexicana* (3) **11** (2005), no. 2, 191-203.
- [4] M. Bras-Amors, Fibonacci-like behavior of the number of numerical semigroups of a given genus, *Semigroup Forum*, **76** (2008), 379-384.
- [5] C. Delorme, Sous-monoides d'intersection complète de \mathbb{N} , *Ann. Scient. École Norm. Sup. (4)*, **9** (1976), 145-154.
- [6] M. Delgado, Experiments with numerical semigroups, in preparation.
- [7] M. Delgado, P.A. García-Sánchez, and J. Morais, "numericalsgps": a gap package on numerical semigroups, (<http://www.gap-system.org/Packages/numericalsgps.html>).
- [8] The GAP Group, *GAP - Groups, Algorithms, and Programming, Version 4.4*; 2004, (<http://www.gap-system.org>).
- [9] P. A. García-Sánchez, I. Ojeda, J. C. Rosales, Affine semigroups having a unique Betti element, *J. Algebra Appl.* **12** (2013), 1250177 (11 pages).
- [10] J. Herzog, E. Kunz, Die Wertehalbgruppe eines lokalen Rings der Dimension 1, *S. B. Heidelberger Akad. Wiss. Math. Natur. Kl* (1971), 2767.
- [11] S. M. Johnson, A linear Diophantine problem, *Can. J. Math.*, **12** (1960), 390-398.
- [12] C. Kirfel, R. Pellikaan, The minimum distance of codes in an array coming from telescopic semigroups, *IEEE Trans. Inform. Theory*, **41**(1995) 1720-1732. Special issue on algebraic geometry codes.
- [13] J. C. Rosales, On presentations of subsemigroups of \mathbb{N}^n , *Semigroup Forum* **55** (1997), 152-159.
- [14] J. C. Rosales, P. A. García-Sánchez, On numerical semigroups with high embedding dimension, *J. Algebra* **203** (1998), 567-578.
- [15] J. C. Rosales, P. A. García-Sánchez, On free affine semigroups, *Semigroup Forum* **58** (1999), no. 3, 367-385.
- [16] J. C. Rosales and P. A. García-Sánchez, *Numerical Semigroups*, **20**. Springer, New York, 2009.
- [17] J.C. Rosales, P. A. García-Sánchez, J.I. García-García, J.A. Jiménez-Madrid, Fundamental gaps in numerical semigroups, *J. Pure Appl. Alg.* **189** (2004), 301-313.
- [18] A. Zhai, Fibonacci-like growth of numerical semigroups of a given genus, arXiv:1111.3142
- [19] O. Zariski, Le problème des modules pour les courbes planes, Hermann, 1986.

UNIVERSITÉ D'ANGERS, DÉPARTEMENT DE MATHÉMATIQUES, LAREMA, UMR 6093, 2 BD LAVOISIER, 49045
ANGERS CEDEX 01, FRANCE

E-mail address: `assi@univ-angers.fr`

DEPARTAMENTO DE ÁLGEBRA, UNIVERSIDAD DE GRANADA, E-18071 GRANADA, ESPAÑA

E-mail address: `pedro@ugr.es`