



HAL
open science

Structure of Arbitrarily Partitionable Graphs with cutset of size k

Olivier Baudon, Florent Foucaud, Jakub Przybylo, Mariusz Woźniak

► **To cite this version:**

Olivier Baudon, Florent Foucaud, Jakub Przybylo, Mariusz Woźniak. Structure of Arbitrarily Partitionable Graphs with cutset of size k . 2012. <hal-00690253v1>

HAL Id: hal-00690253

<https://hal.science/hal-00690253v1>

Preprint submitted on 22 Apr 2012 (v1), last revised 26 Apr 2013 (v9)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Structure of Arbitrarily Partitionable Graphs with cutset of size k^*

Olivier Baudon^a, Florent Foucaud^a, Jakub Przybyło^b, Mariusz Woźniak^{b, †}

^aLaBRI, Université de Bordeaux

351, cours de la Libération, 33405 Talence Cedex, France

^bFaculty of Applied Mathematics, AGH University of Science and Technology
Al. Mickiewicza 30, 30-059 Kraków, Poland

April 22, 2012

Abstract

A graph $G = (V, E)$ is arbitrarily partitionable if for any sequence τ of positive integers adding up to $|V|$, there is a sequence of vertex-disjoint subsets of V whose orders are given by τ , and which induce connected subgraphs. In this paper, we focus on the number and the sizes of the components of an arbitrarily partitionable graph after removing a cutset.

1 Introduction

1.1 Arbitrarily partitionable graphs

Consider a network of computers that you want to separate into disjoint subnetworks of given sizes. If you want to assume that the sizes of subnetworks may be chosen completely arbitrarily, then the underlying graph, where computers are represented by vertices and links between two computers by edges, must be *arbitrarily partitionable*.

More formally, let n, τ_1, \dots, τ_k be positive integers such that $\tau_1 + \dots + \tau_k = n$. $\tau = (\tau_1, \dots, \tau_k)$ is called a *decomposition* of n .

Let $G = (V, E)$ be a graph of order n and S a subset of V . $G[S]$ denotes the subgraph of G induced by S . Let (τ_1, \dots, τ_k) be a decomposition of n . G is τ -*partitionable* iff it exists a partition of $V : V_1, \dots, V_k$ such that for each $i, 1 \leq i \leq k$

- $|V_i| = \tau_i$
- $G[V_i]$ is connected

In that case, τ is said *realizable* in G and (V_1, \dots, V_k) is a *realization* of τ in G .

A graph $G = (V, E)$ of order n is *arbitrarily partitionable* (*AP* for short) iff for each decomposition τ of n , G is τ -partitionable.

*This research was partially supported by the partnership Hubert Curien Polonium 22658VG and the Polish Ministry of Science and Higher Education.

[†]email addresses: {olivier.baudon, florent.foucaud}@labri.fr, przybylo@wms.mat.agh.edu.pl, mwozniak@agh.edu.pl

1.2 On-line and recursive versions

Consider now that the sizes of the subnetworks are requested one by one instead of all together. Using the graph modeling, that means that you must provide a connected subgraph of given order and assume that the rest of the graph remains "on-line" partitionable or is empty. Graphs which have these property for any sequence of requests are called *on-line arbitrarily partitionable*.

The following characterization of such a graphs has been given in [HTW07] :

A connected graph $G = (V, E)$ of order n is *on-line arbitrarily partitionable* (OLAP for short) iff for each integer $1 \leq \lambda \leq n - 1$, there exists a subset V_λ of V such that

- $|V_\lambda| = \lambda$,
- $G[V_\lambda]$ is connected,
- $G[V \setminus V_\lambda]$ is OL-AVD.

Another family of arbitrarily partitionable graphs has also been considered in [BGW11] : *recursively arbitrarily partitionable* graphs. In that case, we want not only to provide connected subgraphs, but also that these subgraphs are themselves partitionables.

A graph $G = (V, E)$ of order n is *recursively arbitrarily partitionable* (RAP for short) iff

- $G = K_1$ or
- G is connected and for each decomposition $\tau = (\tau_1, \dots, \tau_k)$ of n , $k \geq 2$, there exists a partition of $V : V_1, \dots, V_k$ such that for all $i, 1 \leq i \leq k$,
 - $|V_i| = \tau_i$,
 - $G[V_i]$ is RAP.

In [BGW11], it has been shown that $AP \Rightarrow OLAP \Rightarrow RAP$ and that it exists AP graphs that are not OLAP and OLAP graphs that are not RAP.

1.3 Previous results

Most of the papers on AP, OLAP and RAP graphs consider quite simple classes of graphs like trees or a class of 2-connected graphs called balloons. Among the previous results, we will focus on those which help us to determine how many components may be obtained after removing a cutset and the sizes of these components.

In [BF06], the following result gives an upper bound on the degree of AP trees :

Theorem 1 [BF06] *If a tree T is AP, then its maximum degree is 4. Moreover, every vertex of degree 4 in T is adjacent to a leaf.*

In [HTW07] and [BGW11], OLAP and RAP-trees have been completely characterized. For that, we need the following notations :

- A *k-pode* $T_k(t_1, \dots, t_k)$ is a tree of order $1 + \sum_{i=1}^k t_i$ composed by k paths of respective orders t_1, \dots, t_k , connected to a unique node, called the *root* of the *k-pode* (cf. Figure 1a).
- Let a and b be two positive integers. A *caterpillar* $Cat(a, b)$ is a tree of order $a + b$, composed by three paths of order a , b and 2, sharing exactly one node, called the *root* of the caterpillar. $Cat(a, b)$ is isomorphic to $T_3(a - 1, b - 1, 1)$ (cf. Figure 1b).




Figure 1: Examples of Graphs

Theorem 2 [HTW07] *A tree T is OLAP if and only if either T is a path or T is a caterpillar $Cat(a, b)$ with a and b given in Table 1 or T is the 3-pode $T_3(2, 4, 6)$.*

a	b	a	b
2, 4	$\equiv 1 \pmod{2}$	7	8, 9, 11, 13, 15
3	$\equiv 1, 2 \pmod{3}$	8	11, 19
5	6, 7, 9, 11, 14, 19	9, 10	11
6	$\equiv 1, 5 \pmod{6}$	11	12

Table 1: Values a, b ($b \geq a$), such that $Cat(a, b)$ is OLAP

Theorem 3 [BGW11] *A tree T is RAP if and only if either T is a path or T is a caterpillar $Cat(a, b)$ with a and b given in Table 2 or T is the 3-pode $T_3(2, 4, 6)$.*

a	b
2, 4	$\equiv 1 \pmod{2}$
3	$\equiv 1, 2 \pmod{3}$
5	6, 7, 9, 11, 14, 19
6	7
7	8, 9, 11, 13, 15

Table 2: Values a, b ($b \geq a$), such that $Cat(a, b)$ is RAP

We have also some results on the class of 2-connected graphs called k -balloons. Let b_1, \dots, b_k be positive integers. A k -balloon $B(b_1, \dots, b_k)$ is a graph of order $2 + \sum_{i=1}^k b_i$ composed by two vertices (called *roots*) linked by k paths (called *branches*) of *widths* (the number of internal vertices) b_1, \dots, b_k (cf. Figure 1c).

Theorem 4 [BGW11] *If a k -balloon is RAP, then $k \leq 5$. This bound is tight.*

This result has been extended to OLAP- k -balloons :

Theorem 5 [BBFP12] *If a k -balloon is OLAP, then $k \leq 5$.*

Upper bounds for the size of the smallest branch of a RAP or OLAP- k -balloon have also been given :

Theorem 6 [BBFP12] Let $B(b_1, \dots, b_k)$ be a k -balloon, with $k \geq 4$ and $b_1 \leq \dots \leq b_k$. If $B(b_1, \dots, b_k)$ is OLAP, then $b_1 \leq 11$. If $B(b_1, \dots, b_k)$ is RAP, then $b_1 \leq 7$.

2 Size and number of components after removing a cutset of size at most 2

Results given in subsection 1.3 allow us to easily deduce some bounds on the number of components and their sizes, after removing a cutset of size 1 or 2, using the following remarks :

Remark 7 If G contains a spanning subgraph which is AP (resp. OLAP, RAP), then G is AP (resp. OLAP, RAP). In particular, if G is traceable (containing an hamiltonian path), then G is RAP (and then OLAP and AP).

Remark 8 Let $G = (V, E)$ be a graph with a cutset S and let V_1, \dots, V_k the connected components of $G[V \setminus S]$:

- if $|S| = 1$ and G is AP (resp. OLAP, RAP), then the k -pode $T_k(|V_1|, \dots, |V_k|)$ is AP (resp. OLAP, RAP) ;
- if $|S| = 2$ and G is AP (resp. OLAP, RAP), then the k -balloon $B(|V_1|, \dots, |V_k|)$ is AP (resp. OLAP, RAP) ;

Using Remark 8 and Theorems 1, 2, 3, 4, 5, 6, we obtain the following theorem :

Theorem 9 Let G be a graph with a cutset S , and V_1, \dots, V_k the connected components of $G[V \setminus S]$, with $|V_1| \leq \dots \leq |V_k|$:

- if $|S| = 1$ then
 - if G is AP, then $k \leq 4$ and if $k = 4$, then $|V_1| = 1$;
 - if G is OLAP (resp. RAP), then $k \leq 3$, and if $k = 3$, then either $(|V_1|, |V_2|, |V_3|) = (1, a, b)$, with values a and b taken in Table 1 (resp. 2) , or $(|V_1|, |V_2|, |V_3|) = (2, 4, 6)$;
- if $|S| = 2$ then
 - if G is OLAP, then $k \leq 5$;
 - if G is OLAP (resp. RAP) and $k \in \{4, 5\}$ then $|V_1| \leq 11$ (resp. ≤ 7).

3 Number of components after removing a cutset of size $k \geq 2$ in AP graphs

In the previous section, we have proved that if we remove a cutset of size 1 to a graph G , then the number of the remaining connected components is at most 4 if G is AP, and 3 if G is OLAP. Similarly, if we remove a cutset of size 2 to an OLAP graph, then the number of components is at most 5.

In this section, we will prove that for any $k \geq 2$, we may find an AP graph G such that if we remove a cutset of size k to G , the number of components is as big as we want.

Theorem 10 For any integer $k \geq 2$ and $c \geq 2$ it exists an AP graph $G = (V, E)$ with a cutset S of size k such that the number of connected components of $G[V \setminus S]$ is c .

Proof.

The proof of this theorem is based on examples of AP graphs having a cutset S of size k and such that the number of the remaining connected components after deleting vertices of S is c .

We first consider the case where $c \leq k$. Let $G = (S_1 \cup P \cup S_2, E)$ the graph with $2k$ vertices constructed as follow :

- $G[S_1]$ is a stable set with $c - 1$ vertices ;
- $G[P]$ is a path with $k - c + 1$ vertices ;
- $G[S_2]$ is a stable set with k vertices ;
- every vertex of S_1 or P is adjacent to all the vertices of S_2 .

Clearly, S_2 is a cutset of size k . The components of $G[V \setminus S_2]$ are the $c - 1$ vertices of S_1 plus the path induced by P . Because G admits the complete bipartite graph $K_{k,k}$ as partial graph and $K_{k,k}$ is hamiltonian, then G is AP.

Now, we assume that $c > k$.

We denote by $K_k(a_1, \dots, a_c)$ the graph formed by $c + 1$ cliques, one of size k , the others of size a_1, \dots, a_c . Moreover, we add all the edges between the vertices of the clique of size k and the vertices of the other cliques (see Figure 2). Clearly, the clique K_k is a cutset of $K_k(a_1, \dots, a_c)$.




Figure 2: Graph $K_2(2, 3, 4)$

We consider the case of the graph $K_k(a_1, \dots, a_c)$ where values a_1, \dots, a_c are chosen as follow :

- $1 \leq a_1 \leq \dots \leq a_c$
- $\forall i, 1 \leq i \leq c$, we denote $n_i = i + \sum_{1 \leq j \leq i} a_j$.
- $\forall i, 1 \leq i \leq c - 1$, choose a_{i+1} such that
 1. $\forall j, 2 \leq j \leq n_i, a_{i+1} \equiv 0 \pmod{j}$
 2. $a_{i+1} \geq n_i a_i$

For example, $a_i \cdot n_i!$ is a possible value for a_{i+1} .

Let $\tau = (\tau_1, \dots, \tau_l)$ be a decomposition of $n = k + \sum_{1 \leq i \leq c} a_i$, with $\tau_1 \leq \dots \leq \tau_l$. We consider two cases.

First case: $\tau_l \geq a_{c-1}$

In that case, $\tau_l \geq n_{c-2} = c - 2 + \sum_{1 \leq i \leq c-2} a_i \geq k - 1 + \sum_{1 \leq i \leq c-2} a_i$. Thus the part of size τ_l may cover all the cliques K_{a_i} with $i \leq c - 2$, plus $k - 1$ vertices of K_k and eventually some vertices of the cliques $K_{a_{c-1}}$ and K_{a_c} . The remaining graph is formed by at most two cliques joined by the last vertex of K_k and is traceable.

Second case: $\tau_l < a_{c-1}$

We denote by q_i the number of values i in τ , for each $i, 1 \leq i \leq \tau_l$.

We have $n = \sum_{1 \leq i \leq \tau_l} i \cdot q_i$

Then it exists a value $a, 1 \leq a \leq \tau_l$ such that $a \cdot q_a \geq \frac{n}{\tau_l}$. Thus, $a \cdot q_a \geq \frac{n}{a_{c-1}} \geq \frac{n_c}{a_{c-1}} \geq \frac{n_{c-1} a_{c-1}}{a_{c-1}} = n_{c-1}$

We denote by s the index such that for all $i \leq s, a_i \not\equiv 0 \pmod{a}$ and for all $i > s, a_i \equiv 0 \pmod{a}$. Note that s may be equal to 0.

Because $a_s \not\equiv 0 \pmod{a}$, we have $a \geq n_{s-1}$. That means that one part of size a may cover all the cliques $K_{a_1}, \dots, K_{a_{s-1}}$ plus a vertex of K_k and may be some vertices of K_{a_s} .

Because $\forall i > s, a_i \equiv 0 \pmod{a}$, and $a \cdot q_a \geq n_{k-1}$, we may cover also the cliques $K_{a_{s+1}}, \dots, K_{a_{c-1}}$ with parts of size a .

Thus the remaining graph is formed by K_{a_c} , remaining vertices of K_{a_s} and $k-1$ vertices of K_k . This graph is traceable. \square

Balloons The previous result (Theorem 10) can be adapted to the special case of balloons. The interest of this result is that it gives (for the case of $k=2$) an example of graph with order n , and number of edges in $O(n)$.

Theorem 11 $\forall k \geq 1$, it exists a k -balloon AP.

Proof. We consider a k -balloon $B(b_1, \dots, b_k)$ where branches have the same size than the cliques of $K_k(a_1, \dots, a_c)$ given in the proof of Theorem 10 :

$b_1 \leq \dots \leq b_k$ and we denote $n_i = 1 + \sum_{1 \leq j \leq i} b_j$. $B(b_1, \dots, b_k)$ is constructed as follow:

- $b_1 \geq 1$
- $\forall i \leq k-1$, choose b_{i+1} such that
 1. $\forall j, 2 \leq j \leq n_i, b_{i+1} \equiv 0 \pmod{j}$
 2. $b_{i+1} \geq n_i b_i$

Using same arguments than those used for $K_k(a_1, \dots, a_c)$, it is easy to see that such a balloon is AP. \square

4 Size of components

In this section, we will show that, if the number of the connected components, after removing a cutset of size at least 2, may be as large as wanted, the size of these components grows exponentially with their number.

Theorem 12 Let $G = (V, E)$ be a graph with n vertices, S a cutset of G of size k , c_1, \dots, c_l the orders of the connected components of $G_{V \setminus S}$, such that $l > k$ and $1 \leq c_1 \leq c_2 \leq \dots \leq c_l$. Then the values of the sequence $(c_i)_{i \geq 1}$ grow exponentially with i .

To prove this theorem, we will first prove the following lemma :

Lemma 13 Let $G = (V, E)$ be a graph with n vertices, S a cutset of G of size k , c_1, \dots, c_l the orders of the connected components of $G_{V \setminus S}$, such that $l > k$ and $1 \leq c_1 \leq c_2 \leq \dots \leq c_l$.

Let $a, q_1, \dots, q_l, r_1, \dots, r_l$ be integers such that

- $2 \leq a \leq n-1$
- $\forall i, 1 \leq i \leq l, c_i = q_i a + r_i$ with $r_i < a$

If G is AP, then

$$\sum_{1 \leq i \leq l} r_i \leq (k+1) \cdot (a-1)$$

Proof.

Let G be a partial graph of $K_k(c_1, \dots, c_l)$. If a decomposition of n $\tau = (\tau_1, \dots, \tau_p)$ is realizable in G , and V_1, \dots, V_p is a realization of τ in G , then τ is realizable in $K_k(c_1, \dots, c_l)$ and V_1, \dots, V_p is a realization of τ in $K_k(c_1, \dots, c_l)$. Moreover, if G is AP, then $K_k(c_1, \dots, c_l)$ is AP.

Then, to prove Lemma 13, we may consider only graphs of the form $K_k(c_1, \dots, c_l)$.

First, we remark that in a decomposition of $K_k(c_1, \dots, c_l)$, the vertices of K_k may be covered by the greatest parts of a decomposition τ (note that a same part may contain more than one vertex of K_k). In fact, suppose that we have a part V_1 which does not cover any vertex of K_k and a vertex v in K_k covered by a part V_2 such that $|V_2| < |V_1|$. Then we may always exchange $|V_1| - |V_2|$ vertices from V_1 to V_2 inside the clique containing V_1 .

We consider a decomposition of n , $\tau = (a, \dots, a, r)$ with $r < a$ and its realization in $K_k(c_1, \dots, c_l)$. Because $a > r$, we may suppose that all the parts covering the vertices of K_k are of size a . The cliques K_{c_1}, \dots, K_{c_l} are also covered mainly by parts of size a . The remaining vertices must be covered by the at most k parts of size a which cover also the k vertices of K_k , and the part of size r . Because $r \leq a - 1$, we obtain $\sum_{1 \leq i \leq k} r_i + k \leq k \cdot a + a - 1$. \square

Corollary 14 *Let $G = (V, E)$ be a graph with n vertices, S a cutset of G of size k , c_1, \dots, c_l the orders of the connected components of $G_{V \setminus S}$, such that $l > k$ and $1 \leq c_1 \leq c_2 \leq \dots \leq c_l$. If G is AP, then $\forall i, 1 \leq i \leq k$,*

$$c_i \geq \frac{1}{k} \sum_{1 \leq j < i} c_j$$

Proof. We consider $a = c_i + 1$. In that case, for all $j \leq i$, we have $r_j = c_j$. Thus, Lemma 13 becomes $\sum_{i < j \leq k} r_j + c_i + \sum_{1 \leq j < i} r_j \leq (k + 1) \cdot c_i$. Because $\sum_{i < j \leq k} r_j > 0$, we obtain the result. \square

Proof of the Theorem 12

We can now conclude the proof of the Theorem 12 :

$\forall i, j$ such that $k \cdot i > j \geq k(i - 1)$, we have $c_j \geq c_{k(i-1)}$. Thus,

$$c_{k \cdot i} \geq \frac{1}{k} \cdot (k \cdot c_{k(i-1)} + k \cdot c_{k(i-2)} + \sum_{1 \leq j < k(i-2)} c_j)$$

Thus, $c_{k \cdot i} \geq c_{k(i-1)} + c_{k(i-2)}$, and the sequence $(c_{ki})_{i \geq 1}$ grows at least as a Fibonacci sequence. \square

References

- [BBFP12] O. Baudon, J. Bensmail, F. Foucaud, and M. Pilśniak. On some properties of arbitrarily partitionable balloons with five branches. Technical report, 2012. oai:hal.archives-ouvertes.fr:hal-00656232.
- [BF06] D. Barth and H. Fournier. A degree bound on decomposable trees. *Discrete Applied Mathematics*, 306:469–477, 2006.
- [BFP11] O. Baudon, F. Foucaud, and M. Pilśniak. On on-line and recursive arbitrarily partitionable balloons. Technical report, 2011. oai:hal.archives-ouvertes.fr:hal-00656232.
- [BGW11] O. Baudon, F. Gilbert, and M. Woźniak. Recursively arbitrarily vertex-decomposable graphs. *to appear in Opuscula Mathematica*, 2011.
- [HTW07] M. Hornák, Z. Tuza, and M. Woźniak. On-line arbitrarily vertex decomposable trees. *Discrete Applied Mathematics*, 155:1420–1429, 2007.