



HAL
open science

A Rescaling Velocity Method for Dissipative Kinetic Equations - Applications to Granular Media

Francis Filbet, Thomas Rey

► **To cite this version:**

Francis Filbet, Thomas Rey. A Rescaling Velocity Method for Dissipative Kinetic Equations - Applications to Granular Media. 2012. hal-00690598v1

HAL Id: hal-00690598

<https://hal.science/hal-00690598v1>

Preprint submitted on 23 Apr 2012 (v1), last revised 16 Apr 2013 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A RESCALING VELOCITY METHOD FOR DISSIPATIVE KINETIC EQUATIONS APPLICATIONS TO GRANULAR MEDIA

FRANCIS FILBET AND THOMAS REY

ABSTRACT. We present a new numerical algorithm based on a relative energy scaling for collisional kinetic equations allowing to study numerically their long time behavior, without the usual problems related to the change of scales in velocity variables. Several applications are presented for Boltzmann like equations. This method is particularly efficient for numerical simulations of the granular gases equation with dissipative energy: it allows to study accurately the long time behavior of this equation and is very well suited for the study of clustering phenomena.

CONTENTS

1. Introduction	1
2. The Rescaling Velocity Method	2
3. Applications of the Rescaling Velocity Method	4
4. Boundary Conditions	8
5. Discretization of the Problem	10
6. Numerical Simulations	14
Acknowledgment	22
References	24

1. INTRODUCTION

We are interested in numerical simulations of the long time behavior of collisional kinetic equations such as the Boltzmann equation for granular gases. To this end, we introduce a new technique which is based on the information provided by the hydrodynamic fields computed from a macroscopic model corresponding to the original kinetic equation. Then we simply rescale the kinetic equation according to these hydrodynamic quantities. The reason to do so is that the change of scales in velocity is a challenging numerical problem when one wants to deal with solutions to dissipative kinetic equations on a fixed grid. Indeed, most of the usual deterministic methods fails to capture the correct long time behavior because of concentration or spreading over the velocity space.

Recently, F. Filbet and G. Russo proposed in [22] a rescaling method for space homogeneous kinetic equations on a fixed grid. This idea is mainly based on the self-similar behavior of the solution to the kinetic equation. However for space non homogeneous case, the situation is much more complicated and this method cannot be applied since the transport operator and the boundary conditions break down this self-similar behavior. Here we propose a very simple idea based on a rescaling of the kinetic equation according to its hydrodynamic limit. We will see that this approach can actually be applied to various type of collisional kinetic equations such as elastic and inelastic Boltzmann equation (also known as the *granular gases equation*).

Among popular methods for numerical simulations of collisional gases, the most widely used are meshless. On the one hand, the Molecular Dynamics algorithm is a deterministic method up to the random initialization of the particles which is valid in a dense regime [27]. This method has been used to describe

2010 *Mathematics Subject Classification*. Primary: 76P05, 82C40, Secondary: 65N08, 65N35 .

Key words and phrases. Boltzmann equation, spectral methods, boundary values problem, large time behavior, granular gases.

the apparition of shocks in supersonic sand and reproduces very accurately experimentation results [35]. On the other hand, Direct Simulation Monte Carlo (DSMC) methods are stochastic algorithms working either in dense or rarefied regimes. We refer for instance to [5] for a complete review of this topic. These methods are really efficient in term of computational cost since their complexity grows linearly with the number N of particles but they are rather inaccurate since the order of accuracy is about $\mathcal{O}(1/\sqrt{N})$. Another approach consists in a direct resolution of the Boltzmann operator on a phase space grid. For instance, deterministic and highly accurate methods based on a spectral discretization of the collisional operator have been proposed by F. Filbet, G. Naldi, L. Pareschi, G. Toscani and G. Russo in [33, 31, 21] for the space-homogeneous setting. Although being of complexity $\mathcal{O}(N^2)$, they are spectrally accurate and then need very few points to be precise. Another spectral method, inspired of the direct fast Fourier transform approach of A.V. Bobylev and S. Rjasanow [7, 9, 8] using Lagrange multiplier to improve the conservations was also used for space-homogeneous simulations by I. Gamba and S.H. Tharkabhushanam in [23].

Due to the large number of particles involved in the study of rarefied gas dynamics, we shall adopt a statistical physics' point of view, by the use of Boltzmann-like kinetic equations. For a given nonnegative initial condition f_0 , we will consider a particle distribution function $f^\varepsilon = f^\varepsilon(t, x, v)$, for $t \geq 0$, $x \in \Omega \subset \mathbb{R}^d$ and $v \in \mathbb{R}^d$, solution to the initial-boundary value problem

$$(1.1) \quad \begin{cases} \frac{\partial f^\varepsilon}{\partial t} + v \cdot \nabla_x f^\varepsilon = \frac{\mathcal{Q}(f^\varepsilon, f^\varepsilon)}{\varepsilon}, \\ f^\varepsilon(0, x, v) = f_0(x, v). \end{cases}$$

where the collision operator \mathcal{Q} is a Boltzmann-like operator, which preserves at least mass and momentum. This equation describes numerous models such as the Boltzmann equation for elastic and inelastic collisions or Fokker-Planck-Landau type equations. The parameter $\varepsilon > 0$ is the dimensionless Knudsen number, that is the ratio between the mean free path of particles before a collision and the length scale of observation. It measures the rarefaction of the gas: the gas is *rarefied* if $\varepsilon \sim 1$ and *dense* if $\varepsilon \ll 1$. The open set Ω is a bounded Lipschitz-continuous domain of \mathbb{R}^d , which means that the model (1.1) has to be supplemented with boundary conditions described later.

The article is organized as follows. We present in Section 2 the rescaling velocity method and give two possible choices of scaling functions, the coupled Filbet-Russo scaling and the new uncoupled macroscopic one. We choose to use the latter in the rest of the article, and apply it to three different collision operators in Section 3: the Boltzmann operator, the full granular gases operator and a simplified BGK-like granular gases operator. Subsequently, we introduce in Section 4 the spatial boundary conditions we have to impose on Ω : the so-called *Maxwell* boundary conditions, a convex combination of specular and diffusive reflections at the boundary. We present these conditions for the kinetic model (in classical and rescaled variables) and for the macroscopic equations. In Section 5, we introduce the numerical methods we shall use for the discretization of each term of the problem: the collision operator, the conservative transport term and the system of conservation laws with source term. Finally, we present in Section 6 some numerical results concerning the full and simplified granular gases models, in space homogeneous and inhomogeneous settings, with different geometries.

2. THE RESCALING VELOCITY METHOD

This section is devoted to the presentation of a new scaling for equation (1.1) allowing to follow the change of scales in velocity. It is an extension to the space-dependent setting of the method first introduced in [22], using the relative kinetic energy as a scaling function. Moreover, the idea is very close to the classical one used by J.A. Carrillo and G. Toscani in [13] and J.A. Carrillo, M. Di Francesco and G. Toscani in [11] for nonlinear diffusion equations. It was also extended by J.A. Carrillo and J. Vázquez in [14] to show the apparition of chaotic behavior for a precise (constructive) nonlinearity : the self-similar profile of this equation “oscillates” between Gaussian (heat equation) and Zel’dovich-Kompaneets-Barenblatt (porous medium equation) profiles.

Let us drop for simplicity the ε -dependence in f . For a given positive function ω , we introduce a new distribution $g(t, x, \xi)$ by setting

$$(2.1) \quad f(t, x, v) = \frac{1}{\omega(t, x)^d} g\left(t, x, \frac{v}{\omega(t, x)}\right),$$

where the function ω is assumed to be an accurate measure of the “support” or scale of the distribution f in velocity variables. Then according to this scaling, the distribution g should naturally “follow” either the concentration or the spreading in velocity of the distribution f .

Let us now derive the kinetic equation verified by the distribution g . Differentiating relation (2.1) with respect to time yields

$$\frac{\partial f}{\partial t} = \frac{1}{\omega^d} \left[\frac{\partial g}{\partial t} - \frac{1}{\omega} \frac{\partial \omega}{\partial t} \operatorname{div}_\xi(\xi g) \right].$$

Provided that $v = \omega \xi$, one also has

$$v \cdot \nabla_x f = \frac{\xi}{\omega^{d-1}} \cdot \left[\nabla_x g - \frac{1}{\omega} \nabla_x \omega \operatorname{div}_\xi(\xi g) \right].$$

Then, if f is solution to (1.1), the distribution g given by (2.1) is solution to the following equation:

$$\frac{\partial g}{\partial t} + \omega \xi \cdot \nabla_x g - \frac{1}{\omega} \left[\left(\frac{\partial \omega}{\partial t} + \omega \xi \cdot \nabla_x \omega \right) \operatorname{div}_\xi(\xi g) \right] = \frac{\omega^d}{\varepsilon} \tilde{\mathcal{Q}}(g, g),$$

where $\tilde{\mathcal{Q}}$ is such that $\tilde{\mathcal{Q}}(g, g) = \mathcal{Q}(f, f)$. This equation can actually be written in the more convenient conservative form:

$$(2.2) \quad \frac{\partial g}{\partial t} + \operatorname{div}_x(\omega \xi g) - \operatorname{div}_\xi \left[\left(\frac{1}{\omega} \frac{\partial \omega}{\partial t} \xi + \xi \otimes \xi \nabla_x \omega \right) g \right] = \frac{\omega^d}{\varepsilon} \tilde{\mathcal{Q}}(g, g).$$

In order to make this rescaling efficient, the main difficulty is now to choose an appropriate scaling function ω to define completely the distribution g .

2.1. The Generalized Filbet-Russo Scaling. The first and natural idea follows from the work [22]. It consists in computing the function ω directly from the distribution function f , by setting

$$(2.3) \quad \omega := \sqrt{\frac{2 E_f}{d \rho_f}},$$

where ρ_f and E_f are respectively the local mass and kinetic energy of f

$$(2.4) \quad \rho_f := \int_{\mathbb{R}^d} f(v) dv, \quad E_f := \int_{\mathbb{R}^d} f(v) \frac{|v|^2}{2} dv$$

We also define the mean velocity field \mathbf{u}_f and temperature T_f by

$$(2.5) \quad \mathbf{u}_f := \frac{1}{\rho_f} \int_{\mathbb{R}^d} f(v) v dv, \quad T_f := \frac{1}{d \rho_f} \left(2 E_f - \rho_f |\mathbf{u}_f|^2 \right).$$

Assuming that f is nonnegative, the quantity ω will then provide correct information on its support: if f is concentrated, ω will be small, whereas it will be large for scattered distributions. This approach has been shown to be very accurate for the space-homogeneous setting in [22], but it is difficult to extend to our case because the definition of ω yields very restrictive constraints on the moments of g .

Indeed using the change of variable $v \rightarrow \omega \xi$ in (2.4), we find that

$$\omega^2(t, x) = \frac{2 E_f(t, x)}{d \rho_f(t, x)} = \frac{2 E_g(t, x)}{d \rho_g(t, x)} \omega^2(t, x)$$

and then for a distribution function g given by (2.1), the condition (2.3) imposes that

$$\rho_g(t, x) = \frac{2}{d} E_g(t, x), \quad \forall (t, x) \in \mathbb{R}_+ \times \Omega.$$

Stated in another form, it means that

$$\int_{\mathbb{R}^d} g(t, x, \xi) d\xi = \frac{1}{d} \int_{\mathbb{R}^d} g(t, x, \xi) |\xi|^2 d\xi, \quad \forall (t, x) \in \mathbb{R}_+ \times \Omega.$$

This condition gives a very strong constraint on g since it has to be satisfied locally both in time and space. From a numerical point of view, it means that the time and space discretizations have to be done in such a way that this condition is also preserved at the discrete level, which becomes very restrictive on the choice of the numerical method. Moreover, given that ω depends on the distribution g itself, the linear free transport equation satisfied by f becomes a strongly nonlinear transport equation in g , which can be difficult to solve numerically.

2.2. The Macroscopic Scaling. In order to relax this constraint on the distribution function g , we propose a different choice of the function ω . Moreover, to avoid a strong coupling between ω and g leading to nonlinear terms, we do not want to compute ω directly from the distribution function g itself but only from macroscopic quantities which are assumed to be close enough to the one computed from f . Therefore, a good candidate will be a solution to the system of macroscopic equations obtained from a closure of the kinetic model (1.1).

More precisely, assume that in the asymptotic limit $\varepsilon \rightarrow 0$, the distribution function f solution to (1.1) converges towards a steady state $\mathcal{M}_{\rho, \mathbf{u}, T}$, where ρ is the *mass*, \mathbf{u} the *velocity* and T the *temperature*. Then, a first order approximation with respect to ε would be

$$f^\varepsilon = \mathcal{M}_{\rho, \mathbf{u}, T} + \mathcal{O}(\varepsilon),$$

where the macroscopic quantities are solutions to [25]

$$(2.6) \quad \begin{cases} \partial_t \rho + \operatorname{div}_x(\rho \mathbf{u}) = 0, \\ \partial_t(\rho \mathbf{u}) + \operatorname{div}_x(\rho \mathbf{u} \otimes \mathbf{u} + p \mathbf{I}) = 0, \\ \partial_t E + \operatorname{div}_x(\mathbf{u}(E + p)) = G(\rho, \mathbf{u}, E). \end{cases}$$

In this last system, E is the kinetic energy given by

$$2E = d\rho T + \rho |\mathbf{u}|^2,$$

$p > 0$ is the pressure and $G \in \mathbb{R}$ the “energy” of the collision operator. A natural choice for ω is to set

$$(2.7) \quad \omega := \sqrt{\frac{2E}{d\rho}}.$$

The scaling function ω becomes independent of the original distribution but still follows the change of scales in velocity, provided that the closure has been done properly. The system (2.2)-(2.6) is now uncoupled which is better both for a numerical and mathematical point of view. Moreover, it will be easier to solve thanks to the hyperbolic structure of (2.6).

In the following section we give several examples of application of such a method.

3. APPLICATIONS OF THE RESCALING VELOCITY METHOD

We start with the simple example of the classical Boltzmann equation and then present different applications to inelastic kinetic models for granular gases.

3.1. Application to the Boltzmann Equation. The Boltzmann equation describes the behavior of a dilute gas of particles when the only interactions taken into account are binary elastic collisions. It is given by equation (1.1) where $f^\varepsilon(t, x, v)$ is the time-dependent particle distribution function in the phase space and the Boltzmann collision operator \mathcal{Q}_B is a quadratic operator, local in (t, x)

$$(3.1) \quad \mathcal{Q}_B(f, f)(v) = \int_{\mathbb{R}^d} \int_{\mathbb{S}^{d-1}} B(|v - v_*|, \cos \theta) [f'_* f' - f_* f] d\sigma dv_*,$$

where we used the shorthand $f = f(v)$, $f_* = f(v_*)$, $f' = f(v')$, $f'_* = f(v'_*)$. The velocities of the colliding pairs (v, v_*) and (v', v'_*) are related by

$$v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2} \sigma, \quad v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2} \sigma.$$

The *collision kernel* B is a non-negative function which by physical arguments of invariance only depends on $|v - v_*|$ and $\cos \theta = \hat{u} \cdot \sigma$, where $\hat{u} = (v - v_*)/|v - v_*|$.

Boltzmann's collision operator has the fundamental properties of conserving mass, momentum and energy

$$\int_{\mathbb{R}^d} \mathcal{Q}_{\mathcal{B}}(f, f) \varphi(v) dv = 0, \quad \varphi(v) = 1, v, |v|^2$$

and satisfies the well-known Boltzmann's H -theorem (for the space homogeneous case)

$$-\frac{d}{dt} \int_{\mathbb{R}^d} f \log f dv = - \int_{\mathbb{R}^d} \mathcal{Q}_{\mathcal{B}}(f, f) \log(f) dv \geq 0.$$

The functional $-\int f \log f dv$ is the entropy of the solution. Boltzmann H -theorem implies that any equilibrium distribution function, *i.e.* any function which is a maximum of the entropy, has the form of a locally Maxwellian distribution [16, 25]

$$(3.2) \quad \mathcal{M}_{\rho, u, T}(v) = \frac{\rho}{(2\pi T)^{d/2}} \exp\left(-\frac{|u - v|^2}{2T}\right),$$

where the density, velocity and temperature of the gas ρ , u and T are computed from the distribution function as in (2.4)-(2.5). For further details on the physical background and derivation of the Boltzmann equation we refer to [16, 43].

Assuming that the collision kernel is given by the *variable hard spheres* model, that is,

$$B(|z|, \cos \theta) = |z|^\lambda b(\cos \theta)$$

for $\lambda \in [0, 1]$ and b bounded, then for a distribution g given by (2.1), we have using (3.1)

$$\mathcal{Q}_{\mathcal{B}}(f, f) = \omega^{\lambda-d} \mathcal{Q}_{\mathcal{B}}(g, g)$$

and the rescaled equation (2.2) reads

$$(3.3) \quad \frac{\partial g}{\partial t} + \operatorname{div}_x(\omega \xi g) - \operatorname{div}_\xi \left[\left(\frac{1}{\omega} \frac{\partial \omega}{\partial t} \xi + \xi \otimes \xi \nabla_x \omega \right) g \right] = \frac{\omega^\lambda}{\varepsilon} \mathcal{Q}_{\mathcal{B}}(g, g).$$

Moreover, applying a standard closure of the Boltzmann equation by a Maxwellian distribution (3.2), we find that the macroscopic quantities and then ω thanks to (2.7), are given by

$$(3.4) \quad \begin{cases} \partial_t \rho + \operatorname{div}_x(\rho \mathbf{u}) = 0, \\ \partial_t(\rho \mathbf{u}) + \operatorname{div}_x(\rho \mathbf{u} \otimes \mathbf{u} + \rho T \mathbf{I}) = 0, \\ \partial_t E + \operatorname{div}_x(\mathbf{u}(E + \rho T)) = 0. \end{cases}$$

We shall now apply the rescaling method to the granular gases equation.

3.2. Application to the Granular Gases Equation. A granular gas is a set of particles which interact by energy dissipating collisions leading to inelastic collisions. This inelasticity is characterized by a collision mechanics where mass and momentum are conserved and kinetic energy is dissipated. Thus, the collision phenomenon is a non-*microreversible* process. The velocities of the colliding pairs (v, v_*) and (v', v'_*) are related by

$$v' = v - \frac{1+e}{2} (u \cdot \sigma) \sigma, \quad v'_* = v_* + \frac{1+e}{2} (u \cdot \sigma) \sigma,$$

where $u := v - v_*$ is the *relative velocity*, $\sigma \in \mathbb{S}^{d-1}$ the *impact direction* and $e \in [0, 1]$ the dissipation parameter, known as *restitution coefficient*. It means that the energy is dissipated in the impact direction only and will be chosen constant. The parametrization of post-collisional velocities can also be found by using some properties of the model. Indeed, it is equivalent to the conservation of impulsions and dissipation of energy:

$$(3.5) \quad \begin{cases} v' + v'_* = v + v_*, \\ |v'|^2 + |v'_*|^2 - |v|^2 - |v_*|^2 = -\frac{1-e^2}{2} |u \cdot \sigma|^2 \leq 0. \end{cases}$$

This microscopic mechanism allows us to describe the granular collision operator $\mathcal{Q}_{\mathcal{I}}$. If the nonnegative particles distribution function f depends only on $v \in \mathbb{R}^d$, the collision operator $\mathcal{Q}_{\mathcal{I}}(f, g)$ can be expressed in the following weak form: given a smooth test function ψ ,

$$(3.6) \quad \int_{\mathbb{R}^d} \mathcal{Q}_{\mathcal{I}}(f, f) \psi \, dv := \frac{1}{2} \int_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{S}^{d-1}} |v - v_*| f_* f (\psi' + \psi'_* - \psi - \psi_*) b(\cos \theta) \, d\sigma \, dv \, dv_*,$$

where we have used the shorthand $\psi' := \psi(v')$, $\psi'_* := \psi(v'_*)$, $\psi := \psi(v)$ and $\psi_* := \psi(v_*)$.

Taking $\psi(v) = 1$, v and $|v|^2$ in (3.6), the relations (3.5) yield conservation of mass and momentum at the kinetic level and dissipation of kinetic energy

$$\int_{\mathbb{R}^d} \mathcal{Q}_{\mathcal{I}}(f, f) |v|^2 \, dv = -D(f).$$

where $D(f)$ is the *energy dissipation functional*, given by

$$(3.7) \quad D(f) = C(1 - e^2) \iint_{\mathbb{R}^d \times \mathbb{R}^d} f f_* |v - v_*|^3 \, dv \, dv_* \geq 0,$$

for C an explicit, nonnegative constant depending on the cross section b and the dimension. The decrease of the energy, together with conservation of mass and momentum imply that the equilibria of the collision operator are Dirac distributions $\rho \delta_{\mathbf{u}}(v)$ where the density ρ and momentum $\rho \mathbf{u}$ are prescribed. This is the major problem for the numerical study of equation (1.1) with deterministic algorithms because of this concentration in velocity variables. In that case, the rescaling method (2.1) becomes an essential tool to follow the concentration in velocity of the distribution function.

For the hard sphere kernel (3.6) and for a rescaled distribution function g given by (2.1), we have

$$\mathcal{Q}_{\mathcal{I}}(f, f) = \omega^{1-d} \mathcal{Q}_{\mathcal{I}}(g, g).$$

The rescaled equation (2.2) then reads

$$(3.8) \quad \frac{\partial g}{\partial t} + \operatorname{div}_x(\omega \xi g) - \operatorname{div}_\xi \left[\left(\frac{1}{\omega} \frac{\partial \omega}{\partial t} \xi + \xi \otimes \xi \nabla_x \omega \right) g \right] = \frac{\omega}{\varepsilon} \mathcal{Q}_{\mathcal{I}}(g, g).$$

To complete the rescaling procedure, it remains to derive a set of equations for ρ and E to define the rescaling function ω . To this end, we will assume that the restitution coefficient e is close to 1, which corresponds to the weak inelasticity case, in order to write the collision operator as a perturbation of the classical Boltzmann operator and then write the system (2.6) associated to granular gases. Let us mention that there are alternative methods for deriving hydrodynamic equations from dissipative kinetic models, which can be found in the review paper of M. Bisi, J.A. Carrillo and G. Spiga [6]. Here, we focus on the approach developed by G. Toscani [40, 41]: we write the granular gases operator as a perturbation of the elastic Boltzmann operator

$$(3.9) \quad \mathcal{Q}_{\mathcal{I}}(f, f) = \mathcal{Q}_{\mathcal{B}}(f, f) + \varepsilon \mathcal{I}(f, f) + \mathcal{O}(\varepsilon^2),$$

where \mathcal{I} is a quadratic, energy dissipative nonlinear friction operator given by

$$\mathcal{I}(f, f)(v) = \frac{1}{2} \operatorname{div}_v \left(\int_{\mathbb{R}^d \times \mathbb{S}^{d-1}} |u \cdot \sigma| (u \cdot \sigma) \sigma f_* f b(\cos \theta) \, d\sigma \, dv_* \right).$$

Using the expansion (3.9) of the operator with respect to ε , it was shown in [40] that we can derive the macroscopic system (2.6) associated to $\mathcal{Q}_{\mathcal{I}}$. Assuming that the inelasticity is of the same order of the Knudsen number:

$$(3.10) \quad 1 - e = \varepsilon,$$

then, up to the second order in ε , the macroscopic quantities are solution to

$$(3.11) \quad \begin{cases} \partial_t \rho + \operatorname{div}_x(\rho \mathbf{u}) = 0, \\ \partial_t(\rho \mathbf{u}) + \operatorname{div}_x(\rho(\mathbf{u} \otimes \mathbf{u}) + \rho T \mathbf{I}) = 0, \\ \partial_t E + \operatorname{div}_x(\mathbf{u}(E + \rho T)) = -K_d \rho^2 T^{3/2}, \end{cases}$$

for an explicit and nonnegative constant K_d [40]. The solution of this system will give the scaling function ω in (2.7).

3.3. Application to a Simplified Granular Gases Model. The rescaling velocity method can be also applied to a simplified model designed to mimic the behavior of the granular gases equation with a simpler collision operator. Indeed, A. Astillero and A. Santos derived in [2, 3] a very simple approximation of the granular gases equation, containing its main features. It consists in replacing the *bilinear* collision operator $\mathcal{Q}_{\mathcal{I}}$ by a *nonlinear* relaxation operator, supplemented with a drag force in the velocity space, depending on the moment of the particle distribution function itself, in order to match the correct macroscopic dissipation of kinetic energy of the granular gases equation.

The relaxation operator we will use is the so-called Ellipsoidal Statistical BGK operator (ES-BGK) [1]. To this aim we first define some macroscopic quantities of the particle distribution function f such as the opposite of the *stress tensor*

$$\Theta_f(t, x) = \frac{1}{\rho_f} \int_{\mathbb{R}^d} (v - \mathbf{u}_f) \otimes (v - \mathbf{u}_f) f(t, x, v) dv.$$

Therefore the *translational temperature* is related to the stress tensor as $T_f = \text{tr}(\Theta_f)/d$. We finally introduce the corrected tensor

$$\mathcal{T}_f(t, x) = [(1 - \nu) T_f \mathbf{I} + \nu \Theta_f](t, x),$$

which can be viewed as a linear combination of the initial stress tensor Θ_f and of the isotropic stress tensor $T_f \mathbf{I}$ developed by a Maxwellian distribution. The parameter $-\infty < \nu < 1$ is used to modify the value of the Prandtl number through the formula

$$0 \leq \text{Pr} = \frac{1}{1 - \nu} \leq +\infty \quad \text{for } \nu \in (-\infty, 1).$$

Thus we introduce a corrected Gaussian $\mathcal{G}[f]$ defined by

$$\mathcal{G}[f] = \frac{\rho_f}{\sqrt{\det(2\pi \mathcal{T}_f)}} \exp\left(-\frac{(v - \mathbf{u}_f) \mathcal{T}_f^{-1} (v - \mathbf{u}_f)}{2}\right)$$

and the corresponding collision operator is now

$$(3.12) \quad \mathcal{Q}_{\mathcal{S}}(f) = \tau(P_f) (\mathcal{G}[f] - f) + C \rho_f T_f^{1/2} \text{div}_v((v - \mathbf{u}_f)f),$$

where τ is a given function depending on the *kinetic pressure* $P_f := \rho_f T_f$ and C is given by

$$C = (1 - e) \frac{K_d}{d}.$$

The coefficient e corresponds to the restitution coefficient of the granular operator $\mathcal{Q}_{\mathcal{I}}$ and will be assumed equal to $1 - \varepsilon$ as in (3.10). This expression yields the correct granular energy dissipation (3.7).

This new operator replaces the inelastic collision operator by an “equivalent” elastic operator and the drift term acts like an external drag force which mimics the cooling effect of the particles. In particular, the hydrodynamic equations associated to the rescaled equation (2.2) with operator (3.12) are exactly given by the system (3.11). The operator $\tilde{\mathcal{Q}}_{\mathcal{S}}$ such that $\tilde{\mathcal{Q}}_{\mathcal{S}}(g) = \mathcal{Q}_{\mathcal{S}}(f)$ is given by

$$\tilde{\mathcal{Q}}_{\mathcal{S}}(g) = \frac{1}{\omega^d} \left[\tau_g (\mathcal{G}[g] - g) + C \omega \rho_g T_g^{1/2} \text{div}_{\xi}((\xi - \mathbf{u}_g) g) \right],$$

using the shorthand $\tau_g := \tau(\omega^2 P_g)$. Equation (2.2) then simply reads

$$\frac{\partial g}{\partial t} + \text{div}_x(\omega \xi g) = \frac{\tau_g}{\varepsilon} (\mathcal{G}[g] - g) + \text{div}_{\xi} \left[\left(C \omega \rho_g T_g^{1/2} (\xi - \mathbf{u}_g) + \frac{1}{\omega} \frac{\partial \omega}{\partial t} \xi + \xi \otimes \xi \nabla_x \omega \right) g \right],$$

where the function ω is given by (2.7) using the closed system (3.11).




FIGURE 1. Schematic representation of Maxwell conditions at the boundary.

4. BOUNDARY CONDITIONS

In order to define completely the mathematical problem for the kinetic equation (1.1), one has to set boundary conditions on $\partial\Omega$. The most simple (yet physically relevant) are the so-called Maxwell boundary conditions [28], where a fraction $1 - \alpha \in [0, 1]$ of the particles is *specularly* (elastically) reflected by a wall, whereas the remaining fraction α is *thermalized*, leaving the wall (moving at a velocity \mathbf{u}_w) at a temperature $T_w > 0$ (see Figure 1). The parameter α is called *accommodation* coefficient, because it expresses the tendency of the gas to accommodate to the state of the wall. More details on the subject can be found in the textbook of Cercignani [15]. The main difficulty consists to apply appropriate boundary conditions on the macroscopic model which are consistent with the kinetic boundary value problem.

Let us consider a distribution function $f(t, x, v)$ solution to (1.1) for $(t, x, v) \in \mathbb{R}_+ \times \Omega \times \mathbb{R}^d$ with Ω a bounded Lipschitz-continuous domain of \mathbb{R}^d . We assume the boundary $\partial\Omega$ to have a unit outer normal \mathbf{n}_x for all $x \in \partial\Omega$. Then, for an *incoming* velocity $v \in \mathbb{R}^d$ (namely $v \cdot \mathbf{n}_x \leq 0$), we set

$$f(t, x, v) = \alpha \mathcal{R}f(t, x, v) + (1 - \alpha) \mathcal{M}f(t, x, v), \quad \forall x \in \partial\Omega, v \cdot \mathbf{n}_x \leq 0,$$

with

$$(4.1) \quad \begin{cases} \mathcal{R}f(t, x, v) = f(t, x, v - 2(\mathbf{n}_x \cdot v) \mathbf{n}_x), \\ \mathcal{M}f(t, x, v) = \mu(t, x) \mathcal{M}_{1, \mathbf{u}_w, T_w}(v). \end{cases}$$

In this last expression, $\mathcal{M}_{1, \mathbf{u}_w, T_w}$ is the wall Maxwellian

$$\mathcal{M}_{1, \mathbf{u}_w, T_w}(v) := \frac{1}{(2\pi T_w)^{d/2}} \exp\left(-\frac{|v - \mathbf{u}_w|^2}{2T_w}\right)$$

and μ insures global mass conservation:

$$(4.2) \quad \int_{v \cdot \mathbf{n}_x \geq 0} f(t, x, v) v \cdot \mathbf{n}_x dv = -\mu(t, x) \int_{v \cdot \mathbf{n}_x < 0} \mathcal{M}_{1, \mathbf{u}_w, T_w}(v) v \cdot \mathbf{n}_x dv.$$

Thanks to these definitions, we have the following result:

Proposition 4.1. *Let f be a smooth solution to the kinetic equation (1.1) with Maxwell boundary conditions (4.1), $\Psi \in C^1(\mathbb{R}_+, \mathbb{R}_+)$ and $\varphi \in C^1(\mathbb{R}^d, \mathbb{R})$. If φ is symmetric with respect to the normal component*

of the velocity, then we have for any $x \in \partial\Omega$ and $t \in \mathbb{R}_+$:

$$(4.3) \quad \begin{cases} \int_{v \cdot \mathbf{n}_x \leq 0} \Psi(\mathcal{R}f(t, x, v)) v \cdot \mathbf{n}_x \varphi(v) dv = - \int_{v \cdot \mathbf{n}_x \geq 0} \Psi(f(t, x, v)) v \cdot \mathbf{n}_x \varphi(v) dv, \\ \int_{v \cdot \mathbf{n}_x \leq 0} \Psi(\mathcal{R}f(t, x, v)) v \cdot \boldsymbol{\tau}_x \varphi(v) dv = \int_{v \cdot \mathbf{n}_x \geq 0} \Psi(f(t, x, v)) v \cdot \boldsymbol{\tau}_x \varphi(v) dv, \end{cases}$$

where $\boldsymbol{\tau}_x$ belongs to \mathbf{n}_x^\perp . Moreover, we also have

$$(4.4) \quad \int_{v \cdot \mathbf{n}_x \leq 0} \mathcal{M}f(t, x, v) v \cdot \mathbf{n}_x dv = - \int_{v \cdot \mathbf{n}_x \geq 0} f(t, x, v) v \cdot \mathbf{n}_x dv.$$

Proof. Concerning (4.3), for any $x \in \partial\Omega$, we integrate on the set $\{v \cdot \mathbf{n}_x \leq 0\}$ the function $v \mapsto \Psi(\mathcal{R}f(t, x, v)) v \cdot \mathbf{n}_x \varphi(v)$. The transformation $v' = v - 2(\mathbf{n}_x \cdot v) \mathbf{n}_x$ being isometric, its Jacobian is equal to 1 and the change of variable $v \mapsto v'$ preserves the Lebesgue measure dv . Moreover, we have the identities $v' \cdot \mathbf{n}_x = -v \cdot \mathbf{n}_x$ and $v' \cdot \boldsymbol{\tau}_x = v \cdot \boldsymbol{\tau}_x$. Hence, writing $v = (v \cdot \mathbf{n}_x) \mathbf{n}_x + (v \cdot \boldsymbol{\tau}_x) \boldsymbol{\tau}_x$ yields using the symmetry of φ

$$\begin{aligned} \int_{v \cdot \mathbf{n}_x \leq 0} \Psi(\mathcal{R}f(t, x, v)) v \cdot \mathbf{n}_x \varphi(v) dv &= \int_{v \cdot \mathbf{n}_x \leq 0} \Psi(f(t, x, v')) v \cdot \mathbf{n}_x \varphi(v) dv \\ &= - \int_{v' \cdot \mathbf{n}_x \geq 0} \Psi(f(t, x, v')) v' \cdot \mathbf{n}_x \varphi(v') dv', \end{aligned}$$

which proves the first part of (4.3). The second part is obtained exactly the same way.

The relation (4.4) is straightforward using the definition (4.2) of μ . \square

Thanks to this proposition, we can state some of the properties of equation (1.1) under Maxwell boundary conditions:

Corollary 4.2. *Let f be a smooth solution to the kinetic equation (1.1) with Maxwell boundary conditions (4.1). Then,*

- (1) *the mass of f is globally conserved for any accommodation coefficient $\alpha \in [0, 1]$;*
- (2) *if \mathcal{Q} preserves the kinetic energy, then in the specular case $\alpha = 1$ the energy of f is globally conserved;*
- (3) *if \mathcal{Q} produces entropy, namely*

$$- \int_{\mathbb{R}^d} \mathcal{Q}_{\mathcal{B}}(f, f) \log(f) dv \geq 0$$

then in the specular case $\alpha = 1$ the entropy functional $\int f \log f dv dx$ is dissipated.

Proof. (1) We start by multiplying equation (1.1) by φ and integrating in velocity and space variables. We have made the general assumption that \mathcal{Q} preserves the mass. Then, using the Stokes formula and taking $\Psi(f) = f$ and $\varphi(v) = 1$ (which is symmetric with respect to the normal component of v) in (4.3) is enough to guarantee the mass conservation for the specular boundary conditions. The relation (4.4) allows to conclude in the diffusive case.

(2) In the pure specular case, the elastic collisions at the boundary yield the microscopic conservation of kinetic energy: $|v'|^2 = |v|^2$. Then, taking $\Psi(f) = f$ and $\varphi(v) = |v|^2$ (which is symmetric with respect to the normal component of v) in (4.3) and using the Stokes formula gives the conservation of kinetic energy.

(3) Multiplying equation (1.1) by $\log f + 1$ and taking $\Psi(f) = f \log f$, $\varphi(v) = 1$ in (4.3) yields the dissipation of entropy, thanks to Stokes formula. \square

We shall now present the boundary conditions associated to the Maxwell conditions for the rescaled distribution g and the hydrodynamics quantities $(\rho, \rho \mathbf{u}, E)$ solution to the Euler' system. We split in two parts, pure specular and pure diffusive boundary conditions.

4.1. Specular Boundary Conditions. The natural translation of a specular reflection of a particle of velocity v in the physical space, at a point x of the boundary $\partial\Omega$ is a specular reflection at the point x , but with a rescaled velocity $\xi = v\omega$. More precisely, for an incoming velocity ξ (namely $\xi \cdot \mathbf{n}_x \leq 0$), we set

$$\mathcal{R}g(t, x, \xi) = g(t, x, [\xi - 2(\mathbf{n}_x \cdot \xi) \mathbf{n}_x]).$$

Concerning the macroscopic quantities, the relation (4.3) of Proposition 4.1 imposes the so-called *slip boundary* condition [39]. More precisely, it corresponds to a zero-flux boundary conditions for the mean velocity:

$$\mathbf{u} \cdot \mathbf{n}_x = 0.$$

This condition yields in particular the global conservation of mass and kinetic energy that we observed at the kinetic level in Corollary 4.2. It then remains to fix a condition for the momentum flux. To mimic the specular reflection, we reverse the velocity at the boundary, as presented in [4].

4.2. Diffusive Boundary Conditions. We want to impose a velocity \mathbf{u}_w and a temperature T_w at a point x of the boundary, using a wall maxwellian $\mathcal{M}_{1, \mathbf{u}_w, T_w}$ in the physical space. Then, for an incoming velocity ξ in the rescaled space, we set

$$\mathcal{M}g(t, x, \xi) = \mu(t, x) \mathcal{M}_{1, \tilde{\mathbf{u}}_w, \tilde{T}_w}(\xi).$$

In this last expression, $\mathcal{M}_{1, \tilde{\mathbf{u}}_w, \tilde{T}_w}$ is the ‘‘rescaled’’ wall Maxwellian, with $\tilde{\mathbf{u}}_w = \omega \mathbf{u}_w$, $\tilde{T}_w := \omega^2 T_w$ and μ insures the global mass conservation:

$$\int_{\xi \cdot \mathbf{n}_x > 0} g(t, x, \xi) \xi \cdot \mathbf{n}_x d\xi = -\mu(t, x) \int_{\xi \cdot \mathbf{n}_x < 0} \mathcal{M}_{1, \tilde{\mathbf{u}}_w, \tilde{T}_w}(\xi) \xi \cdot \mathbf{n}_x d\xi.$$

At the macroscopic level, the relation (4.4) of Proposition 4.1 imposes a zero-flux boundary conditions for the mean velocity:

$$\mathbf{u} \cdot \mathbf{n}_x = 0.$$

This yields the global conservation of mass for the macroscopic model, corresponding to the result of Corollary 4.2. We also want to fix the temperature $T_w > 0$ of the wall. We set

$$E(x) = \frac{d}{2} \rho(x) T_w.$$

These boundary conditions allow to solve the boundary value problems (3.4) or (3.11).

5. DISCRETIZATION OF THE PROBLEM

We will present in this Section the numerical schemes we used for the discretization of the uncoupled models (2.2) and (2.6). We need to distinguish three independent problems: the treatment of the free transport term, the collision term and the system of conservation laws.

5.1. The Finite Volume Method for a Conservative Transport Equation. In this subsection, we focus on the space discretization of the conservative transport equation

$$(5.1) \quad \begin{cases} \frac{\partial g}{\partial t} + \operatorname{div}_X (a(t, X) g) = 0, \quad \forall (t, X) \in \mathbb{R}_+ \times \mathcal{O}, \\ g(0, X) = g_0(X), \end{cases}$$

for a smooth function $a : \mathbb{R}_+ \times \mathcal{O} \rightarrow \mathbb{R}^d$ and a Lipschitz-continuous domain $\mathcal{O} \subset \mathbb{R}^d$. Our approach of the problem will be made in the framework of finite volume schemes. We shall present the spatial discretization of (5.1) in one space dimension for simplicity purposes. The extension for Cartesian grid in the multidimensional case will then be straightforward.

In the one dimensional setting, the domain $\mathcal{O} = (a, b)$ is a finite interval of \mathbb{R} . We define a mesh of \mathcal{O} , not necessarily uniform, by introducing a sequence of N_x control volume $K_i := (x_{i-\frac{1}{2}}, x_{i+\frac{1}{2}})$ for $i \in \llbracket 0, N_x \rrbracket$ with $x_i := (x_{i-\frac{1}{2}} + x_{i+\frac{1}{2}}) / 2$ and

$$a = x_{\frac{1}{2}} < x_1 < \dots < x_{i-\frac{1}{2}} < x_i < x_{i+\frac{1}{2}} < \dots < x_{N_x} < x_{N_x+\frac{1}{2}} = b.$$

The Lebesgue measure of the control volume is then simply $m(K_i) = x_{i+\frac{1}{2}} - x_{i-\frac{1}{2}}$. Let $g_i = g_i(t)$ be an approximation of the mean value of g over a control volume K_i . By integrating the transport equation (5.1) over K_i , we get the space discretized free transport equation

$$(5.2) \quad \begin{cases} m(K_i) \frac{\partial g_i}{\partial t}(t) + \mathcal{F}_{i+\frac{1}{2}} - \mathcal{F}_{i-\frac{1}{2}} = 0, \quad \forall t \in \mathbb{R}_+, \quad i \in \llbracket 1, N_x \rrbracket, \\ g_i(0) = \frac{1}{m(K_i)} \int_{K_i} g_0(x) dx. \end{cases}$$

In the equation (5.2), the quantity $\mathcal{F}_{i+\frac{1}{2}}$ is an approximation of the flux $x \mapsto a(t, x) g(t, x)$ at the cell interface $x_{i+\frac{1}{2}}$. We choose to use in the following upwind fluxes with the second order Van Leer's slope limiter [42]. If we denote by $A_i(t) := a(t, x_i)$ the velocity at the cell center, then the slope limited upwind flux is given by

$$\mathcal{F}_{i+\frac{1}{2}} = A_i^+ g_{i+\frac{1}{2},-} - A_i^- g_{i+\frac{1}{2},+}$$

where $x^+ := \max(x, 0)$ and $x^- := \max(-x, 0)$ are respectively the positive and negative part of a real number x and $g_{i+\frac{1}{2},\pm}$ is the slope limited reconstruction of g at the cell interface, namely

$$\begin{cases} g_{i+\frac{1}{2},-} = g_i + \frac{1}{2} \phi(\theta_i) (g_{i+1} - g_i), \\ g_{i+\frac{1}{2},+} = g_i - \frac{1}{2} \phi(\theta_i) (g_{i+2} - g_{i+1}). \end{cases}$$

In this last expression, θ_i is the *slope*

$$\theta_i = \frac{g_i - g_{i-1}}{g_{i+1} - g_i}$$

and ϕ is the so-called *Van Leer's limiter*

$$\phi(\theta) := \frac{\theta + |\theta|}{1 + \theta}.$$

5.2. The Spectral Method for the Collision Operator. We now focus on the numerical resolution of

$$(5.3) \quad \frac{\partial g}{\partial t} = \mathcal{Q}(g, g),$$

where $\mathcal{Q} = \mathcal{Q}_B$ or \mathcal{Q}_L . These Boltzmann-like operators will be computed using a spectral method. Fourier techniques for the resolution of the Boltzmann equation have been first introduced independently by L. Pareschi and B. Perthame in [32] and by A. Bobylev and S. Rjasanow in [7]. It has since been investigated by a lot of authors. One can find numerous results about this method in the article [33] of L. Pareschi and G. Russo. It has been recently shown to be convergent by F. Filbet and C. Mouhot in [19] for the elastic case and allows spectral accuracy [10]. It has then been derived for the granular gases operator (3.6) by F. Filbet, L. Pareschi and G. Toscani in [21]. Let us present the main features of this method.

Any deterministic numerical method for the computation of the Boltzmann operator requires to work on a *bounded* velocity space. Our approach consists in adding some non physical binary collisions by periodizing the particle distribution function and the collision operator. This implies the loss of some local invariant, but a careful periodization allows at least the preservation of mass. This periodization is the basis of spectral methods.

We thus consider a bounded velocity domain $\mathcal{D}_V := [-V, V]^d$. If g is compactly supported in \mathcal{D}_V , the collision operator spreads the support of g by a factor at most equal to $\sqrt{2}$. This was shown in Lemma 3.1 of [21] using the weak formulation (3.6) of $\mathcal{Q}_{\mathcal{I}}$ and can actually be improved in dimension 1. In this latter case, G. Naldi and G. Toscani showed in [31] that the size of the support is conserved by $\mathcal{Q}_{\mathcal{I}}$. Using these results, simple geometric considerations [33] shows that one can truncate the weak form of $\mathcal{Q}_{\mathcal{I}}$ to made him act on the set $\mathcal{U} := \mathcal{D}_V \times B(0, 2\lambda V) \times \mathbb{S}^{d-1}$: for any smooth test function ψ , the truncated collision operator $\mathcal{Q}_{\mathcal{I}}^V$ is given by

$$(5.4) \quad \int_{\mathbb{R}^d} \mathcal{Q}_{\mathcal{I}}^V(g, g) \psi \, dv := \frac{1}{2} \int_{\mathcal{U}} |v - v_*| g_* g (\psi' - \psi) b(\cos \theta) \, d\sigma \, dv \, dv_*.$$

The coefficient $\lambda := (3 + \sqrt{2})/2$ is an *anti-aliasing* parameter used when periodizing the distribution g .

The approximate solution to the periodized equation is represented as a truncated Fourier sum g_{N_v} , defined for $N_v \in \mathbb{N}$ by

$$g_{N_v}(v) = \sum_{|k| \leq N_v} \hat{g}_k \psi_k(v), \quad \psi_k(v) := e^{i \frac{\pi}{V} k \cdot v}.$$

In this expression, the sum is taken over the multi-index $k = (k_1, \dots, k_d) \in \mathbb{Z}^d$ and \hat{g}_k stands for the k^{th} Fourier coefficient of g , namely

$$\hat{g}_k := \frac{1}{(2V)^d} \int_{\mathcal{D}_V} g(v) \psi_{-k}(v) \, dv.$$

In a Fourier-Galerkin spectral method, the unknowns are the coefficients $(\hat{g}_k(t))_{k \in \llbracket -N_v, N_v \rrbracket^d}$. They are obtained for any $k \in \llbracket -N_v, N_v \rrbracket^d$ by the solution to the Fourier transformed, truncated equation (5.3)

$$\frac{\partial \hat{g}_k}{\partial t} = \left\langle \mathcal{Q}_{\mathcal{I}}^V(g_{N_v}, g_{N_v}), \psi_{-k} \right\rangle_{L^2(\mathcal{D}_V)}.$$

The Fourier coefficients of the truncated collision operator $\mathcal{Q}_{\mathcal{I}}^V$ are given using (5.4) by

$$(5.5) \quad \begin{aligned} \left\langle \mathcal{Q}_{\mathcal{I}}^V(g_{N_v}, g_{N_v}), \psi_{-k} \right\rangle_{L^2(\mathcal{D}_V)} &= \frac{1}{2(2V)^d} \int_{\mathcal{U}} |u| g_{N_v}(v) g_{N_v}(v-u) \left(e^{-i \frac{\pi}{V} k \cdot v'} - e^{-i \frac{\pi}{V} k \cdot v} \right) \, d\sigma \, dv \, du, \\ &= \sum_{\substack{l, m = -N_v \\ l+m=k}}^{N_v} \hat{g}_l \hat{g}_m \left(\hat{B}(l, m) - \hat{\gamma}(m) \right), \end{aligned}$$

where $\hat{B}(l, m)$ and $\hat{\gamma}(m)$ are defined for $l, m \in \llbracket -N_v, N_v \rrbracket^d$ by

$$\begin{cases} \hat{B}(l, m) = \frac{1}{2} \int_{\mathcal{B}(0, 2\lambda V)} |u| e^{-i \frac{\pi}{V} \left(\left(\frac{3-e}{4} \right) u \cdot m - \left(\frac{1+e}{4} \right) u \cdot l \right)} \left(\int_{\mathbb{S}^{d-1}} e^{-i \frac{\pi}{V} \left(\frac{1+e}{4} \right) |u|(l+m) \cdot \omega} \, d\omega \right) \, du, \\ \hat{\gamma}(m) = \frac{|\mathbb{S}^{d-1}|}{2} \int_{\mathcal{B}(0, 2\lambda V)} |u| e^{-iu \cdot m} \, du. \end{cases}$$

The computation of $\mathcal{Q}_{\mathcal{I}}(g_{N_v}, g_{N_v})$ in Fourier variables is then reduced to the evaluation of a sum, which has the same structure (bilinear in g and written as the difference of two terms, the gain and loss one) than the integral in classical variables.

Remark 5.1. *Notice that the kernel modes have only to be computed once, which is of great interest for the numerical implementation. Moreover, it can be done explicitly for $d = 1$ and $d = 3$ [21]. The case $d = 2$ needs a numerical integration, which can be made as precise as one wants and then stored for further use.*

It then remains to project the Fourier collision operator (5.5) in the space \mathbb{P}_{N_v} of trigonometric polynomials of degree at most N_v in each direction, using the orthogonal projection of $L^2(\mathcal{D}_V)$ onto \mathbb{P}_{N_v} .

5.3. The WENO Approach for a System of Conservation Laws. The fluid equations (3.11) can be written as a system of nonlinear hyperbolic conservation laws with source term

$$\frac{\partial U}{\partial t} + \operatorname{div}_x F(U) = G(U),$$

where $U : \mathbb{R}_+ \times \Omega \rightarrow \mathbb{R}^n$, $F, G : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $n = d + 2$. There exists a large number of numerical methods to solve this type of problem, and we choose to use a finite difference approach.

The reconstruction of the fluxes will be made by a high-order Weighted Essentially Non-Oscillatory (*WENO*) method with a 5 points stencil. Indeed, it is very well adapted to the treatment of shocks in fluid system. The reader can consult the lecture notes [38] of C.W. Shu for most of the details about this method. Let us present it in the one dimensional setting and on an uniform grid.

We will approximate the solution U at time t and position $x_i = i\Delta x$, $\Delta x > 0$, by the vector $(U_i)_{i \in \llbracket 1, N_x \rrbracket}$ for $N_x \in \mathbb{N}$, solution to

$$\frac{d}{dt} U_i(t) + \frac{\mathcal{F}_{i+\frac{1}{2}} - \mathcal{F}_{i-\frac{1}{2}}}{\Delta x} = G(U_i), \quad \forall t \in \mathbb{R}_+, \quad i \in \llbracket 1, N_x \rrbracket,$$

The vector $(\mathcal{F}_{i+\frac{1}{2}})_{i \in \llbracket 1, N_x \rrbracket}$ is the numerical flux. We will adopt a characteristic-wise *WENO* method for the reconstruction of this flux. It can be summarized in the following way. Given some divided differences of the cell averages \bar{U}_i of $(U_i)_{i \in \llbracket 1, N_x \rrbracket}$ at time t :

- (1) Compute an average (also called intermediate) state $U_{i+\frac{1}{2}}$ using arithmetic mean.
- (2) Compute the right (resp. left) eigenvectors matrix $R(U_{i+\frac{1}{2}})$ (resp. $R^{-1}(U_{i+\frac{1}{2}})$) and the eigenvalues $\lambda_l(U_{i+\frac{1}{2}})$ of $F'(U_{i+\frac{1}{2}})$, then set

$$R = R(U_{i+\frac{1}{2}}), \quad R^{-1} = R^{-1}(U_{i+\frac{1}{2}}), \quad \Lambda = \operatorname{Diag}(\lambda_l(U_{i+\frac{1}{2}})).$$

- (3) Project the divided differences lying in the 5 points stencil of the reconstruction onto the local characteristics fields:

$$\bar{V}_j := R^{-1} \bar{U}_j, \quad \bar{H}_j := R^{-1} F(\bar{U}_j), \quad \forall j \in \llbracket i-5, i+5 \rrbracket.$$

- (4) Perform the scalar reconstruction of the local characteristic variables, namely
 - 4.1. Obtain the 5th order reconstructed values $V_{i+\frac{1}{2},+}$ and $V_{i+\frac{1}{2},-}$ of \bar{V}_i using the *WENO* procedure described in [38, Procedure 2.3];
 - 4.2. Compute the Lax-Friedrichs flux

$$H_{i,\pm} := \frac{1}{2} \left(\bar{H}_i + \max_{i,l} |\lambda_l(U_{i+\frac{1}{2}})| V_{i+\frac{1}{2},\pm} \right).$$

- (5) Go back to the physical space, thanks to the right eigenvectors

$$F_{i,\pm} := R H_{i,\pm}$$

and finally compute the numerical flux

$$\mathcal{F}_{i+\frac{1}{2}} = F_{i,+} + F_{i,-}.$$

The rescaling function ω (and its time and space derivatives) given by the macroscopic scaling (2.7) is then evolved in time thanks to the *WENO* fluxes $\mathcal{F}_{i+\frac{1}{2}}$.

We shall now present numerical simulations of the rescaling velocity method for energy dissipative equations.

6. NUMERICAL SIMULATIONS

6.1. Convergence toward a Dirac mass. In this section, we will deal with a space homogeneous distribution $f = f(t, v)$, solution to the granular gases equation

$$(6.1) \quad \begin{cases} \frac{\partial f}{\partial t} = \mathcal{Q}_{\mathcal{I}}(f, f), \\ f(t=0) = f_0, \end{cases}$$

with an initial condition chosen as the sum of two gaussian distributions of total temperature equal to 1:

$$f_0(v) := \frac{1}{2(2\pi\sigma^2)^{d/2}} \left(\exp\left(-\frac{|v - \sigma \mathbf{e}|^2}{2\sigma^2}\right) + \exp\left(-\frac{|v + \sigma \mathbf{e}|^2}{2\sigma^2}\right) \right), \quad \forall v \in \mathbb{R}^d,$$

where $\mathbf{e} = (1, \dots, 1)^\top$ and $\sigma = 1/10$. With this choice, we have

$$(6.2) \quad \int_{\mathbb{R}^d} f_0(v) \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} dv = \begin{pmatrix} 1 \\ 0 \\ d \end{pmatrix}.$$

The corresponding scaled distribution $g = g(t, \xi)$ given by (2.1) is then solution to the homogeneous granular equation with a time-dependent drift term

$$(6.3) \quad \begin{cases} \frac{\partial g}{\partial t} - \nabla_{\xi} \cdot \left(\frac{1}{\omega} \frac{d\omega}{dt} \xi g \right) = \omega \mathcal{Q}_{\mathcal{I}}(g, g), \\ g(t=0) = g_0. \end{cases}$$

The initial condition g_0 is chosen by $g_0 = f_0$ because $\omega(0) = 1$ according to (6.2).

To study the evolution of ω , we propose to compare the rescaling approach proposed by F. Filbet and G. Russo in [22] and the macroscopic one based on the closure of the kinetic model. In the space homogeneous setting, the mass $\rho = 1$ is preserved. Hence, we have

$$\omega(t) = \sqrt{\frac{2}{d} E(t)}$$

and it is enough to compute the time evolution of the kinetic energy E to obtain ω . Then on the one hand, following [22], we will study the distribution g solution to (6.3) coupled with the exact kinetic energy

$$(6.4) \quad \frac{dE}{dt} = \omega^3 \int_{\mathbb{R}^d} \mathcal{Q}_{\mathcal{I}}(g, g) \frac{|\xi|^2}{2} d\xi.$$

On the other hand, from the result of Subsection 3.2, we replace the ordinary differential equation (6.4) by

$$(6.5) \quad \frac{dE}{dt} = -K_d \rho^2 T^{3/2},$$

where $dT(t) = 2E(t)$ because of the conservation of momentum. Let us emphasize that (6.3)-(6.5) is an uncoupled model. We will then compare it with the solution to (6.1) in classical variables.

Description of the results. We start with a two dimensional test, designed to compare the results obtained with both coupled (6.3)-(6.4) and uncoupled models (6.3)-(6.5) and intended to show that the latter one will give better results. We choose the computational domain in velocity as the box $\mathcal{D}_V = [-V, V]^2$ with $V = 9$ to avoid some possible loss of mass due to the drift term. We take successively $N_v = 32$ and $N_v = 48$ half Fourier modes in each direction of the velocity space.

We present in Figure 2 the time evolution of the first marginal distribution

$$g^{(1)}(t, \xi_1) := \int_{\mathbb{R}} g(t, \xi_1, \xi_2) d\xi_2.$$

We can see that both models give close results for short time in the coarse and fine grids, but some oscillations eventually appear in the coupled model, leading to the breakout of the scheme. This is due to the wrong computation of the kinetic energy of g , because of the heavy tails of this distribution [29].




FIGURE 2. Comparison of the first marginal distribution $g^{(1)}$ for coupled model (6.3)-(6.4) (points) and uncoupled model (6.3)-(6.4) (dashed line), at time $t = 0.5, 10$ and 50 for $N_v = 32$ (left) and $N_v = 48$ (right).

This phenomenon does not occur anymore when we refine the grid. In this case, the results given by both models are in very good agreement, especially for the long time behavior: the equilibrium distribution is very well captured by the new uncoupled model.

We then compare the result of this test for the uncoupled model with a solution to equation (6.1) computed in classical variables with the spectral scheme. We take $V = 8$ and $N_v = 32$ half modes in each directions of the classical velocity space.

We observe in Figure 3 that both models give the same result for short time. Then, on the one hand, the solution obtained by the spectral scheme in classical variables breaks down for intermediate time ($t \sim 1$) because of the concentration which yields spurious oscillations. We can see in particular the loss of symmetry of the solution at time $t = 1$. On the other hand, the rescaled variables prevent concentration and thus allows the spectral scheme to achieve its full accuracy: the Dirac mass is very well captured for large time. This is also due to the correct change of scales in the support of the scaled distribution, thanks to the expression of ω .




FIGURE 3. Comparison of the first marginal $f^{(1)}$ of the solution to (6.1) computed in classical (dashed line) and rescaled variables (solid line), at time $t = 0.3, 1$ and 20 .

6.2. Trend to Equilibrium in the Non Homogeneous Case. We consider now the full inelastic Boltzmann equation in dimension d on the torus

$$\frac{\partial f}{\partial t} + v \cdot \nabla_x f = \frac{\mathcal{Q}_{\mathcal{I}}(f, f)}{\varepsilon}, \quad x \in \mathbb{T}^d, v \in \mathbb{R}^d$$

with periodic boundary conditions in x . We first introduce the hydrodynamical fields associated to a kinetic distribution $f(t, x, v)$. These are the $(d+1)$ fields of density ρ (scalar) and mean velocity \mathbf{u} (vector valued) defined by the formulas (2.4)-(2.5). Whenever $f(t, x, v)$ is a smooth solution to the granular gases equation with periodic boundary conditions, one has the global conservation laws for mass and momentum

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{T}^d \times \mathbb{R}^d} f(t, x, v) dx dv &= 0, \\ \frac{d}{dt} \int_{\mathbb{T}^d \times \mathbb{R}^d} f(t, x, v) v dx dv &= 0. \end{aligned}$$

Therefore, without loss of generality we shall impose

$$\int_{\mathbb{T}^d \times \mathbb{R}^d} f(t, x, v) dx dv = 1, \quad \int_{\mathbb{T}^d \times \mathbb{R}^d} f(t, x, v) v dx dv = 0,$$

where we normalized the measure on the torus: $m(\mathbb{T}^d) = 1$. These conservation laws are then enough to uniquely determine the stationary state of the inelastic Boltzmann equation: the normalized global Dirac distribution in velocity and constant in space

$$(6.6) \quad f_{\infty}(v) = \delta_0(v).$$

Our goal here is to investigate numerically the long time behavior of the solution f . If f is any reasonable solution of the inelastic Boltzmann equation, satisfying certain *a priori* bounds of compactness, then it is expected that f does converge to the global distribution function f_{∞} as t goes to $+\infty$. The point is that on the opposite to the elastic Boltzmann equation, the solution does not satisfy any entropy theorem. Therefore, to measure the convergence to equilibrium we compute the evolution of the quantity

$$\mathcal{L}(t) = \|\rho(t) - 1\|_{L^1}.$$




FIGURE 4. Comparison of the global inhomogeneous cooling process in original and scaled variables ($\varepsilon = 0.1$).

We present numerical results for the full granular gases equation (1.1), using the uncoupled rescaled model (3.8)-(3.11). The initial condition $g_0(x, \xi) = \omega(0, x)^d f_0(x, \omega(0, x) \xi)$ is set as

$$\begin{cases} f_0(x, v) = \frac{\rho_0(x)}{(2\pi)^{d/2}} \exp\left(-\frac{|v|^2}{2}\right), & \forall (x, v) \in \mathbb{T}^d \times \mathbb{R}^d, \\ \omega(0, x) = \sqrt{\frac{2}{d\rho_0(x)}}, \end{cases}$$

with $\rho_0(x) = 1 + 0.1 \cos(\pi x)$ and $\mathbb{T} = [0, L]$.

Description of the results. We start with a simple one dimensional test (in space and velocity), intended to show that the uncoupled model is a good approximation of equation (1.1) in the space inhomogeneous case. We take $L = 1$ and $N_x = 25$ points in space variable, a rescaled velocity box \mathcal{D}_V of size $V = 10$ and $N_v = 32$ half Fourier modes in rescaled variables. We also perform computations in classical velocity variables, with a velocity box of size $V = 8$ and $N_v = 128$ half modes, in order to achieve a good accuracy and avoid oscillations in short time.

We compare the evolution of the global kinetic energy $E(t)$ of both simulations. We also plot the theoretical asymptotic behavior of the kinetic energy (namely, the Haff's Law [26]), given formally by

$$E(t) \simeq \frac{E(0)}{(1 + \sqrt{2}K_1 t)^2}.$$

We can see in Figure 4 that the Haff's Law is very well captured by the rescaling method. This is also the case for short time in the classical method, but due to the fast apparition of spurious oscillations (even with $N_v = 128$ half modes), a strong divergence from the expected global behavior occurs in this case. Once more, the efficiency of the rescaling method for the long time behavior of the solution becomes clear, since the Haff's Law is perfectly described.

Let us investigate more precisely the long time behavior of the numerical solution obtained from the rescaled method. In Figure 5, we present the time evolution in log-log scale of $\mathcal{L}(t)$ which describes the relaxation towards global equilibrium of the local mass $\rho(t, x)$ in L^1 norm.

Numerical investigations on the elastic Boltzmann equation showed some evidences of oscillations during the relaxation process, for example in the local relative entropy [20], as it was conjectured by L. Desvillettes

and C. Villani in [17]. In the inelastic case, the situation is different. On the one hand the solution $f(t)$ does not converge to a smooth steady state since f_∞ is the Dirac mass (6.6), but hopefully our rescaling method allows to follow the concentration in velocity. On the other hand, on the evolution of $\mathcal{L}(t)$ we still observe the apparition of oscillations, but now the oscillation frequency is constant with respect to the logarithmic time scale, which is a big difference with the elastic case (see Figure 5). Let us mention that we can also observe this oscillation phenomenon with pure specular boundary conditions.

We also remark that the scheme breaks down when $\mathcal{L}(t)$ becomes too small (this is particularly clear for the case $\varepsilon = 0.05$ in Figure 5). This can be avoided by refining the space and velocity grids.

Interpretation of the results. Here, we performed simulations on the full inelastic Boltzmann equation in a simplified geometry (one dimension of space and velocity, periodic and specular boundary conditions, fixed Knudsen number) with the spectral method to observe the evolution of the quantity $\mathcal{L}(t)$ and to check numerically if such oscillations occur. It is in fact possible to give a simple interpretation of these oscillations thanks to the work [18]. Since this effect is observed near the global equilibrium, which is homogeneous in space, one can consider that the space inhomogeneous granular gases equation is a small perturbation of the homogeneous equation, which exhibits a self-similar behavior and the Haff's Law. Therefore, we rescale this equation as

$$(6.7) \quad \begin{cases} f(t, x, v) = V(t)^d h(s(t), x, V(t)v), \\ V(t) = 1 + (1 - e)t, \\ s(t) = \frac{\log(1 + (1 - e)t)}{(1 - e)}. \end{cases}$$

Then, one can show after straightforward computations that the distribution h is solution to the granular Boltzmann equation with an anti-drift term in the usual set (see the paper [29] and the references therein) of self similar variables $(s, x, \nu) := (s(t), x, V(t)v)$

$$(6.8) \quad \frac{\partial h}{\partial s} + \nu \cdot \nabla_x h + (1 - e) \nabla_\nu (\nu h) = \frac{\mathcal{Q}_{\mathcal{I}}(h, h)}{\varepsilon}.$$

Due to the convergence toward an equilibrium distribution H_e , we can replace the collision operator by its linearization \mathcal{L}_e around H_e in (6.8). The initial condition we took actually amounts to study the long time behavior of the first spatial Fourier mode of h . This behavior is formally given by the one of $t \mapsto e^{\lambda_j t}$ when $t \rightarrow \infty$ for any $j \in \{0, \dots, d + 1\}$, where λ_j is one of the $d + 2$ hydrodynamic eigenvalues of the linearized operator \mathcal{L}_e .

At least for weak inelasticity, we have seen in Subsection 3.2 that the inelastic collision operator can be understood as a perturbation of the elastic Boltzmann operator. In [36], we used this expansion and results from [30] to perform a spectral analysis of \mathcal{L}_e , as it has already been done for the elastic Boltzmann operator in [18]. This allows to compute the first terms of the Taylor expansion of the hydrodynamic eigenvalues λ_j . We give this expansion with respect to $\eta = 2\pi/L$ and $1 - e$:

$$(6.9) \quad \lambda_j = i\eta \lambda_j^{(1)} + \eta^2 \lambda_j^{(2)} - (1 - e) + \mathcal{O}(\eta^3) + \mathcal{O}((1 - e)^2),$$

where $\lambda_j^{(1)}$ is the first order expansion in η of the elastic hydrodynamic eigenvalues:

$$\lambda_j^{(1)} = \begin{cases} \pm \sqrt{1 + \frac{2}{d}}, & j \in \{0, 1\}, \\ 0, & j \in \{2, \dots, d + 1\}. \end{cases}$$




FIGURE 5. Return to equilibrium of $\mathcal{L}(t) = \|\rho - 1\|_{L^1}$.

Restitution coefficient e	Damping rate β	$-(1-e)\beta$
0.8	-1.05	0.21
0.85	-1.4	0.21
0.9	-2.05	0.205
0.95	-4.1	0.205

TABLE 1. Influence of the restitution coefficient e on the damping rate in the time evolution of the quantity \mathcal{L} .

The second order expansion in η is given by

$$\lambda_j^{(2)} = \begin{cases} -\frac{\kappa}{d+2} - \frac{\mu}{2}, & j \in \{0, 1\}, \\ -\left(1 + \frac{2}{d}\right)\kappa, & j = 2, \\ -\frac{d}{2(d-1)}\mu, & j \in \{3, \dots, d+1\}, \end{cases}$$

with μ the viscosity coefficient and κ the heat conductivity of the Navier-Stokes limit of the elastic Boltzmann equation.

The expansion (6.9) can explain the oscillations and the damping observed in Figure 5. Indeed, the first order term in η is imaginary and will generate oscillations during the return toward equilibrium, of period $\eta\sqrt{1+2/d}$. Then, the period of oscillation ζ will be proportional to L^{-1} , as in [20]. In our case, the time has not been scaled, and then according to the definition (6.7) of $s(t)$, the oscillations will appear in a logarithmic time. The second order term in η and the term in $(1-e)$ are nonpositive and then will exhibit a damping effect of magnitude $\eta^2\alpha + (1-e)\beta$ for α, β positive, according to (6.9). We can see in Table 1 that this effect can be observed numerically: when the restitution coefficient varies, the quantity $(1-e)\beta$ remains constant, when measured in the logarithmic scale.

6.3. Inelastic Shear Flow Problem. The uniform shear flow problem is perhaps one of the most widely studied nonequilibrium problem for granular gases, both experimentally and theoretically [3]. It can be described as follows: consider a gas enclosed between two infinite planes located at $y = -1/2$ and $y = 1/2$, in opposite relative motion with respective velocities $-1/2$ and $1/2$ in the x -direction. The planes are not described as realistic thermalized, reflecting walls (*i.e.* Maxwell boundary conditions) but as Lee-Edwards boundaries: when a particle of velocity v hits one of the planes, it is reemitted through the opposite one with a velocity v' such that the relative velocity of the particle with respect to the plane is preserved before and after this exchange. In consequence, the energy of the system increases during this process.

Thus, the particles gain energy at the boundary while other particles lose energy due to the inelasticity of the collisions. This problem has been solved numerically using Direct Simulation Monte Carlo methods in [3, 37]. We shall consider it using the full deterministic inelastic Boltzmann equation. The gas is taken initially at thermal equilibrium in the two dimensional infinite tube

$$f_0(x, v) = \frac{1}{2\pi} \exp\left(-\frac{|v|^2}{2}\right), \quad \forall x \in [-1/2, 1/2] \times \mathbb{R}, \quad \forall v \in \mathbb{R}^2.$$

The evolution of the system will be induced by the inelastic collisions as well as the flow created by the energy input at the boundaries.

Description of the results. The simulations are made thanks to the rescaled model (3.8), with the granular gases collision operator $\mathcal{Q}_{\mathcal{I}}$, on a simplified geometry: one dimension of space and velocity with $\Omega = [-1/2, 1/2]$. The shear flow problem is mimicked by using diffusive boundary conditions: we impose constant temperature $T_w = 1$ and velocity $u_w = 0$ on the boundary $x = \pm 1/2$. The restitution coefficient of the inelastic collisions is $e = 0.9$. We take $N_x = 100$ and 150 points for the space grid and $N_v = 32$ half Fourier modes on a rescaled velocity box $\mathcal{D}_V = [-V, V]$ of size $V = 12$. We compare these results with computations made on the model (1.1) in classical variables. We take in this case $N_x = 100$ spatial points and $N_v = 32$ half Fourier modes on a velocity box \mathcal{D}_V with $V = 8$.




FIGURE 6. Normalized density $\rho(t, x)/\bar{\rho}$ and temperature $T(t, x)/\bar{T}(t)$ of the shear flow problem computed in classical and rescaled variables, at time $t = 0.25, 0.75$ and 2 .

We present in Figure 6 the evolution of the normalized density $\rho(t, x)/\bar{\rho}$ and temperature $T(t, x)/\bar{T}(t)$, where $\bar{\rho}$ and \bar{T} are the space averaged quantities

$$\bar{T}(t) := \frac{1}{|\Omega|} \int_{\Omega} T(t, x) dx, \quad \bar{\rho} := \frac{1}{|\Omega|} \int_{\Omega} \rho(t, x) dx.$$

On the one hand, we observe that the computation in classical and rescaled variables agree very well for short time. Then, the oscillations due to the concentration in velocity variable for the classical case quickly break the result of this computation. On the other hand, the solution obtained in rescaled variables can be computed for large time, thanks to the change of scales induced by the choice (2.7) of ω . We also notice the stability of the scheme with respect of the space grid.

The short time behavior of this quantity is close to the one presented in [3] using DSMC simulations on the more realistic $2d_x \times 2d_v$ geometry. Indeed, we can see that the thermalized walls heat the system

at the boundary and the collisions cool it inside the domain. This induces very strong gradients near the boundaries for the macroscopic quantities. When time evolves, the temperature will eventually become uniform inside the domain with bigger and bigger gradients near the boundary. This large time behavior is different of the one described in [3] where the temperature reaches a space uniform steady state. This might be due to the use of a simplified geometry. Indeed, in their case, the heating of the boundary induced by the Lee-Edwards conditions will stop because of the uniformization of the gas.

6.4. Cluster Formation in Granular Gases. We consider now the kinetic equation (1.1) with the simplified granular gases operator \mathcal{Q}_S in dimension 2 of space and velocity, on the torus $\mathbb{T}^2 = [0, 1]^2$:

$$\frac{\partial f}{\partial t} + v \cdot \nabla_x f = \frac{\mathcal{Q}_S(f)}{\varepsilon}, \quad \forall (x, v) \in \mathbb{T}^2 \times \mathbb{R}^2$$

with periodic boundary conditions in x . We shall present some numerical evidences of *clustering* for this simplified granular gases model.

The problem of cluster formation in a force-free granular media is a particularly difficult test case of numerical methods for granular gases simulations. It has been shown to occur numerically in both microscopic and macroscopic simulations. More precisely, in the classical Molecular Dynamics (MD) simulations of [24], clusters arise from an initially homogeneous gas of particles. It was also observed using MD simulations in [34] that clustering is a transient phenomenon: an initially homogeneous system of particles interacting by inelastic collisions will tend to form clusters and then will eventually come back after some time to the homogeneous cooling state. Clusters were also observed using deterministic numerical simulations for the macroscopic quantities, thanks to the energy dissipative Navier-Stokes model of [12].

We will show that this phenomenon also occurs at the mesoscopic scale, namely using a kinetic description. The initial condition for the problem is taken as a Maxwellian distribution of homogeneous, isotropic momentum and temperature and uniform density:

$$f_0(x, v) = \frac{U(x)}{(2\pi)^{d/2}} \exp\left(-\frac{|v|^2}{2}\right), \quad \forall (x, v) \in \mathbb{T}^2 \times \mathbb{R}^2$$

with U an uniformly distributed random variable over $[0, 1]^2$.

Description of the results. The simulations are made using the simplified granular gases operator \mathcal{Q}_S presented in Subsection 6.4 in the rescaled model (2.2)-(2.6). The macroscopic scaling function ω , given by (2.7), is computed thanks to the energy dissipative Euler system (3.11). Due to the correct energy dissipation, this model mimics the behavior of the kinetic granular gases equation and is then consistent with the MD simulations of [24]. Moreover, the ES-BGK operator used in the definition of \mathcal{Q}_S gives the good Prandtl number for the Navier-Stokes limit of the model, which is also consistent with the macroscopic model of [12].

We take $N_x = 50$ points in each direction of space on the torus and $N_v = 50$ points in each direction of a velocity box of size $V = 7$. The restitution coefficient is $e = 0.99$ (and then $\varepsilon = 0.01$ thanks to the weak inelasticity assumption (3.10)). We show in Figure 7 the density contour plot given by the kinetic model and by the dissipative Euler system (3.11) used to compute the scaling function. We see that both models give the same results for short time, which is consistent with the fact that the hydrodynamic limit of the kinetic equation (1.1) with the Fokker-Planck operator (3.12) is given at the first order in ε by Euler system (2.6). Then, for larger time, clusters start to appear even with a coarse grid in space. It is interesting to notice that the clusters remains bounded through time, even without taking into account the dense gas effects through a ‘‘pair correlation function’’ depending of a maximal packing fraction, as in [12].

ACKNOWLEDGMENT

The authors would like to thank ICERM at Brown University for the kind hospitality during the 2011 Fall Semester Program on ‘‘Kinetic Theory and Computation’’ where part of this work has been carried on. TR also like to thanks Lorenzo Pareschi for interesting discussions about the BGK-like simplified




FIGURE 7. Contour plot of the density given by the simplified granular gases model (top) and the dissipative Euler system (bottom), at time $t = 0, 0.25, 0.5, 1, 5$ and 10 . High values are displayed in red, low in blue.

granular gases model. This work is partially supported by the European Research Council ERC Starting Grant 2009, project 239983-NuSiKiMo.

REFERENCES

- [1] ANDRIES, P., LE TALLEC, P., PERLAT, J.-P., AND PERTHAME, B. The Gaussian-BGK model of Boltzmann equation with small Prandtl number. *European Journal of Mechanics - B/Fluids* 19, 6 (Nov. 2000), 813–830.
- [2] ASTILLERO, A., AND SANTOS, A. A granular fluid modeled as a driven system of elastic hard spheres. In *The Physics of Complex Systems. New Advances and Perspectives*, F. Mallamace and H. E. Stanley, Eds., no. 2. IOS Press, Amsterdam, Sept. 2004, pp. 475–480.
- [3] ASTILLERO, A., AND SANTOS, A. Uniform shear flow in dissipative gases: Computer simulations of inelastic hard spheres and frictional elastic hard spheres. *Physical Review E* 72, 3 (Sept. 2005), 1–23.
- [4] BESSE, C., BORGHOL, S., GOUDON, T., LACROIX-VIOLET, I., AND DUDON, J.-P. Hydrodynamic Regimes, Knudsen Layer, Numerical Schemes: Definition of Boundary Fluxes. *Adv. Appl. Math. Mech.* 3, 5 (2011), 519–561.
- [5] BIRD, G. *Molecular Gas Dynamics and the Direct Simulation of Gas Flows*, 2nd ed. Oxford University Press, 1994.
- [6] BISI, M., CARRILLO, J., AND SPIGA, G. Some alternative methods for hydrodynamic closures to dissipative kinetic models. *Eur. Phys. J. Special Topics* 179, 1 (May 2010), 165–178.
- [7] BOBYLEV, A., AND RJASANOW, S. Difference scheme for the Boltzmann equation based on the fast Fourier transform. *Eur. J. Mech. B Fluids* 16, 2 (1997), 293–306.
- [8] BOBYLEV, A. V., AND RJASANOW, S. Fast deterministic method of solving the Boltzmann equation for hard spheres. *Eur. J. Mech. B Fluids* 18, 5 (1999), 869–887.
- [9] BOBYLEV, A. V., AND RJASANOW, S. Numerical solution of the boltzmann equation using a fully conservative difference scheme based on the fast fourier transform. *Transport Theory Statist. Phys.* 29, 3-5 (2000), 289–310.
- [10] CANUTO, C., HUSSAINI, M., QUARTERONI, A., AND ZANG, T. *Spectral methods in fluid dynamics*. Springer Series in Computational Physics. Springer-Verlag, New York, 1988.
- [11] CARRILLO, J., FRANCESCO, M. D., AND TOSCANI, G. Intermediate Asymptotics Beyond Homogeneity and Self-Similarity: Long Time Behavior for $u_t = \Delta\phi(u)$. *Arch. Ration. Mech. Anal.* 180, 1 (Jan. 2006), 127–149.
- [12] CARRILLO, J., POESCHEL, T., AND SALUENA, C. Granular hydrodynamics and pattern formation in vertically oscillated granular disk layers. *J. Fluid Mech.* 597 (2008), 119–144.
- [13] CARRILLO, J., AND TOSCANI, G. Asymptotic L^1 -decay of solutions of the porous medium equation to self-similarity. *Indiana Univ. Math. J.* 49, 1 (2000), 113–142.
- [14] CARRILLO, J. A., AND VÁZQUEZ, J. Asymptotic Complexity in Filtration Equations. *J. Evol. Equ.* 7, 3 (Feb. 2007), 471–495.
- [15] CERCIGNANI, C. *The Boltzmann Equation and its Applications*. Springer, 1988.
- [16] CERCIGNANI, C., ILLNER, R., AND PULVIRENTI, M. *The mathematical theory of dilute gases*, vol. 106 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1994.
- [17] DESVILLETTES, L., AND VILLANI, C. On the Trend to Global Equilibrium for Spatially Inhomogeneous Kinetic Systems: the Boltzmann Equation. *Invent. Math.* 159, 2 (2005), 245–316.
- [18] ELLIS, R., AND PINSKY, M. The First and Second Fluid Approximations to the Linearized Boltzmann Equation. *J. Math. Pures Appl.* 54, 9 (1975), 125–156.
- [19] FILBET, F., AND MOUHOT, C. Analysis of Spectral Methods for the Homogeneous Boltzmann Equation. *Trans. Amer. Math. Soc.* 363 (2011), 1947–1980.
- [20] FILBET, F., MOUHOT, C., AND PARESCHI, L. Solving the Boltzmann Equation in $N \log^2 N$. *SIAM J. Sci. Comput.* 28, 3 (2007), 1029–1053.
- [21] FILBET, F., PARESCHI, L., AND TOSCANI, G. Accurate Numerical Methods for the Collisional Motion of (Heated) Granular Flows. *J. Comput. Phys.* 202, 1 (2005), 216–235.
- [22] FILBET, F., AND RUSSO, G. A Rescaling Velocity Method for Kinetic Equations: the Homogeneous Case. In *Proceedings Modelling and Numerics of Kinetic Dissipative Systems* (Lipari, 2004), Nova-Science, p. 11.
- [23] GAMBA, I. M., AND THARKABHUSHANAM, S. H. Spectral-Lagrangian methods for collisional models of non-equilibrium statistical states. *J. Comput. Phys.* 228, 6 (Apr. 2009), 2012–2036.
- [24] GOLDBIRSHCH, I., AND ZANETTI, G. Clustering instability in dissipative gases. *Physical Review Letters* 70, 11 (Mar. 1993), 1619–1622.
- [25] GOLSE, F. The Boltzmann equation and its hydrodynamic limits. In *Handbook of Differential Equations: Evolutionary Equations Vol. 2*, C. Dafermos and E. Feireisl, Eds. North-Holland, 2005, pp. 159–303.
- [26] HAFF, P. Grain flow as a fluid-mechanical phenomenon. *J. Fluid Mech.* 134 (1983), 401–30.
- [27] HAILE, J. *Molecular dynamics simulation: elementary methods*. Wiley professional paperback series. Wiley, 1992.
- [28] MAXWELL, J. C. On the dynamical theory of gases. *Philos. Trans. Roy. Soc. London* 157 (1867), pp. 49–88.
- [29] MISCHLER, S., AND MOUHOT, C. Cooling process for inelastic Boltzmann equations for hard spheres, Part II: Self-similar solutions and tail behavior. *J. Stat. Phys.* 124, 2 (2006), 703–746.
- [30] MISCHLER, S., AND MOUHOT, C. Stability, convergence to self-similarity and elastic limit for the Boltzmann equation for inelastic hard spheres. *Commun. Math. Phys.* 288, 2 (2009), 431–502.

- [31] NALDI, G., PARESCHI, L., AND TOSCANI, G. Spectral methods for one-dimensional kinetic models of granular flows and numerical quasi elastic limit. *ESAIM Math. Model. Numer. Anal.* 37, 1 (Mar. 2003), 73–90.
- [32] PARESCHI, L., AND PERTHAME, B. A Fourier Spectral Method for Homogeneous Boltzmann Equations. *Transport Theory Statist. Phys.* 25, 3 (1996), 369–382.
- [33] PARESCHI, L., AND RUSSO, G. Numerical Solution of the Boltzmann Equation I : Spectrally Accurate Approximation of the Collision Operator. *SIAM J. Numer. Anal.* 37, 4 (2000), 1217–1245.
- [34] PÖSCHEL, T., BRILLIANTOV, N. V., AND SCHWAGER, T. Transient clusters in granular gases. *Journal of Physics: Condensed Matter* 17, 24 (June 2005), S2705–S2713.
- [35] RERICHA, E. C., BIZON, C., SHATTUCK, M. D., AND SWINNEY, H. L. Shocks in supersonic sand. *Phys. Rev. Lett.* 88 (Dec 2001), 014302.
- [36] REY, T. First and second order fluid approximations to the granular gases equation. In preparation.
- [37] RUIZ-MONTERO, M., AND BREY, J. J. On the development of inhomogeneities in freely evolving granular gases: The shear state and beyond. *The European Physical Journal Special Topics* 179, 1 (May 2010), 249–262.
- [38] SHU, C.-W. High order ENO and WENO schemes for computational fluid dynamics. In *High-order methods for computational physics*, vol. 9 of *Lect. Notes Comput. Sci. Eng.* Springer, Berlin, 1999, pp. 439–582.
- [39] SONE, Y. *Molecular Gas Dynamics: Theory, Techniques, And Applications*. Modeling And Simulation in Science, Engineering And Technology. Birkhäuser, 2007.
- [40] TOSCANI, G. Kinetic and Hydrodynamic Models of Nearly Elastic Granular Flows. *Monatsh. Math.* 142, 1 (2004), 179–192.
- [41] TOSCANI, G. Hydrodynamics from the dissipative Boltzmann equation. In *Mathematical models of granular matter*, vol. 1937 of *Lecture Notes in Math.* Springer, Berlin, 2008, pp. 59–75.
- [42] VAN LEER, B. Towards the ultimate conservative difference scheme III. Upstream-centered finite-difference schemes for ideal compressible flow. *Journal of Computational Physics* 23, 3 (Mar. 1977), 263–275.
- [43] VILLANI, C. *A review of mathematical topics in collisional kinetic theory*. Elsevier Science, 2002.

FRANCIS FILBET, UNIVERSITÉ DE LYON, CNRS UMR 5208, UNIVERSITÉ LYON 1, INSTITUT CAMILLE JORDAN, 43 BLVD. DU 11 NOVEMBRE 1918, F-69622 VILLEURBANNE CEDEX, FRANCE.

E-mail address: `filbet@math.univ-lyon1.fr`

THOMAS REY, UNIVERSITÉ DE LYON, CNRS UMR 5208, UNIVERSITÉ LYON 1, INSTITUT CAMILLE JORDAN, 43 BLVD. DU 11 NOVEMBRE 1918, F-69622 VILLEURBANNE CEDEX, FRANCE.

E-mail address: `rey@math.univ-lyon1.fr`