



**HAL**  
open science

## Parameter sensitivity of CIR process

Sidi Mohamed Ould Aly

► **To cite this version:**

Sidi Mohamed Ould Aly. Parameter sensitivity of CIR process. *Electronic Communications in Probability*, 2013, 18 (34), pp.1-6. 10.1214/ECP.v18-2035 . hal-00690856v2

**HAL Id: hal-00690856**

**<https://hal.science/hal-00690856v2>**

Submitted on 27 May 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Parameter sensitivity of CIR process

**S. M. OULD ALY**

Université Paris-Est, Laboratoire d'Analyse et de Mathématiques Appliquées

5, boulevard Descartes, 77454 Marne-la-Vallée cedex 2, France

email: sidi-mohamed.ouldali@univ-paris-est.fr

May 27, 2013

## **Abstract**

We study the differentiability of the CIR process with respect to its parameters. We give a stochastic representation for these derivatives in terms of the paths of  $V$ .

## **1 Introduction**

The CIR process is defined as the unique solution of the following stochastic differential equation:

$$dV_t = (a - bV_t)dt + \sigma\sqrt{V_t}dW_t, \quad V_0 = v, \quad (1.1)$$

where  $a, \sigma, v \geq 0$  and  $b \in \mathbb{R}$  (see [8] for the existence and uniqueness of the solution of the SDE). This process is widely used in finance to model short term interest rate (see [3]) but also used to model stochastic volatility in the Heston stochastic volatility model. The option prices in these models depend in the values of the parameters of CIR process. On the other hand, these parameters are often calibrated to market prices of derivatives, so they tend to change their values regularly. The knowledge of the derivatives of the CIR process with respect to its parameters is therefore crucial for the study the sensitivities of prices in these models.

The most common approach to study the sensitivity of stochastic differential equation with respect to its parameters is to use the Malliavin calculus, especially for the sensitivity

with respect to the initial value. The Malliavin derivative gives a stochastic representation of the sensitivity of process with respect to its initial value. We note that the coefficients of (1.1) are neither differentiable in 0 nor globally Lipschitz, so the standard results (see e.g [9],[5]) cannot be used here. Nevertheless, for the special case of CIR process, Alòs and Ewald ([1]) show the existence of Malliavin derivative of the CIR process under assumption ( $2a > \sigma^2$ ). In mathematical finance, the sensitivities of option prices with respect to not only the initial point, but also other parameters, need to be studied very carefully.

In this article, we study the differentiability of the solution of (1.1) with respect to the parameters  $a$ ,  $b$  and  $\sigma$  in  $\mathcal{L}_p$  sense (see next section). We show that, under some assumptions, this process is differentiable with respect to these parameters and give a stochastic representation of its derivatives.

## 2 Differentiability

For technical reasons, we will rather consider the square root of  $V^v$ , denoted  $X^v$ . Throughout this paper, we assume that

$$2a \geq \sigma^2 \tag{2.1}$$

Under this assumption, we have for any  $T, v > 0$ ,  $\mathbb{P}(\forall t \in [0, T] : V_t^v > 0) = 1$ . The process  $X^v$  is the unique solution of the following stochastic differential equation

$$dX_t^v = \left( \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t^v} - \frac{b}{2} X_t^v \right) dt + \frac{\sigma}{2} dW_t, \quad X_0^v = \sqrt{v}. \tag{2.2}$$

We start by studying the differentiability of  $X$  with respect to the parameter  $a$ . We consider here the  $\mathcal{L}_p$ -differentiability of the function  $a \mapsto X^v(a)$ , i.e the existence of a process  $\dot{X}_a$  so that

$$\lim_{\epsilon \rightarrow 0} \left\| \sup_{s \leq t} \left| \frac{X_s^v(a + \epsilon) - X_s^v(a)}{\epsilon} - \dot{X}_a(s) \right| \right\|_p = 0 \tag{2.3}$$

We have the following result

**Proposition 2.1.** *Let  $b \in \mathbb{R}$  and  $\sigma, x \geq 0$ . For every  $a \in ]\sigma^2, +\infty[$ , let  $X_a$  be the unique*

solution of the SDE :

$$dX_t = \left( \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t} - \frac{b}{2} X_t \right) dt + \frac{\sigma}{2} dW_t, \quad X(0) = x$$

and let  $a_0 > \sigma^2$ . Then the function  $a \mapsto X_a$  is  $\mathcal{L}_p$ -differentiable at  $a_0$ , for any  $1 \leq p \leq \frac{2a_0}{\sigma^2} - 1$  and its derivative ( $\dot{X}_a$ ) is given by

$$\dot{X}_a(t) = \int_0^t \frac{1}{2X_s} \exp \left( -\frac{b}{2}(t-u) - \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \int_s^t \frac{du}{X_u^2} \right) ds. \quad (2.4)$$

**Proof:** Let  $X^\epsilon$  be the unique solution of the stochastic differential equation

$$dX_t^\epsilon = \left( \left( \frac{a+\epsilon}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t^\epsilon} - \frac{b}{2} X_t^\epsilon \right) dt + \frac{\sigma}{2} dW_t, \quad X_0^\epsilon = \sqrt{v}.$$

For  $\epsilon > 0$ , define  $R_0^\epsilon(t) := X_t^\epsilon - X_t$ . We can easily see that  $R_0^\epsilon$  is given by

$$R_0^\epsilon(t) = \epsilon U_t^\epsilon \int_0^t (U_s^\epsilon)^{-1} \frac{1}{2X_s} ds,$$

where

$$U^\epsilon = \exp \left( - \int_0^t \alpha_s^\epsilon ds \right), \quad \text{with } \alpha_t^\epsilon = \left( \frac{a+\epsilon}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_s^\epsilon X_s} + \frac{b}{2}.$$

We have, using the fact that for any  $s \leq t$ ,  $e^{-\int_s^t \alpha_u^\epsilon du} \leq e^{-bt/2} \vee 1$  a.s.,

$$\frac{|R_0^\epsilon(t)|}{\epsilon} \leq \frac{t(e^{-bt/2} \vee 1)}{2} \sup_{s \leq t} \frac{1}{X_s^v}.$$

On the other hand, we have, using Lemma 2.3.2 of [4],

$$\forall p < 2 \left( \frac{2a}{\sigma^2} - 1 \right), \quad \mathbb{E} \left[ \sup_{s \leq t} \frac{1}{X_s^p} \right] < +\infty. \quad (2.5)$$

In particular, we have for any  $p \in [1, 2 \left( \frac{2a}{\sigma^2} - 1 \right) [$ ,

$$\|R_0^\epsilon\|_p \leq C\epsilon.$$

Let's now set

$$\dot{X}_a(t) := \lim_{\epsilon \rightarrow 0} \frac{R_0^\epsilon}{\epsilon}(t) = U_t^0 \int_0^t (U_s^0)^{-1} \frac{1}{2X_s} ds.$$

We have  $\|\dot{X}_a\|_p \leq C$ . Furthermore,  $\dot{X}_a$  is solution of the stochastic differential equation:

$$d\dot{X}_a(t) = - \left( \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t^2} + \frac{b}{2} \right) \dot{X}_a(t) dt + \frac{1}{2X_t} dt.$$

Let  $R_1^\epsilon(t) = X_t^\epsilon - X_t - \epsilon \dot{X}_a(t)$ . The process  $R_1^\epsilon$  is a solution of the stochastic differential equation

$$dR_1^\epsilon(t) = \left( -\alpha_t^\epsilon R_1^\epsilon(t) - \epsilon \dot{X}_a(t) \left( \alpha_t^\epsilon - \left[ \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t^2} + \frac{b}{2} \right] \right) \right) dt.$$

On the other hand, we have

$$\alpha_t^\epsilon - \left( \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t^2} + \frac{b}{2} \right) = - \left( \frac{\alpha_t^\epsilon}{X_t} - \frac{b}{2X_t} \right) R_0^\epsilon(t) + \frac{\epsilon}{2X_t^2}.$$

It follows that  $R_1^\epsilon$  can be written as

$$R_1^\epsilon(t) = U_t^\epsilon \int_0^t (U_s^\epsilon)^{-1} \left( \dot{X}_a(t) \left( \epsilon \left( \frac{\alpha_t^\epsilon}{X_t} - \frac{b}{2X_t} \right) R_0^\epsilon(t) - \frac{\epsilon^2}{2X_t^2} \right) \right) ds,$$

Using (2.5) and the fact that for any  $s \leq t$ , we have  $e^{-\int_s^t \alpha_u^\epsilon du} \leq 1 \vee e^{-bt/2}$  and  $\int_0^t \alpha_s^\epsilon e^{-\int_s^t \alpha_u^\epsilon du} ds = 1 - e^{-\int_0^t \alpha_u^\epsilon du}$ , we get

$$\forall 1 \leq p < \frac{2a}{\sigma^2} - 1, \quad \|R_1^\epsilon\|_p \leq C\epsilon^2. \quad \square$$

The differentiability with respect to  $b$  is obtained in the same. The proof of the next Proposition is almost identical to Proposition 2.1.

**Proposition 2.2.** *Let  $x, a, \sigma \geq 0$  so that  $4a > 3\sigma^2$ . For every  $b \in \mathbb{R}$ , let  $X_b$  be the unique solution of the SDE :  $dX_t = \left( \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t} - \frac{b}{2} X_t \right) dt + \frac{\sigma}{2} dW_t$ ,  $X_0 = x$  and let  $b_0 \in \mathbb{R}$ . The function  $b \mapsto X_b$  is  $\mathcal{L}_p$ -differentiable at  $b_0$ , for any  $1 \leq p < 2(\frac{2a}{\sigma^2} - 1)$  and its derivative  $\dot{X}_b$  is given by*

$$\dot{X}_b(t) = - \int_0^t \frac{X_s}{2} \exp \left( -\frac{b}{2}(t-u) - \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \int_s^t \frac{du}{X_u^2} \right) ds \quad (2.6)$$

We now consider the differentiability of  $X$  with respect to the parameter  $\sigma$ . We propose an extension of the result of Benhamou et al (cf. [2]) who show that  $\sigma \mapsto X$  is  $\mathcal{C}^2$  in neighborhood of 0. We will show that this function is  $\mathcal{C}^1$  in  $[0, \sqrt{a}[$  and  $C^\infty$  around 0.

**Proposition 2.3.** *For any  $\sigma \in [0, \sqrt{a}[$ , the function  $\sigma \mapsto X$  is  $\mathcal{C}^1$  at  $\sigma$  in  $\mathcal{L}_p$ -sense, for every  $p \in [1, \frac{2a}{\sigma^2} - 1[$  and its derivative is the unique solution of the SDE :*

$$d\dot{X}_\sigma(t) = \left( -\frac{\sigma}{4X_t} - \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{\dot{X}_\sigma(t)}{X_t} - \frac{b}{2} \dot{X}_\sigma(t) \right) dt + \frac{1}{2} dW_t. \quad (2.7)$$

**Proof:** Let  $X^\epsilon$  be the unique solution of the SDE :

$$dX_t^\epsilon = \left( \left( \frac{a}{2} - \frac{(\sigma + \epsilon)^2}{8} \right) \frac{1}{X_t^\epsilon} - \frac{b}{2} X_t^\epsilon \right) dt + \frac{\sigma + \epsilon}{2} dW_t, \quad X_0^\epsilon = \sqrt{v}.$$

Let set  $R_0^\epsilon(t) = X_t^\epsilon - X_t$ . In particular,  $R_0^\epsilon$  solves the stochastic differential equation:

$$\begin{aligned} dR_0^\epsilon(t) &= \left( \left( \frac{a}{2} - \frac{(\sigma + \epsilon)^2}{8} \right) \frac{1}{X_t^\epsilon} - \frac{b}{2} X_t^\epsilon - \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t} + \frac{b}{2} X_t \right) dt + \frac{\epsilon}{2} dW_t \\ &= \left( - \left[ \left( \frac{a}{2} - \frac{(\sigma + \epsilon)^2}{8} \right) \frac{1}{X_s^\epsilon X_s} + \frac{b}{2} \right] R_0^\epsilon(t) - \frac{2\epsilon\sigma + \epsilon^2}{8X_t} \right) dt + \frac{\epsilon}{2} dW_t. \end{aligned}$$

It follows that  $R_0^\epsilon$  can be written as

$$R_0^\epsilon(t) = U_t^\epsilon \int_0^t (U_s^\epsilon)^{-1} \left( -\frac{2\epsilon\sigma + \epsilon^2}{8X_s} ds + \frac{\epsilon}{2} dW_s \right),$$

where  $U^\epsilon$  is given by

$$U_t^\epsilon = \exp \left( - \int_0^t \alpha_s^\epsilon ds \right) \quad (2.8)$$

and

$$\alpha_s^\epsilon = \left( \frac{a}{2} - \frac{(\sigma + \epsilon)^2}{8} \right) \frac{1}{X_s^\epsilon X_s} + \frac{b}{2}. \quad (2.9)$$

Applying the Itô formula to the product  $(U_t^\epsilon)^{-1} W_t$ , we have

$$R_0^\epsilon(t) = -\frac{2\epsilon\sigma + \epsilon^2}{8} U_t^\epsilon \int_0^t (U_s^\epsilon)^{-1} \frac{ds}{X_s} + \frac{\epsilon}{2} W_t + U_t^\epsilon \int_0^t W_s d(U^\epsilon)_s^{-1}.$$

On the other hand, using the fact that  $\alpha_t^\epsilon \geq b/2$ , *a.s.*, we know that for any  $s \leq t$ , we have  $0 \leq U_t^\epsilon (U_s^\epsilon)^{-1} \leq 1 \vee e^{-bt/2}$ , *a.s.* It follows that

$$\begin{aligned} |R_0^\epsilon(t)| &\leq c(t) \int_0^t \frac{ds}{X_s} + \frac{\epsilon}{2} \left( \sup_{s \leq t} W_s + \sup_{s \leq t} W_s (1 - U_t^\epsilon) U_t^\epsilon \right) \\ &\leq c(t) \sup_{s \leq t} \frac{1}{X_s} + \epsilon \sup_{s \leq t} W_s (1 + U_t^\epsilon) U_t^\epsilon. \end{aligned}$$

Using (2.5), we have, for any  $1 \leq p < 2 \left( \frac{2a}{\sigma^2} - 1 \right)$ ,

$$\|R_0^\epsilon\|_p \leq C\epsilon. \quad (2.10)$$

Let's now set

$$\dot{X}_\sigma(t) := U_t^0 \int_0^t (U_s^0)^{-1} \left( -\frac{\sigma}{4X_s} ds + \frac{1}{2} dW_s \right).$$

We have  $\|\dot{X}_\sigma\|_p \leq C$ . Furthermore, we can easily see that  $\dot{X}_\sigma$  is solution to the stochastic differential equation:

$$d\dot{X}_\sigma(t) = - \left( \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t^2} + \frac{b}{2} \right) \dot{X}_\sigma(t) dt - \frac{\sigma}{4X_t} dt + \frac{1}{2} dW_t.$$

Set  $R_1^\epsilon(t) = X_t^\epsilon - X_t - \epsilon \dot{X}_\sigma(t)$ . The process  $R_1^\epsilon$  solves the stochastic differential equation:

$$dR_1^\epsilon(t) = \left( -\alpha_t^\epsilon R_1^\epsilon(t) - \epsilon \dot{X}_\sigma(t) \left( \alpha_t^\epsilon - \left[ \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t^2} + \frac{b}{2} \right] \right) - \frac{\epsilon^2}{8X_t} \right) dt.$$

On the other hand, we can easily see that

$$\alpha_t^\epsilon - \left( \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{1}{X_t^2} + \frac{b}{2} \right) = - \left( \frac{\alpha_t^\epsilon}{X_t} - \frac{b}{2X_t} \right) R_0^\epsilon(t) - \frac{2\epsilon\sigma + \epsilon^2}{8X_t^2}.$$

It follows that  $R_1^\epsilon$  can be written as

$$R_1^\epsilon(t) = U_t^\epsilon \int_0^t (U_s^\epsilon)^{-1} \left( -\frac{\epsilon^2}{8X_s} ds + \epsilon \dot{X}_\sigma(s) \left( \left( \frac{\alpha_s^\epsilon}{X_s} - \frac{b}{2X_s} \right) R_0^\epsilon(s) + \frac{2\epsilon\sigma + \epsilon^2}{8X_s^2} \right) \right) ds.$$

We have

$$\begin{aligned}
|R_1^\epsilon(t)| &\leq \int_0^t U_t^\epsilon(U_s^\epsilon)^{-1} \left( \frac{\epsilon^2}{8X_s} ds + \epsilon |\dot{X}_\sigma(s)| \left( \left( \frac{\alpha_t^\epsilon}{X_s} + \frac{b}{2X_s} \right) |R_0^\epsilon(s)| + \frac{2\epsilon\sigma + \epsilon^2}{8X_s^2} \right) \right) ds \\
&\leq \int_0^t U_t^\epsilon(U_s^\epsilon)^{-1} \left( \frac{\epsilon^2}{8X_s} ds + \epsilon |\dot{X}_\sigma(t)| \left( \frac{b}{2X_s} |R_0^\epsilon(s)| + \frac{2\epsilon\sigma + \epsilon^2}{8X_s^2} \right) \right) ds + \\
&\quad \epsilon \int_0^t U_t^\epsilon(U_s^\epsilon)^{-1} \frac{\alpha_t^\epsilon}{X_s} |\dot{X}_\sigma(s)| |R_0^\epsilon(s)| ds \\
&\leq c(t) \int_0^t \left( \frac{\epsilon^2}{8X_s} ds + \epsilon |\dot{X}_\sigma(t)| \left( \frac{b}{2X_s} |R_0^\epsilon(s)| + \frac{2\epsilon\sigma + \epsilon^2}{8X_s^2} \right) \right) ds \\
&\quad + \epsilon c_2(t) \sup_{s \leq t} \left( \frac{|\dot{X}_\sigma(s)| |R_0^\epsilon(s)|}{X_s} \right).
\end{aligned}$$

Finally, using (2.5), we have, for any  $1 \leq p < (\frac{2a}{\sigma^2} - 1)$ ,

$$\|R_1^\epsilon\|_p \leq C\epsilon^2. \quad \square$$

**Proposition 2.4.** *Under the assumptions of Propositions 2.3, 2.1, 2.2, the solution of the SDE (1.1) is differentiable with respect to the parameters  $a$ ,  $b$  and  $\sigma$ . Its derivatives, denoted by  $\dot{V}_a$ ,  $\dot{V}_b$  and  $\dot{V}_\sigma$  respectively, are given as*

$$\begin{aligned}
\dot{V}_a(t) &= \sqrt{V}_t \int_0^t \frac{1}{\sqrt{V}_s} \exp \left( -\frac{b}{2}(t-u) - \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \int_s^t \frac{du}{V_u} \right) ds, \\
\dot{V}_b(t) &= -\sqrt{V}_t \int_0^t \sqrt{V}_s \exp \left( -\frac{b}{2}(t-u) - \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \int_s^t \frac{du}{V_u} \right) ds, \\
\dot{V}_\sigma(t) &= \frac{2}{\sigma} V_t - \frac{2}{\sigma} \sqrt{V}_t \left( \sqrt{v} e^{-\frac{b}{2}t - (\frac{a}{2} - \frac{\sigma^2}{8}) \int_0^t \frac{dr}{V_r}} + a \int_0^t \frac{e^{-\frac{b}{2}(t-u) - (\frac{a}{2} - \frac{\sigma^2}{8}) \int_u^t \frac{dr}{V_r}}}{\sqrt{V}_u} du \right) \quad (2.11)
\end{aligned}$$

**Proof:** As  $V_t = X_t^2$ ,  $V$  is differentiable with respect to the parameters  $a$ ,  $b$  and  $\sigma$  under the assumptions of Propositions 2.3, 2.1, 2.2. The derivatives  $\dot{V}_\sigma$  is given as solution of the SDE :

$$d\dot{V}_\sigma(t) = -b\dot{V}_\sigma(t)dt + \sqrt{V}_t dW_t^2 + \sigma \frac{\dot{V}_\sigma(t)}{2\sqrt{V}_t} dW_t, \quad \dot{V}_\sigma(0) = 0.$$



One can see that the process  $Z_t := \dot{V}_\sigma(t) - \frac{2}{\sigma}V_t$  is solution of the SDE :

$$dZ_t = \left( -\frac{2a}{\sigma} - bZ_t \right) dt + \sigma \frac{Z_t}{2\sqrt{V_t}} dW_t^2, \quad Z_0 = -\frac{2}{\sigma}x.$$

On the other hand, applying Itô formula to the process  $ZV^\alpha$ , for  $\alpha \in \mathbb{R}^*$ , we have

$$d(ZV^\alpha)(t) = \left( -\frac{2a}{\sigma}V_t^\alpha - b(1+\alpha)Z_tV_t^\alpha + (\alpha a + \frac{\alpha^2}{2}\sigma^2)Z_tV_t^{\alpha-1} \right) dt + (\alpha + \frac{1}{2})Z_tV_t^{\alpha-\frac{1}{2}}dW_t^2.$$

It follows that, for  $\alpha = -\frac{1}{2}$ , the process  $Y = ZV^{-\frac{1}{2}}$ ,  $Y$  has finite variation and is given as solution of

$$dY_t = \left( -\frac{2a}{\sigma}V_t^{-\frac{1}{2}} - \frac{b}{2}Y_t - \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \frac{Y_t}{V_t} \right) dt, \quad Y_0 = -\frac{2}{\eta}\sqrt{v}.$$

We can easily solve this equation, we get

$$Y_t := \frac{V_\sigma(t) - \frac{2}{\sigma}V_t}{\sqrt{V_t}} = -\frac{2}{\sigma}\sqrt{v}e^{-\gamma t} - \frac{2a}{\sigma} \int_0^t \frac{e^{-(\gamma t - \gamma u)}}{\sqrt{V_u}} du, \quad a.s.,$$

where

$$\gamma_t := \frac{b}{2}t + \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \int_0^t \frac{dr}{V_r}. \quad (2.12)$$

Thus

$$\dot{V}_\sigma(t) = \frac{2}{\sigma}V_t - \frac{2}{\sigma}\sqrt{V_t} \left( \sqrt{v}e^{-\frac{b}{2}t - \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \int_0^t \frac{dr}{V_r}} + a \int_0^t \frac{e^{-\frac{b}{2}(t-u) - \left( \frac{a}{2} - \frac{\sigma^2}{8} \right) \int_u^t \frac{dr}{V_r}}}{\sqrt{V_u}} du \right), \quad a.s. \quad \square$$

## References

- [1] Alòs, E. and Ewald, C.-O.: Malliavin differentiability of the Heston volatility and applications to option pricing. *Advances in Applied Probability*, 40 (1). pp. 144-162, 2008.
- [2] Benhamou, E., Gobet, E. and Miri, M.: Times dependent Heston model, *SIAM Journal on Financial Mathematics*, Vol.1, pp.289-325, 2010
- [3] Cox, J, Ingersoll, J, and Ross, J. A.: Theory of the Term structure of Interest Rates. *Econometrica*, 53:385–407, 1985

- [4] De Marco, S.: On Probability Distributions of Diffusions and Financial Models with non-globally smooth coefficients, PhD dissertation, Université Paris Est et Scuola Normale Superiore di Pisa, 2010.
- [5] Detemple, J., Garcia, R. and Rindisbacher, M.: Representation formulas for Malliavin derivatives of diffusion processes, *Finance and Stochastics* 9 (3), 2005.
- [6] Dufresne, D.: The integrated square-root process. Research Paper no. 90, Centre for Actuarial Studies, University of Melbourne, 2001.
- [7] Heston, S.: A closed-form solution for options with stochastic volatility with applications to bond and currency options, *Review of Financial Studies* (1993)
- [8] Ikeda, N. and Watanabe, S.: Stochastic Differential Equations and Diffusion Processes. *North-Holland/Kodansha*, 1989.
- [9] Nualart, D.: The Malliavin Calculus and Related Topics, Probability and Its Applications *Springer*, New York 1985.