



**HAL**  
open science

## Tail behaviour of -TARCH models

Péter Elek, László Márkus

► **To cite this version:**

Péter Elek, László Márkus. Tail behaviour of -TARCH models. *Statistics and Probability Letters*, 2010, 10.1016/j.spl.2010.07.020 . hal-00691789

**HAL Id: hal-00691789**

**<https://hal.science/hal-00691789>**

Submitted on 27 Apr 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Accepted Manuscript


Tail behaviour of  $\beta$ -TARCH models

Péter Elek, László Márkus

PII: S0167-7152(10)00214-2  
DOI: 10.1016/j.spl.2010.07.020  
Reference: STAPRO 5758

To appear in: *Statistics and Probability Letters*

Received date: 8 August 2009  
Revised date: 28 July 2010  
Accepted date: 28 July 2010



Please cite this article as: Elek, P., Márkus, L., Tail behaviour of  $\beta$ -TARCH models. *Statistics and Probability Letters* (2010), doi:10.1016/j.spl.2010.07.020

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Tail behaviour of $\beta$ -TARCH models

Péter Elek<sup>a,1,\*</sup>, László Márkus<sup>a</sup>

<sup>a</sup>Department of Probability Theory and Statistics, Eötvös Loránd University, Hungary  
H-1117 Budapest, Pázmány Péter sétány 1/c.

---

### Abstract

It is now common knowledge that the simple quadratic ARCH process has regularly varying tail even when generated by a normally distributed noise, and the tail behaviour is well-understood under more general conditions as well. Much less studied is the case of  $\beta$ -ARCH-type processes, i.e. when the conditional variance is a  $2\beta$ -power function with  $0 < \beta < 1$ . Being a little more general and allowing for asymmetry, we consider threshold  $\beta$ -ARCH models, driven by noises with Weibull-like tails. (Special cases include the Gaussian or the Laplace distributions.) We show that the generated process has approximately Weibull like tail, too, albeit with different exponent:  $1-\beta$  times that of the noise, in the sense that the tail can be bounded from both sides by Weibull distributions of this exponent but slightly different constants. The proof is based on taking an appropriate auxiliary sequence and then applying the general result of Klüppelberg and Lindner (2005) for the tail of infinite MA sequences with light-tailed innovations. *Keywords:* ARCH-type model, conditional heteroscedasticity, extreme value theory, tail behaviour

---

### 1. Introduction

In this paper we examine the tail behaviour of the stationary distribution of certain ARCH-type models defined by the equation

$$X_t = \left( \omega + \alpha_+ (X_{t-1}^+)^{2\beta} + \alpha_- (X_{t-1}^-)^{2\beta} \right)^{1/2} Z_t, \quad (1)$$

---

\*Corresponding author

Email addresses: [elekpeti@cs.elte.hu](mailto:elekpeti@cs.elte.hu) (Péter Elek), [markus@cs.elte.hu](mailto:markus@cs.elte.hu) (László Márkus)

<sup>1</sup>Phone number: +36-1-327-5630

where we apply the usual notations  $x^+ = \max(x, 0)$  and  $x^- = -\min(x, 0)$ . The model parameters  $\omega$ ,  $\alpha_+$  and  $\alpha_-$  satisfy  $\min(\alpha_+, \alpha_-) \geq 0$ ,  $\max(\alpha_+, \alpha_-) > 0$  and  $\omega > 0$ , and  $Z_t$  is an i.i.d. sequence with zero mean and finite variance.

An important feature of this process is that – if the autocorrelation function is defined at all – it is an uncorrelated but not an independent sequence because its conditional variance is changing over time as a function of the lagged values (conditional heteroscedasticity). If  $\beta = 1$  and  $\alpha_+ = \alpha_- > 0$ , we obtain the well-known ARCH (autoregressive conditionally heteroscedastic) model (Engle, 1982), where the conditional variance has a quadratic functional form. Since this process can reproduce the stylised facts (e.g. uncorrelatedness, conditional heteroscedasticity, nonnormality) of financial time series in an easily estimable way, it has become a basic tool in financial econometrics in the past two decades, and gave rise to various generalisations. For instance, in order to model the fact that the variance of stock returns responds more strongly to negative shocks than to positive ones, Glosten et al. (1993) defined the TARCH (threshold ARCH) process by allowing  $\alpha_+ \neq \alpha_-$  in the equation with  $\beta = 1$ . (Hence  $\alpha_+ < \alpha_-$  generally holds in financial applications.) For a broad overview of the various generalisations of ARCH models and some of their properties we refer the reader to Terasvirta (2009).

Due to the popularity of the quadratic ARCH models in finance, their probabilistic properties are quite much studied and well understood. It is a well-known fact for  $\beta = 1$  (see e.g. Embrechts et al. (1997)) that not all choices of  $(\omega, \alpha_+, \alpha_-)$  and of the distribution of  $Z_t$  permit a stationary solution of equation (1). For instance, if  $Z_t$  is normally distributed, the quadratic ARCH model (i.e. the case  $\alpha_+ = \alpha_-$ ) has a stationary solution if and only if  $\alpha_+ = \alpha_- < 2 \exp(\delta) \approx 3.562$ , where  $\delta$  is the Euler-constant. (A different choice for the noise distribution yields a different domain of stationarity.) Also, much is known about the tail behaviour of the stationary distribution if  $\beta = 1$ . It was proven two decades ago (Goldie, 1991) that the simple ARCH process has regularly varying tail (roughly speaking: polynomially decaying tail) even when  $Z_t$  is normally distributed. This phenomenon is often summarised as: "light-tailed input can cause heavy-tailed output". More generally, Borkovec and Klüppelberg (2001) proved that the AR(1) model driven by a quadratic ARCH(1) innovation has regularly varying

tail for a very general class of noise distributions. Using the concepts of extreme value theory (EVT) it follows that the stationary distribution of quadratic ARCH processes belongs to the maximum domain of attraction of the Frechet extreme value distribution or, equivalently, their tail can be approximated by a generalised Pareto distribution (GPD) with shape parameter  $\xi > 0$ . (For an introduction into EVT we refer the reader to Embrechts et al. (1997).)

The  $0 < \beta < 1$  case – where the conditional variance is increasing slower than a quadratic function of the lagged values – is very different from the usual  $\beta = 1$  parameter choice, and is much less studied in the literature. The model may then be called the  $\beta$ -TARCH process and was analysed e.g. by Guegan and Diebolt (1994). It follows relatively easily from the drift condition for Markov chains (Meyn and Tweedie, 1993) that in the  $0 < \beta < 1$  case the  $X_t$  process defined by (1) is stationary irrespective of the choice for the parameters and for the distribution of  $Z_t$  (provided that the latter has a finite second moment and some basic conditions for its density are fulfilled). Moreover, if the  $m$ th moment of  $Z_t$  is finite, the  $m$ th moment of the stationary distribution of  $X_t$  will be finite, too (see Guegan and Diebolt (1994), or in a more general setting Elek and Márkus (2008)). Hence, if all moments of  $Z_t$  is finite and its distribution has infinite support, the distribution of  $X_t$  may only belong to the maximum domain of attraction of the Gumbel law and, equivalently, the shape parameter of the GPD fitted to it may only be zero – if the distribution belongs to the maximum domain of attraction of an extreme value law at all.

This result already yields that the  $\beta$ -TARCH model is lighter tailed than the usual, quadratic specification: for light-tailed  $Z_t$  noises the tail of  $X_t$  decays faster to zero than a polynomial function. The finding, however, does not determine the exact tail behaviour: the maximum domain of attraction of the Gumbel law contains many different types of distributions (e.g. normally, exponentially or lognormally decaying ones). In this paper we give a more precise estimate for the tail decay by showing that  $X_t$  has approximately Weibull-like tail provided that  $Z_t$  has a Weibull-like distribution. Our research is motivated by the fact that  $\beta$ -TARCH models proved useful to model conditional heteroscedasticity in areas where the quadratic ARCH model was considered too heavy-tailed, such as in the analysis of water discharge series of rivers with large

catchments. (See e.g. Elek and Márkus (2008) or in a broader context Szilágyi et al. (2006).)

Throughout the paper we will use the notations  $\bar{F}_X(u) = 1 - F_X(u)$  for the survival function and  $f_X(u)$  for the density function of the random variable  $X$ .

## 2. Tail behaviour

To examine the tail behaviour of  $X_t$  let us introduce an assumption on the tail of  $Z_t$ :

**Assumption 1.**  $Z_t$  is an i.i.d. sequence with a symmetric, absolutely continuous probability distribution. Moreover, there exist  $u_0 > 0$ ,  $\gamma > 0$ ,  $\kappa > 0$ ,  $K_1 > 0$  and  $K_2$  such that its probability density satisfies

$$f_{Z_t}(u) = K_1 |u|^{K_2} \exp(-\kappa |u|^\gamma) \quad (2)$$

for every  $|u| > u_0$ , and  $f_{Z_t}(u)$  is bounded away from zero on  $[-u_0, u_0]$ .

According to this assumption,  $Z_t$  has a Weibull-like tail with exponent  $\gamma$ . The Gaussian ( $\gamma = 2$ ) or the Laplace ( $\gamma = 1$ ) distributions are obtained as special cases.

Guegan and Diebolt (1994) showed under the assumption  $\min(\alpha_+, \alpha_-) > 0$  that if  $\beta > (\gamma - 1)/\gamma$ ,  $X_t$  has no exponential moment (i.e. it is heavier tailed than the exponential distribution) while if  $\beta < (\gamma - 1)/\gamma$ ,  $X_t$  has a moment generating function defined around the neighbourhood of zero. This finding already suggests that  $X_t$  may possess (approximately) a Weibull-like tail with exponent  $\gamma(1 - \beta)$ . Assuming a normally distributed noise (i.e.  $\gamma = 2$ ),  $\alpha_+ = \alpha_-$  and  $1/2 < \beta < 1$ , Robert (2000) argued that this is indeed the case: under his assumptions  $X_t$  has Weibull-like tail with exponent  $2(1 - \beta)$ . Although the proof of his conjectures seems to be incomplete,<sup>2</sup>

---

<sup>2</sup>He derives a functional equation for the logarithm of the moment generating function  $L_Y(s) = E(\exp(sX_t))$  of  $Y_t = \log X_t^2$  and estimates the tail of  $Y_t$  based on the behaviour of  $L_Y(s)$  around  $\infty$ . During the calculations he assumes (see Appendix 1 of his paper) that if a function  $g$  satisfies  $g(x) - g(\alpha x) = O(1/x)$  as  $x \rightarrow \infty$ , then  $g(x) = O(1/x)$ . However, this is not the case: if e.g.  $g(x) = \sin(2\pi \log x / \log \alpha)$  then  $g(x) - g(\alpha x) = 0$ .

some of his ideas are useful to prove that  $X_t$  has approximately Weibull-like tail even if we consider the more general case, i.e.  $\alpha_+ \neq \alpha_-$ ,  $\gamma \neq 2$  and  $0 < \beta < 1$ .

**Theorem 1.** *Assume that  $X_t$  satisfies equation (1), Assumption 1 holds, and  $\omega > 0$ ,  $\min(\alpha_+, \alpha_-) > 0$ ,  $0 < \beta < 1$ . Then, using the notation  $\alpha = \max(\alpha_+, \alpha_-)$ , the survival function of the stationary distribution of  $X_t$  satisfies*

$$\begin{aligned} \exp\left(-\frac{\alpha^{-\gamma/2} \kappa \gamma \beta^{-\frac{\beta}{1-\beta}}}{2} u^{\gamma(1-\beta)} + O\left(u^{\gamma(1-\beta)/2}\right)\right) &\leq \bar{F}_{X_t}(u) \\ &\leq \exp\left(-\frac{(\alpha + \omega)^{-\gamma/2} \kappa \gamma \beta^{-\frac{\beta}{1-\beta}}}{2} u^{\gamma(1-\beta)} + O\left(u^{\gamma(1-\beta)/2}\right)\right). \end{aligned} \quad (3)$$

Clearly, since  $(K - \epsilon)z < Kz + O(z^{1/2}) < (K + \epsilon)z$  as  $z \rightarrow \infty$  for any  $K > 0$  and  $\epsilon > 0$ , the exponent and the bounds for the multiplier of  $u^{\gamma(1-\beta)}$  are exact in the above given Weibull like approximation, but (even if the constants were the same in the upper and lower bound) there still remains room for a deviation from an asymptotically exact Weibull limit distribution.

*Proof.* We may assume without loss of generality that  $\alpha = \alpha_+ \geq \alpha_-$ . Let  $Y_t = \log(X_t^2)$ ,  $U_{t,1} = \log(\alpha_+ Z_t^2)$ , and  $U_{t,2} = \log(\alpha_- Z_t^2)$ . Furthermore, let us introduce the functions

$$h_1(y) = \log(\omega/\alpha_+ + \exp(\beta y)),$$

$$h_2(y) = \log(\omega/\alpha_- + \exp(\beta y))$$

and the random variables  $V_{t,i} = h_i(Y_{t-1}) - \beta Y_{t-1}$  ( $i = 1, 2$ ). Then

$$Y_t = h_1(Y_{t-1}) + U_{t,1} = \beta Y_{t-1} + U_{t,1} + V_{t,1} \quad \text{if } Z_{t-1} > 0,$$

$$Y_t = h_2(Y_{t-1}) + U_{t,2} = \beta Y_{t-1} + U_{t,2} + V_{t,2} \quad \text{if } Z_{t-1} \leq 0.$$

Since  $h_i(y) \geq \beta y$  ( $i = 1, 2$ ),  $V_{t,i} \geq 0$  a.s. Moreover, since  $Z_t$  is a symmetrically distributed i.i.d. sequence,  $Y_t$  can be written as

$$Y_t = \beta Y_{t-1} + U_t + V_t,$$

where  $U_t = U_{t,1}\chi_{\{Z_{t-1}>0\}} + U_{t,2}\chi_{\{Z_{t-1}\leq 0\}}$  is an independent 1/2-1/2 mixture of  $U_{t,1}$  and  $U_{t,2}$ , and is itself an i.i.d. process. Similarly  $V_t = V_{t,1}\chi_{\{Z_{t-1}>0\}} + V_{t,2}\chi_{\{Z_{t-1}\leq 0\}}$  is an independent 1/2-1/2 mixture of  $V_{t,1}$  and  $V_{t,2}$ .

Let us introduce the auxiliary autoregressive sequence

$$Y_t^* = \beta Y_{t-1}^* + U_t = \sum_{i=0}^{\infty} \beta^i U_{t-i}.$$

As we use log in the definition of  $U_t$ , the convergence of the sum cannot be taken for granted. However, the distribution of  $Z_t$  is absolutely continuous (at zero as well) and it has finite variance, hence  $E(U_t^2) < \infty$ , therefore  $Y_t^*$  has a stationary distribution and  $E(Y_t^*)^2 < \infty$ .

The importance of this process comes from the fact that it is stochastically smaller than  $Y_t$ , i.e.  $\bar{F}_{Y_t^*}(u) \leq \bar{F}_{Y_t}(u)$ . To see this, let  $\hat{Y}_0 = Y_0$  and define  $\hat{Y}_t$  recursively as

$$\hat{Y}_t = \beta \hat{Y}_{t-1} + U_t. \quad (4)$$

Using that  $V_t \geq 0$  a.s., we can prove by induction that  $\hat{Y}_t \leq Y_t$ :

$$\hat{Y}_t = \beta \hat{Y}_{t-1} + U_t \leq \beta Y_{t-1} + U_t \leq Y_t. \quad (5)$$

Since  $\hat{Y}_t$  tends in distribution to the stationary distribution of  $Y_t^*$  as  $t \rightarrow \infty$ ,  $Y_t^*$  is indeed stochastically smaller than  $Y_t$ . Therefore, by examining the tail behaviour of  $Y_t^*$  we can obtain a lower bound for the tail of  $Y_t$  as well. (A similar majorisation technique is used in a different context in Lemma 2.2. of Brachner et al. (2009).)

To determine the tail of  $Y_t^*$ , we will apply the framework of Klüppelberg and Lindner (2005) who examined the tail behaviour of linear moving average processes with increments lighter tailed than the exponential distribution. Let  $\sum_{-\infty}^{\infty} c_i W_{t-i}$  be the examined process and assume that the probability density of the i.i.d. sequence  $W_t$  satisfies

$$f(u) = \nu(u) \exp(-\psi(u)), \quad u \geq u_0 \quad (6)$$

for some  $u_0$ , and  $\psi(u)$  is  $C^2$ ,  $\psi'(u_0) = 0$ ,  $\psi'(\infty) = \infty$  and  $\psi''$  is strictly positive on  $[u_0, \infty]$ . Moreover, using the notation  $\phi = 1/\sqrt{\psi''}$ , we assume that the function  $\nu$  is flat for  $\phi$ , i.e.

$$\lim_{u \rightarrow \infty} \frac{\nu(u + x\phi(u))}{\nu(u)} = 1$$

uniformly on bounded  $x$ -intervals, and also that  $\phi$  is self-neglecting, i.e. flat for itself. (Roughly speaking, these assumptions require that  $\psi(u)$  should be strictly convex and  $\nu(u)$  should behave approximately as a constant as  $u \rightarrow \infty$ .)



Furthermore, following the original notations of Klüppelberg and Lindner (2005), define  $q(\tau) = \psi'^{-1}(\tau)$  and

$$Q(\tau) = \sum_{i=-\infty}^{\infty} c_i q(c_i \tau), \quad \sigma_i^2(\tau) = c_i^2 q'(c_i \tau), \quad \sigma_{\infty}^2(\tau) = \sum_{i=-\infty}^{\infty} \sigma_i^2(\tau).$$

It follows from the conditions that  $Q$  is a strictly increasing function. Then, provided that  $c_i$  is a summable sequence of non-negative real numbers, not all zero, and assuming that the two conditions below hold:

$$\lim_{m \rightarrow \infty} \limsup_{\tau \rightarrow \infty} \frac{\sum_{|j| > m} \sigma_j^2(\tau)}{\sigma_{\infty}^2(\tau)} = 0, \quad (7)$$

$$\lim_{m \rightarrow \infty} \limsup_{\tau \rightarrow \infty} \frac{\sum_{|j| > m} \sigma_j(\tau)}{\sigma_{\infty}^2(\tau)} = 0, \quad (8)$$

the following theorem is true:

**Theorem 2.** (Klüppelberg and Lindner, 2005) *Under the above conditions, as  $u \rightarrow \infty$ ,*

$$P\left(\sum_{i=-\infty}^{\infty} c_i W_{t-i} > u\right) \sim \frac{1/\sqrt{2\pi}}{Q^{-1}(u) \sigma_{\infty}(Q^{-1}(u))} \exp\left(-\int_{u_0}^u \sum_{\Sigma c_i} (Q^{-1}(v) + \rho(Q^{-1}(v))) dv\right)$$

where  $\rho(\tau) = o(1/\sigma_{\infty}(\tau))$ . It is also true that  $1/\sigma_{\infty}(\tau) = o(\tau)$  so the first term in the integral is the leading term.

In our case, this theorem will be used with the choice  $U_t = W_t$  so the conditions of the theorem should be checked first. Here,

$$f_{U_t}(u) = \frac{1}{2} \left( K_1 \exp(K_2 u) \exp\left(-\kappa \alpha_+^{-\gamma/2} e^{\frac{\gamma u}{2}}\right) \right) + \frac{1}{2} \left( K_3 \exp(K_4 u) \exp\left(-\kappa \alpha_-^{-\gamma/2} e^{\frac{\gamma u}{2}}\right) \right) = \nu(u) \exp(-\psi(u))$$

with appropriate constants  $K_1 > 0$ ,  $K_3 > 0$  and  $K_2, K_4$ . To satisfy the necessary assumptions with  $u_0 = 0$  in (6),  $\psi(u)$  can be defined as

$$\begin{aligned} \psi(u) &= \kappa \alpha^{-\gamma/2} e^{\frac{\gamma u}{2}} - e \kappa \alpha^{-\gamma/2} / 2 & \text{if } u \geq 2/\gamma, \\ \psi(u) &= e \kappa \alpha^{-\gamma/2} (\gamma/2)^2 u^2 / 2 & \text{if } u < 2/\gamma. \end{aligned}$$

Then it is a matter of routine to check that the resulting  $\nu(u)$  function is flat for  $\phi(u)$  and that  $\phi(u)$  is self-neglecting (see also Example 2.4. (c) in Klüppelberg and Lindner (2005)), so the tail of  $Y_t^*$  can in principle be approximated using  $c_i = \beta^i$  for  $i \geq 0$  and  $c_i = 0$  for  $i < 0$ . (Conditions (7)-(8) will be checked, see below.) Using the notation  $\tau_0 = \epsilon \kappa \alpha^{-\gamma/2} \gamma/2$ , we obtain

$$\begin{aligned}\psi'(u) &= \kappa \alpha^{-\gamma/2} (\gamma/2) e^{\frac{\gamma u}{2}} = e^{-1} \tau_0 e^{\frac{\gamma u}{2}} && \text{if } u \geq 2/\gamma, \\ \psi'(u) &= \epsilon \kappa \alpha^{-\gamma/2} (\gamma/2)^2 u = \tau_0 (\gamma/2) u && \text{if } u < 2/\gamma\end{aligned}$$

and hence

$$\begin{aligned}q(\tau) &= 2\gamma^{-1} \log(e\tau/\tau_0) = 2\gamma^{-1} (\log \tau - \log \tau_0 + 1) && \text{if } \tau \geq \tau_0, \\ q(\tau) &= 2\gamma^{-1} \tau/\tau_0 && \text{if } \tau < \tau_0.\end{aligned}$$

Then

$$\begin{aligned}Q(\tau) &= \sum_{j=0}^{\infty} \beta^j q(\beta^j \tau) = 2\gamma^{-1} \sum_{j=0}^{\infty} \beta^j \log(\beta^j e\tau/\tau_0) \\ &\quad + 2\gamma^{-1} \sum_{j: \beta^j \tau < \tau_0} \beta^j (\beta^j \tau/\tau_0 - \log(\beta^j e\tau/\tau_0)). \quad (9)\end{aligned}$$

For any  $0 < \theta < 1$  (we shall choose  $\theta$  appropriately from this range later), the sum in the second term can be written as

$$\begin{aligned}&\sum_{j: \beta^j \tau < \tau_0} \beta^j (\beta^j \tau/\tau_0 - \log(\beta^j e\tau/\tau_0)) \\ &= (e\tau/\tau_0)^{-\theta} \sum_{j: \beta^j \tau < \tau_0} (\beta^{1-\theta})^j \left( e^\theta (\beta^j \tau/\tau_0)^{1+\theta} - (\beta^j e\tau/\tau_0)^\theta \log(\beta^j e\tau/\tau_0) \right).\end{aligned} \quad (10)$$

For  $0 < \theta < 1$  the function  $g(x) = x^\theta \log x$  is bounded on  $(0, e]$ , hence both  $(\beta^j e\tau/\tau_0)^\theta \log(\beta^j e\tau/\tau_0)$  and  $e^\theta (\beta^j \tau/\tau_0)^{1+\theta}$  are bounded if  $0 < \beta^j \tau/\tau_0 < 1$ . This way we have for  $\tau \rightarrow \infty$

$$\begin{aligned}&\sum_{j: \beta^j \tau < \tau_0} (\beta^{1-\theta})^j \left( e^\theta (\beta^j \tau/\tau_0)^{1+\theta} - (\beta^j e\tau/\tau_0)^\theta \log(\beta^j e\tau/\tau_0) \right) \\ &= O(1) \sum_{j: \beta^j \tau < \tau_0} (\beta^{1-\theta})^j = o(1)\end{aligned}$$

because  $\tau \rightarrow \infty$  implies that  $j \rightarrow \infty$  in the summation condition.

Therefore the sum in (10) (and so the second term in (9)) is  $o(\tau^{-\theta})$ . As a consequence, using the notations

$$A = 2\gamma^{-1}(1-\beta)^{-1} \quad \text{and} \quad B = \beta(1-\beta)^{-1} \log \beta + 1 - \log \tau_0,$$

we obtain for  $\tau \rightarrow \infty$

$$\begin{aligned} Q(\tau) &= 2\gamma^{-1} \sum_{j=0}^{\infty} \beta^j (\log(\beta^j \tau) + 1 - \log \tau_0) + o(\tau^{-\theta}) \\ &= 2\gamma^{-1}(1-\beta)^{-1} (\log \tau + \beta(1-\beta)^{-1} \log \beta + 1 - \log \tau_0) + o(\tau^{-\theta}) \\ &= A(\log \tau + B) + o(\tau^{-\theta}). \end{aligned}$$

Trivially,  $Q^{-1}(u) \rightarrow \infty$  and hence

$$\exp\left(-o\left((Q^{-1}(u))^{-\theta}\right)\right) = 1 + o\left((Q^{-1}(u))^{-\theta}\right) = 1 + o(1) \quad (11)$$

as  $u \rightarrow \infty$ , therefore

$$Q^{-1}(u) = \exp(A^{-1}u - B) \exp\left(-o\left((Q^{-1}(u))^{-\theta}\right)\right) = \exp(A^{-1}u - B) (1 + o(1)). \quad (12)$$

Using again (11) and then (12) we obtain a better estimate for  $Q^{-1}(u)$ :

$$\begin{aligned} Q^{-1}(u) &= \exp(A^{-1}u - B) \left(1 + o\left((Q^{-1}(u))^{-\theta}\right)\right) \\ &= \exp(A^{-1}u - B) \left(1 + o(\exp(-\theta A^{-1}u))\right) \\ &= \frac{\kappa\gamma\alpha^{-\gamma/2}\beta^{-\frac{\beta}{1-\beta}}}{2} \exp\left(\frac{\gamma(1-\beta)}{2}u\right) + o\left(\exp\left(\frac{\gamma(1-\beta)(1-\theta)}{2}u\right)\right). \end{aligned}$$

Let us also check conditions (7)-(8). We obtain that  $q'(\tau) = 2\gamma^{-1}/\tau_0$  if  $\tau < \tau_0$  and  $q'(\tau) = 2\gamma^{-1}/\tau$  if  $\tau \geq \tau_0$ , hence  $\sigma_i^2(\tau) = 2\gamma^{-1}\beta^{2i}/\tau_0$  if  $\beta^i\tau < \tau_0$  and  $\sigma_i^2(\tau) = 2\gamma^{-1}\beta^i/\tau$  if  $\beta^i\tau \geq \tau_0$ . Thus

$$\sigma_{\infty}^2(\tau) = 2\gamma^{-1} \left( \sum_{j: \beta^j\tau \geq \tau_0} \beta^j/\tau + \sum_{j: \beta^j\tau < \tau_0} \beta^{2j}/\tau_0 \right) \sim 2\gamma^{-1}(1-\beta)^{-1}/\tau,$$

so (7)-(8) are easily seen for the  $c_i = \beta^i$  sequence, consequently Theorem 2 can be applied. It also follows that  $\rho(\tau) = O(\tau^{1/2})$  and  $u_0 = 0$  in that Theorem. If we

choose  $\theta > 1/2$ , we obtain

$$\begin{aligned}\bar{F}_{Y_t^*}(u) &= \exp\left(-\log Q^{-1}(u) - \log \sigma_\infty(Q^{-1}(u)) - \int_0^u \left(Q^{-1}(v) + O\left(e^{\frac{\gamma(1-\beta)}{4}v}\right)\right) dv\right) \\ &= \exp\left(-\frac{\kappa\alpha^{-\gamma/2}\beta^{-\frac{\beta}{1-\beta}}}{1-\beta} e^{\frac{\gamma(1-\beta)}{2}u} + O\left(e^{\frac{\gamma(1-\beta)}{4}u}\right)\right).\end{aligned}\quad (13)$$

Taking into account that  $Y_t^*$  is stochastically smaller than  $Y_t = \log(X_t^2)$ , the lower bound is obtained for  $\bar{F}_{X_t}(u)$  in (3).

To show the upper bound for the tail, we can first prove with a similar majorisation technique as used in the proof of  $\bar{F}_{Y_t^*}(u) \leq \bar{F}_{Y_t}(u)$  (equations (4)-(5)) that the increase of either  $\alpha_+$  or  $\alpha_-$  does not make the tail of  $Y_t$  any lighter. Therefore, we can assume that  $\alpha = \alpha_+ = \alpha_-$  and get an upper bound for the tail of this restricted model. In this case, let us introduce for each  $t$  a random variable  $U_t^{**} \geq 0$  such that  $U_t \leq U_t^{**}$  a.s. and  $f_{U_t^{**}}(u) = K f_{U_t}(u)$  for all  $u > 0$  with an appropriate  $K > 0$ . (Such a variable can easily be constructed.) Define also  $h(y) = \beta y \chi_{\{y \geq 0\}} + \log(1 + \omega/\alpha)$ . It follows from  $\alpha_+ = \alpha_-$  that  $h_i(y) \leq h(y)$  ( $i = 1, 2$ ) and thus it can be shown again with a majorisation argument that  $\bar{F}_{Y_t^{**}}(u) \geq \bar{F}_{Y_t}(u)$  holds for the stationary distribution of the model defined by

$$Y_t^{**} = h(Y_{t-1}^{**}) + U_t^{**}.$$

Indeed, let  $\tilde{Y}_0 = Y_0$  and define  $\tilde{Y}_t$  recursively as

$$\tilde{Y}_t = h(\tilde{Y}_{t-1}) + U_t^{**}.$$

Using  $U_t \leq U_t^{**}$  we can prove by induction that  $Y_t \leq \tilde{Y}_t$ :

$$Y_t \leq h(Y_{t-1}) + U_t \leq h(\tilde{Y}_{t-1}) + U_t \leq h(\tilde{Y}_{t-1}) + U_t^{**} = \tilde{Y}_t.$$

Since the distribution of  $\tilde{Y}_t$  tends to the stationary distribution of  $Y_t^{**}$  as  $t \rightarrow \infty$ ,  $Y_t$  is stochastically smaller than  $Y_t^{**}$ .

As  $h(y) \geq 0$  for all  $y$  and  $U_t^{**} \geq 0$  a.s., an alternative definition for  $Y_t^{**}$  is

$$Y_t^{**} = \beta Y_{t-1}^{**} + U_t^{**} + \log(1 + \omega/\alpha) = \sum_{i=0}^{\infty} \beta^i U_{t-i}^{**} + \frac{\log(1 + \omega/\alpha)}{1 - \beta}.\quad (14)$$

The result of Klüppelberg and Lindner (2005) again gives that the tail of  $\sum_{i=0}^{\infty} \beta^i U_{t-i}^{**}$  has the same form as the tail of  $Y_t^*$  (equation (13)). Completing the above reasoning with the effect of the last constant term in (14) it is easy to derive the upper bound for the tail of  $X_t$  in (3).  $\square$

In some applications (e.g. Elek and Márkus (2008)) the  $\alpha_- = 0$  restriction is used. Then the upper bound in (3) certainly holds and a slightly weaker lower bound can also be proven easily:

**Proposition 1.** *Assume the assumptions of Theorem 1 but allow  $\alpha_- = 0$ . Then for every  $\delta > 0$  there exists a  $K > 0$  such that*

$$\exp\left(-Ku^{(1+\delta)\gamma(1-\beta)}\right) \leq \bar{F}_{X_t}(u). \quad (15)$$

*Proof.* Let us use the same notations as in the proof of Theorem 1 and let  $Y_t^{***}$  be defined by

$$\begin{aligned} Y_t^{***} &= \beta Y_{t-1}^{***} + U_t^+ && \text{if } Z_t \geq 0 \\ Y_t^{***} &= U_t^0 && \text{if } Z_t < 0 \end{aligned}$$

where  $U_t^0 = \log(\alpha_0 Z_t^2)$ . It is easily shown that  $Y_t^{***} \leq Y_t$  stochastically in this case. Moreover, as  $Z_t$  is symmetrically distributed, for every  $n \in \mathbb{Z}^+$  with probability  $2^{-n}$

$$Y_t^{***} = \sum_{i=0}^{n-1} \beta^i U_{t-i}^+ + \beta^n Y_{t-n}^{***}.$$

Therefore, using the notations  $q = \bar{F}_{Y_{t-n}^{***}}(0)$  and  $Y_{t,n} = \sum_{i=0}^{n-1} \beta^i U_{t-i}^+$ ,

$$\bar{F}_{Y_t^{***}}(u) \geq q2^{-n} \bar{F}_{Y_{t,n}}(u).$$

Moreover, similarly to the derivation of the tail of  $Y_t^*$ , it follows again from Theorem 2 that

$$\bar{F}_{Y_{t,n}}(u) = \exp\left(-Ke^{\frac{\gamma(1-\beta)}{2(1-\beta^n)}} + O\left(e^{\frac{\gamma(1-\beta)}{4(1-\beta^n)}}\right)\right)$$

with a suitable  $K > 0$ . Choosing  $n$  such that  $1/(1-\beta^n) < 1+\delta$  we obtain

$$\bar{F}_{Y_t^{***}}(u) \geq \exp\left(-Ke^{\frac{(1+\delta)\gamma(1-\beta)}{2}}\right)$$

with a possibly different  $K > 0$ , and transforming  $Y_t$  to  $X_t$  gives the statement of the proposition.  $\square$

### 3. Conclusions

In this paper we showed that the tail of a  $\beta$ -TARCH model can be approximated by Weibull-like distributions with exponent  $\gamma(1 - \beta)$  if the generating noise has Weibull-like tail with exponent  $\gamma$ . We conjecture that the lower bound in Theorem 1 gives the precise asymptotics for the tail of the model but we could not prove this conjecture yet.

It is a natural question to ask how the tail behaviour is modified when AR- or MA-terms are added to the simple uncorrelated  $\beta$ -TARCH model, as e.g. in Elek and Márkus (2008). This question is not yet settled but we conjecture that the more general ARMA- $\beta$ -TARCH model has approximately a Weibull-like tail with the same exponent as the more restricted model examined in this paper.

### References

- Borkovec, M., Klüppelberg, K., 2001. The tail of the stationary distribution of an autoregressive process with ARCH(1) errors. *Ann. Appl. Probab.* 11, 1220-1241.
- Brachner, C., Fasen, V., Lindner, A., 2009. Extremes of autoregressive threshold processes. *Adv. in Appl. Probab.* 41, 428-451.
- Elek, P., Márkus, L., 2008. A light-tailed conditionally heteroscedastic time series model with an application to river flows. *J. Time Ser. Anal.* 29, 14-36.
- Embrechts, P., Klüppelberg, C., Mikosch, T., 1997. *Modelling Extremal Events for Insurance and Finance*. Springer, Berlin.
- Engle, R. F., 1982. Autoregressive conditional heteroskedasticity with estimates of the variance of the United Kingdom inflation. *Econometrica* 50, 987-1007.
- Glosten, L. R., Jagannathan, R., Runkle, D., 1993. On the relation between the expected value and the volatility of the normal excess return on stocks. *J. Finance* 48, 1779-1801.
- Goldie, C. M., 1991. Implicit renewal theory and tails of solutions of random equations. *Ann. Appl. Probab.* 1, 126-166.

- Guegan, D., Diebolt, J., 1994. Probabilistic properties of the  $\beta$ -ARCH-model. *Statist. Sinica* 4, 71-87.
- Klüppelberg, C., Lindner, A., 2005. Extreme value theory for moving average processes with light-tailed innovations. *Bernoulli* 11(3), 381-410.
- Meyn, S. P., Tweedie, R. L., 1993. *Markov Chains and Stochastic Stability*. Springer, London.
- Robert, C., 2000. Extremes of alpha-ARCH models. In: *Measuring Risk in Complex Stochastic Systems*, Ed.: Franke, J., Hardle, W. and Stahl, G., 219-251. Springer, Berlin.
- Szilágyi, J., Bálint, G., Csík, A., 2006. Hybrid, Markov chain-based model for daily streamflow generation at multiple catchment sites. *J. Hydrologic Eng.* 11, 245-256.
- Terasvirta, T., 2009. An introduction to univariate GARCH models. In: *Handbook of Financial Time Series* (eds.: Andersen, T. G., Davis, R. A., Kreiss, J-A., Mikosch, T.). Springer, Berlin.