



HAL
open science

Confinement of knotted polymers in a slit

Richard James Matthews, Ard Louis, Julia Yeomans

► **To cite this version:**

Richard James Matthews, Ard Louis, Julia Yeomans. Confinement of knotted polymers in a slit. *Molecular Physics*, 2011, 109 (07-10), pp.1289-1295. 10.1080/00268976.2011.556094 . hal-00692125


HAL Id: hal-00692125

<https://hal.science/hal-00692125>

Submitted on 28 Apr 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Confinement of knotted polymers in a slit

Journal:	<i>Molecular Physics</i>
Manuscript ID:	TMPH-2010-0417.R1
Manuscript Type:	Special Issue paper - In honour of Bob Evans
Date Submitted by the Author:	03-Jan-2011
Complete List of Authors:	Matthews, Richard; Oxford University Louis, Ard; Oxford University Yeomans, Julia; Oxford University
Keywords:	knotted polymers, confinement, Langevin dynamics
<p>Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.</p> <p>Knot_slit_paper_Resubmit.zip</p>	

SCHOLARONE™
Manuscripts

RESEARCH ARTICLE

Confinement of knotted polymers in a slit

R. Matthews*, A.A. Louis and J.M. Yeomans

*Rudolf Peierls Centre for Theoretical Physics, 1 Keble Road, Oxford OX1 3NP, England**(v4.5 released September 2009)*

We investigate the effect of knot type on the properties of a ring polymer confined to a slit. For relatively wide slits, the more complex the knot, the more the force exerted by the polymer on the walls is decreased compared to an unknotted polymer of the same length. For more narrow slits the opposite is true. The crossover between these two regimes is, to first order, at smaller slit width for more complex knots. However, knot topology can affect these trends in subtle ways. Besides the force exerted by the polymers, we also study other quantities such as the monomer-density distribution across the slit and the anisotropic radius of gyration.

Keywords: knotted polymers; confinement; Langevin dynamics

1. Introduction

Whilst knots are generic, and any polymer of sufficient length in solution is very likely to be knotted [1], they have proven to have particular relevance in biological systems. The DNA contained in the cells of living organisms is typically both long and confined: the genome of *E. coli*, a well studied bacterium [2], is about 3×10^4 persistence lengths [2] ($\sim 1.5\text{mm}$) and is contained in a cell with a typical volume $\sim \mu\text{m}^3$ [3]. Although the DNA is organized using supercoiling and proteins (particularly in eukaryotes) [4], confinement makes entanglements likely [5, 6], particularly during processes such as replication [7]. Knots in DNA can prevent transcription [8], a process necessary for the production of proteins. Indeed, there is a family of enzymes, the topoisomerases [9], one of whose functions is to control knotting. Evidence of highly confined DNA knotting has also been found in bacteriophages [10], viruses that infect bacteria. These knots may affect the ejection speed of a bacteriophage's DNA [11, 12]. Given its relevance in biology, it is important to understand the interplay of knotting and confinement.

The confinement of linear polymers in a slit, between two infinite, parallel walls, has been well studied. The simplest model is to consider a single chain with walls that provide only geometric constraints. For $D > R$, where D and R are slit and polymer size respectively, the chain will maintain a shape similar to that seen without confinement. For $D < R$, however, significant deformation occurs. A scaling form for the free energy of such a chain for $D < R$ was given by Daoud and de Gennes [13] using a blob picture, in which the polymer is taken to form a series of independent blobs of size determined by D , within each of which the polymer

*Corresponding author. Email: r.matthews1@physics.ox.ac.uk

behaves as if it is free. They predicted

$$\frac{F_{conf}}{k_B T} \approx N \left(\frac{a}{D} \right)^{1/\nu}, \quad (1)$$

where N is the number of monomers in the chain, a is monomer size and ν is a three dimensional exponent $\approx 3/5$. Predictions made by Daoud and de Gennes for the polymer size were verified by Webman, Lebowitz and Kalos using simulations [14]. Later, it was predicted that the ratio between the monomer density by the wall and the force exerted on the wall is universal, independent of microscopic details [15]. Again, attempts have been made to verify this prediction by simulation, see ref. [16] and references therein. Whilst the majority of simulation work has used Monte Carlo methods, a recent study has also compared the results of Molecular Dynamics simulations to scaling predictions [17]. Other work has considered issues such as attractive walls [18].

By comparison to the case of linear polymers, there has been little work on knotted polymers in slits and basic issues remain open. Tesi *et al.* [6] looked at the probability that a knot will be found in a slit-confined polymer. They found that for relatively wide slits the probability increased as the slit width decreased. However, for narrower slits, the probability peaked and eventually became less than the value for wide slits. Janse van Rensburg [19] applied both analytical calculations and a Monte Carlo approach that samples configurations with the same topology but varying N . The change of the average value of N with slit width was investigated. Knotted polymers were found to expand as the width increased, in contrast to unknotted polymers whose size showed a plateau after a certain width was reached. In ref. [20], the ends of knotted and unknotted polymers were attached to parallel plates, separated so as to stretch the chains. It was found that the forces exerted on knotted polymers deviated from scaling predictions. The results were used to deduce an estimate of how the knot size scales with polymer length.

Motivated by the common occurrence of confined knotted polymers in biology, we use a coarse-grained model to gain insight into the behaviour of knotted ring polymers in a slit, concentrating on the force exerted on the walls. We then go on to look at other quantities, such as the radius of gyration and the monomer density distribution, that help us to interpret the trends seen.

We find that the effect of knotting differs depending on the degree of confinement of the polymer. We find that for narrow slits, more complex knots exert higher forces on the walls, whereas for relatively wide slits, the opposite is true. We relate the forces to the monomer densities near the walls and interpret the results in terms of the effect of knots on the ability of the polymer to spread out in the slit.

2. Model

We use a standard, bead-spring, polymer model [21]. Excluded volume is included through a truncated Lennard-Jones potential:

$$\begin{aligned} V_{EV}(x) &= 4\epsilon \left[\left(\frac{\sigma}{x} \right)^{12} - \left(\frac{\sigma}{x} \right)^6 + \frac{1}{4} \right], & x \leq 2^{1/6}\sigma, \\ V_{EV}(x) &= 0, & x > 2^{1/6}\sigma, \end{aligned} \quad (2)$$




Figure 1. Knot topology for 8_{18} , based on a diagram from ref. [26].

where σ is the size of a bead, ϵ is an energy scale and x is the bead separation. Neighbouring beads are connected with FENE springs [22]

$$V_{FENE} = -\frac{kX_0^2}{2} \ln \left[1 - \left(\frac{x}{X_0} \right)^2 \right], \quad (3)$$

where X_0 , the maximum extension, is set to 1.5σ and k is chosen as $30\epsilon/\sigma^2$.

The polymer motion is simulated using the Langevin equation [23]:

$$m\ddot{\mathbf{x}}_i = \mathbf{f}_i - \zeta\dot{\mathbf{x}}_i + \sqrt{2\zeta k_B T} \mathbf{r}_i(t), \quad (4)$$

where \mathbf{x}_i is the position of the i^{th} bead, \mathbf{f}_i is the total force acting on it, m is the bead mass, ζ is a friction constant and $k_B T$ is the temperature, which we set to ϵ . $\mathbf{r}_i(t)$ are random vectors that satisfy

$$\begin{aligned} \langle \mathbf{r}_i(t) \rangle &= 0, \\ \langle \mathbf{r}_i(t) \mathbf{r}_j(t') \rangle &= \delta_{ij} \delta(t - t') \mathcal{I}, \end{aligned} \quad (5)$$

where \mathcal{I} is the identity matrix. The second and third terms in Eq. (4) represent the drag due to the solvent and Brownian noise respectively. The parameters may be combined to give a convenient unit of simulation time, $t_0 = \sqrt{(\sigma^2 m / \epsilon)}$. We integrate the motion using a velocity Verlet algorithm [24] with a timestep of $\Delta t = 0.01$.

The polymer is placed in a slit between two parallel plane walls, modelled by the same excluded volume potential as for the bead-bead interactions, Eq. (2). The surfaces of the walls are taken as lying at the point at which the potential is truncated. We consider slit widths from $D = 4\sigma$ to $D = 18\sigma$. This range was chosen, through initial exploratory simulations, to show the crossover between the regimes where more complex knots are harder or easier to confine. The simulations resemble those of Dimitrov *et al.* [17], who considered confined, linear polymers.

We choose a polymer length, $N = 300$, which is sufficiently short that the knots will significantly affect the polymer properties. It should be borne in mind that, since knots in polymers in a good solvent are weakly localized [25], different results would be obtained for other values of N . The results do, however, give a qualitative picture of how a knot, *which is of significant length compared to N* , will affect a polymer in a slit.

We consider linear and unknotted ring polymers (denoted 0_1) and also polymers with knot types 3_1 , 6_1 and 9_1 . C_p is a standard notation [27], where C gives the minimum number of crossings in any projection onto a plane, the essential crossings. p is used to distinguish knots with equal C . 3_1 and 9_1 are torus knots; they both belong to a group of knots with similar topology. 6_1 is an even-twist knot. In addition, we consider two other knots. Firstly, we include a twelve-crossing even-twist knot. We label it 12_1 , although the standard notation only extends up to $C = 10$. Secondly, we also look at 8_{18} , which is chosen because in initial simulations

1 it was found to have an unusually spherical average instantaneous shape [28]. In
 2 Fig. 1. we show a knot diagram [26] for 8_{18} . The polymer model used should
 3 prevent chain crossings [21] and we verified that the knot type remained constant
 4 by calculating the Alexander polynomial [27], $A_k(t)$, at $t = -2$.

5 The radius of gyration of the unconfined linear polymer, $R_G(\text{linear}) = 14.2 \pm 0.1\sigma$.
 6 For the ring polymers with different knots, $R_G(0_1) = 10.65 \pm 0.03\sigma$, $R_G(3_1) =$
 7 $9.01 \pm 0.02\sigma$, $R_G(6_1) = 7.78 \pm 0.01\sigma$, $R_G(8_{18}) = 6.86 \pm 0.01\sigma$, $R_G(9_1) = 7.28 \pm 0.01\sigma$
 8 and $R_G(12_1) = 6.90 \pm 0.01\sigma$. However, it should be kept in mind that the instantane-
 9 ous configurations of linear polymers are not spherical, but rather one can define
 10 3 axes that characterize a prolate shape [28]. We investigated the anisotropic instan-
 11 aneous distribution of knotted polymer shapes, which tend to be more spherical
 12 than linear polymers [29], although here there is no universal behaviour because it
 13 depends on the size of the knot compared to the length of the polymer.

14 We note that for some of the more complex knots, an increase in the internal
 15 energy of the chain larger than the standard error was observed: for narrow slits and
 16 complex knots the probability of collision increases and so the amount of overlap
 17 between beads is higher, leading to an internal energy increase due to the excluded
 18 volume interaction. The largest such increase, $\Delta U = 0.56 \pm 0.06k_B T$, was seen for
 19 12_1 between $D = 8\sigma$ and $D = 4\sigma$. In the middle of this range, the force on the
 20 confining walls was $7.734 \pm 0.003k_B T/\sigma$. This suggests that the internal energy
 21 change accounts for $\lesssim 1\%$ of the free energy increase of the chains, with the rest
 22 being due to entropy. We are thus confident that this effect, which depends on the
 23 specific potential chosen, does not affect our conclusions, which are qualitative.

24 The starting configurations for the polymers were created by performing runs
 25 in which D was reduced at a rate of $10^{-4}\sigma/t_0$ from a wide slit, $D = 25\sigma$, to the
 26 narrowest slit we considered, $D = 4\sigma$. The final configuration was then taken as the
 27 starting configuration for all the simulations at different D for that particular knot
 28 type. By performing extra compression runs, we checked that the configurations
 29 used were not unusual: all those seen at end of the additional compression runs
 30 were qualitatively similar. When viewed from the direction across the slit, they
 31 showed a number of chain crossings that was typically equal to, or only slightly
 32 larger than, the number of essential crossings of the knot.

33 Simulations for each D were performed separately. Starting from a polymer in
 34 the configuration obtained from the compression run, an initialisation period of
 35 $10^5 t_0$ was allowed before results were recorded for $10^7 t_0$. Relaxation was slower in
 36 the two free directions than the confined direction and also depended on D and
 37 knot type [30]: depending on the particular system, the timescale may vary with
 38 knot complexity in different ways [31]. The relaxation time in the free directions,
 39 taken as the time for the auto-correlation function of the relevant component of
 40 the radius of gyration to decay to $1/e$, was typically on the order of $10^2 t_0$ for the
 41 knotted polymers and even for the slowest case ($D = 4\sigma$, linear polymer) was
 42 clearly less than $10^4 t_0$.

43 Errors were estimated from the variance of the relevant quantities, initially as-
 44 suming that all data points were independent. During the simulations, data was
 45 recorded every $100t_0$. If the estimate of the relevant correlation time, t_C , was seen
 46 to be significantly greater than $100t_0$ then the estimated error was increased by a
 47 factor of $\sqrt{2t_C/100t_0}$ [24].

48 **3. Results**

49 We study the average force, f , exerted on the confining walls by the polymer as
 50 function of D for different knot types. The average force for knotted and unknotted
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60




Figure 2. The average force, f , exerted on the walls of a confining slit by polymers with different knot types – 0_1 (\times), 3_1 (\circ), 6_1 (\triangle), 8_{18} (\square), 9_1 ($+$) and 12_1 (\diamond) – divided by the values for a linear chain, as a function of slit width, D . Errorbars are plotted but for most points they are much smaller than the symbols.

rings is plotted as a ratio to the value for a linear chain in Fig. 2.

We first consider the forces exerted by linear and unknotted ring polymers, which are very similar over the range of slit widths investigated. The scaling picture of a confined polymer is based on it forming a series of blobs of dimension $\sim D$ [13]. For small widths, both linear and ring polymers should form the same number of blobs. However, since linear polymers have a larger radius of gyration, it would be expected that, for wide enough slits ($D \gg R_G$), the forces exerted by the unknotted ring would decrease below those for linear polymers. At $D = 30\sigma$, the component of the radius of gyration in the direction of confinement was $5.24 \pm 0.01\sigma$ for a linear polymer, compared to $5.14 \pm 0.01\sigma$ for an unknotted ring, and the force exerted by the unknotted polymer was 0.886 ± 0.006 of the linear polymer value. The difference reflects the fact that $R_G(0_1) < R_G(\text{linear})$.

The topological constraints of a knot that is relatively large compared to the polymer prevent the formation of independent blobs so it is expected that results for knotted rings will deviate from those for linear and unknotted polymers. Fig. 2 shows that, for narrow slits, polymers with more complex knots exert a higher force, whilst for wide slits the opposite is true. The crossover occurs at smaller slit widths for more complex knots, likely because they are more compact [30]. 8_{18} , picked because it is unusually spherical, shows, uniquely amongst the knots studied, non-monotonic behaviour for the force ratio as a function of D . The force itself always showed monotonic behaviour for each knot type. We conjecture that the non-monotonicity in the ratio for 8_{18} reflects a qualitatively different behaviour for this knot compared to the others due to its more spherical shape. A polymer which typically has an extended shape may respond to increasing confinement by losing rotational degrees of freedom for its overall configuration. A highly spherical polymer, such as one with an 8_{18} knot, on the other hand, cannot do this and must deform. This leads to a more rapid increase in the force at relatively large D .

Although the force exerted by 8_{18} is the highest at intermediate slit widths, for narrower D it is very close to 6_1 , a knot with less essential crossings. Once the polymer is significantly deformed, its unconfined shape would not necessarily be expected to further influence its behaviour. Polymers in narrower slits should show behaviour like two-dimensional, adsorbed polymers and, as shown in ref. [25] for example, knots in such polymers show quite different behaviour to those in three-dimensional polymers.

We next compare the radius of gyration of the polymers, focussing on the com-




Figure 3. The radius of gyration, in the confined direction, R_{perp} , of polymers in a slit with different knot types – 0_1 (\times), 3_1 (\circ), 6_1 (\triangle), 8_{18} (\square), 9_1 ($+$) and 12_1 (\diamond) – divided by the values for a linear chain, as a function of slit width, D .




Figure 4. Monomer densities as a function of $\zeta = z/(D + \delta a)$, where z is the distance from the centre of the slit and δ is a correction necessary to achieve agreement with scaling formulae when D is comparable to a [16, 17]. (a) Monomer density across the slit for $D = 4\sigma$ (\square), $D = 12\sigma$ (\circ) and $D = 18\sigma$ (\times) for linear polymers. The solid line is a scaling formula [16, 17]. We use a scaling exponent $\nu = 0.588$ [17] and a correction, $\delta = 1.38$, found to give the best fit to the data. (b)-(d) Monomer density as function of the position across the slit for various knot types: 0_1 (\times), 3_1 (\circ), 6_1 (\triangle), 8_{18} (\square), 9_1 ($+$) and 12_1 (\diamond). Results, for (b) $D = 4\sigma$, (c) $D = 12\sigma$ and (d) $D = 18\sigma$, are plotted as ratios to the values for linear polymers.

ponent in the confined direction, R_{perp} . In Fig. 3, results are plotted as ratios to the values for a linear polymer. The results for R_{perp} show very similar trends to those seen for the forces: as may be appreciated by comparing Figs. 2 and 3, the ordering of the different knot types is almost identical for all D .

To help interpret the results for f and R_{perp} , we examine the monomer density as a function of position across the slit for various D . The results are shown in Fig. 4. Firstly, in Fig. 4(a), we compare the results for linear polymers to the following

1 scaling formula for monomer density [16, 17]:

$$2 \rho(\zeta) = \frac{\Gamma(2 + 2/\nu)}{(\Gamma(1 + 1/\nu))^2} (1/4 - \zeta^2)^{1/\nu}, \quad (6)$$

3 where ζ is the position across the slit divided by $D + \delta a$. δ is a correction, necessary
 4 when D is comparable to a [16, 17]. Assuming a scaling exponent, $\nu = 0.588$ [17], we
 5 find the best fit to the data using $\delta = 1.38$. The results show reasonable agreement
 6 with the scaling formula, although clear differences are seen, particularly for $D =$
 7 4σ . Given that, for this width, D is only a few times bigger than the bead size,
 8 it is unsurprising that such deviations are seen. We also checked our results by
 9 comparing the forces for linear polymers to a scaling formula derived from Eq. (1).
 10 As in the work of Dimitrov *et al.* [17], we found good agreement using the same
 11 value of δ as for the monomer density.

12 In Figs. 4(b)-(d), we compare monomer densities across the slit for different knot
 13 types for $D = 4\sigma$, 12σ and 18σ , by plotting them as ratios to the results for linear
 14 polymers. These simulations show that knots change how the polymer is distributed
 15 across the slit in a different way for different slit widths. For wide slits, $D \gtrsim 14\sigma$,
 16 the constraints of more complex knots make the polymer more compact [30] leading
 17 to a lower monomer density at the walls. For narrower slits, $D \lesssim 8\sigma$, however, the
 18 more complex the knot the higher the wall density. The value of the monomer
 19 density is expected to control the force exerted by the polymer and comparison of
 20 Figs. 2 and 4 show a close correlation between the two quantities.

21 Similarly, the trends for the radius of gyration across the slit shown in Fig. 3 can
 22 be explained in terms of the results in Fig. 4: for narrower slits the monomers tend
 23 to lie closer to the walls for more complex knots leading to an increase in R_{perp} ,
 24 whereas for wider slits the reverse is true.

25 Although it is intuitively reasonable that for wider slits more complex knots lead
 26 to a more compact polymer and hence a lower monomer density at the walls, it
 27 is less obvious why the monomer density at the walls should increase with knot
 28 complexity in narrower slits. A suggestion as to why this may be the case is found
 29 by looking at the distribution of (instantaneous) R_{para} , the radius of gyration in
 30 the two unconfined directions, for a slit with $D = 18\sigma$, the widest we consider.
 31 As shown in Fig. 5, for more complex knots the distribution is narrower and has
 32 a maximum at lower R_{para} indicating that the more complex knots prevent the
 33 polymer from exploring configurations where they are spread out along the slit.
 34 Hence this suggests that when the slit becomes narrow and significantly distorts
 35 the polymer, the polymer cannot so easily distribute its monomers further out in
 36 the free directions and instead their density near the wall increases. As shown in
 37 Fig. 6 the increase in the average R_{para} as D is changed from 18σ to 4σ is less for
 38 more complex knots.

3.1. Summary and Discussion

39 To summarize, we have investigated the effect of knot type on the properties of
 40 a ring polymer confined in a slit of width D . We found that, for wide slits, more
 41 complex knots exert lower forces than a linear polymer on the walls. For more
 42 narrow slits the opposite occurs: knotting leads to larger forces. The forces are
 43 seen to correlate with the monomer densities near the walls. The crossover occurs
 44 at different slit widths for different knots. For larger slit width, when the polymers
 45 are not significantly deformed, those with more complex knots have a smaller radius
 46 of gyration [30], reducing the monomer density by the walls and correspondingly




Figure 5. Distribution of the (instantaneous) radius of gyration in the two free directions, R_{para} , for ring polymers in a slit of $D = 18\sigma$ with different knots: 0_1 (\times), 3_1 (\circ), 6_1 (\triangle), 8_{18} (\square) and 12_1 (\diamond). Results for 9_1 are omitted to improve the clarity of the plot but are seen to fit with the general trend.




Figure 6. Change in the radius of gyration in the two free directions, R_{para} , between slits of width $D = 18\sigma$ and $D = 4\sigma$ for different knot types. Knot type is indicated on the horizontal axis by C , the essential crossings.

the force exerted on them. For smaller slits, the knot prevents the polymer from deforming as much in the free directions as an unknotted polymer can, and so the monomer densities, and consequently forces, are increased.

The example of 8_{18} , which has an unusual spherical shape and shows non-monotonic behaviour of the force ratio (to that for a linear polymer) as a function of D , demonstrates that the particular topology of the knot plays a role. We conjecture that the maximum for 8_{18} occurs because, due to their shape, polymers with 8_{18} cannot respond to decreasing D by losing rotational degrees of freedom for their overall configuration, as more elongated polymers can. Instead, they begin to deform at a point, relative to their size, where other polymers still maintain their preferred shape, if with reduced orientational freedom.

Although we exclusively consider one polymer length, the trends we have uncovered should be generic: except in the large N limit, knots make polymers more compact and will decrease their ability to spread out as they are confined. Therefore we expect that there is a wide range of polymer lengths for which similar qualitative trends will be observed. An interesting issue to be explored in the future is the crossover from weak to strong knot localization as the polymer is confined: polymers in a good solvent in three-dimensions are weakly localized [25] whilst those in two-dimensions are strongly localized. It would also be useful to look at the crossover from the type of behaviour we observe to the large N limit, where, due to localization, knot type will not affect polymer properties.

Acknowledgements

We are very pleased to contribute to this Special Issue of Molecular Physics dedicated to Professor Bob Evans in recognition of his important contributions to the field. We wish him much success in the coming years. We would also like to thank Dirk Aarts and Friederike Schmid for helpful discussions.

References

- [1] D. Sumners and S. Whittington, *J. Phys. A: Math. Gen.* **21**, 1689 (1988).
- [2] J. Berg, J. Tymoczko and L. Stryer, *Biochemistry*, 5th ed. (W. H. Freeman and Co., New York, 2002).
- [3] H. Kubitschek, *J. Bacteriol.* **172** (1), 94 (1990).
- [4] C. Woldringh, *Mol. Microbiol.* **45** (1), 17 (2002).
- [5] J. Michels and F. Wiegel, *Proc. Roy. Soc. London A* **403**, 269 (1986).
- [6] M. Tesi, E. Janse van Rensburg, E. Orlandini and S. Whittington, *J. Phys. A: Math. Gen.* **27**, 347 (1994).
- [7] J. Sogo, A. Stasiak, M. Martínez-Robles, D. Krimer, P. Hernández and J. Schwartzman, *J. Mol. Biol.* **286** (3), 637 (1999).
- [8] J. Portugal and A. Rodríguez-Campos, *Nucleic Acid Res.* **24** (24), 4890 (1996).
- [9] P. Watt and I. Hickson, *Biochem. J.* **303** (3), 681 (1994).
- [10] J. Arsuaga, M. Vázquez, S. Trigueros, D. Sumners and J. Roca, *Proc. Nat. Acad. Sci.* **99** (8), 5373 (2002).
- [11] R. Matthews, A. Louis and J. Yeomans, *Phys. Rev. Lett.* **102** (8), 88101 (2009).
- [12] D. Marenduzzo, E. Orlandini, A. Stasiak, D. Sumners, L. Tubiana and C. Micheletti, *Proc. Nat. Acad. Sci.* **106** (52), 22269 (2009).
- [13] M. Daoud and P. de Gennes, *J. Physique* **38** (1), 85 (1977).
- [14] I. Webman, J. Lebowitz and M. Kalos, *J. Physique* **41** (6), 579 (1980).
- [15] E. Eisenriegler, *Phys. Rev. E* **55** (3), 3116 (1997).
- [16] H. Hsu and P. Grassberger, *J. Chem. Phys.* **120**, 2034 (2004).
- [17] D. Dimitrov, A. Milchev, K. Binder, L. Klushin and A. Skvortsov, *J. Chem. Phys.* **128**, 234902 (2008).
- [18] R. Brak, A. Owczarek, A. Rechnitzer and S. Whittington, *J. Phys. A: Math. Gen.* **38**, 4309 (2005).
- [19] E. Janse van Rensburg, *J. Stat. Mech: Theor. Exp.* **3**, 1 (2007).
- [20] O. Farago, Y. Kantor and M. Kardar, *Europhys. Lett.* **60**, 53 (2002).
- [21] G. Grest and K. Kremer, *Phys. Rev. A* **33** (5), 3628 (1986).
- [22] H. Warner Jr., *Ind. Eng. Chem. Fund.* **11** (3), 379 (1972).
- [23] D. Lemons and A. Gythiel, *Amer. J. Phys.* **65**, 1079 (1997).
- [24] D. Frenkel and B. Smit, *Understanding Molecular Simulation: from Algorithms to Applications* (Academic Press, London, 2002).
- [25] E. Orlandini, A. Stella and C. Vanderzande, *Phys. Biol.* **6**, 025012 (2009).
- [26] C. Livingston, *Knot Theory* (The Mathematical Association of America, Washington, 1993).
- [27] E. Orlandini and S. Whittington, *Rev. Mod. Phys.* **79** (2), 611 (2007).
- [28] J. Aronovitz and D. Nelson, *J. Physique* **47** (9), 1445 (1986).
- [29] E. Rawdon, J. Kern, M. Piatek, P. Plunkett, A. Stasiak and K. Millett, *Macromolecules* **41** (21), 8281 (2008).
- [30] S. Quake, *Phys. Rev. Lett.* **73** (24), 3317 (1994).
- [31] R. Matthews, A. Louis and J. Yeomans, *EPL* **89**, 20001 (2010).