



**HAL**  
open science

## Ecological assessment of *Phragmites australis* wetlands using multi-season SPOT-5 scenes

Brigitte Poulin, Aurélie Davranche, Gaëtan Lefebvre

► **To cite this version:**

Brigitte Poulin, Aurélie Davranche, Gaëtan Lefebvre. Ecological assessment of *Phragmites australis* wetlands using multi-season SPOT-5 scenes. *Remote Sensing of Environment*, 2010, 114 (7), pp.1602-1609. 10.1016/j.rse.2010.02.014 . hal-00692539

**HAL Id: hal-00692539**

**<https://hal.science/hal-00692539>**

Submitted on 16 May 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1  
2 **Ecological assessment of *Phragmites australis* wetlands using multi-season**  
3 **SPOT-5 scenes**

4  
5  
6 **Brigitte Poulin<sup>1</sup>, Aurélie Davranche<sup>1,2</sup> and Gaëtan Lefebvre<sup>1</sup>.**

7 <sup>1</sup>Tour du Valat Research Center, Le Sambuc, 13200 Arles, France

8 <sup>2</sup>Institut für Geographie, Friedrich-Alexander Universität Erlangen-Nürnberg, Kochstraße 4/4,  
9 91054 Erlangen, Germany

10

11

12

13

14

15

16

17 **Running title : Wetland ecological assessment**

18 **Word count : 6594**

19

20

21

22 \*Corresponding author: Tel: +33 (0) 4 90 97 29 58, FAX +33 (0) 4 90 97 20 19

23 E-mail address: [poulin@tourduvalat.org](mailto:poulin@tourduvalat.org)

24

25

26 **ABSTRACT**

27  
28 Ecologists and conservationists need accurate and replicable tools for monitoring wetland  
29 conditions in order to develop and implement adaptive management strategies efficiently. The  
30 Rhone Delta (Camargue) in southern France encloses 9200 ha of fragmented reed marshes  
31 actively managed for reed harvesting, waterfowl hunting or cattle grazing, and holding significant  
32 numbers of vulnerable European birds. We used multi-season SPOT-5 data in conjunction with  
33 ground survey to assess the predictive power of satellite imagery in modelling indicators of reed  
34 structure (height, diameter, density and cover of green/dry stems) relevant to ecosystem  
35 management and bird ecology. All indicators could be predicted accurately with a combination of  
36 bands (SWIR, NIR) and indices (SAVI, OSAVI, NDWI, DVI, DVW, MSI) issued from scenes of  
37 March, June, July, September or December and subtraction between these. All models were  
38 robust when validated with an independent set of satellite and field data. The high spatial  
39 resolution of SPOT-5 scenes (pixel of 10 X 10 m) permits the monitoring of detailed attributes  
40 characterizing the reed ecosystem across a large spatial extent, providing a scientifically-based,  
41 replicable tool for managers, stakeholders and decision-makers to follow wetland conditions in  
42 the short and long-term. Combined with models on the ecological requirements of vulnerable bird  
43 species, these tools can provide maps of potential species ranges at spatial extents that are  
44 relevant to ecosystem functioning and bird populations.

45  
46 **KEY-WORDS:** Camargue, ecosystem health; GLM modelling; multispectral indices; multitemporal  
47 imagery; *Phragmites australis*; vegetation structure, SPOT-5 satellite; state indicators; wetland  
48 monitoring.

## 50 **1. Introduction**

51  
52 Although wetlands support high biodiversity and provide an extensive range of public  
53 goods and services to humankind, they are among the most threatened habitats in the world  
54 (Williams, 1993; Dudgeon et al., 2006). Their high rate of disappearance has prompted the  
55 development of remotely-sensed techniques for mapping their distribution, but little has been  
56 done for monitoring their state of health, especially at extensive spatial extents that are relevant to  
57 ecosystem functioning and species populations. Ecologists and conservationists need accurate  
58 and replicable tools for monitoring wetland conditions in order to develop and implement  
59 adaptive-management strategies efficiently (Ostendorp et al., 1995; Kerr & Ostrovsky, 2003).

60 Common reed *Phragmites australis* (Cav. Trin. ex Steudel) is the most widely distributed  
61 flowering plant on earth (Marks et al., 1994; Güsewell & Klötzli, 2000). It typically grows in or  
62 near freshwater, brackish, and alkaline wetlands along a gradient from deep water (> 2m) to  
63 terrestrial (< 1m below substrate) conditions (Clevering, 1998). It often dominates the area it  
64 occupies to form dense stands in floodplains, lowland shallow lakes and along natural river  
65 channels or irrigation canals. Reed stands are considered as undesirable invaders in some areas of  
66 North America where non-endemic genotypes are proliferating, but have high conservation and  
67 socio-economic value throughout in Europe due to specific vulnerable bird species and various  
68 traditional, recreational and commercial activities (Güsewell & Klötzli, 2000; Ludwig et al.,  
69 2003; Valkamaa et al., 2008).

70 Common reed can resist fire, frost, high pH, water deficit and salt, but has a low tolerance  
71 for wave and current action (Marks et al., 1994; Pagter et al., 2005). Each spring, annual shoots  
72 emerge from perennial underground rhizomes, growing up to 3-4 m tall in optimal conditions.  
73 These vertical stems produce leaves, flower, and eventually set seed. The stems die in early

74 winter but stand as rigid canes for several months (Burgess & Evans, 1989). Density may reach  
75 200 (wet stands) or 300 (dry stands) shoots per meter square (Hara et al., 1993), leading to a rapid  
76 accumulation of decaying matter, which contributes to drying out of the reedbed that eventually  
77 evolves towards scrub and woodland (Granéli, 1989; Cowie et al., 1992). Common reed is  
78 relatively intolerant of summer mowing and cattle grazing (van Deursen & Drost, 1990), but  
79 cutting or burning the dry stems in winter will slow down this hydroseral process (Burgess &  
80 Evans, 1989; Bedford, 2005). Common reed can tolerate a constant salinity of up to 22.5 g/L, but  
81 shoot height, diameter and density will start decreasing above 5 g /L (Lissner & Schierup, 1997).  
82 Decreased reed density and height also result from water deficit (Engloner, 2009). Optimal  
83 conditions for reed growth are freshwater bodies exhibiting a seasonal fluctuations of 30-cm in  
84 water levels (Deegan et al., 2007). A vigorous reedbed will have homogeneous vegetation  
85 cover, tall green stems and a 2:1 ratio of dry to green stems. Permanent flooding without water  
86 renewal will result in lower shoots, a higher dry-to-green-stem ratio, and eventually a clumpy  
87 distribution of reeds following the death of rhizome buds and roots (Armstrong et al., 1996).  
88 Winter burning or cutting will have a positive impact on shoot density and diameter, while  
89 reducing shoot height and increasing plant richness the next spring (Granéli, 1989; Cowie et al.,  
90 1992). A recent review of the factors influencing reed structure, growth and biomass is provided  
91 by Engloner (2009).

92 Habitat selection by breeding birds is tightly related to structural components of the reed  
93 ecosystem throughout Europe (Leisler et al., 1989; Boar, 1992; Jedraszko-Dabrowska, 1992;  
94 Graveland, 1999; Martinez-Vilalta et al., 2002; Gilbert et al., 2005; Polak et al., 2008).  
95 Ecological requirements of reed bird species in southern France have been well identified  
96 (Barbraud et al., 2002; Poulin & Lefebvre, 2002; Poulin et al., 2002; 2005; 2009): the great reed  
97 warbler (GRW) *Acrocephalus arundinaceus* breeds in harvested or non-harvested reedbeds

98 where shoot diameter is above 6 mm (or 195 cm in height); the moustached warbler (MW) *A.*  
99 *melanopogon* prefers areas with a high proportion of stems with flower head (one-year old reed)  
100 intermingled with other emergent plants; the bearded tit (BT) *Panurus biarmicus* is most  
101 abundant in reedbeds having high densities of dry and thin shoots, which often reflects a stress  
102 response to permanent flooding; the colonial purple heron (PH) *Ardea purpurea* nests in flooded  
103 reedbeds having tall and thick shoots with a 2:1 dry:green stem ratio; the Eurasian bittern (EB)  
104 *Botaurus stellaris* is either found in harvested areas (no dry stems) with homogeneous vegetation  
105 cover or in non-harvested reedbeds characterized by numerous small open-water areas used for  
106 foraging, both including small patches of reed cut two winters ago (1:1 dry:green stem ratio).

107 The Rhone delta (Camargue) in southern France encompasses 145 000 ha including 9200  
108 ha of reed marshes that are actively water managed for various socio-economic uses (Mathevet et  
109 al., 2007). These marshes enclose over 50% of the French population of three vulnerable reed  
110 birds in Europe: the moustached warbler, the purple heron and the Eurasian bittern. Conflicts  
111 over water management among users, the presence of salt in the water table together with the  
112 foreseen impacts of climate change justify the development of replicable and robust tools for  
113 monitoring the state of health of these reedbeds. Remote sensing appeared as the most  
114 appropriate approach for monitoring this quasi monospecific and fragmented habitat spread over  
115 a large area and partially located on private properties with difficult access (Davranche et al.,  
116 2009b). In this study, we used multispectral and multi-seasonal data in conjunction with ground  
117 surveys to assess the predictive power of SPOT-5 scenes in modelling reedbed features that are  
118 relevant to bird ecological requirements and management practices. The ultimate goal is to  
119 produce replicable maps of reed-stand conditions to orient management and conservation actions  
120 over the long term.

121

122 **2. Methods**

123

124 *2.1 Study area*

125

126 The Camargue reedbeds are found within water bodies of various types (marsh, pond,  
127 lagoon, river, canal), size (from < 1 to > 2000 ha), hydrology (permanent or temporary flooding  
128 with stable or fluctuating water levels) and underground salinity (from 0.5 to 30 g/L). Spatial  
129 distribution of reedbeds was previously assessed with SPOT-5 scenes and field data using a  
130 binary classification-tree algorithm (Fig. 1). The resulting maps for two successive years lead to  
131 an overall accuracy of 99% and 98% respectively (Davranche et al., 2009b). Water levels in  
132 most of these reedbeds are managed to improve yield of socio-economic activities such as  
133 waterfowl hunting, winter-reed cutting, cattle grazing, fishing and nature conservation (Mathevet  
134 et al., 2007). For instance, harvested reedbeds are typically flooded during the growing season  
135 (March to June) and drained in winter to facilitate access and reduce impact of cutting engines on  
136 the rhizomes, whereas hunting marshes are permanently flooded or drained in spring only. These  
137 activities have a direct impact on reed structure through the creation of open-water areas for  
138 hunting (by cattle grazing or removal of the rhizomes) and the withdrawing of dry stems by  
139 cutting. They also affect reed structure indirectly through water management which, combined  
140 with rainfall and the presence of salt in the water table, will affect reed growth and the overall  
141 state of health of the plant formation. The climate is Mediterranean with mild and windy winters  
142 and hot and dry summers. Mean annual rainfall over the last 30 years is  $579 \pm 158$  (SD) mm,  
143 being concentrated in spring and autumn, with large intra- and inter-annual variations  
144 (Chauvelon, 2009). Total rainfall was 664 and 411 mm in 2005 and 2006, respectively.

145

146 *2.2 Indicators of reed condition*

147

148         Hawke & José (1996) have suggested four indicators of reedbed health: height, diameter,  
149 and density of reed along with presence of shrubs. We have selected five additional criteria easily  
150 measurable in the field that are indicative of stand condition and associated with ecological  
151 requirements of birds and reed harvesters (Table 1). Height and diameter of green stems, as well  
152 as density of flower heads, can vary on a yearly basis and are tightly related to hydrological  
153 conditions (fluctuations in water levels and salinity). Density of reeds and the ratio of dry-to-  
154 green reeds are also affected by hydrology, but more directly by management practices such as  
155 reed cutting, burning or grazing. Low and high reed density can both reflect ecosystem  
156 degradation and this parameter must be interpreted along with reed height or diameter for  
157 condition assessment. Reed-cover homogeneity and scrub encroachment are indicative of a  
158 degraded reedbed evolving towards open water or woodland, respectively. Two of these criteria,  
159 plant richness and shrub encroachment, could not be predicted in this study following their low  
160 levels in our study area.

161

162 *2.3 Field data*

163

164         Water and vegetation measurements were taken in June and July at 39 reedbed sites  
165 (training sample) in 2005 and 21 sites (validation sample) in 2006 (Fig.1). Selection of study sites  
166 resulted in a compromise between admittance, accessibility, and getting a representative sample  
167 of reedbeds based on aerial photographs and videos collected during aerial surveys. At each site,  
168 one sampling plot of 20 X 20 m corresponding to four pixels of a SPOT-5 scene was located  
169 within a homogeneous area representative of a larger zone and located at least 70 m from the



170 border to reduce edge effects in spectral responses. Sampling plots were geolocated with a GPS  
171 (Holux GR-230XX) of 2-5 m accuracy using the average value obtained from the centre of the  
172 plot at three meters above ground to limit the echo caused by tall reeds. Water level, plant cover  
173 and floristic composition were estimated along two diagonals 28-m long crossing the entire plot.  
174 Water levels were systematically estimated with a rule every 4 m along each diagonal and in the  
175 centre of the plot ( $N = 17$ ). We recorded whether this measure was taken above reeds or bare  
176 ground and used the proportion of readings taken in reeds as a degree of reed-cover homogeneity.  
177 This distinction was not made in 2006, hence we used half the 2005 plots for the training sample  
178 and applied the other half on the 2006 scenes for model validation, assuming that this  
179 environmental parameter varied little over one year. Reed density, height and diameter were  
180 measured within four quadrats of 50 X 50 cm per plot located at seven meters from the centre of  
181 the plot in each cardinal direction. Reed density corresponds to the total number of green and dry  
182 stems inside the quadrat. Among dry reeds, we distinguished whole stems with flower heads  
183 (one-year reed) from flowerless or broken stalks. Whole stems with or without flowers were  
184 aggregated during the 2006 field survey, hence we used half the 2005 data for validation on the  
185 2005 scenes. Reed height and diameter were measured on two green and four dry ‘average’ stems  
186 inside the quadrats on both years.

187

#### 188 *2.4 SPOT-5 data*

189

190 SPOT-5 scenes centred on the Rhone delta were acquired on 30 December 2004, 17 March,  
191 19 May, 18 June, 31 July and 21 September 2005 through the SPOT Image Programming Service  
192 (Copyright CNES). These periods were selected based on key events in the phenology and water  
193 management of reedbeds (Davranche et al., 2009b). For model validation, we used images from

194 18 December 2005, 16 March, 29 May, 23 June, 24 July and 15 October 2006, as no scene was  
195 available in September. SPOT-5 scenes have 10-m pixel resolution and four spectral bands: B1  
196 (green: 0.50-0.59  $\mu$  m), B2 (red: 0.61-0.68  $\mu$  m), B3 (near infrared NIR: 0.79-0.89  $\mu$  m) and B4  
197 (short-wave infrared SWIR: 1.58-1.75  $\mu$  m). Radiometric corrections were performed using, the  
198 Second Simulation of the Satellite Signal in the Solar Spectrum (6S), developed by Vermote et al.  
199 (1997). This atmospheric model predicts the sensor signal assuming cloudless atmosphere, taking  
200 into account the main atmospheric effects (gaseous absorption by water vapour, carbon dioxide,  
201 oxygen and ozone; scattering by molecules and aerosols) and lead to a mean error of 0.7% per  
202 band (Davranche et al., 2009a). The corrected scenes were projected to Lambert conformal conic  
203 projection datum NTF (Nouvelle Triangulation Française) using a second-order transformation  
204 and nearest-neighbour re-sampling (Davranche et al., 2009a), and georeferenced to a topographic  
205 map at 1:25 000 scale using ground control points (RMSE < one pixel). Mean reflectance values  
206 at the locations of the field plots were extracted for each band using the ‘Spatial Analyst’ of  
207 ArcGIS version 9.2 (Environmental Systems Research Institute). Eleven multispectral indices  
208 among the most commonly found in the literature (Table 2) were calculated for each scene, as  
209 well as multitemporal indices obtained by subtracting two monthly values from a same indice.

210

### 211 *2.5 Model calibration and validation*

212

213 The six scenes provided 15 possible combinations of subtractions between two dates and  
214 yielded 315 explanatory variables when multiplied by the four bands and 11 indices. A pre-  
215 selection of conceptually meaningful explanatory variables based on environmental resilience of  
216 reeds was made based on the following assumptions: (1) under similar abiotic conditions, reed  
217 parameters should not vary annually and the remotely-sensed data useful for describing these

218 parameters should have a similar constancy; (2) reed parameters are mostly influenced by human  
219 intervention, which is site specific and unlikely to affect distinctly a large number of marshes on  
220 a given year. We compared the degree of relationships ( $R^2$ ) and the mean value ( $t$ -test) of each  
221 predictive variable from the 39 study plots in 2005 and 2006, and conservatively used  $P > 0.01$  as  
222 threshold value for variable selection with both tests.

223 We built Generalized Regression Models with a forward-stepwise procedure in *Statistica*  
224 version 8.0 (StatSoft Inc.) to predict reed parameters from the remotely-sensed data. For each  
225 computed model, the regression analysis was rerun excluding any selected variable, one by one,  
226 to verify that model fit was not improved when the variables selected automatically were  
227 replaced by two or more other variables. Goodness-of-fit of the model was assessed by  
228 calculating the coefficient of determination ( $R^2$ ) and the normalized root-mean-square error  
229 (NRMSE) between the predicted and observed 2005 values (training sample). Predictive  
230 accuracy of the model was assessed by calculating  $R^2$  and NRMSE between the predicted and  
231 observed 2006 values when applying the 2005 model. This validation approach, based on a  
232 dataset independent from the training sample, is considered as the most compelling  
233 demonstration of model usefulness (Mac Nally & Fleishman, 2004; Thomson et al., 2007).  
234 Means of predicted and observed values from the validation sample were further compared using  
235 Student's  $t$ -tests (or Welch's  $t$ -test if unequal variances) to determine whether field calibration  
236 would be required in future model application.

237

238

239

240

241

242 **3. Results**

243

244 *3.1 Pre-selection of predictive variables*

245

246 Fifty-three variables did not differ significantly between 2005 and 2006 and were selected  
247 for model building. Their distribution according to each month and band/index is shown in Figs.  
248 2-3. March and July were the most frequently selected months, followed by June and December  
249 with May showing the least consistent value (Fig. 2). The NIR and SWIR bands and the DVI,  
250 SAVI, OSAVI were the most stable, with many multispectral indices showing less consistent  
251 reflectance values than single bands (Fig. 3).

252

253 *3.2 Modelling of reed indicators*

254

255 All seven reed indicators were modelled with two bands and six indices issued from three to  
256 five scenes (dates), involving subtraction of bands and/or indices between these (Table 3). SAVI  
257 and NDWI were the most commonly selected indices, with NIR and SWIR being the only  
258 selected bands. DVI was selected for modelling dry- and green-reed density and DVW for  
259 modelling reed-cover homogeneity. All these models were accurate when applied to the 2006  
260 independent dataset (Table 3, Fig. 4-6). Reed height was particularly easy to predict, as revealed  
261 by the high correlation between the predicted and observed values (Fig. 4) and by the numerous  
262 significant single- and two-date models that arose during the model selection process. Density of  
263 green and dry stems and their ratio showed significantly or nearly significantly different means in  
264 2006, suggesting that these parameters might require a field calibration (Table 3, Fig. 5).  
265 Actually, a similar reed density can exhibit different spatial patterns from the regular distribution

266 of shoots to sparsely-distributed dense tussocks, which probably explains the lower predictive  
267 power of these models. Reed-cover homogeneity could be accurately and repeatedly predicted, in  
268 spite of the low sample size and the fact that these data were originally meant to reflect  
269 bathymetric variation rather than vegetation coverage (Fig. 6). Modelling of this parameter  
270 would, however, certainly benefit from an increased sampling effort with measurements taken  
271 every meter.

272

### 273 *3.3 Model application*

274

275 Height of reed stem is a good indicator of stand condition. It is correlated with stem  
276 diameter and leaf size (respectively  $R = 0.82$  and  $0.72$ , d.f. = 39,  $P < 0.001$ ), as well as flower  
277 production and resistance to breakage (van der Toorn & Mook 1982; Boar, 1992). It evolves  
278 rapidly following modifications in water management, especially under brackish conditions. The  
279 largest continuous reed area in the Camargue, the Charnier-Scamandre site, is located along a  
280 salinity gradient in its northernmost part, with mean underground salinity varying from 27 to 3  
281 g/L from west to east (Fig. 7). Mapped reed height in 2005 covaries with this salinity gradient.  
282 Predicted reed height ranged from 29 cm to 309 cm per pixel, and from 132 to 222 cm per  
283 hydrological unit. Overall, these values were relatively similar in 2006 (respectively 19-309, 117-  
284 220), although some hydrological units exhibited a different pattern between years owing to a  
285 different water management (Fig. 7). This spatio-temporal variation in reed height can help orient  
286 management options whether in terms of defining optimal hydrological regimes, protecting areas  
287 for vulnerable birds (e.g. great reed warbler and purple heron) or selecting areas for reed  
288 harvesting.

289

#### 290 4. Discussion

291

292 We demonstrate that multi-spectral and seasonal imagery can be a powerful tool for  
293 monitoring fine-scale variations in reedbed attributes relevant to bird ecology and habitat  
294 management. Studies using scenes from multiple dates often rely on a single value (eg: mean  
295 reflectance or principal-component axes) for model building (Suárez-Seoane et al., 2002;  
296 Wiegand et al., 2008). Monthly subtraction between bands and indices issued from different  
297 images associated with specific phenological stages of reed was a key feature of the models  
298 developed in this study, allowing these bands and indices to carry a lot more information than  
299 what they were originally created for.

300 Although SPOT-5 scenes were radiometrically corrected, many bands and indices showed  
301 inconsistent values between years. Their pre-selection, based on their reliability for describing the  
302 selected reed parameters under different environmental and hydrological conditions, appears as a  
303 necessary step of model development. For instance, NDVI was stable in July only and was not  
304 selected in any of the final models, highlighting the usefulness of exploring the potential of other,  
305 less well-known indices. This statement certainly holds for any remote-sensing application in  
306 ecological studies that are currently largely restricted to the use of NDVI (Pettorelli et al., 2005).  
307 The most useful indices for modelling reed attributes in this study were the SAVI, NDWIs  
308 (NDWIF, MNDWI), and DVI for height, diameter and density of green reeds, in addition to MSI  
309 for the ratio dry/green reeds and DVW for reed-cover homogeneity. SAVI accounts for the  
310 spectral contribution of soil and is recommended for predicting biomass when soil exposure is  
311 high relative to vegetation cover (Zhang *et al*, 1997). This index, designed to eliminate soil-  
312 induced variation, was mostly selected in December and March when vegetation coverage of reed  
313 stands is minimal. SAVI was always combined with indices (NDWIs and MSI) or bands

314 involving the NIR and/or SWIR in our models. These bands provide a pigment-independent  
315 quantitative estimate of vegetation water content, and are influenced by leaf structure, leaf dry  
316 matter, canopy matter, canopy structure and leaf area index (Ceccato et al, 2002; Cheng et al,  
317 2006). DVI has been shown to be a good predictor of deciduous plantations and is considered  
318 more sensitive to vegetation density than other indices (Franklin, 2001). In our model, DVI was  
319 most useful for predicting density of green and dry stems and their ratio. DWIs and MSI probably  
320 reflect dry matter and water content, with the advantage of the NDWI not being saturated during  
321 the growing period like other vegetation indices. Gao (1996) suggests that the NDWI, which  
322 increases from dry soil to free water, should be used in combination with NDVI. The DVW  
323 combines both indices to reinforce the perception of free water bodies (Gond *et al*, 2004), and is  
324 then particularly well-suited to model reed-cover homogeneity which reflects the proportion of  
325 open-water areas inside reedbeds.

326       Thorough field campaigns to develop accurate and robust remote-sensing tools that do not  
327 require field sampling when re-applied (or limited field work for calibration), is a desirable  
328 approach for ecosystem long-term monitoring. Application of the models developed in this study  
329 will allow detecting local or regional reedbed degradation in order to orient stakeholders toward  
330 more sustainable management practices. In cases of conflicts around the water resource among  
331 landowners, users and nature conservationists, they will provide an unbiased source of  
332 information to address the impact of various management options for collective-decision making.  
333 Combined with models on the ecological requirements of vulnerable-bird species, they have the  
334 potential to provide precise estimates of potential species ranges and their evolution in the long  
335 term. Human activities are increasing in intensity and extent, ensuing habitat loss and degradation  
336 that impair ecosystem function and reduce the value of ecosystem services for humans. The need

337 to detect and predict changes in natural ecosystems in general and wetlands in particular has  
338 never been greater.

339

## 340 **5. Conclusion**

341

342 Our study shows that stand condition of reed marshes can be mapped accurately over a  
343 major river delta, providing a powerful and robust monitoring tool for evidence-based habitat  
344 management. This study highlights the potentialities of multiseasonal and multi-spectral data in  
345 ecological applications. The increased availability of powerful statistical techniques, geographic  
346 information systems (GIS) and satellite sensors is opening a new field for monitoring the health  
347 of ecosystems across large geographic areas (Guisan & Zimmermann, 2000; Kerr & Ostrovsky,  
348 2003). The high spatial resolution of SPOT-5 scenes makes it possible to obtain detailed  
349 attributes of ecosystem characterization that can be modelled across large spatial extents (Wulder  
350 et al., 2005) abolishing the recurrent scale mismatch between field and remotely sensed-data.

351

352

## 353 **Acknowledgements**

354

355 We are indebted to the Centre National d'Études Spatiales (CNES) for largely funding the  
356 programming and acquisition of SPOT scenes (ISIS Projects 698 & 795), to the Office National  
357 de la Chasse et de la Faune Sauvage (ONCFS) for providing a thesis grant to A. Davranche, and  
358 to the Foundation Tour du Valat for logistic support. Thanks are extended to A. Sandoz for his  
359 technical advice and to E. Duborper, C. Nourry, A. Diot, J. Desgagné, M. Gobert & S. Didier, for  
360 their contribution to the field work. We are also grateful to the Marais du Vigueirat, commune de



361 Vauvert, Syndicat Mixte de Camargue Gardoise and the various stakeholders who gave us access  
362 to their reed marshes.

363

364 **References**

365

366 Armstrong, J., Armstrong, W., & van der Putten, W. H. (1996). Phragmites die-back: bud and  
367 root death, blockages within the aeration and vascular systems and the possible role of  
368 phytotoxins. *New Phytologist*, 133, 399-414.

369 Barbraud, C., Lepley, M., Mathevet, R., & Mauchamp, A. (2002). Reedbed selection and colony  
370 size of breeding purple herons *Ardea purpurea* in Southern France. *Ibis*, 144, 227-235.

371 Bedford, A. P. (2005). Decomposition of *Phragmites australis* litter in seasonally flooded and  
372 exposed areas of a managed reedbed. *Wetlands*, 25, 713-720.

373 Boar, R. R. (1992). Factors which affect the growth of *Phragmites australis*. In D. Ward (Ed.),  
374 *Reedbeds for wildlife* (pp. 21-30). Oxford: The Royal Society for the Protection of Birds &  
375 University of Bristol Information Press.

376 Burgess, N. D., & Evans, C. E. (1989). *The management of reedbeds for birds*. Sandy: The Royal  
377 Society for the Protection of Birds.

378 Ceccato, P., Gobron, N., Flasse, S., Pinty, B., & Tarantola, S. (2002). Designing a spectral index  
379 to estimate vegetation water content from remote sensing data: Part 1 – Theoretical approach.  
380 *Remote Sensing of Environment* 82, 188–197.

381 Chauvelon, P. (2009). *Gestion Intégrée d'une Zone humide littorale méditerranéenne aménagée :*  
382 *contraintes, limites et perspectives pour l'Ile de CAMargue (GIZCAM)*. Programme LITEAU  
383 2, Ministère de l'Ecologie, de l'Energie, du Développement durable et de l'Aménagement du  
384 Territoire, Final report, Tour du Valat, 84 pp.

385 Cheng, Y. B., Zarco-Tejada, P. J., Riano, D., Rueda, C. A., & Ustin, S. L. (2006). Estimating  
386 vegetation water content with hyperspectral data for different canopy scenarios: Relationships  
387 between AVIRIS and MODIS indexes. *Remote Sensing of Environment* 105, 354-366.

388 Clevering, O. A. (1998). Effects of litter accumulation and water table on morphology and  
389 productivity of *Phragmites australis*. *Wetlands Ecology and Management*, 5, 275-287.

390 Cowie, N. R., Sutherland, W. J., Dithogo, M. K. M., & James, R. (1992). The effects of  
391 conservation management of reed beds II. The flora and litter disappearance. *Journal of*  
392 *Applied Ecology*, 29, 277-284.

393 Davranche, A., Lefebvre, G., & Poulin, B. (2009a). Radiometric normalization of SPOT 5  
394 scenes: 6S atmospheric model versus pseudo-invariant features. *Photogrammetric*  
395 *Engineering and Remote Sensing*, 75, 723-728.

396 Davranche, A., Lefebvre, G., & Poulin, B. (2009b) Wetland monitoring using classification trees  
397 and SPOT-5 seasonal time series. *Remote Sensing of Environment*.

398 Deegan, B. M., White, S. D., & Ganf, G. G. (2007). The influence of water level fluctuations on  
399 the growth of four emergent macrophyte species. *Aquatic Botany*, 86, 309-315.

400 Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z., Knowler, D. J., Lévêque, C.,  
401 Naiman, R. J., Prieur-Richard, A. H., Soto, D., Stiassny, M. L., & Sullivan, C. A. (2006).  
402 Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological*  
403 *Reviews of the Cambridge Philosophical Society*, 81, 163-182.

404 Engloner, A. I. (2009). Structure, growth dynamics and biomass of reed (*Phragmites australis*) –  
405 A review. *Flora*, 204, 331-346.

406 Franklin, S.E, (2001). *Remote sensing for sustainable forest management*. Lewis Publishers, New  
407 York.

408 Gao, B. G. (1996). NDWI - a normalized difference water index for remote sensing of vegetation  
409 liquid water from space. *Remote Sensing of Environment*, 58, 257-266.

410 Gilbert, G., Tyler, G. A., Dunn, C. J., & Smith, K. W. (2005). Nesting habitat selection by  
411 bitterns *Botaurus stellaris* in Britain and the implications for wetland management. *Biological*  
412 *Conservation*, 124, 547-553.

413 Gond, V., Bartholome, E., Ouattara, F., Nonguierma, A. & Bado, L. (2004). Surveillance et  
414 cartographie des plans d'eau et des zones humides et inondables en régions arides avec  
415 l'instrument VEGETATION embarqué sur SPOT-4. *International Journal of Remote*  
416 *Sensing*, 25, 987–1004.

417 Granéli, W. (1989). Influence of standing litter on shoot production in reed, *Phragmites australis*  
418 (Cav. Trin. ex Steudel). *Aquatic Botany*, 35, 99-109.

419 Graveland, J. (1999). Effects of reed cutting on density and breeding success of reed warbler  
420 *Acrocephalus scirpaceus* and sedge warbler *Acrocephalus schoenobaenus*. *Journal of Avian*  
421 *Biology*, 30, 469-482.

422 Guisan, A., & Zimmermann, N. E. (2000). Predictive habitat distribution models in ecology.  
423 *Ecological Modelling*, 135, 147–186.

424 Güsewell, S., & Klötzli, F. (2000). Assessment of aquatic and terrestrial reed (*Phragmites*  
425 *australis*) stands. *Wetlands Ecology and Management*, 8, 367-373.

426 Hanqiu, X. (2006). Modification of Normalized Difference Water Index (NDWI) to enhance  
427 open water features in remotely sensed imagery. *International Journal of Remote Sensing*, 27,  
428 3025 – 3033.

429 Hara, T., van der Toorn, J., & Mook, J. H. (1993). Growth dynamics and size structure of shoots  
430 of *Phragmites australis*, a clonal plant. *Journal of Ecology*, 81, 47-60.

- 431 Hawke, C. J., & Jose, P. V. (1996). *Reedbed management for commercial and wildlife interests*,  
432 The Royal Society for the Protection of Birds, Sandy.
- 433 Huete, A. R. (1988). A Soil-Adjusted Vegetation Index (SAVI). *Remote sensing of Environment*,  
434 25, 295-309.
- 435 Hunt, E. R., & Rock, B. N. (1989). Detection of changes in leaf water content using near- and  
436 middle-infrared reflectances. *Remote Sensing of Environment*, 30, 43-54.
- 437 Jedraszko-Dabrowska, D. (1992). Reeds as construction supporting great reed warbler  
438 (*Acrocephalus arundinaceus* L.) and reed warbler (*Acrocephalus scirpaceus* Herm.) nests.  
439 *Ekologia Polska*, 39, 229-242.
- 440 Kerr, J. T., & Ostrovsky, M. (2003). From space to species: ecological applications for remote  
441 sensing. *Trends in Ecology and Evolution*, 18, 299-305.
- 442 Leisler, B., Ley, H. W., & Winkler, H. (1989). Habitat, behaviour and morphology of  
443 *Acrocephalus* warblers: An integrated analysis. *Ornis Scandinavica*, 20, 181-186.
- 444 Lillesand, T. M., & Kiefer, R. W. (1987). *Remote sensing and image interpretation, 2nd edition*.  
445 John Wiley and Sons, New York.
- 446 Lissner, J., & Schierup, H. H. (1997). Effects of salinity on the growth of *Phragmites australis*.  
447 *Aquatic Botany*, 55, 247-260.
- 448 Ludwig, D. E., Iannuzzi, T. J., & Esposito, A. N. (2003). *Phragmites* and environmental  
449 management: a question of values. *Estuaries*, 26 (2B), 624-630.
- 450 Mac Nally, R., & Fleishman, E. (2004) A successful predictive model of species richness using  
451 'indicator' species. *Conservation Biology*, 18, 646-54.
- 452 Marks, M., Lapin, B., & Randall, J. (1994). *Phragmites australis* (*P. communis*): threats,  
453 management, and monitoring. *Natural Areas Journal*, 14, 285-294.

454 Martinez-Vilalta, J., Bertolero, A., Bigas, D., Paquet, J. Y., & Martinez-Vitalta, A. (2002).  
455 Habitat selection of passerine birds nesting in the Ebro Delta reedbeds (NE Spain):  
456 management implications. *Wetlands*, 22, 318-325.

457 Mathevet, R., Le Page, C., Etienne, M., Lefebvre, G., Poulin, B., Gigot, G., & Mauchamp, A.  
458 (2007). ButorStar: a role-playing game for collective awareness of reedbed wise use.  
459 *Simulation & Gaming*, 38, 233-262.

460 Ostendorp, W., Iscli, C., Krauss, M., Krumscheid-Plankert, P., Moret, J. L., Rollier, M., &  
461 Schanz, F. (1995). Lake shore deterioration, reed management and bank restoration in some  
462 central European lakes. *Ecological Engineering*, 5, 51-75.

463 Pagter, M., Bragato, C., & Brix, H. (2005). Tolerance and physiological responses of *Phragmites*  
464 *australis* to water deficit. *Aquatic Botany*, 81, 285-299.

465 Pearson, R. L., & Miller, L. D. (1972). Remote mapping of standing crop biomass for estimation  
466 of the productivity of the short-grass Prairie, Pawnee National Grasslands, Colorado.  
467 Processing of the 8th International Symposium on Remote Sensing of Environment, ERIM,  
468 Ann Arbor, MI, 1357-1381.

469 Pettorelli, N., Vik, J. O., Mysterud, A., Gaillard, J.-M., Tucker, C. J., & Stenseth, N. C. (2005).  
470 Using the satellite-derived NDVI to assess ecological responses to environmental change.  
471 *Trends in Ecology and Evolution*, 20, 503-510.

472 Polak, M., Kasprzykowski, Z., & Kucharczyk, M. (2008). Micro-habitat nest preferences of the  
473 great bittern, *Botaurus stellaris*, on fishponds in central-eastern Poland. *Annales Zoologici*  
474 *Fennici*, 45, 102-108.

475 Poulin, B., & Lefebvre, G. (2002). Effect of winter cutting on the passerine breeding assemblage  
476 in French Mediterranean reedbeds. *Biodiversity & Conservation*, 11, 1567-1581.

- 477 Poulin, B., Lefebvre, G., & Mauchamp, A. (2002). Habitat requirements of passerines and  
478 reedbed management in southern France. *Biological Conservation*, 107, 315-325.
- 479 Poulin, B., Lefebvre, G., Allard, S., & Mathevet, R. (2009). Reed harvest and summer drawdown  
480 enhance bittern habitat in the Camargue. *Biological Conservation*, 142, 689-695.
- 481 Poulin, B., Lefebvre, G., & Mathevet, R. (2005). Habitat selection by booming bitterns *Botaurus*  
482 *stellaris* in French Mediterranean reed-beds. *Oryx*, 39, 265-274.
- 483 Richardson, A. J., & Everitt, J. H. (1992). Using spectral vegetation indices to estimate rangeland  
484 productivity. *Geocarto International*, 1, 63-69.
- 485 Rondeaux, G., Steven, M., & Baret, F. (1996). Optimization of Soil-Adjusted Vegetation Indices.  
486 *Remote Sensing of Environment*, 55, 95-107.
- 487 Rouse, J. W., Haas, R. H., Schell, J. A., & Deering, D. W. (1973). Monitoring vegetation systems  
488 in the great plains with ERTS, Third ERTS Symposium, *NASA SP-351*, 1, 309-317.
- 489 Suárez-Seoane, S., Osborne, P. E., & Alonso, J. C. (2002). Large-scale habitat selection by  
490 agricultural steppe birds in Spain: identifying species-habitat responses using generalised  
491 additive models. *Journal of Applied Ecology*, 39, 755-771.
- 492 Thomson, J., Mac Nally, R., Fleishman, E., & Horrocks, G. (2007). Predicting bird species  
493 distributions in reconstructed landscapes. *Conservation Biology*, 21, 752-66.
- 494 Valkamaa, E., Lyytinen, S., & Koricheva, J. (2008). The impact of reed management on  
495 wildlife: a meta-analytical review of European studies. *Biological Conservation*, 141, 364-  
496 374.
- 497 van der Toorn, J., & Mook, J. H. (1982). The influence of environmental factors and management  
498 on stands of *Phragmites australis*. I. Effects of burning, frost, and insect damage on shoot  
499 density and shoot size. *Journal of Applied Ecology*, 19, 477-499.

- 500 van Deursen, E. J. M., & Drost, H. J. (1990). Defoliation and treading by cattle of reed  
501 *Phragmites australis*. *Journal of Applied Ecology*, 27, 284-297.
- 502 Vermote, E., Tanre, D., Deuze, J. L., Herman, M., & Morcrette, J. J. (1997). Second simulation  
503 of the satellite signal in the Solar Spectrum, 6S: An overview. *IEEE Transactions on*  
504 *Geoscience and Remote Sensing*, 35, 675-686.
- 505 Wiegand, T., Naves, J., Garbulsky, M. F., & Fernández, N. (2008). Animal habitat quality and  
506 ecosystem functioning: exploring seasonal patterns using NDVI. *Ecological Monographs*, 78,  
507 87–103.
- 508 Williams, M. (1993). *Wetlands: A Threatened Landscape*, Blackwell Publishers, Oxford.
- 509 Wulder, M. A., Hall, R. J., Coops, N. C., & Franklin, S. E. (2005). High spatial resolution  
510 remotely sensed data for ecosystem characterization. *BioScience*, 54, 511-521.
- 511 Zhang, M., Ustin, S. L., Rejmankova, E., & Sanderson, E. W. (1997). Monitoring Pacific coast  
512 salt marshes using remote sensing. *Ecological Applications*, 7, 1039-1053.

**Table 1**

Selected indicators of reed conditions and their relationship (+ or –) with reedbed health, requirements of birds and reed harvesters, and management practices. See text for bird species' abbreviations.

Indicator	Reedbed health	Ecological needs		Ecosystem impacts		
		Bird	Harvester	Cutting	Burning	Grazing
Green reed height	+	+ <i>PH</i>	+	–	–	–
Green reed diameter	+	+ <i>GRW, PH</i>	+	+	+	–
Green reed density	+/-	– <i>EB</i>	+	+	+	+
Dry reed density	–	+ <i>BT</i>	–	–	–	–
Flower head density	+	+ <i>MW</i>	+	–	+	–
Ratio dry/green shoots	–	<i>EB, PH</i>	–	–	–	–
Reed cover homogeneity	+	+ <i>EB</i>	+			–
Plant richness	–	+ <i>MW</i>	–	+	+	+
Scrub encroachment	–		–	–	–	–



**Table 2**


Multispectral indices used in this study.

Index	Formula	References
SR - Simple Ratio	$B2/B3$	Pearson & Miller, 1972
VI - vegetation index	$B3/B2$	Lillesand & Kiefer, 1987
DVI - Differential Vegetation Index	$B3-B2$	Richardson & Everitt, 1992
MSI - Moisture Stress Index	$B4/B3$	Hunt & Rock, 1989
NDVI - Normalized Difference Vegetation Index	$(B3-B2)/(B3+B2)$	Rouse et al., 1973
SAVI - Soil Adjusted Vegetation Index	$1.5*(B3-B2)/(B3+B2+0.5)$	Huete, 1988
OSAVI – Optimized SAVI	$(B3-B2)/(B3+B2+0.16)$	Rondeaux <i>et. al.</i> , 1996
NDWI – Normalized Difference Water Index	$(B3-B4)/(B3+B4)$	Gao, 1996
NDWIF – Normalized Difference Water Index of Mc Feeters	$(B1-B3)/(B1+B3)$	Mc Feeters, 1996
MNDWI – Modified Normalized Difference Water Index	$(B1-B4)/(B1+B4)$	Hanqiu, 2006
DVW – Difference between Vegetation and Water	NDVI - NDWI	Gond <i>et al</i> , 2004


**Table 3**

Best models for each reed indicator with their goodness-of-fit (2005) and predictive accuracy (2006) estimated from the correlations between predicted and observed values (\*:  $P \leq 0.05$ , \*\*:  $P \leq 0.01$ , \*\*\*:  $P \leq 0.001$ ) and the normalized root-mean-square error.  $P$  values from  $t$ -tests comparing the predicted and observed mean values are also provided for 2006. Variables are labeled as follow: B4/09 = reflectance of the SWIR band in September; SAVI/12-03 = difference between the SAVI of December and March.


Indicator	Best model	2005		2006		
		$R^2$	NRMSE	$R^2$	NRMSE	$t$ -test
Green-reed height	$141.71 + 414.50 * \text{SAVI}/12-03 - 224.01 * \text{B4}/03-07 - 145.43 * \text{MSI}/06 + 613.57 * \text{B4}/09$	0.67***	12%	0.56***	16%	0.611
Green-reed diameter	$4.02 + 10.60 * \text{OSAVI}/12-03 + 4.91 * \text{MNDWI}/06 + 7.26 * \text{SAVI}/06$	0.45***	19%	0.28*	23%	0.722
Green-reed density	$310.45 - 2258.65 * \text{DVI}/12 + 1356.02 * \text{B3}/03-06$	0.50***	15%	0.18*	29%	0.001
Dry-reed density	$-83.10 + 1885.33 * \text{B3}/12-07 - 826.50 * \text{DVI}/03-06 - 1054.36 * \text{NDWIF}/03-07 - 3247.56 * \text{B3}/03-07$	0.67***	17%	0.35***	28%	0.052
Dry/green-reed ratio	$-2.01 + 52.39 * \text{SAVI}/12-03 - 19.48 / \text{OSAVI}/12-07 - 9.02 * \text{DVI}/03-06 - 24.67 * \text{NDWIF}/03-07$	0.71***	13%	0.52***	32%	0.017
Flowered-reed density	$-13.05 + 168.62 * \text{SAVI}/12 - 233.99 * \text{B3}/03-06$	0.56***	22%	0.56***	22%	0.829
Reed-cover homogeneity	$152.81 - 103.79 * \text{B4}/12-03 + 117.85 * \text{DVW}/03-07 + 202.70 * \text{NDWIF}/06$	0.88***	10%	0.23*	22%	0.645




**Fig 1.** Reedbed distribution in the Camargue with the location of the training (2005) and validation (2006) field plots.




**Fig.2.** Contribution of each scene to the 53 pre-selected predictive variables.




**Fig. 3.** Contribution of each band and indice to the 53 pre-selected predictive variables. Black columns refers to single-date variables and white columns to two-date variables.



**Fig. 4.** Correlation between the predicted and observed values for green-stem height in 2005 (o) and 2006 ( $\Delta$ ) using the best 2005 model.




**Fig. 5.** Correlation between the predicted and observed values for the dry/green-stem ratio in 2005 (O) and 2006 ( $\Delta$ ) using the best 2005 model.



**Fig. 6.** Correlation between the predicted and observed values for reed-cover heterogeneity in 2005 (O) and 2006 (Δ) using the best 2005 model.





**Fig. 7.** Mapping of green-reed height (in cm) at the Charnier-Scamandre site in 2005 and 2006 using natural breaks for color scheme. Water or land areas free of reed are shown in white. Hydrological units either refer to large water bodies or to embanked reed marshes having independent water management.