



**HAL**  
open science

## Optimisation of a Multivalent Strep-tag for Protein Detection

Michael Busby, Lukas Kurt Josef Stadler, Paul Ko Ferrigno, Jason J. Davis

► **To cite this version:**

Michael Busby, Lukas Kurt Josef Stadler, Paul Ko Ferrigno, Jason J. Davis. Optimisation of a Multivalent Strep-tag for Protein Detection. *Biophysical Chemistry*, 2010, 152 (1-3), pp.170. 10.1016/j.bpc.2010.09.005 . hal-00694277

**HAL Id: hal-00694277**

**<https://hal.science/hal-00694277>**

Submitted on 4 May 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Accepted Manuscript


## Optimisation of a Multivalent Strep-tag for Protein Detection

Michael Busby, Lukas Kurt Josef Stadler, Paul Ko Ferrigno, Jason J. Davis

PII: S0301-4622(10)00229-2  
DOI: doi: [10.1016/j.bpc.2010.09.005](https://doi.org/10.1016/j.bpc.2010.09.005)  
Reference: BIOCHE 5431

To appear in: *Biophysical Chemistry*

Received date: 2 August 2010  
Revised date: 20 September 2010  
Accepted date: 20 September 2010



Please cite this article as: Michael Busby, Lukas Kurt Josef Stadler, Paul Ko Ferrigno, Jason J. Davis, Optimisation of a Multivalent Strep-tag for Protein Detection, *Biophysical Chemistry* (2010), doi: [10.1016/j.bpc.2010.09.005](https://doi.org/10.1016/j.bpc.2010.09.005)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Optimisation of a Multivalent Strep-tag for Protein Detection****Michael Busby<sup>a,1</sup>, Lukas Kurt Josef Stadler<sup>a,2</sup>, Paul Ko Ferrigno<sup>2,\*</sup>, Jason J Davis<sup>1,\*</sup>**

<sup>1</sup>Department of Chemistry, University of Oxford,  
Mansfield Road, Oxford OX1 3QR.

<sup>2</sup>Section of Experimental Therapeutics, Leeds Institute of Molecular Medicine, St  
James's University Hospital, Beckett St, Leeds LS9 7TF.

\*Jason.davis@chem.ox.ac.uk

<sup>a</sup>: LKJS and MB made equal contributions to this work


**Keywords - Strep tag, Multivalency, Microarray, Surface Plasmon Resonance, FRET**

**Abstract**

The Strep tag is a peptide sequence that is able to mimic biotin's ability to bind to streptavidin. Sequences of Strep tags from 0-5 have been appended to the N-terminus of a model protein, the Stefin A Quadruple Mutant (SQM) peptide aptamer scaffold, and the recombinant fusion proteins expressed. The affinities of the proteins for streptavidin have been assessed as a function of the number of tags inserted using a variety of labelled and label free bioanalytical and surface based methods (western blots, microarray assays and surface plasmon resonance spectroscopy). The binding affinity increases with the number of tags across all assays, reaching nanomolar levels with 5 inserts, an observation assigned to a progressive increase in the probability of a binding interaction occurring. In addition a novel interfacial FRET based assay has been developed for generic Strep tag interactions, which utilises a conventional microarray scanner and bypasses the requirement for expensive lifetime imaging equipment. By

labelling both the tagged StrepX-SQM<sub>2</sub> and streptavidin targets, the conjugate is primed for label free FRET based displacement assays.

### Graphical abstract



### 1. Introduction

The use of protein affinity tags to facilitate the purification and detection of recombinant proteins has become indispensable to the field of protein chemistry and beyond. The most common classes of tags include antibody epitope tags, such as Myc and haemagglutinin (HA) [1,2], fusion protein tags such as glutathione S-transferase (GST) and maltose binding protein [3,4], and short affinity tags such as the 6xhistidine tag and poly-arginine tag [5,6]. The interaction between streptavidin and biotin possesses one of the lowest dissociation constants in biology and is of particular interest because it occurs even when the biotin is covalently attached to proteins [7]. Each streptavidin monomer binds one biotin molecule with femtomolar affinity, one of the strongest non-covalent interactions observed in nature [8]. Over the past 15 years several groups have developed peptide binders to the tetrameric streptavidin protein [9-13] and a variety of protein engineering approaches have been applied to tailor affinity for specific applications [14-16]. This provides a simple and clean system for protein purification as a streptavidin binding peptide can be selectively eluted from a streptavidin affinity column using appropriate concentrations of biotin. A further advantage of streptavidin as an affinity reagent is its relative low production cost. This in turn has led to the commercialisation of a wide variety of streptavidin-derived materials such as enzymes, plates, microarray slides, beads and more. Schmidt and Skerra were among the first to develop a streptavidin binding peptide when they identified the Strep-tag in a peptide library screen [11]. The Strep-tag, although useful for the one-step purification of recombinant proteins [17], could only be used in the C-terminus of the protein, as a free carboxy-terminus is required for binding. This deficit motivated the development of the StrepII-tag, which is similar in length and can be included anywhere within the protein sequence [10]. Both these tags contain the conserved HPQ amino acid motif, which is found in many streptavidin binding peptides and is chiefly involved in mediating the interaction with streptavidin. Although the shortness of the Strep-tags makes them very attractive as biochemical tools, both bind streptavidin with relatively low affinity (37 $\mu$ M and 72 $\mu$ M, respectively [10]). This poses a problem in cases where higher affinity is required e.g. in recombinant protein microarrays, or in the detection of

Strep-tagged proteins in Western Blots. Consequently, several groups have set out to find higher affinity alternatives to the Strep-tags [9, 13]. Using a mRNA display library, Wilson and coworkers, for example, have identified a 38 amino acid long peptide (streptavidin-binding peptide, SBP), which binds with nanomolar affinity [13].

Juntilla and co-workers have explored the possibility of Strep-tag multivalency by using two tandem StrepII-tags separated by a 12 amino acid linker, a configuration named a StrepIII-tag, in the purification of protein complexes with streptavidin coated beads [18]. An improved detection of the relevant protein complex when using the StrepIII-tag compared to the StrepII-tag was observed on a Western blot assay. However, it remains to be shown whether the increased effectiveness of multiple combinations of StrepII-tags results in multivalency effects, where the measured binding affinity is determined by a combination of stable interactions, or simply by a probability effect, where low affinity interactions occur at high frequency because the binding sites are present in excess. Multivalency [19] effects are commonly observed in nature, where they have evolved to strengthen a diverse array of biochemical interactions. These include polyvalent antibodies binding to an array of antigens on a pathogenic microorganism [20,21], or indeed the attachment of a virus particle coated in binding proteins (e.g. haemagglutinin) to multiple receptors on a host cell [22]. In the laboratory, recent exploitations of polyvalency include the development of a self-assembling peptide polymer, used to improve the attachment rate of nanoparticles to human cells [23], or the development of a dual 6xHistidine tag (two 6xHis tags in tandem) in order to improve binding of recombinant proteins to Ni<sup>2+</sup> coated microarray slides [24].

In the present study we have taken further the development of the strepIII-tag [18] in setting out to investigate the role of binding affinity effects in the performance of the StrepII-tag. We have combined up to 5 StrepII-tags (each of sequence WSHPGF EK) separated by three copies of a regular spacer peptide(GGGS) (Fig. 1A). The tags herein have been appended to the N-terminus of a model protein using the Stefin A Quadruple

Mutant (SQM) peptide aptamer scaffold protein [25]. Peptide aptamers are engineered proteins designed to possess the binding specificities and affinities of antibodies, without the requirement for a host animal or complex post-translational modifications for their production. In our system, the SQM protein is expressed in *E coli* with an amino-terminal tail designed to contain a variety of functional domains that enable subsequent purification or modification of the expressed protein. One of these domains introduces a unique cysteine residue[26] that allows the dimerisation of the SQM proteins under oxidising conditions. We use a subscript “2” to indicate where the SQM protein (and any attached Strep tags) are observed to be largely dimerised (Fig 1C). The StrepII-tag, which for simplicity will from now on only be referred to as the Strep-tag, was introduced either once (strep1-SQM) or up to 5 times (strep5-SQM<sub>2</sub>) into the N-terminal tail of the SQM scaffold, with each tag separated by the 12 amino acid (GGGS)<sub>3</sub> linker. The enhanced binding efficacies afforded by multiple Strep-tags have been assessed by western blot, surface plasmon resonance and fluorescent microarrays. In addition to obvious application to chromatographic purification, the basic format of a surface microarray based FRET assay [27] is also described for the peptide aptamer scaffold. The advantages of a FRET assay over a single (target labelled) fluorophore equivalent are the inherent reductions in artefactual noise (such as protein autofluorescence) and the confirmation that the two binding partners are in an intimate sub 10 nm binding sphere [28,29]. Such a platform is also inherently “label free”, as has shown to be useful for DNA microarrays [30,31].

## 2. Materials and Methods

### 2.1. Plasmid and DNA manipulation for bacterial expression

The streptavidin binding tags were cloned into a modified pET30a+ vector (Novagen, USA; residue[26]), amino-terminal to the SQM peptide aptamer scaffold (pET30a+-SQM, [25]). The Strep-tag (WSHPGF EK) was generated by annealing the primers

F: 5'-CTGGAGCCACCCGCGAGTTCGAAAAG-3'

R: 5'-CTTTTCGAACTGCGGGTGGCTCCAGGTAC-3',

followed by ligation into *KpnI* cut pET30a+-SQM vector. Strep-GGGS(x3)-Strep (strep2) was generated with flanking *KpnI* sites by PCR-amplifying the Strep containing pET30a+ vector using the following primers:

F: 5'-GATCTGGGTACCTGGAGCCACC-3'

R: 5'-

ACGCGGTACCCTTTTCGAACTGCGGGTGGCTCCAAGAACCGCCACCAGAACCGCCACCAGAACC  
GCCACCCTTTTCGAACTGCGGGTG-3'.

The PCR product was purified, digested with *KpnI* and ligated into the *KpnI* cut pET30a+-SQM vector.

A mutagenesis PCR reaction was carried out in order to remove the upstream *KpnI* restriction site flanking the Strep-tag insert using the primers

F: 5'-CCAGATCTGAGTACCTGG-3'

R: 5'-CCAGGTA CT CAGATCTGG-3'

In order to make 3, 4, and 5 Strep tags, we first annealed the primers:

F: 5'-TCTGGGTACCGGTTCTGGTGGCGGTTCTGGTGGCGGTTCTGGAGC-3'

R: 5'-CGTCGGTACCCTTTTCGAACTGCGGGTGGCTCCAAGAACCGCC-3',

followed by a 'fill-in' reaction using Phusion polymerase (NEB, UK). This yields the element GTGSGGSGGGS-Strep (i.e. spacer plus tag), flanked by *KpnI* restriction sites. The amino acid sequence GTG instead of GGG at the beginning of the spacer is necessary as the encoding DNA sequence contains the upstream *KpnI* restriction site. After the PCR reaction the DNA was digested with *KpnI*, followed by a ligation reaction with *KpnI* cut Strep-pET30a+-SQM containing a 10-fold excess of the insert. This yields a Strep-pET30a+-SQM vector with varying additional copies of the Strep-tag (i.e. a total of



3, 4, 5 and more Strep-tags). Finally, to reverse the GTGS part of the spacer (generated by the *KpnI* restriction site) back to the desired GGGs sequence a PCR mutagenesis reaction was performed on the new constructs using the following oligonucleotides:

F: 5'-GCAGTTCGAAAAGGGTGGCGGTTCTG-3'

R: 5'-CAGAACCGCCACCCTTTTCGAACTGC-3'

This reaction will yield a mix of plasmids, some containing the desired reverting mutation to encode the amino acids GGGs, others having some GTGS and some GGGs. Consequently several colonies have to be screened until the desired plasmid encoding all GGGs is found. This was achieved with Strep3- and Strep4-pET30a+-SQM, but not with Strep5-pET30a+-SQM, which consequently remained with the amino acid sequence GTGS at the beginning of 3 of its spacers. A schematic diagram of the constructs used in this study is shown (Fig 1A).

## *2.2. Gel electrophoresis, gel staining and Western Blotting*

StrepX-SQM<sub>1</sub> was separated on a 15% polyacrylamide SDS gel followed by either transfer onto nitrocellulose membrane (Pierce, UK) for western blotting, or incubation in Coomassie blue stain (40% ethanol, 10% isopropanol, 10% acetic acid, 0.25% w/v Coomassie brilliant blue) for 1h at room temperature, followed by incubation in destain (the staining solution without Coomassie blue) overnight at 4°C. For western blotting the membrane was blocked in 3% fat-free powdered milk in TBST (Tris buffered saline pH 8.5 with 0.05% Tween-20) at 4°C overnight and then briefly rinsed in TBST. The membrane was then incubated with 1 µg/ml streptavidin-HRP (Horseradish peroxidase; Invitrogen, USA) for 1 hour at room temperature and washed 3 times for 10 minutes each in TBST. A chemi-luminescent HRP substrate (Perbio, UK) was then applied, excess rinsed off in TBS and the membrane air-dried and exposed to photographic film. For the loading controls the membrane was re-probed using an anti-S-tag antibody (1:5000;

Novagen, USA), which recognises a unique S-tag encoded by the pET30 vector upstream of the Strep tags. A secondary anti-mouse antibody (Jackson Immunolabs) was used at 1:12,000 in TBST. Western blots were quantified using the Quantity One Software (Biorad) and data processed using Excel for Macintosh (Microsoft).

### 2.3. Surface Plasmon Resonance (SPR)

**Surface Activation:** SPR experiments (Autolab ESPRIT, Ecochemie) were performed on gold surfaces initially cleaned in hot piranha solution (warning, piranha solution is a strong oxidant and reacts violently with many organic materials) and then subsequently functionalised overnight with a monolayer of COOH-OEG thiol ( $\text{HS}(\text{CH}_2)_{10}(\text{OCH}_2\text{CH}_2)_3\text{OCH}_2\text{COOH}$ , Prochimia Surfaces), 625  $\mu\text{M}$  in ethanol. SPR chips were then equilibrated to a stable time dependant signal under 10 mM phosphate buffer, pH 7.4 (running buffer). Terminal acid groups were activated with 100  $\mu\text{l}$  of a 1:1 0.2 M EDC and 0.05 M NHS solution for 15 minutes. Depending on the required surfaces, the following final immobilisation protocols were used. **Streptavidin Surface:** A SAM (Self-Assembling Monolayer) consisting of a biotin termination coupled to an underlying gold surface via three repeating ethylene glycol groups was formed by coupling the commercially available EZ-Link Biotin-PEO-LC-Amine, 0.01 M (Pierce) in 0.1 M Bicarbonate buffer pH 8 with the active NHS ester for 20 min. Subsequently, streptavidin was injected at a concentration of 10  $\mu\text{M}$  in running buffer until the absorption had reached a plateau and the SPR sensorgram indicated a large change in surface mass, within 5mDeg. **StrepX-SQM<sub>2</sub> Surface:** The relevant StrepX-SQM<sub>2</sub> protein was injected at a concentration of 10  $\mu\text{M}$  in running buffer until a stable baseline was reached. The complementary target group, either streptavidin or the StrepX-SQM<sub>2</sub> was then injected at variable concentrations and absorption/desorption profiles were measured. Surfaces were regenerated using 0.1 M glycine, pH 2.4, for 2 minutes.

Surface coverages in  $\text{ng mm}^{-2}$  were calculated from the relationship  $1\text{mDeg} = 122\text{ng mm}^{-2}$  [32]. Kinetic parameters were calculated with Kinetic Evaluation Software (Ecochemie).

#### 2.4. Microarray and FRET Assays

All proteins were labelled using standard protocols supplied by the dye manufacturer, Atto-tec GmbH. Concentrations were checked using a Nanodrop spectrophotometer (Thermo Scientific). Microarray assays were performed by spotting (BioOdessey, Biorad Corp) the StrepX-SQM<sub>2</sub> samples on amine coated glass slides (Genetix) using 100  $\mu\text{m}$  capillary pins. Samples were printed at concentrations of 10  $\mu\text{M}$  in print buffer which comprised of PBST (phosphate buffered saline + 3% Tween 20) and 10% glycerol. Array features were printed in repeats of four and the entire array was repeated three times across the slide. Spotted volumes were allowed to incubate on the slide for 60 min prior to a blocking step in 1% BSA in PBST for 60 min. Atto 647N-NHS ester labelled streptavidin (1:1 labelling ratio) was then incubated at a concentration of 7 nM in 1% BSA in PBST for 40 min using 50  $\mu\text{l}$  volume Lifterslips (Thermo Scientific). Slides were washed in PBST twice for 10 min then twice in deionised water for 2 min. The slides were dried under a stream of nitrogen and scanned at a resolution of 5  $\mu\text{m}$  with 543 nm lasers under the Cy3 detection protocol or 633 nm lasers under the Cy5 detection protocol (ScanArray Express, Perkin Elmer). Spot intensities were analyzed with the integrated ScanArray software. FRET assays were performed under identical conditions to microarray binding assays, *vide supra*, though now both capture and target molecule are labelled, with donor (Atto 550) and acceptor (Atto 647N) fluorophores, respectively. For these assays, array features were printed in repeats of three and at two concentrations (5  $\mu\text{M}$  and 10  $\mu\text{M}$ ).

### 3. Results

#### 3.1. *Strep-tag/streptavidin interaction in a Western Blot format*

Initially the relative changes in binding affinity between one and multiple Strep-tags for streptavidin were assessed in a western blot format. Heat denaturation and SDS stabilisation of the StrepX-SQM proteins in the presence of DTT produce linear, monomeric forms of the proteins containing 0-5 Strep-tags that are separated electrophoretically and transferred onto nitrocellulose. Therefore in this assay, the interactions occur between reduced monomeric StrepX-SQM immobilised on a membrane and soluble HRP-conjugated streptavidin, although some re-folding can occur during the transfer step. HRP metabolises a chemi-luminescent substrate and relative binding affinities are measured via signal intensity on photographic film (Fig. 2A). The use of 2 Strep-tags increases binding to streptavidin by approximately 5-fold compared to a single Strep tag according to this method. A further significant increase in affinity is observed when 3 tags are used (12-fold compared to a single Strep-tag and 2.5 fold compared to Strep2). Further small improvements are observed with Strep4 and Strep5 (Fig. 2B,  $p > 0.05$ ; Student's t-test used).

### 3.2. *Surface Plasmon resonance measurements*

A more quantitative analysis of relative binding affinities of the StrepX-SQM<sub>2</sub> systems in their native state (oxidised, dimeric, with no denaturation) was sought through surface plasmon resonance analysis. The series of StrepX-SQM<sub>2</sub> samples were flowed at 1  $\mu$ M over a pre-formed streptavidin-Biotin-SAM (Fig. 3A). The maximum equilibrium coverage taken from sensorgrams was plotted after subtraction of any non-specific binding contributions measured using Strep0-SQM<sub>2</sub> and the relative affinities were displayed as  $\mu\text{M cm}^{-2}$ . Consistent with the western blot data, binding is observed to increase with additional Strep-tags from 1-5, though the rate of increase is observed to progressively fall. The percentage increase in the number of moles of streptavidin immobilised between Strep1-SQM<sub>2</sub> and Strep2-SQM<sub>2</sub> was 124% whereas between Strep4-SQM<sub>2</sub> and Strep5-SQM<sub>2</sub> the increase was much smaller at 13%. This observation is consistent with the onset of steric saturation as the surface density of Strep tags

increases (Fig. 3B) [33]. Analyses repeated under reducing conditions were equivalent (data not shown).

These quantitative trends were observed to be fully consistent on inverting the interfacial configuration, that is on exposing immobilised StrepX-SQM<sub>2</sub> on a COOH-OEG terminated SAM to solution phase streptavidin (0.05 to 2  $\mu$ M). Equilibrium dissociation constants,  $K_D$ , obtained for Strep1-SQM<sub>2</sub> and Strep5-SQM<sub>2</sub> (Fig. 3C) were  $1.49 \pm 0.01 \mu$ M and  $260 \pm 0.18$ nM respectively (Fig. 3D). The associated  $R_{max}$  values for the interaction were  $186.61 \pm 0.70$  mDeg for the Strep1-SQM<sub>2</sub> surface and  $877.07 \pm 0.17$  mDeg for that presented by Strep5-SQM<sub>2</sub>. When comparing the  $R_{max}$  for the initial StrepX-SQM<sub>2</sub> surface and the  $R_{max}$  for the saturated binding data it becomes apparent that, although capture ratios scale with the number of Strep-tags, they remain below 1:1 for both conditions. Thus, the capture ratio for Strep1-SQM<sub>2</sub> = 0.18:1 while that for Strep5-SQM<sub>2</sub> = 0.43:1. This is consistent with both expected steric limitations and observed electrophoretic mobility shift assays carried out in solution where only 1:1 complexation was observed (data not shown). These analyses, of course, do not shed light on how many tags per SQM molecule are involved in the binding to any one streptavidin but prior work, at typical surface presentations of biotin, have confirmed that streptavidin binds predominantly through two of its sites [34,35,36], a configuration demonstrably useful in building up surface structures through accessing the initially unused streptavidin sites pointing away from the surface.

### 3.3. *Affinity measurements in a protein microarray format*

In order to further study the binding behaviour of the multiple Strep-tags and pave the way for a surface based FRET sensor, the binding efficiencies of the Strep-tag systems were investigated in a microarray format. StrepX-SQM<sub>2</sub> samples were spotted from 10  $\mu$ M solutions onto amine modified glass slides in repeated arrays to compensate for potential slide inhomogeneities. An excessive concentration was used to ensure that all

the active surface binding sites were saturated with the StrepX-SQM<sub>2</sub> protein. Incubation with fluorescently labelled streptavidin and subsequent fluorescence intensity analysis confirmed a broadly linear increase in binding affinity with tag number (Fig. 4B), tailing off slightly at higher tag numbers. Interestingly, in this format, the strep0-SQM<sub>2</sub> gave a zero background response, indicating that the microarray surface was entirely specific.

#### 3.4. *Streptavidin FRET Assay*

A microarray streptavidin FRET assay, was developed using the microarray surface described previously. In this configuration, the capture and target biomolecules are labelled with complementary (FRET pairing) dye molecules. In the proximity afforded by binding, donor excitation-dependant fluorescence is detected at the acceptor emission wavelength. In a comparative analysis of binding interactions between streptavidin and Strep0-SQM<sub>2</sub> and Strep5-SQM<sub>2</sub> previously discussed, one would expect that “no FRET” and “strong FRET” behaviours will be observed respectively (if nonbonded dye-dye interactions are negligible, Fig. 5A). The dye pair choice was optimised so as to both maximise spectral overlap at convenient frequencies and generate a Förster radius large enough to accommodate the dimensions of the proteins used herein (Atto550/Atto647N,  $R_0 = 6.5$  nm, Atto-tec.com). In all experiments, dye-to-protein labelling ratios were maintained at 1:1, so as to remove potential contributions from labelling variance to any observed variations in emission.

The first microarray lines (Fig. 5B) show Strep0-SQM<sub>2</sub>-Atto550 spots used as donor only leakage controls, to assess whether the donor contributes fluorescence at the measured emission wavelength of the acceptor. As noted earlier, streptavidin-Atto647N binding is not expected here. The absence of acceptor dye is confirmed by the strong donor signal present, confirming no FRET and no crosstalk. The second line of the array confirms the presence of streptavidin bound acceptor (streptavidin-Atto647N) bound to Strep5-SQM<sub>2</sub>-Atto550, as excitation of the SQM bound donor results in emission from Atto647.

The bottom microarray line is a Strep5-SQM<sub>2</sub>/streptavidin-Atto647N pairing. Here, only the Atto647N dye is present which acts as the acceptor leakage control; again no signal is seen in either the donor or the FRET channel. These results confirm that the dyes are acting orthogonally to each other and the assay is not detecting crosstalk. It also confirms that the FRET signal is due to binding interactions between the two labelled proteins (see discussion for further details).

#### 4. Discussion

The first experiments on the multivalent effects of multiple Strep-tag units enhancing streptavidin binding affinity came from Hengsakul and Cass, who showed that, when coupling the original Strep-tag [10,11] to the C-terminus of dimerising alkaline phosphatase, a dissociation constant of 0.5  $\mu$ M was measured [37]. Though  $K_d$  values obtained were lower than reported with a single “original” Strep-tag (72  $\mu$ m) [10], an observation assigned to the presence of two tags in the dimeric phosphatase, direct comparisons cannot be made as experimental conditions in the two studies were distinct. Though the Streplll-tag, comprised of two StreplI-tags, is reported to engender improved streptavidin binding affinity, reported results are wholly qualitative [18]. We have demonstrated herein that combining 2 StreplI-tags separated by a 12 amino acid linker and fused to an empty peptide aptamer scaffold protein does improve streptavidin binding, by approximately five fold, compared to that of the single tag. Furthermore, the addition of a third, fourth and fifth StreplI-tag leads to further sequential binding increases, though the advantage of this progressively diminishes. These observations have been reproducibly confirmed under carefully controlled conditions across three assay formats. They have also been confirmed in either “inversion” of the surface binding assay (the SQM fusion protein on the surface capturing solution phase streptavidin or *vice versa*).

The physical basis of the increase in target pair binding can be explained in terms of multivalence [19,21]. The increase in tag number results, specifically, in both a higher local concentration of binding points, and a greater receptive surface area at which the large 52.8 kDa streptavidin molecules can be accommodated. The added advantage of streptavidin having four binding pockets increases the potential network of interactions that may occur, though it is likely that only 1-2 binding pockets per streptavidin are bound to any one SQM receptor at any single point in time [34-36]. An approximately linear increase in affinity is achieved on adding more tags through the Strep1-Strep5 series and associated  $K_d$  values progressively fall into the nanomolar regime. Based on the linear behaviour, the effect is deemed to be probabilistic resulting from greater capture of individual streptavidin molecules. The discrepancy observed herein between the Strep1-SQM<sub>2</sub> and original calculated StrepII-tag  $K_d$  values may be reflective of interactions involving the scaffold protein. Mammen et al [38], have presented the relationship between poly and monovalent binding constants ( $K_{poly}$  and  $K_{mono}$  respectively) as  $K_{poly} = (K_{mono})\alpha^N$ , where  $\alpha$  quantifies the “degree of cooperativity” and  $N$  is the potential number of individual interactions. Applied here to the data presented in section 3.2 (Strep1-SQM<sub>2</sub> vs Strep5-SQM<sub>2</sub>),  $\alpha$  is resolved as being very close to 1.0, consistent with these observations being probabilistic rather than cooperative. All experimental formats utilised indicate a progressive reduction in the advantage conferred by adding more tags across the StrepX-SQM<sub>2</sub> series. This observation is entirely consistent with a progressive steric saturation of binding sites presented by the StrepX-SQM<sub>2</sub> protein (the steric footprint of streptavidin covering large portions of the active surface), preventing any further streptavidin molecules from binding and is thus related to similar observations made on increasing biotin surface density at planar monolayer surfaces [33]. Such behaviour has been quantitatively defined herein by SPR, where the maximum molar ratio of streptavidin that may bind to the Strep5-SQM<sub>2</sub> is observed to be 0.43:1. For the Strep1-SQM<sub>2</sub> surface the maximum streptavidin binding is 0.18:1.



Finally, the strong tagged StrepX-SQM<sub>2</sub> streptavidin association can be utilised in establishing robust interfacial FRET interactions. Though numerous examples of the applications of FRET based bioassays exist [39-43] few have translated to solid surfaces [27,44,45] due to the complexity of experiment design. In addition, the majority of interfacial FRET based assays have relied on the use of life time imaging instrumentation [27,45], which although convenient and sensitive is also highly specialised. Examples which rely on photoluminescence intensity have utilised quantum dot/ gold nanoparticle pairs, which due to their size may interfere with complex protein-protein interactions [44]. The work described herein presents a self consistent methodology for configuring surface based assays between two protein pairs that can be performed on a conventional microarray scanner, with small molecule modification of the capture and target proteins. The data also confirms that the two binding partners are in an intimate sub 10 nm binding sphere. These factors contribute favourably to the potential applications of the assay towards the capture of more complex recombinantly expressed biomolecules with appended Strep-tag tails or towards the development of more sophisticated and sensitive assays based with potential label free applications [30,31].

## 5. Conclusion

A progressive improvement in binding affinity between the StrepII-tag and streptavidin can be engineered in association with a StrepX-SQM<sub>2</sub> scaffold. On increasing the number of Strep-tags to 5, nanomolar  $K_d$ 's are afforded and can be replicated across several interfacial assay formats. A surface based microarray based on FRET between the capture molecule and the target has also been demonstrated, highlighting this new highly sensitive, selective assay platform. Future work should explore the use of different length spacers between the tags, which might further improve affinity. Applications of the FRET assay can be progressed with the introduction of FRET assays that detect more complex and physiologically relevant targets. In addition even higher

sensitivity and limits of detection are obtainable based on lifetime dependant analysis [27].

### Acknowledgements

This work was funded through BBSRC Grant BB/F011296/1 to JJD and PKF. LKJS and PKF gratefully acknowledge the support of AstraZeneca.

### References


- [1] P.A. Kolodziej, R.A. Young, Epitope tagging and protein surveillance, *Methods Enzymol.*, 194 (1991) 508-519.
- [2] T.N. Tai, W.A. Havelka, S. Kaplan, A broad-host-range vector system for cloning and translational lacZ fusion analysis, *Plasmid*, 19 (1988) 175-188.
- [3] P. Duplay, H. Bedouelle, A. Fowler, I. Zabin, W. Saurin, M. Hofnung, Sequence of male gene and of its product, the maltosebinding protein of *Escherichia coli* K12, *J. Biol. Chem.*, 259 (1988) 10606-10613.
- [4] D.B. Smith, K. Johnson, Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase, *Gene*, 67 (1988) 31-40.
- [5] J. Porath, J. Carlsson, I. Olsson, G. Belfrage, Metal chelate affinity chromatography, a new approach to protein fractionation, *Nature*, 258 (1975) 598-599.
- [6] K. Terpe, Overview of tag protein fusions: from molecular and biochemical fundamentals to commercial systems, *Appl. Microbiol. Biotechnol.*, 60 (2003) 523-533.
- [7] E.P. Diamandis, T.K. Christopoulos, The biotin-(strept)avidin system: principles and applications in biotechnology, *Clin. Chem.*, 37 (1991) 625-636.
- [8] N.M. Green, Avidin and streptavidin, *Methods Enzymol.*, 184 (1990) 51-67.
- [9] T. Lamla, V.A. Erdmann, Searching sequence space for high-affinity binding peptides using ribosome display, *J. Mol. Bio.*, 329 (2003) 381-388.
- [10] T.G.M. Schmidt, J. Koepke, R. Frank, A. Skerra, Molecular interaction between the strep-tag affinity peptide and its cognate target, streptavidin, *J. Mol. Bio.*, 225 (1996) 753-766.
- [11] T.G.M. Schmidt, A. Skerra, The random peptide library-assisted engineering of a C-terminal affinity peptide, useful for the detection and purification of a functional Ig Fv fragment, *Protein Eng.*, 6 (1993) 109-122.
- [12] P.C. Weber, M.W. Pantoliano, L.D. Thompson, Crystal structure and ligand-binding studies of a screened peptide complexed with streptavidin, *Biochem. J.*, 31 (1992) 9350-9354.
- [13] D.S. Wilson, A.D. Keefe, J.W. Szostak, The use of mRNA display to select high-affinity protein-binding peptides, *PNAS*, 98 (2001) 3750-3755.
- [14] L.A. Klumb, V. Chu, P.S. Stayton, Energetic roles of hydrogen bonds at the ureido oxygen binding pocket in the streptavidin biotin complex, *Biochem.*, 37 (1998) 7657-7663.

- [15] A. Chilkoti, P.H. Tan, P.S. Stayton, Site-directed mutagenesis studies of the high-affinity streptavidin-biotin complex: contributions of tryptophan residues 79, 108, and 120, *PNAS*, 92 (1995) 1754-1758.
- [16] S. Freitag, V. Chu, J.E. Penzotti, L.A. Klumb, R. To, D. Hyre, I. Le Trong, T.P. Lybrand, R.E. Stenkamp, P.S. Stayton, A structural snapshot of an intermediate on the streptavidin-biotin dissociation pathway, *PNAS*, 96 (1999) 8384-8389.
- [17] T.G.M. Schmidt, A. Skerra, One-step affinity purification of bacterially produced proteins by means of the "Strep tag" and immobilized recombinant core streptavidin, *J. Chromatog. Sect.*, 676 (1994) 337-345.
- [18] M.R. Junttila, S. Saarinen, S. Schmidt, J. Kast, J. Westermarck, Single-step Streptag® purification for the isolation and identification of protein complexes from mammalian cells, *Proteomics*, 5 (2005) 1199-1203.
- [19] M. Mathai, C. Seok-Ki, M.W. George, Polyvalent Interactions in Biological Systems: Implications for Design and Use of Multivalent Ligands and Inhibitors, *Angew. Chem. Int. Ed.*, 37 (1998) 2754-2794.
- [20] N.J. Dimmock, Multiple mechanisms of neutralization of animal viruses, *Trends. Biochem. Sci.*, 12 (1987) 70-74.
- [21] M. Oda, T. Azuma, Reevaluation of stoichiometry and affinity/avidity in interactions between anti-hapten antibodies and mono- or multi-valent antigens, *Mol. Immunol.*, 37 (2000) 1111-1122.
- [22] W.J. Lees, A. Spaltenstein, J.E. Kingery-Wood, G.M. Whitesides, Polyacrylamides bearing pendant alpha-sialoside groups strongly inhibit agglutination of erythrocytes by influenza A virus: multivalency and steric stabilization of particulate biological systems., *J. Med. Chem.*, 37 (1994) 3419-3433.
- [23] A.J. Simnick, C.A. Valencia, R. Liu, A. Chilkoti, Morphing low-affinity ligands into high-avidity nanoparticles by thermally triggered self-assembly of a genetically encoded polymer, *ACS Nano*, 4 (2010) 2217-2227.
- [24] C. Steinhauer, C. Wingren, F. Khan, M. He, M.J. Taussig, C.A.K. Borrebaeck, Improved affinity coupling for antibody microarrays: Engineering of double-(His)<sub>6</sub>-tagged single framework recombinant antibody fragments, *Proteomics*, 6 (2006) 4227-4234.
- [25] T. Hoffmann, L.K.J. Stadler, M. Busby, Q. Song, A.T. Buxton, S.D. Wagner, J.J. Davis, P. Ko Ferrigno, Structure-function studies of an engineered scaffold protein derived from stefin A. I: Development of the SQM variant, *Prot. Eng. Des. Sel.*, 23 (2010) 403-413.
- [26] J.J. Davis, J. Tkac, S. Laurenson, P. Ko Ferrigno, Peptide aptamers in label-free protein detection: 1. Characterization of the immobilized scaffold, *Anal Chem*, 79 (2007) 1089-1096.
- [27] S. Nagl, R. Bauer, U. Sauer, C. Preininger, U. Bogner, M. Schaeferling, Microarray analysis of protein-protein interactions based on FRET using subnanosecond-resolved fluorescence lifetime imaging, *Biosens. Bioelec.*, 24 (2008) 397-402.
- [28] S. Nagl, M. Schaeferling, O.S. Wolfbeis, Fluorescence analysis in microarray technology, *Microchim. Acta*, 151 (2005) 1-21.



- [29] E. Oh, M.-Y. Hong, D. Lee, S.-H. Nam, H.C. Yoon, H.-S. Kim, Inhibition assay of biomolecules based on fluorescence resonance energy transfer (FRET) between quantum dots and gold nanoparticles, *J. Am. Chem.Soc.*, 127 (2005) 3270-3271.
- [30] A.A. Lubin, K.W. Plaxco, Folding-based electrochemical biosensors: The case for responsive nucleic acid architectures, *Acc. Chem. Res.*, 43 (2010) 496-505.
- [31] K.M. Wang, Z.W. Tang, C.Y.J. Yang, Y.M. Kim, X.H. Fang, W. Li, Y.R. Wu, C.D. Medley, Z.H. Cao, J. Li, P. Colon, H. Lin, W.H. Tan, Molecular engineering of DNA: molecular beacons, *Angew. Chem.-Int. Edit.*, 48 (2009) 856-870.
- [32] X. Su, Y.-J. Wu, W. Knoll, Comparison of surface plasmon resonance spectroscopy and quartz crystal microbalance techniques for studying DNA assembly and hybridization, *Biosens. Bioelect.*, 21 (2005) 719-726.
- [33] L.M. Bonanno, L.A. DeLouise, Steric Crowding Effects on Target Detection in an Affinity Biosensor, *Langmuir*, 23 (2007) 5817-5823.
- [34] L.S. Jung, K.E. Nelson, P.S. Stayton, C.T. Campbell, Binding and dissociation kinetics of wild-type and mutant streptavidins on mixed biotin-containing alkylthiolate monolayers, *Langmuir*, 16 (2000) 9421-9432.
- [35] V.H. Pérez-Luna, M.J. O'Brien, K.A. Opperman, P.D. Hampton, G.P. López, L.A. Klumb, P.S. Stayton, Molecular recognition between genetically engineered streptavidin and surface-bound biotin, *Journal of the American Chemical Society*, 121 (1999) 6469-6478.
- [36] L. Haeussling, H. Ringsdorf, F.J. Schmitt, W. Knoll, Biotin-functionalized self-assembled monolayers on gold: surface plasmon optical studies of specific recognition reactions, *Langmuir*, 7 (1991) 1837-1840.
- [37] M. Hengsakul, A.E.G. Cass, Alkaline Phosphatase-Strep tag fusion protein binding to streptavidin: resonant mirror studies, *J. Mol. Bio.*, 266 (1997) 621-632.
- [38] M. Mammen, S.-K. Choi, G.M. Whitesides, Polyvalent interactions in biological systems: implications for design and use of multivalent ligands and inhibitors, *Angew. Chem. Int. Ed.*, 37 (1998) 2754-2794.
- [39] R.M. Clegg, Fluorescence resonance energy transfer, *Curr. Opin. Biotechnol.*, 6 (1995) 103 - 110.
- [40] R. Heim, R.Y. Tsien, Engineering green fluorescent protein for improved brightness, longer wavelengths and fluorescence resonance energy transfer, *Curr Biol*, 6 (1996) 178 - 182.
- [41] J.R. Lakowicz, *Principles of Fluorescence Spectroscopy*, (2006).
- [42] J. Kwan, A. Ling, E. Papp, D. Shaw, J.M. Bradshaw, A fluorescence resonance energy transfer-based binding assay for characterizing kinase inhibitors: Important role for C-terminal biotin tagging of the kinase, *Anal. Biochem.*, 395 (2009) 256-262.
- [43] C.S. Lebakken, S.M. Riddle, U. Singh, W.J. Frazee, H.C. Eliason, Y. Gao, L.J. Reichling, B.D. Marks, K.W. Vogel, Development and applications of a broad-coverage, TR-FRET-based kinase binding assay platform, *J. Biomol. Screen.*, 14 (2009) 924-935.
- [44] Y.P. Kim, Y.H. Oh, E. Oh, S. Ko, M.K. Han, H.S. Kim, Energy transfer-based multiplexed assay of proteases by using gold nanoparticle and quantum dot conjugates on a surface, *Anal. Chem.*, 80 (2008) 4634-4641.

[45] S.H. Kim, M. Jeyakumar, J.A. Katzenellenbogen, Dual-mode fluorophore-doped nickel nitrilotriacetic acid-modified silica nanoparticles combine histidine-tagged protein purification with site-specific fluorophore labeling, *J. Am. Chem. Soc.*, 129 (2007) 13254-13264.

ACCEPTED MANUSCRIPT



**Fig. 1. A.** Multiple cloning site region of the pET30a<sup>+</sup>-SQM construct used in this study. All streptavidin binding tags were inserted into the KpnI restriction site. Approximate location of the cysteine residue (cys) is indicated. **B.** Purified samples of StrepX-SQM separated on SDS-PAGE with and without DTT followed by staining with Coomassie blue. Dimer formation mediated by the cysteine residue is readily observed as a molecular weight shift (increase) in the oxidised state (-DTT). Numbers 1-5 signify number of Strep-tags (X) in the monomer. **C.** Schematic representation of dimer formation by StrepX-SQM.



C.


No. Strep-tags	1	2	3	4	5
StrepX/Strep1	1	4.8	12.1	13.7	15.6

**Fig. 2. A.** Western Blot probing using streptavidin-HRP and anti-S-tag antibody loading control. Amounts of protein loaded: 1.5, 3, 6, 30 pmoles. The highest amount, 30 pmoles was only used for 0 and 1 Strep-tags, as the signal was too strong at this level with multiple tags. **B.** Densitometry analysis using Strep1-SQM as the baseline. Affinity of Strep2-SQM compared to Strep1-SQM is significantly improved (student's t-test  $p < 0.05$ ), as well as Strep3-SQM compared to Strep2-SQM ( $p < 0.05$ ). Although there is further improvement with 4 and 5 Strep-tags, this is not statistically significant ( $p > 0.05$ ). \*denotes a statistically significant increase in affinity of StrepX compared to Strep(X-1).


C. Fold increase of the affinity of streptavidin for StrepX-SQM normalised to Strep1-SQM.

ACCEPTED MANUSCRIPT






**Fig. 3. A.** Surface Plasmon Resonance response of the target StrepX-SQM<sub>2</sub> (X = 0-5) injected at equal concentrations over a streptavidin-biotin-SAM. **B** Plot of corrected molar surface coverage of the target StrepX-SQM<sub>2</sub> (X = 0-5) obtained from A. In comparing molar surface coverages. A trend in increasing interaction is clearly seen from Strep 1-5. All values have been corrected for molecular mass differences, and nonspecific (Strep0-SQM<sub>2</sub>) binding. **C** Variable concentrations of the target streptavidin injected over a Strep5-SQM<sub>2</sub>-OEG-SAM immobilised surface. **D** Equilibrium analysis over the variable concentrations of the target streptavidin obtained from C, showing the calculated  $k_D = 260 \pm 0.18$  nM and  $R_{max} = 877.07 \pm 0.17$  mDeg.



**Fig. 4. A.** Microarray spotting of StrepX-SQM<sub>2</sub> sample capture of Atto-647N labelled streptavidin. The microarray consists of seven spotted samples: a buffer control (spotted twice) and six StrepX-SQM<sub>2</sub> samples (X= 0-5). Each line contains two samples each, spotted in repeats of four. There are four lines in total, the first two lines cannot be seen due to no signal (buffer control, Strep0-SQM<sub>2</sub>) or low signal (Strep1-SQM<sub>2</sub>). **B** Relative average intensities with standard deviations over four samples of captured streptavidin emission.



**Fig. 5. A.** Experiment and control scenarios which define the specificity of the FRET assay. FRET will only occur when both StrepX-SQM<sub>2</sub> and streptavidin are labelled with donor (green) and acceptor (red) dyes, and when the StrepX-SQM<sub>2</sub> is present to bring the donor and acceptor units together in proximity. StrepX-SQM<sub>2</sub> proteins are in the dimeric state. **B)** Microarray FRET assay measured in different emission channels. i)

donor channel measuring Atto 550 fluorescence, which corresponds to the spotted StrepX-SQM<sub>2</sub> samples. ii) Acceptor channel measuring Atto 647N fluorescence, which corresponds to the bound streptavidin-Atto 647N molecules. iii) FRET channel measuring Atto 647N fluorescence upon Atto 550 excitation. The sample layout is described beneath different lines correspond to different samples spotted in repeats of three times at two concentrations.

ACCEPTED MANUSCRIPT