



Une démonstration du théorème de Catalan-Mihăilescu et généralisation à la conjecture de Pilai

Jamel Ghannouchi

► To cite this version:

Jamel Ghannouchi. Une démonstration du théorème de Catalan-Mihăilescu et généralisation à la conjecture de Pilai. 2012. hal-00695280v19

HAL Id: hal-00695280

<https://hal.science/hal-00695280v19>

Preprint submitted on 4 Nov 2012 (v19), last revised 31 Jan 2013 (v22)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Une démonstration du théorème de Catalan-Mihăilescu et généralisation à la conjecture de Pillai

Jamel Ghanouchi

jamel.ghanouchi@topnet.tn

Abstract

(MSC=11D04) More than one century after its formulation by the Belgian mathematician Eugene Catalan, Preda Mihailescu has solved the open problem. But, is it all ? Mihailescu's solution utilizes computation on machines, we propose here not really a proof as it is intended classically, but a resolution of an equation like the resolution of the polynomial equations of third and fourth degrees. This solution is totally algebraic and does not utilize, of course, computers or any kind of calculation.

(Keywords : Diophantine equations, Catalan equation ; Algebraic resolution)

Introduction

Catalan theorem has been proved in 2002 by Preda Mihailescu. In 2004, it became officially Catalan-Mihăilescu theorem. This theorem stipulates that there are not consecutive pure powers. There do not exist integers strictly greater than 1, $X > 1$ and $Y > 1$, for which with exponents strictly greater than 1, $p > 1$ and $q > 1$,

$$Y^p = X^q + 1$$

but for $(X, Y, p, q) = (2, 3, 2, 3)$. We can verify that

$$3^2 = 2^3 + 1$$

Euler has proved that the equation $X^3 + 1 = Y^2$ has this only solution. We propose in this study a general solution. The particular cases already solved concern $p = 2$, solved by Ko Chao in 1965, and $q = 3$ which has been solved in 2002. The case $q = 2$ has been solved by Lebesgue in 1850. We solve here the equation for the general case.

The resolution

Let Catalan equation

$$Y^p = X^q + 1$$

We pose

$$c = \frac{X^p - 1}{Y^{\frac{p}{2}}}; \quad c' = \frac{7 - X^p}{Y^{\frac{p}{2}}}$$

Thus

$$Y^{\frac{p}{2}} = \frac{6}{c + c'}$$

$$X^p = cY^{\frac{p}{2}} + 1 = \frac{6c}{c + c'} + 1 = \frac{7c + c'}{c + c'}$$

And

$$X^q = Y^p - 1 = \frac{36 - (c + c')^2}{(c + c')^2}$$

We have $X^p > 7$ hence $c' < 0$ and $c > c'$ and $Y^{\frac{p}{2}} \geq 3$ thus $0 < c + c' \leq 2$ Two cases
 $c^2 > 1$ and $c^2 \leq 1$

$c^2 > 1$ We have

$$\begin{aligned} X^q - X^{2p} &= \frac{36 - (c + c')^2 - (7c + c')^2}{(c + c')^2} \\ &< \frac{36 - 36c^2}{(c + c')^2} < 0 \end{aligned}$$

We deduce

$$q + 1 \leq 2p$$

But let

$$Z = \frac{c + 7c'}{c + c'}$$

We have

$$Z + X^p = 8$$

Thus

$$8X^q - (Z + X^p)^2 = 8X^q - 64 \geq 0$$

And

$$8X^{q-2p} \geq (ZX^{-p} + 1)^2$$

If $X^p = 8$ or $(X, p) = (2, 3)$ then $2p - 1 = 5 \geq q$ implies $q \in \{2, 3, 4, 5\}$ and it is impossible. Let

$$n = \frac{-6X^p}{6X^p - 48} = X^p Z^{-1} < 0 \Rightarrow n + 1 = \frac{-48}{6X^p - 48} = 8Z^{-1} < 0$$

$$\Rightarrow \frac{n+1}{n} = \frac{8}{X^p} > 0$$

$$m = \frac{6Z}{-6Z + 48} = ZX^{-p} < 0 \Rightarrow m + 1 = \frac{48}{-6Z + 48} = 8X^{-p} > 0$$

$$\Rightarrow \frac{m+1}{m} = \frac{8}{Z} < 0$$

But, we have

$$\begin{aligned} 8(n-m)(Z^{-1}+X^p)(m+1) &= (n-m)(66+m+n) = 8(n-m)(Z+X^{-p})(n+1) = 64(n-m)(Z^{-1}X^{-p}+1) \\ &= (n-m)((n+1)(m+1)+64) \end{aligned}$$

But it means that

$$X^{-p}Z = \frac{m+1}{n+1} = \frac{8(n-m)(X^{-p}+Z)+64m-n-1}{64n-m-1}$$

And

$$Z^{-1}X^p = \frac{n+1}{m+1} = \frac{8(m-n)(Z^{-1}+X^p)+64n-m-1}{64m-n-1}$$

Hence

$$\begin{aligned} &\frac{(m+1)^2}{(n+1)^2} \\ &= \frac{(8(n-m)(X^{-p}+Z)+64m-n-1)(64m-n-1)}{(8(m-n)(Z^{-1}+X^p)+64n-m-1)(64n-m-1)} \end{aligned}$$

And

$$\begin{aligned} &(8(m-n)(Z^{-1}+X^p)(m+1)+64(n+1)-(m+1)^2)(64(n+1)-(m+1)^2) \\ &= (8(n-m)(X^{-p}+Z)(n+1)+64(m+1)-(n+1)^2)(64(m+1)-(n+1)^2) \\ &= 64(m-n)(Z^{-1}X^{-p}+1)(64(n+1)-(m+1)^2)+(64(n+1)-(m+1)^2)^2 \\ &= 64(n-m)(Z^{-1}X^{-p}+1)(64(m+1)-(n+1)^2)+(64(m+1)-(n+1)^2)^2 \\ &\quad (m-n)(n+m+66)(64(n-m)+(n-m)(n+m+2)) \\ &= (64(m+1)-(n+1)^2)^2-(64(n+1)-(m+1)^2)^2 \\ &= (64(m-n)+(m-n)(n+m+2))(64(m+n+2)-(n+1)^2-(m+1)^2) \end{aligned}$$

Thus

$$(n-m)^2(n+m+66)^2 = (n-m)(m+n+66)(64(m+n+2)-(n+1)^2-(m+1)^2)$$

It means that $X^p = Z$ (because $n = m$) or

$$ZX^{-p} = 1$$

Thus

$$X^p = Z \Rightarrow 8X^{q-2p} > (ZX^{-p}+1)^2 = 4$$

Hence

$$4X^{q+1} \geq 8X^q > 4X^{2p} \geq 4X^{q+1}$$

Or

$$q+1 = 2p$$

And $X = 2$ and

$$\begin{aligned} 7c+c' &= 7c'+c \Rightarrow (c-c')Y^{\frac{p}{2}} = 0 = X^p - 1 - 7 + X^p \\ &\Rightarrow (X,p) = (2,2) \Rightarrow (Y,q) = (\pm 3, 3) \end{aligned}$$

$$c^2 \leq 1$$

$$\begin{aligned} 2X^q - X^{2p} &= \frac{72 - 2(c+c')^2 - (7c+c')^2}{(c+c')^2} \\ &> \frac{64 - 49c^2}{(c+c')^2} > 0 \end{aligned}$$

Thus

$$X^q \geq X^{2p-1}$$

Or

$$q+1 \geq 2p$$

But let

$$Z = 8 - X^q$$

If we suppose $X^q \neq 8$. We have

$$Z + X^q = 8$$

$$4X^{2p} - (Z + X^q)^2 = 4X^{2p} - 64 \geq 0$$

And

$$4X^{2p-q} \geq X^q(ZX^{-q} + 1)^2$$

Let

$$\begin{aligned} n &= \frac{-6X^q}{6X^q - 48} < 0 \Rightarrow n+1 = \frac{-48}{6X^q - 48} < 0 \\ &\Rightarrow \frac{n+1}{n} = \frac{8}{X^q} > 0 \\ m &= \frac{6Z}{-6Z + 48} < 0 \Rightarrow m+1 = \frac{48}{-6Z + 48} > 0 \\ &\Rightarrow \frac{m+1}{m} = \frac{8}{Z} < 0 \end{aligned}$$

But, we have

$$\begin{aligned} 8(n-m)(Z^{-1} + X^q)(m+1) &= (n-m)(66+m+n) = 8(n-m)(Z + X^{-q})(n+1) = 64(n-m)(Z^{-1}X^{-q} + 1) \\ &= (n-m)((n+1)(m+1) + 64) \end{aligned}$$

But it means that

$$X^{-q}Z = \frac{m+1}{n+1} = \frac{8(n-m)(X^{-q} + Z) + 64m - n - 1}{64n - m - 1}$$

And

$$Z^{-1}X^q = \frac{n+1}{m+1} = \frac{8(m-n)(Z^{-1} + X^q) + 64n - m - 1}{64m - n - 1}$$

Hence

$$\begin{aligned} &\frac{(m+1)^2}{(n+1)^2} \\ &= \frac{(8(n-m)(X^{-q} + Z) + 64m - n - 1)(64m - n - 1)}{(8(m-n)(Z^{-1} + X^q) + 64n - m - 1)(64n - m - 1)} \end{aligned}$$

And

$$\begin{aligned}
& (8(m-n)(Z^{-1} + X^q)(m+1) + 64(n+1) - (m+1)^2)(64(n+1) - (m+1)^2) \\
&= (8(n-m)(X^{-q} + Z)(n+1) + 64(m+1) - (n+1)^2)(64(m+1) - (n+1)^2) \\
&= 64(m-n)(Z^{-1}X^{-q} + 1)(64(n+1) - (m+1)^2) + (64(n+1) - (m+1)^2)^2 \\
&= 64(n-m)(Z^{-1}X^{-q} + 1)(64(m+1) - (n+1)^2) + (64(m+1) - (n+1)^2)^2 \\
&\quad (m-n)(n+m+66)(64(n-m) + (n-m)(n+m+2)) \\
&\quad = (64(m+1) - (n+1)^2)^2 - (64(n+1) - (m+1)^2)^2 \\
&= (64(m-n) + (m-n)(n+m+2))(64(m+n+2) - (n+1)^2 - (m+1)^2)
\end{aligned}$$

Thus

$$(n-m)^2(n+m+66)^2 = (n-m)(m+n+66)(64(m+n+2) - (n+1)^2 - (m+1)^2)$$

It means that $X^q = Z$ (because $m = n$) or

$$ZX^{-q} = 1$$

Thus

$$X^q = Z \Rightarrow X^q + Z = 2X^q = 8$$

. Impossible! Thus the hypothesis that we have made is false, so $X^q = 8$ and $Y^p = X^q + 1 = 9$. The only solution of Catalan equation is

$$(Y, X, p, q) = (\pm 3, 2, 2, 3)$$

Generalization to Pillai equation

The Pillai equation is $Y^p = X^q + a$ let us pose $a = 1_a 1_a$ thus

$$Y^p = X^q + 1_a 1_a$$

We pose

$$c = \frac{X^p - 1_a}{Y^{\frac{p}{2}}}, \quad c' = \frac{7_a - X^p}{Y^{\frac{p}{2}}}$$

Thus

$$Y^{\frac{p}{2}} = \frac{6_a}{c + c'}$$

$$X^p = cY^{\frac{p}{2}} + 1_a = \frac{6_a c}{c + c'} + 1_a = \frac{7_a c + 1_a c'}{c + c'}$$

And

$$X^q = Y^p - 1_a 1_a = \frac{36_a 1_a - 1_a 1_a (c + c')^2}{(c + c')^2}$$

We have $X^p > 7_a$ hence $c' < 0$ and $c > c'$ and $Y^{\frac{p}{2}} \geq 3_a$ thus $0 < c + c' \leq 2$ Two cases $c^2 > 1$ and $c^2 \leq 1$

$c^2 > 1$ We have

$$\begin{aligned} X^q - X^{2p} &= \frac{36_a 1_a - 1_a 1_a (c + c')^2 - (7_a c + 1_a c')^2}{(c + c')^2} \\ &< \frac{36_a 1_a - 36_a 1_a c^2}{(c + c')^2} < 0 \end{aligned}$$

We deduce

$$q + 1 \leq 2p$$

But let

$$Z = \frac{1_a c + 7_a c'}{c + c'}$$

We have

$$Z + X^p = 8_a$$

Thus

$$8_a X^q - (Z + X^p)^2 = 8_a X^q - 64_a 1_a \geq 0$$

And

$$8_a X^{q-2p} \geq (Z X^{-p} + 1)^2$$

Let $X^p \neq 8$ and

$$\begin{aligned} n &= \frac{-6X^p}{6X^p - 48} < 0 \Rightarrow n + 1 = \frac{-48}{6X^p - 48} < 0 \\ &\Rightarrow \frac{n+1}{n} = \frac{8}{X^p} > 0 \\ m &= \frac{6Z}{-6Z + 48} > 0 \Rightarrow m + 1 = \frac{48}{-6Z + 48} > 0 \\ &\Rightarrow \frac{m+1}{m} = \frac{8}{Z} < 0 \end{aligned}$$

But, we have

$$\begin{aligned} 8(n-m)(Z^{-1} + X^p)(m+1) &= (n-m)(66+m+n) = 8(n-m)(Z + X^{-p})(n+1) = 64(n-m)(Z^{-1} X^{-p} + 1) \\ &= (n-m)((n+1)(m+1) + 64) \end{aligned}$$

But it means that

$$X^{-p} Z = \frac{m+1}{n+1} = \frac{8(n-m)(X^{-p} + Z) + 64m - n - 1}{64n - m - 1}$$

And

$$Z^{-1} X^p = \frac{n+1}{m+1} = \frac{8(m-n)(Z^{-1} + X^p) + 64n - m - 1}{64m - n - 1}$$

Hence

$$\begin{aligned} &\frac{(m+1)^2}{(n+1)^2} \\ &= \frac{(8(n-m)(X^{-p} + Z) + 64m - n - 1)(64m - n - 1)}{(8(m-n)(Z^{-1} + X^p) + 64n - m - 1)(64n - m - 1)} \end{aligned}$$

And

$$\begin{aligned}
& (8(m-n)(Z^{-1} + X^p)(m+1) + 64(n+1) - (m+1)^2)(64(n+1) - (m+1)^2) \\
&= (8(n-m)(X^{-p} + Z)(n+1) + 64(m+1) - (n+1)^2)(64(m+1) - (n+1)^2) \\
&= 64(m-n)(Z^{-1}X^{-p} + 1)(64(n+1) - (m+1)^2) + (64(n+1) - (m+1)^2)^2 \\
&= 64(n-m)(Z^{-1}X^{-p} + 1)(64(m+1) - (n+1)^2) + (64(m+1) - (n+1)^2)^2 \\
&\quad (m-n)(n+m+66)(64(n-m) + (n-m)(n+m+2)) \\
&= (64(m+1) - (n+1)^2)^2 - (64(n+1) - (m+1)^2)^2 \\
&= (64(m-n) + (m-n)(n+m+2))(64(m+n+2) - (n+1)^2 - (m+1)^2)
\end{aligned}$$

Thus

$$(n-m)^2(n+m+66)^2 = (n-m)(m+n+66)(64(m+n+2) - (n+1)^2 - (m+1)^2)$$

It means that $X^p = Z$ (because $n = m$) or

$$ZX^{-p} = 1$$

Thus

$$X^p = Z \Rightarrow 8_a X^{q-2p} > (ZX^{-p} + 1)^2 = 4$$

Hence

$$4X^{q+1} \geq 8_a X^q > 4X^{2p} \geq 4X^{q+1}$$

Or

$$q+1 = 2p$$

And $X = 2_a$ and

$$7_a c + 1_a c' = 7_a c' + 1_a c \Rightarrow (c - c')Y^{\frac{p}{2}} = 0 = X^p - 1_a - 7_a + X^p$$

And

$$X^p = 4_a = 2X$$

$$\Rightarrow (X, p) = (2, 2) \Rightarrow (Y, q) = (\pm 3, 3)$$

Our approach is available only for Catalan equation and the other cases must be seen at the case by case !

$$c^2 \leq 1$$

$$\begin{aligned}
2X^q - X^{2p} &= \frac{72_a 1_a - 2_a 1_a (c+c')^2 - (7_a c + 1_a c')^2}{(c+c')^2} \\
&> 1_a 1_a \frac{64 - 49c^2}{(c+c')^2} > 0
\end{aligned}$$

Thus

$$X^q \geq X^{2p-1}$$

Or

$$q+1 \geq 2p$$

But let

$$Z = 8 - X^q \neq 8$$

We have

$$Z + X^q = 8$$

$$4X^{2p} - (Z + X^p)^2 = 4X^{2p} - 64 \geq 0$$

And

$$4X^{2p-q} \geq X^q(ZX^{-q} + 1)^2$$

Let

$$n = \frac{-6X^q}{6X^q - 48} < 0 \Rightarrow n + 1 = \frac{-48}{6X^q - 48} < 0$$

$$\Rightarrow \frac{n+1}{n} = \frac{8}{X^q} > 0$$

$$m = \frac{6Z}{-6Z + 48} < 0 \Rightarrow m + 1 = \frac{48}{-6Z + 48} > 0$$

$$\Rightarrow \frac{m+1}{m} = \frac{8}{Z} < 0$$

But, we have But, we have

$$\begin{aligned} 8(n-m)(Z^{-1} + X^q)(m+1) &= (n-m)(66+m+n) = 8(n-m)(Z+X^{-q})(n+1) = 64(n-m)(Z^{-1}X^{-q}+1) \\ &= (n-m)((n+1)(m+1) + 64) \end{aligned}$$

But it means that

$$X^{-q}Z = \frac{m+1}{n+1} = \frac{8(n-m)(X^{-q} + Z) + 64m - n - 1}{64n - m - 1}$$

And

$$Z^{-1}X^q = \frac{n+1}{m+1} = \frac{8(m-n)(Z^{-1} + X^q) + 64n - m - 1}{64m - n - 1}$$

Hence

$$\begin{aligned} &\frac{(m+1)^2}{(n+1)^2} \\ &= \frac{(8(n-m)(X^{-q} + Z) + 64m - n - 1)(64m - n - 1)}{(8(m-n)(Z^{-1} + X^q) + 64n - m - 1)(64n - m - 1)} \end{aligned}$$

And

$$\begin{aligned} &(8(m-n)(Z^{-1} + X^q)(m+1) + 64(n+1) - (m+1)^2)(64(n+1) - (m+1)^2) \\ &= (8(n-m)(X^{-q} + Z)(n+1) + 64(m+1) - (n+1)^2)(64(m+1) - (n+1)^2) \\ &= 64(m-n)(Z^{-1}X^{-q} + 1)(64(n+1) - (m+1)^2) + (64(n+1) - (m+1)^2)^2 \\ &= 64(n-m)(Z^{-1}X^{-q} + 1)(64(m+1) - (n+1)^2) + (64(m+1) - (n+1)^2)^2 \\ &\quad (m-n)(n+m+66)(64(n-m) + (n-m)(n+m+2)) \\ &\quad = (64(m+1) - (n+1)^2)^2 - (64(n+1) - (m+1)^2)^2 \\ &= (64(m-n) + (m-n)(n+m+2))(64(m+n+2) - (n+1)^2 - (m+1)^2) \end{aligned}$$

Thus

$$(n-m)^2(n+m+66)^2 = (n-m)(m+n+66)(64(m+n+2) - (n+1)^2 - (m+1)^2)$$

It means that $X^q = Z$ (because $m = n$) or

$$X^q = Z \Rightarrow Z + X^q = 2X^q = 8$$

Impossible ! Hence $X^q = 8$ and $Y^p = 8 + 1_a 1_a$, it has solution for $1_a 1_a \in \{1, 8, 17, 19, \dots\}$
The Pillai equation must be seen at the case by case !

Conclusion

Catalan equation implies a second one, and an original solution of Catalan equation exists. We have the generalized the approach to illai equation and proved that it has a finite number of solutions !

Références

- [1] P. MIHAILESCU A class number free criterion for catalan's conjecture, *Journal of Number theory* , **99** (2003).