



**HAL**  
open science

## A proof of Pillai conjecture

Jamel Ghannouchi

► **To cite this version:**

| Jamel Ghannouchi. A proof of Pillai conjecture. 2012. hal-00698687v14

**HAL Id: hal-00698687**

**<https://hal.science/hal-00698687v14>**

Preprint submitted on 24 Jul 2012 (v14), last revised 28 Mar 2013 (v22)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A proof of Pillai conjecture

## Jamel Ghanouchi

jamel.ghanouchi@topnet.tn

### Abstract

(MSC=11D04) More than one century after its formulation by the Belgian mathematician Eugene Catalan, Preda Mihailescu has solved the open problem. But, is it all? Mihailescu's solution utilizes computation on machines, we propose here not really a proof as it is intended classically, but a resolution of an equation like the resolution of the polynomial equations of third and fourth degrees. This solution is totally algebraic and does not utilize, of course, computers or any kind of calculation. Then, we generalize the proof to Pillai conjecture.

(Keywords : Diophantine equations ; Catalan equation ; Pillai conjecture ; Algebraic resolution)

### Introduction

Catalan theorem has been proved in 2002 by Preda Mihailescu. In 2004, it became officially Catalan-Mihailescu theorem. This theorem stipulates that there are not consecutive pure powers. There do not exist integers strictly greater than 1,  $X > 1$  and  $Y > 1$ , for which with exponents strictly greater than 1,  $p > 1$  and  $q > 1$ ,

$$Y^p = X^q + 1$$

but for  $(X, Y, p, q) = (2, 3, 2, 3)$ . We can verify that

$$3^2 = 2^3 + 1$$

Euler has proved that the equation  $X^3 + 1 = Y^2$  has this only solution. We propose in this study a general solution. The particular cases already solved concern  $p = 2$ , solved by Ko Chao in 1965, and  $q = 3$  which has been solved in 2002. The case  $q = 2$  has been solved by Lebesgue in 1850. We solve here the equation for the general case. We generalize the proof to Pillai conjecture

$$Y^p = X^q + a$$

And prove that it has always a finite number of solutions for a fixed  $a$ .

### The approach

Let

$$c = \frac{X^p - 1}{Y^{\frac{p}{2}}}, \quad c' = \frac{7 - X^p}{Y^{\frac{p}{2}}}$$

But

$$(c + c')Y^{\frac{p}{2}} = X^p - 1 + 7 - X^p = 6 \Rightarrow Y^{\frac{p}{2}} = \frac{6}{c + c'}$$

And

$$X^p = cY^{\frac{p}{2}} + 1 = \frac{6c}{c + c'} + 1 = \frac{7c + c'}{c + c'}$$

And

$$X^q = Y^p - 1 = \frac{36 - (c + c')^2}{(c + c')^2}$$

And

$$(c + c')Y^{\frac{p}{2}} = 6 > 0$$

Thus  $c + c' > 0$ . Also  $cY^{\frac{p}{2}} = X^p - 1 > 0$  Thus  $c > 0$ . But  $X^p \geq 4$ , hence

$$cY^{\frac{p}{2}} = X^p - 1 \geq 7 - X^p = c'Y^{\frac{p}{2}}$$

hence  $c \geq c'$  and

$$7 - X^p = 7 - \frac{7c + c'}{c + c'} = \frac{6c'}{c + c'}$$

If  $c' > 0$  Thus  $X^p < 7$  and  $X^p = 4$ , it means that  $c' < 0$  and

$$(c + c')Y^{\frac{p}{2}} = 6 > 0$$

$$(c + c')X^p = 7c + c' > 0$$

thus  $c + c' > 0$  and

$$Y^{\frac{p}{2}} = \frac{6}{c + c'} \geq 3 \Rightarrow c + c' \leq 2$$

**main results** What we must retain is

1)  $c' < 0$

2)  $0 < c + c' < 2$

3)  $0 < 7c + c'$

We will discuss two cases I)  $c^2 > 1$

II)  $c^2 \leq 1$

$c^2 > 1$  We have

$$X^q - X^{2p} = \frac{36 - (c + c')^2 - (7c + c')^2}{(c + c')^2} < \frac{(6 - 6c)(6 + 6c) - (c + c')^2}{(c + c')^2} < 0$$

Hence

$$q \leq 2p - 1$$

And

$$c^2 X^q - X^{2p} = \frac{36c^2 - c^2(c + c')^2 - (7c + c')^2}{(c + c')^2} = \frac{-(c + c')(13c + c') - c^2(c + c')^2}{(c + c')^2} < 0$$

And

$$\begin{aligned} 2c^2X^q - X^{2p} &= \frac{72c^2 - 2c^2(c+c')^2 - (7c+c')^2}{(c+c')^2} \\ &> \frac{64c^2 - (7c+c')^2}{(c+c')^2} = \frac{(c-c')(15c+c')}{(c+c')^2} > 0 \end{aligned}$$

But with  $uc = u'c' = 1$

$$X^p = \frac{7c+c'}{c+c'} = \frac{7u'+u}{u+u'}$$

Let

$$Z = \frac{7c'+c}{c'+c} = \frac{7u+u'}{u+u'}$$

We have

$$8X^q - (Z + X^p)^2 = \frac{288 - 8(c+c')^2 - 64(c+c')^2}{(c+c')^2} = \frac{288 - 72(c+c')^2}{(c+c')^2} > 0$$

Thus

$$8X^{q-2p} - (ZX^{-p} + 1)^2 > 0$$

But

$$\begin{aligned} X^{-2p}Z^2 &= \frac{(7c'+c)(7u+u')}{(7u'+u)(7c+c')} \\ &= \frac{49uc' + u'c + 14}{49u'c + uc' + 14} = \frac{49c'^2 + c^2 + 14cc'}{49c^2 + c'^2 + 14cc'} \end{aligned}$$

Let  $v$  verifying

$$\begin{aligned} vZ^2X^{-2p} - 1 &= \frac{49vc'^2 + vc^2 + 14vcc'}{49c^2 + c'^2 + 14cc'} - 1 \\ &= \frac{(49v-1)c'^2 + (v-49)c^2 + 14cc'(v-1)}{49c^2 + c'^2 + 14cc'} > 0 \end{aligned}$$

It means

$$\begin{aligned} &(49v-1)(X^p-7)^2 + (v-49)(X^p-1)^2 + 14(X^p-1)(7-X^p)(v-1) \\ &= (49v-1+v-49-14v+14)X^{2p} + (-14(49v-1)-2(v-49)+112(v-1))X^p + 49(49v-1)+(v-49)-98(v-1) \\ &= (36v-36)X^{2p} - 576vX^p + (49^2-97)v > 0 \end{aligned}$$

For  $v = 1 + \frac{1}{n} = \frac{X^p}{2X^p-16}$  or  $n = \frac{12X^p-96}{-6X^p+96}$  Thus

$$36X^{2p} + (n+1)(-576X^p + 2304) = 36X^{2p} + \left(\frac{-6X^p}{6X^p-96}\right)(-576X^p + 2304) > 0$$

We have then

$$v = \frac{n+1}{n} \Rightarrow vZ^2 > X^{2p}$$

Hence

$$\begin{aligned} Z^2 &> \frac{X^{2p}n}{n+1} \\ Z^2X^{-2p} + 2ZX^{-p} + 1 &> \frac{2n+1}{n+1} + \frac{14c'+2c}{7c+c'} \end{aligned}$$

$$= \frac{(16n+9)c + (16n+15)c'}{(n+1)(7c+c')} = 1$$

Because  $(9n+2)c + (15n+14)c' = 0$  and

$$(9n+2)(X^p-1) + (15n+14)(7-X^p) = (-6n-12)X^p + (96n+96) = 0$$

or

$$n(-6X^p+96) - (12X^p-96) = 0$$

Then

$$8X^{q-2p} > (ZX^{-p}+1)^2 > 1$$

Or

$$X^{q+1} \geq 8X^q > X^{2p}$$

And

$$q+1 \geq 2p \geq q+1 \Rightarrow q+1 = 2p$$

Let  $m$  and  $v$  verifying  $v = \frac{(13-10^{-m})c + (5-10^{-m-1})c'}{32(c^2-c'^2)}$  : we have

$$7c - c' = 32v(c+c')(c-c') > 4(c+c')$$

Because

$$7c - c' = (13 - 10^{-m})c + (5 - 10^{-m-1})c' > 4c + 4c'$$

Or

$$c(7 - 32v(c+c')) = (1 - 32v(c+c'))c'$$

And

$$(X^p-1)(7-32v(c+c')) = (32v(c+c')-1)(X^p-7)$$

$$0 < X^p(64v(c+c')-8) = 256v(c+c')-14$$

Because

$$\begin{aligned} 64v(c+c')-8 &= \frac{2((13-10^{-m})c + (5-10^{-m-1})c') - 8(c-c')}{c-c'} \\ &= \frac{(18-2(10^{-m}))c + (18-2(10^{-m-1}))c'}{c-c'} > \frac{6(c+c')}{c-c'} > 0 \end{aligned}$$

So

$$\begin{aligned} X^p &= \frac{128v(c+c')-7}{32v(c+c')-4} \\ &= \frac{(45-4(10^{-m}))c + (13-10^{-m-1})c'}{(13-10^{-m})c + (5-10^{-m-1})c'} < 16 \end{aligned}$$

Because

$$\begin{aligned} &(163-12(10^{-m}))c + (67-15(10^{-m-1}))c' \\ &= -\left(\frac{815-163(10^{-m-1})}{6}\right)c' + \frac{163(10^{-m})}{6}c + (67-15(10^{-m-1}))c' \\ &= \frac{-413+155(10^{-m})}{6}c' + \frac{163(10^{-m})}{6}c > 0 \end{aligned}$$

Thus  $X^p \leq 9$  And it means

$$X^p = 4 \Rightarrow (X, p) = (2, 2) \Rightarrow Y^2 = 2^3 + 1 = 9 \Rightarrow (Y, q) = (\pm 3, 3)$$

Another proof : let

$$X^{2p-q} = \frac{n+1}{n}$$

Thus

$$X^{q+2-2p} = \frac{nX^2}{n+1} > \frac{X}{u} \geq 1$$

We have chosen  $u$  such as  $n(uX - 1) > 1$ . Thus

$$0 \geq q + 1 - 2p \geq 0$$

And  $q + 1 = 2p$

$c^2 \leq 1$  We have

$$2X^q - X^{2p} = \frac{72 - 2(c+c')^2 - (7c+c')^2}{(c+c')^2} > \frac{64 - 49c^2}{(c+c')^2} > 0$$

Thus

$$q \geq 2p - 1$$

And

$$c^2 X^q - X^{2p} = \frac{36c^2 - c^2(c+c')^2 - (7c+c')^2}{(c+c')^2} = \frac{-(c+c')(13c+c') - c^2(c+c')^2}{(c+c')^2} < 0$$

Or

$$\begin{aligned} 2c^2 X^q - X^{2p} &= \frac{72c^2 - 2c^2(c+c')^2 - (7c+c')^2}{(c+c')^2} \\ &> \frac{64c^2 - (7c+c')^2}{(c+c')^2} = \frac{(c-c')(15c+c')}{(c+c')^2} > 0 \end{aligned}$$

Let

$$Z' = \frac{9(c+c')^2 - 36}{(c+c')^2} = 9 - Y^p$$

We have

$$Z' + X^q = 8$$

And

$$16X^{2p} - (Z' + X^q)^2 = 16X^{2p} - 64 > 0$$

Hence

$$16X^{2p-q} > (Z'X^{-q} + 1)^2$$

Let  $v = \frac{Y^p-1}{Y^p-9} = \frac{n+1}{n}$  with  $n = \frac{Y^p-9}{8}$  thus

$$Z'^2 X^{-2q} = \frac{(Y^p-9)^2}{(Y^p-1)^2} = \frac{n^2}{(n+1)^2}$$

And

$$\begin{aligned}
16X^{2p-q} &> X^q \left( \frac{2n^2 + 2n + 1}{n^2 + 2n + 1} + \frac{18(c + c')^2 - 72}{36 - (c + c')^2} \right) \\
&= X^q \left( \frac{2n^2 + 2n + 1}{n^2 + 2n + 1} + \frac{18 - 2Y^p}{Y^p - 1} \right) \\
&= X^q \left( \frac{-2nY^p - Y^p + 16n^2 + 34n + 17}{Y^p - 1} \right) = X^q > 1
\end{aligned}$$

Because

$$16n^2 + 34n + 18 = 2(8n^2 + 17n + 9) = 2Y^p + 2nY^p = 2Y^p(n + 1)$$

And

$$\frac{8n^2 + 17n + 9}{n + 1} = 8n + 9 = Y^p$$

We deduce

$$\begin{aligned}
16 &> X^{q-2p} \\
1 &\geq 16X > X^{q-1-2p}
\end{aligned}$$

And as  $q$  is odd

$$q + 1 \geq 2p \geq q + 1 \Rightarrow q + 1 = 2p$$

And

$$\begin{aligned}
0 > (c^2 - 1)Y^p &= X^{2p} - 2X^p - X^q = X^{2p} - 2X^p - X^{2p-1} > 0 \\
X^{p-1} - X^{q-p-1} &= \frac{2}{X} \in \mathbb{N} \Rightarrow X = 2 \\
2^{p-1} = 2^{q-p-1} + 1 &\Rightarrow q = p + 1 = 3 \Rightarrow Y = \pm 3
\end{aligned}$$

Another proof : let

$$X^{q-2p} = \frac{n+1}{n}$$

Thus

$$X^{2p+2-q} = \frac{nX^2}{n+1} > \frac{X}{u} \geq 1$$

We have chosen  $u$  such as  $n(uX - 1) > 1$ . Thus

$$0 \geq q + 1 - 2p \geq 0$$

And  $q+1 = 2p$  Thus we have Catalan solution !

### Generalization to Pillai equation

Pillai equation is  $Y^p = X^q + a$ . We will define Pillai numbers : we note them :  $b_a$  they depend of  $a$  and have the same proprieties than  $b$  for  $a = 1$ . Thus, we write  $1_a + 7_a = 8_a$  and  $7_a - 1_a = 6_a$  : our goal is to follow the approach as we have done for Catalan equation. Of course, we have  $1_a = a_1$  et  $0_a = 0$ . The equation becomes  $Y^p = X^q + 1_a$ . Let us pose for  $p, q, X, Y, a$  integers stricly greater to 1. Let

$$c = \frac{X^p - 1_a}{Y^{\frac{p}{2}}}, \quad c' = \frac{7_a - X^p}{Y^{\frac{p}{2}}}$$

And

$$(c + c')Y^{\frac{p}{2}} = X^p - 1_a + 7_a - X^p = 6_a \Rightarrow Y^{\frac{p}{2}} = \frac{6_a}{c + c'}$$

So

$$X^p = cY^{\frac{p}{2}} + 1_a = \frac{6_ac}{c + c'} + 1_a = \frac{7_ac + 1_ac'}{c + c'}$$

And

$$X^q = Y^p - 1_a = \frac{36_a 1_a - 1_a(c + c')^2}{(c + c')^2}$$

Also

$$(c + c')Y^{\frac{p}{2}} = 6_a > 0$$

Thus  $c + c' > 0$ . And

$$cY^{\frac{p}{2}} = X^p - 1_a > 0$$

So  $c > 0$ . But  $X^p \geq 4_a$ , then

$$cY^{\frac{p}{2}} = X^p - 1_a \geq 7_a - X^p = c'Y^{\frac{p}{2}}$$

And  $c \geq c'$  and

$$7_a - X^p = 7_a - \frac{7_ac + 1_ac'}{c + c'} = \frac{6_ac'}{c + c'}$$

If  $c' > 0$  Then  $X^p < 7_a$  and we have a finite number of solutions, we have  $c' < 0$ , with

$$(c + c')Y^{\frac{p}{2}} = 6_a > 0$$

and  $(c + c')X^p = 7_ac + 1_ac' > 0$  with  $c + c' > 0$ . And

$$Y^{\frac{p}{2}} = \frac{6_a}{c + c'} \geq 3_a \Rightarrow c + c' \leq 2 < 2_a$$

**Main results** We retain

1)  $c' < 0$

2)  $0 < c + c' < 2_a$

3)  $0 < 7_ac + 1_ac'$

Two cases : I)  $c^2 > 1_a$

II)  $c^2 \leq 1_a$

$c^2 > 1_a$  We have

$$1_a X^q - X^{2p} = \frac{36_a 1_a 1_a - 1_a(c + c')^2 - (7_ac + 1_ac')^2}{(c + c')^2} < \frac{36_a 1_a(1_a - c^2) - 1_a(c + c')^2}{(c + c')^2} < 0$$

Hence  $X^{q-2p} < \frac{1}{1_a} < 1$  and

$$q \leq 2p - 1$$

And

$$c^2 X^q - X^{2p} = \frac{36_a 1_a c^2 - c^2(c + c')^2 - (7_ac + 1_ac')^2}{(c + c')^2} = \frac{-1_a 1_a(c + c')(13c + c') - c^2(c + c')^2}{(c + c')^2} < 0$$



And

$$2c^2X^q - X^{2p} = \frac{72_a1_ac^2 - 2c^2(c+c')^2 - (7_ac + 1_ac')^2}{(c+c')^2}$$

$$> \frac{64_a1_ac^2 - (7_ac + 1_ac')^2}{(c+c')^2} = \frac{1_a1_a(c-c')(15c+c')}{(c+c')^2} > 0$$

Thus

$$2c^2 > X^{2p-q} \geq X$$

Thus  $2p \geq q + 1$  But with  $uc = u'c' = 1$ , we have

$$Z = \frac{7_ac' + 1_ac}{c+c'} = \frac{7_a u + 1_a u'}{u+u'}$$

$$X^p = \frac{7_ac + 1_ac'}{c+c'} = \frac{7_a u' + 1_a u}{u+u'}$$

$$8X^q - (Z+X^p)^2 = \frac{288_a1_a - 8(c+c')^2 - 64_a1_a(c+c')^2}{(c+c')^2} > \frac{288_a1_a - 72_a1_a(c+c')^2}{(c+c')^2} > 0$$

Thus

$$8X^{q-2p} - (ZX^{-p} + 1)^2 > 0$$

But

$$X^{-2p}Z^2 = \frac{(7_ac' + 1_ac)(7_a u + 1_a u')}{(7_a u' + 1_a u)(7_ac + 1_ac')}$$

$$= \frac{49_a u c' + 1_a u' c + 14_a}{49_a u' c + 1_a u c' + 14_a} = \frac{49_a c'^2 + 1_a c^2 + 14_a c c'}{49_a c^2 + 1_a c'^2 + 14_a c c'}$$

Let  $v$  verifying

$$vZ^2X^{-2p} - 1 = \frac{49_a v c'^2 + 1_a v c^2 + 14_a v c c'}{49_a c^2 + 1_a c'^2 + 14_a c c'} - 1$$

$$= \frac{(49_a v - 1_a)c'^2 + (1_a v - 49_a)c^2 + 14_a c c'(v - 1)}{49_a c^2 + 1_a c'^2 + 14_a c c'} > 0$$

It means

$$(49_a v - 1_a)(X^p - 7_a)^2 + 1_a(v - 49_a)(X^p - 1_a)^2 + 14_a(X^p - 1_a)(7_a - X^p)(v - 1)$$

$$= (36_a v - 36_a)X^{2p} - 576_a v X^p + 1_a 1_a(49^2 - 97)v > 0$$

For  $v = 1 + \frac{1}{n} = \frac{X^p}{2X^p - 16_a}$  or  $n = \frac{12X^p - 96_a}{-6X^p + 96_a}$  Thus

$$36_a X^{2p} + (n+1)(-576_a X^p + 2304_a 1_a) = 36_a X^{2p} + \left(\frac{-6X^p}{6X^p - 96_a}\right)(-576_a X^p + 2304_a 1_a) > 0$$

We have then

$$v = \frac{n+1}{n} \Rightarrow vZ^2 > X^{2p}$$

Hence

$$Z^2 > \frac{X^{2p}n}{n+1}$$

$$\begin{aligned} Z^2 X^{-2p} + 2ZX^{-p} + 1 &> \frac{2n+1}{n+1} + \frac{14c' + 2c}{7c + c'} \\ &= \frac{(16n+9)c + (16n+15)c'}{(n+1)(7c+c')} = 1 \end{aligned}$$

Because  $(9n+2)c + (15n+14)c' = 0$  and

$$(9n+2)(X^p - 1_a) + (15n+14)(7_a - X^p) = (-6n-12)X^p + 1_a(96n+96) = 0$$

or

$$n(-6X^p + 96_a) - (12X^p - 96_a) = 0$$

Then

$$8X^{q-2p} > (ZX^{-p} + 1)^2 > 1$$

Or

$$X^{q+1} \geq 8X^q > X^{2p}$$

And

$$q+1 \geq 2p \geq q+1 \Rightarrow q+1 = 2p$$

Let  $m$  and  $v$  verifying  $v = \frac{(13_a - 10^{-m})c + (5_a - 10^{-m-1})c'}{32(c^2 - c'^2)}$  : we have

$$7_a c - 1_a c' = 32v(c+c')(c-c') > 4_a(c+c')$$

Because

$$7_a c - 1_a c' = (13_a - 10^{-m})c + (5_a - 10^{-m-1})c' > 4c + 4c'$$

Or

$$c(7_a - 32v(c+c')) = (1_a - 32v(c+c'))c'$$

And

$$\begin{aligned} (X^p - 1_a)(7_a - 32v(c+c')) &= (32v(c+c') - 1_a)(X^p - 7_a) \\ 0 < X^p(64v(c+c') - 8_a) &= 256v(c+c') - 14_a \end{aligned}$$

Because

$$\begin{aligned} 64v(c+c') - 8_a &= \frac{2((13_a - 10^{-m})c + (5_a - 10^{-m-1})c') - 8_a(c-c')}{c-c'} \\ &= \frac{(18_a - 2(10^{-m}))c + (18_a - 2(10^{-m-1}))c'}{c-c'} > \frac{6_a(c+c')}{c-c'} > 0 \end{aligned}$$

So

$$\begin{aligned} X^p &= \frac{128v(c+c') - 7_a}{32v(c+c') - 4_a} \\ &= \frac{(45_a - 4_a(10^{-m}))c + (13_a - 1_a10^{-m-1})c'}{(13_a - 1_a10^{-m})c + (5_a - 1_a10^{-m-1})c'} < 16 \end{aligned}$$

Because

$$\begin{aligned} &(163_a - 12_a(10^{-m}))c + (67_a - 15_a(10^{-m-1}))c' \\ &= -\left(\frac{815 - 163(10^{-m-1})}{6}\right)c' + \frac{163(10^{-m})}{6}c + (67 - 15(10^{-m-1}))c' \end{aligned}$$

$$= \frac{-413 + 155(10^{-m})}{6}c' + \frac{163(10^{-m})}{6}c > 0$$

and there is a finite number of solutions, another proof : let

$$X^{2p-q} = \frac{n+1}{n}$$

Thus

$$X^{q+2-2p} = \frac{nX^2}{n+1} > \frac{X}{u} \geq 1$$

we have chosen  $u$  such as  $n(uX - 1) > 1$ . Thus

$$0 \geq q + 1 - 2p \geq 0$$

And  $q + 1 = 2p$  : There is a finite number of solutions!

$c^2 \leq 1$  We have

$$2_a X^q - X^{2p} = \frac{72_a 1_a 1_a - 2_a(c+c')^2 - (7_a c + 1_a c')^2}{(c+c')^2} > \frac{64_a 1_a 1_a - 49_a 1_a c^2}{(c+c')^2} > 0$$

Thus

$$q \geq 2p - 1$$

And

$$c^2 X^q - X^{2p} = \frac{36_a 1_a c^2 - c^2(c+c')^2 - (7_a c + 1_a c')^2}{(c+c')^2} = \frac{-1_a 1_a (c+c')(13c+c') - c^2(c+c')^2}{(c+c')^2} < 0$$

And

$$\begin{aligned} 2c^2 X^q - X^{2p} &= \frac{72_a 1_a c^2 - 2c^2(c+c')^2 - (7_a c + 1_a c')^2}{(c+c')^2} \\ &> \frac{64_a 1_a c^2 - (7_a c + 1_a c')^2}{(c+c')^2} = \frac{1_a 1_a (c-c')(15c+c')}{(c+c')^2} > 0 \end{aligned}$$

Let

$$Z' = \frac{9(c+c')^2 - 36_a 1_a}{(c+c')^2} = 9 - Y^p$$

We have

$$Z' + X^q = 8$$

And

$$16X^{2p} - 1_a 1_a (Z' + X^q)^2 = 16X^{2p} - 1_a 1_a 64 > 0$$

Hence

$$16X^{2p-q} > 1_a 1_a X^q (Z' X^{-q} + 1)^2$$

Let  $v = \frac{Y^p-1}{Y^p-9} = \frac{n+1}{n}$  with  $n = \frac{Y^p-9}{8}$  thus

$$Z'^2 X^{-2q} = \frac{(Y^p-9)^2}{(Y^p-1)^2} = \frac{n^2}{(n+1)^2}$$

And

$$Z^{p^2} X^{-2q} = \frac{1}{v^2} = \frac{Y^p - 1}{Y^p - 9} = \frac{n}{n+1}$$

Or

$$\begin{aligned} 16X^{2p-q} &> 1_a 1_a X^q \left( \frac{2n^2 + 2n + 1}{n^2 + 2n + 1} + \frac{18(c+c')^2 - 72}{36 - (c+c')^2} \right) \\ &= 1_a 1_a X^q \left( \frac{2n^2 + 2n + 1}{n^2 + 2n + 1} + \frac{18 - 2Y^p}{Y^p - 1} \right) \\ &= 1_a 1_a X^q \left( \frac{-2nY^p - Y^p + 16n^2 + 34n + 17}{Y^p - 1} \right) = 1_a 1_a X^q > 1 \end{aligned}$$

Because

$$16n^2 + 34n + 18 = 2(8n^2 + 17n + 9) = 2Y^p + 2nY^p = 2Y^p(n+1)$$

And

$$\frac{8n^2 + 17n + 9}{n+1} = 8n + 9 = Y^p$$

We deduce

$$\begin{aligned} 16 &> 1_a 1_a X^{q-2p} \\ 1 &\geq \frac{16}{1_a 1_a X} > X^{q-1-2p} \end{aligned}$$

And as  $q$  is odd

$$q+1 \geq 2p \geq q+1 \Rightarrow q+1 = 2p$$

Another proof : let

$$X^{q-2p} = \frac{n+1}{n}$$

Thus

$$X^{2p+2-q} = \frac{nX^2}{n+1} > \frac{X}{u} \geq 1$$

we have chosen  $u$  such as  $n(uX - 1) > 1$ . Thus

$$0 \geq q+1 - 2p \geq 0$$

And  $q+1 = 2p$  : but

$$0 > (c^2 - 1_a)Y^p = X^{2p} - 2_a X^p - 1_a X^q = X^{2p} - 2_a X^p - 1_a X^{2p-1} > 0$$

Hence  $c^2 = 1_a$  and

$$X^{p-1} - 1_a X^{q-p-1} = \frac{2}{X} \in \mathbb{N} \Rightarrow X \leq 2_a$$

It means that there is a finite number of solutions !

### Conclusion

Catalan equation is solved, an original solution exists ! We have generalized the approach to Pillai equation and proved that it always has a finite number of solutions. It is the proof of Pillai conjecture. It seems that many problems of number theory can be solved like this.

## Références

- [1] P. MIHAILESCU A class number free criterion for catalan's conjecture, *Journal of Number theory* , **99** (2003).