



**HAL**  
open science

# Automorphisms of Drinfeld half-spaces over a finite field

Bertrand Rémy, Amaury Thuillier, Annette Werner

► **To cite this version:**

Bertrand Rémy, Amaury Thuillier, Annette Werner. Automorphisms of Drinfeld half-spaces over a finite field. 2012. hal-00703402v2

**HAL Id: hal-00703402**

**<https://hal.science/hal-00703402v2>**

Preprint submitted on 21 Nov 2012 (v2), last revised 27 Nov 2012 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Automorphisms of Drinfeld half-spaces over a finite field

Bertrand Rémy, Amaury Thuillier and Annette Werner

## ABSTRACT

We show that the automorphism group of Drinfeld's half-space over a finite field is the projective linear group of the underlying vector space. The proof of this result uses analytic geometry in the sense of Berkovich over the finite field equipped with the trivial valuation. We also take into account extensions of the base field.

## Introduction

In this note we determine the automorphism group of Drinfeld's half-spaces over a finite field. Given a finite-dimensional vector space  $V$  over a finite field  $k$ , the Drinfeld half-space  $\Omega(V)$  is defined as the complement of all  $k$ -rational hyperplanes in the projective space  $\mathbf{P}(V)$ ; it is an affine algebraic variety over  $k$ . We show that every  $k$ -automorphism of  $\Omega(V)$  is induced by a  $k$ -automorphism of  $\mathbf{P}(V)$ . Hence the automorphism group of  $\Omega(V)$  is equal to  $\mathrm{PGL}(V)$ .

More generally, for an arbitrary field extension  $K$  of  $k$ , we prove that the natural injection of  $\mathrm{PGL}(V)$  into  $\mathrm{Aut}_K(\Omega(V) \otimes_k K)$  is an isomorphism. Our result answers a question of Dat, Orlik and Rapoport [DOR10, p. 338] which was motivated by the analogous statement for Drinfeld half-spaces over a non-Archimedean local field (with non-trivial absolute value).

Drinfeld defined his  $p$ -adic upper half-spaces in [Dri74]. They are the founding examples of the theory of period domains [RZ96]. Analogs of period domains over finite fields have been studied by Rapoport in [Rap97]. They are open subvarieties of flag varieties characterized by a semi-stability condition. Recently, they have been studied by Rapoport, Orlik and others, see e.g. [Orl01], [OR08]. A good introduction is given in the book [DOR10].

Over local non-Archimedean fields with non-trivial absolute value, Drinfeld half-spaces are no longer algebraic varieties and must be defined in the context of analytic geometry. In this setting, it was shown by Berkovich that every automorphism is induced by a projective linear transformation [Ber95]. This was generalized to products of Drinfeld half-spaces by Alon [Alon06], who also pointed out and corrected a discrepancy in Berkovich's proof. Berkovich's strategy is based on the fact that in the case of a local non-Archimedean ground field with non-trivial absolute value, the Bruhat-Tits building of the group  $\mathrm{PGL}(V)$  is contained in  $\Omega(V)$  as the subset of points satisfying a natural maximality condition. This implies that every automorphism of  $\Omega(V)$  induces an automorphism of the Bruhat-Tits building, and with some further work (see [Alon06]) one can show the claim.

One could in fact use a similar strategy in order to determine the automorphism group of  $\Omega(V)$  over a finite field. Indeed, if we endow the finite ground field with the trivial absolute value

and look at the corresponding Berkovich analytic space  $\Omega(V)^{\text{an}}$ , by [Ber90], the *vectorial building* associated to the group  $\text{PGL}(V)$  is contained in  $\Omega(V)^{\text{an}}$ . We believe that one can then follow Berkovich's and Alon's arguments to deduce that this automorphism comes from an element of  $\text{PGL}(V)$ .

However, in this note, we adopt a slightly different, and maybe more natural, viewpoint. Thereby, we want to highlight that the true content of this theorem is about *extension* of automorphisms, and that it has in fact very little to do with buildings, see Remark 2.3. Our approach is the following. We consider the space  $X$  obtained by blowing up all  $k$ -rational linear subspaces of the projective space  $\mathbf{P}(V)$ . Irreducible components of the boundary divisor correspond bijectively to linear subspaces of  $\mathbf{P}(V)$ . Moreover, a family of components has non-empty intersection if and only if the corresponding linear subspaces form a flag. We use Berkovich analytic geometry to prove in Proposition 2.1 that every automorphism of  $\Omega(V)$  preserves the set of discrete valuations on the function field induced by boundary components of  $X$ . Hence by Proposition 1.4 it extends to an automorphism of  $X$ . By taking a closer look at the Chow ring of  $X$  in section 3, we deduce that this automorphism preserves the set of discrete valuations corresponding to hyperplanes, which allows us to conclude that it induces an automorphism of the projective space.

#### ACKNOWLEDGEMENTS

We thank Vladimir Berkovich for acquainting us with the results in [Alon06]. We also thank Carlo Gasbarri for suggesting a more straightforward argument in the last step of the proof, and Stéphane Lamy for an interesting discussion on birational geometry. We are grateful to the referee for a number of helpful corrections and suggestions how to make the paper more accessible.

### 1. Automorphisms of Drinfeld's half-spaces

Let  $k$  be a finite field and let  $V$  be a  $k$ -vector space. We denote by  $\mathbf{P}(V)$  the projective scheme  $\text{Proj}(\text{Sym}^\bullet V)$  and define the  $k$ -scheme  $\Omega(V)$  as the complement of all (rational) hyperplanes in  $\mathbf{P}(V)$ :

$$\Omega(V) = \mathbf{P}(V) - \bigcup_{\substack{W \subset V \\ \dim W = 1}} \mathbf{P}(V/W).$$

For every field extension  $K/k$  we denote by  $V_K = V \otimes_k K$  the induced vector space over  $K$ . Then the base change  $\Omega(V)_K = \Omega(V) \otimes_k K$  is the complement of all  $k$ -rational hyperplanes in  $\mathbf{P}(V_K) = \mathbf{P}(V) \otimes_k K$ .

The main result of this note is the following.

**THEOREM 1.1.** *Let  $V$  be a vector space of finite dimension over a finite field  $k$ .*

(i) *The restriction map*

$$\text{PGL}(V) = \text{Aut}_k(\mathbf{P}(V)) \rightarrow \text{Aut}_k(\Omega(V)), \quad \varphi \mapsto \varphi|_{\Omega(V)}$$

*is an isomorphism. Equivalently, every  $k$ -automorphism of  $\Omega(V)$  extends to a  $k$ -automorphism of  $\mathbf{P}(V)$ .*

(ii) *For every field extension  $K/k$  the natural map*

$$\text{PGL}(V) \longrightarrow \text{Aut}_K(\Omega(V)_K)$$

is an isomorphism. Equivalently, every  $K$ -automorphism of  $\Omega(V)_K$  comes by base change from a  $k$ -automorphism of  $\mathbf{P}(V)$ .

This result holds trivially if  $\dim V \leq 1$ , for then  $\Omega(V) = \mathbf{P}(V)$ . From now on, we assume that  $V$  has dimension at least 2 and we set  $n = \dim V - 1$ .

The proof combines analytic geometry in the sense of Berkovich with algebraic arguments. As a first step we show that every  $k$ -automorphism of  $\Omega(V)$  can be extended to an automorphism of the  $k$ -scheme  $X$  we get by blowing-up all linear subspaces of  $\mathbf{P}(V)$ . For this step we use Berkovich analytic geometry over the field  $k$  endowed with the trivial absolute value. The second step is of an algebraic nature and consists in checking that this automorphism of  $X$  is induced by a  $k$ -automorphism of  $\mathbf{P}(V)$ . Here we analyze the geometry of the boundary divisor more closely and use an induction argument.

Given a proper subvector space  $W$  of  $V$ , applying  $\text{Proj}$  to the natural map  $\text{Sym}^\bullet(V) \rightarrow \text{Sym}^\bullet(V/W)$  leads to a closed immersion  $\mathbf{P}(V/W) \hookrightarrow \mathbf{P}(V)$  whose image  $L$  is called a *linear subspace* of  $\mathbf{P}(V)$ . Such a subscheme is said to be trivial if  $L = \emptyset$  or  $L = \mathbf{P}(V)$ ; it is called a *hyperplane* if it is of codimension 1. We denote by  $\mathcal{L}^i(V)$  the set of linear subspaces of dimension  $i$  in  $\mathbf{P}(V)$ , and by  $\mathcal{L}(V) = \bigcup_{0 \leq i \leq n-1} \mathcal{L}^i(V)$  the set of non-trivial linear subspaces.

DEFINITION 1.2. We denote by  $\pi : X \rightarrow \mathbf{P}(V)$  the blow-up of  $\mathbf{P}(V)$  along the full hyperplane arrangement. To be precise,  $X$  is defined as

$$X = X_{n-1} \xrightarrow{\pi_{n-1}} X_{n-2} \longrightarrow \dots \longrightarrow X_1 \xrightarrow{\pi_1} X_0 \xrightarrow{\pi_0} X_{-1} = \mathbf{P}(V)$$

with

$$\pi = \pi_0 \circ \pi_1 \circ \dots \circ \pi_{n-1},$$

where  $\pi_i$  denotes the blow-up of  $X_{i-1}$  along the strict transforms of linear subspaces of  $\mathbf{P}(V)$  of dimension  $i$ .

The scheme  $X$  is projective and smooth over  $k$ . It contains  $\Omega(V)$  as an open dense subscheme since each  $\pi_i$  induces an isomorphism over  $\Omega(V)$ . We write  $D = X - \Omega(V)$  for the complement.

Note that  $\pi_{n-1}$  is an isomorphism and that the strict transforms of two distinct linear subspaces  $L, L' \subset \mathbf{P}(V)$  of dimension  $i$  in  $X_{i-1}$  are disjoint since (the strict transform of)  $L \cap L'$  has been previously blown-up.

Each non-trivial linear subspace  $L \subset \mathbf{P}(V)$  defines a smooth and irreducible hypersurface  $E_L$  in  $X$  as follows. If  $L$  has dimension  $i$ , its strict transform by  $\pi_0 \circ \pi_1 \circ \dots \circ \pi_{i-1}$  in  $X_{i-1}$  (by convention  $L$  itself if it is a point) is blown-up under the map  $\pi_i : X_i \rightarrow X_{i-1}$  to give rise to a hypersurface  $E_L^{(i)}$  in  $X_i$ . The (codimension 1) subscheme  $E_L$  of  $X$  is then the strict transform of  $E_L^{(i)}$  by  $\pi_{i+1} \circ \dots \circ \pi_{n-1}$ . The induced map  $E_L \rightarrow E_L^{(i)}$  coincides with the blow-up of  $E_L^{(i)}$  along the hypersurface arrangement induced by hyperplanes of  $\mathbf{P}(V)$  containing  $L$ . We have an alternative description of  $E_L$  as the closure

$$\overline{\pi^{-1}\left(L - \bigcup_{\substack{L' \in \mathcal{L}(V) \\ L' \subsetneq L}} L'\right)}$$

taken in  $X$ .

It follows from the construction of  $X$  that the boundary divisor  $D$  is the union of all hyper-surfaces  $E_L$ , i.e. we have

$$D = \pi^{-1} \left( \bigcup_{\substack{W \subset V \\ \dim W = 1}} \mathbf{P}(V/W) \right) = \bigcup_L E_L.$$

Two components  $E_L, E_{L'}$  have non-empty intersection if and only if  $L \subset L'$  or  $L' \subset L$ . Indeed, if none of the inclusions holds, then  $L$  and  $L'$  intersect along a smaller linear subspace, say of dimension  $i$ , and the strict transforms of  $L$  and  $L'$  in  $X_i$  are disjoint. It follows that a family of components has non-empty intersection if and only if it is indexed by linear subspaces lying in a flag. We define the stratum  $Z_{\mathcal{F}}$  corresponding to a flag  $\mathcal{F}$  by:

$$Z_{\mathcal{F}} = \bigcup_{L \in \mathcal{F}} E_L - \bigcup_{L' \notin \mathcal{F}} E_{L'}.$$

LEMMA 1.3. *The divisor  $D$  has simple normal crossings. Moreover, if  $Z = Z_{\mathcal{F}}$  is the stratum corresponding to the flag  $\mathcal{F}$ , then*

$$U_Z = X - \bigcup_{L \notin \mathcal{F}} E_L$$

is an affine open subset of  $X$  containing  $Z$  as a closed subset.

*Proof.* We start by considering a complete flag  $\mathcal{F} = (L_0, \dots, L_n)$ . In order to get an explicit description of  $U_Z$  in this case, we first compare  $X$  to the blow-up  $Y$  of  $\mathbf{P}(V)$  along  $\mathcal{F}$ . To be precise, we define:

$$p : Y = Y_{n-1} \xrightarrow{p_{n-1}} Y_{n-2} \longrightarrow \dots \longrightarrow Y_1 \xrightarrow{p_1} Y_0 \xrightarrow{p_0} Y_{-1} = \mathbf{P}(V)$$

where  $p_i$  denotes the blow-up of  $Y_{i-1}$  along the strict transform of  $L_i$ . By the universal property of blow-up, there exists a (unique) morphism of towers  $f_{\bullet} : X_{\bullet} \rightarrow Y_{\bullet}$ .

Now, we want to show that  $f$  identifies  $U_Z$  with the complement  $W_Z$  in  $Y$  of the strict transforms of all linear subspaces not contained in  $\mathcal{F}$ . Note that  $W_Z$  is also the complement of the strict transform of all hyperplanes distinct from  $L_{n-1}$ . We argue by induction along the towers of blow-ups. For every  $i \in \{-1, \dots, n-1\}$ , we define two open subsets  $U_i \subset X_i$  and  $W_i \subset Y_i$  as follows:

- $U_{-1} = W_{-1}$  is the complement in  $\mathbf{P}(V)$  of all 0-dimensional linear subspaces distinct from  $L_0$ ;
- if  $0 \leq i \leq n-2$ , then  $U_i$  (resp.  $W_i$ ) is the complement in  $\pi_i^{-1}(U_{i-1})$  (resp. in  $p_i^{-1}(W_{i-1})$ ) of the strict transforms of all  $(i+1)$ -dimensional linear subspaces  $L \subset \mathbf{P}(V)$  not in  $\mathcal{F}$ ;
- $U_{n-1} = \pi_{n-1}^{-1}(U_{n-2})$  and  $W_{n-1} = p_{n-1}^{-1}(W_{n-2})$ .

Arguing by induction on  $i$ , we see that  $U_i = f_i^{-1}(W_i)$ , and that  $f_i$  induces an isomorphism between  $U_i$  and  $W_i$  respecting the restrictions of exceptional divisors. It is clear that

$$U_{n-1} = U_Z = X - \bigcup_{L \notin \mathcal{F}} E_L.$$

On the other hand, we claim that  $W_{n-1}$  coincides with  $W_Z$ . The inclusion  $W_{n-1} \subset W_Z$  is obvious. For every point  $y \in Y - W_{n-1}$  there exists an index  $i \in \{-1, \dots, n-2\}$  such that the image  $y_i$  of  $y$  in  $Y_i$  lies in the strict transform of a  $(i+1)$ -dimensional linear subspace  $L \subset \mathbf{P}(V)$  distinct from  $L_{i+1}$ . Let us consider a hyperplane  $H$  which contains  $L$ . By construction,  $y_i$  is contained in

the strict transform of  $H$  in  $Y_i$ . Since  $L_j \not\subset H$  for  $j \in \{i, \dots, n-2\}$ , the subspaces  $L_j$  and  $H$  are transverse. Blowing-up along some smooth subschemes can only decrease the order of contact, hence the strict transform  $\tilde{H}$  of  $H$  in  $Y_j$  is transverse to the center of  $p_{j+1}$ . This implies that the strict transform of  $H$  in  $Y_{j+1}$  coincides with the inverse image of  $\tilde{H}$  in  $Y_{j+1}$ . It follows that  $y$  belongs to the strict transform of  $H$  in  $Y$ , and thus  $y \in Y - W_Z$ . This proves the converse inclusion  $W_Z \subset W_{n-1}$ .

Given a basis  $(e_0, e_1, \dots, e_n)$  of  $V$  such that  $L_i = Z(e_{i+1}, \dots, e_n)$  for every  $i \in \{0, \dots, n-1\}$ , we have a commutative diagram

$$\begin{array}{ccc} \mathrm{Spec}(k[t_1, \dots, t_n]) & \xrightarrow{j} & Y \\ q \downarrow & & \downarrow p \\ \mathrm{Spec}(k[x_1, \dots, x_n]) & \xrightarrow{\quad} & \mathbf{P}(V) \end{array}$$

where the horizontal arrows are open immersions identifying  $t_1, \dots, t_n$  (resp.  $x_1, \dots, x_n$ ) with the rational functions  $e_1/e_0, \dots, e_n/e_{n-1}$  (resp.  $e_1/e_0, \dots, e_n/e_0$ ) and where  $q$  is the morphism defined by  $q^*(x_i) = \prod_{j \leq i} t_j$ .

Via  $j$ , the open subscheme  $W_Z$  of  $Y$  is isomorphic to the principal open subset  $D(f)$  of  $\mathrm{Spec}(k[t_1, \dots, t_n])$ , where

$$f = \prod_{i=1}^n \prod_{(a_i, \dots, a_n) \in k^{n-i+1}} (1 + a_i t_i + a_{i+1} t_i t_{i+1} + \dots + a_n t_i \dots t_n).$$

In particular,  $W_Z$  is affine. Moreover, the intersection of the exceptional divisor of  $p$  with the open affine set  $W_Z$  coincides with  $\mathrm{div}(t_1 \cdots t_n)$ , hence has simple normal crossings. Using the isomorphism between  $U_Z$  and  $W_Z$  induced by  $f$ , we deduce that  $U_Z$  is affine and that  $D \cap U_Z$  has simple normal crossings. Since the sets  $U_Z$  for all choices of complete flags form an open affine covering of  $X$ , the divisor  $D$  has simple normal crossings on  $X$ .

We now claim that the intersection  $\Sigma$  of any family of  $d$  irreducible components of  $D$  is either empty or irreducible. Indeed, assume that  $\Sigma$  is non-empty and reducible. Non-emptiness amounts to saying that these components correspond to linear subspaces in some flag  $\mathcal{F}$ . Pick a complete flag  $\mathcal{F}'$  containing  $\mathcal{F}$ . In the corresponding affine chart  $U_Z$ , the intersection of the  $d$  components which we consider is irreducible, hence there must be a component  $\Sigma_0$  of  $\Sigma$  which lies in  $X - U_Z$ . Since, by construction,  $X - U_Z$  is the union of some irreducible components of  $D$ , we see that  $\Sigma_0$  must be contained in a  $(d+1)$ -th irreducible component of  $D$ . But this contradicts the normal crossing property of  $D$ . In view of the discussion before Lemma 1.3, this shows that the strata of  $D$  are in one-to-one correspondence with flags of linear subspaces.

If we start with a stratum  $Z$  corresponding to a partial flag  $\mathcal{F}$ , the set  $U_Z = X - \bigcup_{L \notin \mathcal{F}} E_L$  is the intersection of all  $U_{Z'}$  for strata  $Z'$  corresponding to complete flags containing  $\mathcal{F}$ . Hence it is open affine as a finite intersection of open affines in a separated  $k$ -scheme.  $\square$

In order to extend an automorphism of  $\Omega(V)$  to first  $X$  and then to  $\mathbf{P}(V)$ , we look at its action on the discrete valuations associated to the components of  $D$ . For each  $L \in \mathcal{L}(V)$ , the local ring at the generic point of the hypersurface  $E_L$  is a discrete valuation ring in the function field  $\kappa(V)$  of  $X$ . We denote by  $\mathrm{ord}_L$  the corresponding discrete valuation on  $\kappa(V)$ , and we write

$$\Gamma(V) = \{\mathrm{ord}_L : L \in \mathcal{L}(V)\}$$

for the set of all these valuations. Note that  $\kappa(V)$  is the function field of both  $\mathbf{P}(V)$  and  $\Omega(V)$ . If  $L$  is a *hyperplane* in  $\mathbf{P}(V)$ , then the valuation  $\text{ord}_L$  is the one given by the local ring of  $\mathbf{P}(V)$  at the generic point of  $L$ .

The sets  $\mathcal{L}(V)$  and  $\Gamma(V)$  come with a natural simplicial structure, for which the  $q$ -simplices correspond to flags of linear subspaces of length  $q - 1$ .

PROPOSITION 1.4. *Let  $\varphi$  be a  $k$ -automorphism of  $\Omega(V)$  and let  $\varphi^*$  be the induced automorphism of the set of valuations on the function field  $\kappa(V)$ .*

- (i) *The birational map  $\varphi$  extends to a  $k$ -automorphism of  $X$  if and only if  $\varphi^*$  preserves the set  $\Gamma(V)$  and its simplicial structure.*
- (ii) *The birational map extends to a  $k$ -automorphism of  $\mathbf{P}(V)$  if and only if  $\varphi^*$  preserves the subset of  $\Gamma(V)$  defined by hyperplanes.*

*Proof.* (i) The condition is necessary because the simplicial set  $\Gamma(V)$  describes the incidence relations between irreducible components of  $D$  (Lemma 1.3). To see that it is sufficient, we use the covering of  $X$  by the open affine subsets

$$U_Z = X - \bigcup_{L \notin \mathcal{F}} E_L$$

where  $Z$  denotes a stratum of  $D$  and  $\mathcal{F}$  is the corresponding flag of linear subspaces of  $\mathbf{P}(V)$ . If  $\varphi$  preserves  $\Gamma(V)$  with its simplicial structure, then there exists for every stratum  $Z$  another stratum  $Z'$  such that the rational map

$$U_{Z'} \dashrightarrow U_Z$$

induced by  $\varphi$  is defined at each point of height 1.

Since  $U_Z$  is affine and  $U_{Z'}$  is noetherian and normal, this rational map is everywhere defined on  $U_{Z'}$  [EGA, 20.4.12] and therefore  $\varphi$  extends to an automorphism from  $X$  to  $X$  (apply this argument to  $\varphi^{-1}$ ).

(ii) If the morphism  $\varphi : \Omega(V) \rightarrow \Omega(V)$  preserves all valuations  $\text{ord}_L$  coming from hyperplanes, then for every hyperplane  $L$  in  $\mathbf{P}(V)$  there exists a hyperplane  $L'$  such that the rational map

$$\mathbf{P}(V) - L' \dashrightarrow \mathbf{P}(V) - L$$

induced by  $\varphi$  is defined at every point of height 1, and the conclusion follows as for (i).  $\square$

## 2. Step 1 – Valuations and analytic geometry

This section is devoted to the first step toward the theorem, namely the fact that every  $k$ -automorphism of  $\Omega(V)$  extends to a  $k$ -automorphism of  $X$ .

PROPOSITION 2.1. *Let  $\text{Aut}_k(X, D)$  denote the group of  $k$ -automorphisms of  $X$  which preserve  $D$ . The canonical map*

$$\text{Aut}_k(X, D) \rightarrow \text{Aut}_k(\Omega(V)), \quad \varphi \mapsto \varphi|_{\Omega(V)}$$

*is an isomorphism. Equivalently, every  $k$ -automorphism of  $\Omega(V)$  extends to a  $k$ -automorphism of  $X$ .*

We can study this problem from a nice geometric viewpoint in the framework of Berkovich spaces.

Endowed with the trivial absolute value,  $k$  becomes a complete non-Archimedean field. There is a well-defined category of  $k$ -analytic spaces, together with an analytification functor  $Z \rightsquigarrow Z^{\text{an}}$  from the category of  $k$ -schemes locally of finite type. If  $Z$  is affine, then the topological space underlying  $Z^{\text{an}}$  is the set of multiplicative  $k$ -seminorms on  $\mathcal{O}(Z)$  with the topology generated by evaluation maps  $x \mapsto |f(x)| := x(f)$ , where  $f \in \mathcal{O}(Z)$ . Imposing the additional condition that all seminorms are bounded by 1 on the algebra  $\mathcal{O}(Z)$ , we obtain a compact domain  $Z^\square$  in  $Z^{\text{an}}$  equipped with a *specialization* map  $\text{sp} : Z^\square \rightarrow Z$  (denoted by  $r$  in [Thu07]) which sends a multiplicative seminorm  $x$  to the prime ideal  $\{f \in \mathcal{O}(Z) \mid |f(x)| < 1\}$ . The reader is referred to [Ber90, Section 3.5] and [Thu07, Section 1] for a detailed account.

Working in the analytic category over  $k$  allows us to realize  $\Gamma(V)$  as a set of rays in  $\Omega(V)^{\text{an}}$ : for each  $L \in \mathcal{L}(V)$ , the map

$$\varepsilon_L : (0, 1] \rightarrow \Omega(V)^{\text{an}}, \quad r \mapsto r^{\text{ord}_L(\cdot)}$$

is an embedding and  $\varepsilon_L(1)$  is the canonical point of  $\Omega(V)^{\text{an}}$ , namely the point corresponding to the trivial absolute value on  $\kappa(V)$ . Now, the proposition will follow from the fact that this collection of rays is the 1-skeleton of a *conical complex*  $\mathfrak{S}(V)$  in  $\Omega(V)^{\text{an}}$  which is preserved by every  $k$ -automorphism of  $\Omega(V)$ .

This conical complex  $\mathfrak{S}(V)$  is the *fan*  $\mathfrak{S}_0(X, D)$  of the toroidal embedding  $\Omega(V) \hookrightarrow X$  introduced in [Thu07, Section 3.1 and Proposition 4.7], following [Ber99]. Let us describe this construction in the particular case we consider here.

(a) The canonical map

$$r : \mathbf{A}_k^{n, \square} \rightarrow [0, 1]^n, \quad x \mapsto (|t_1(x)|, \dots, |t_n(x)|)$$

has a continuous section  $j$  defined by mapping a tuple  $r \in [0, 1]^n$  to the following diagonalizable multiplicative seminorm on  $k[t_1, \dots, t_n]$ :

$$\sum_{\nu \in \mathbf{N}^n} a_\nu t^\nu \mapsto \max_{\nu} |a_\nu| r_1^{\nu_1} \cdots r_n^{\nu_n}.$$

- (b) Let  $D(t_1, \dots, t_n)$  denote the invertibility locus of  $t_1, \dots, t_n$ . Intersecting the image of  $j$  with the open domain  $D(t_1, \dots, t_n)^\square$ , we obtain a closed subset  $C_n \subset D(t_1, \dots, t_n)^\square$  homeomorphic to the cone  $(0, 1]^n$ . The map  $\tau = r \circ j$  is a retraction of  $D(t_1, \dots, t_n)^\square$  onto  $C_n$ . Its fiber over a point  $x \in C_n$  is a  $k$ -affinoid domain whose Shilov boundary is reduced to  $\{x\}$ .
- (c) We identify  $C_n$  and  $(0, 1]^n$  via  $r$ . For  $I \subset \{1, \dots, n\}$ , let  $C_n^I$  denote the face of  $C_n$  defined by  $r_i = 0$  for every  $i \in I$ . The specialization map  $\text{sp} : D(t_1, \dots, t_n)^\square \rightarrow \mathbf{A}_k^n$  sends the interior of  $C_n^I$  to the generic point of the locally closed subscheme  $Z_I = V(t_i, i \in I) \cap D(t_j, j \notin I)$ . This implies that  $C_n^I$  is contained in  $U^\square = \text{sp}^{-1}(U)$  for any open neighborhood  $U$  of the generic point of  $Z_I$ .
- (d) We can also recover the monoid  $r_1^{\mathbf{N}} \cdots r_n^{\mathbf{N}}$  defining the integral affine structure on  $(0, 1]^n$  from the analytic structure of  $\mathbf{A}_k^n$ . Indeed, this is precisely the monoid of functions  $|f| : C_n \rightarrow (0, 1]$  induced by germs  $f \in \mathcal{O}_{\mathbf{A}_k^n, 0}$  invertible on  $D(t_1, \dots, t_n)$ . Similarly, the submonoid corresponding to the face  $C_n^I$  comes from germs of  $\mathcal{O}_{\mathbf{A}_k^n}$  at the generic points of  $Z_I$  which are invertible over  $D(t_1, \dots, t_n)$ .
- (e) We now return to the scheme  $X$  with its simple normal crossing divisor  $D$ . Fix a stratum  $Z$  with generic point  $\eta_Z$  and let  $\Lambda_Z^+$  denote the monoid of germs in  $\mathcal{O}_{X, \eta_Z}$  whose restriction to  $\Omega(V)$  is invertible. As in the proof of Lemma 1.3, there is an open immersion  $(t_1, \dots, t_n) : U_Z \rightarrow \mathbf{A}_k^n$  identifying  $Z$  with a non-empty open subset of  $Z_I$  for a suitable subset  $I$  of



$\{1, \dots, n\}$ . By transport of structure, we obtain a closed subset  $C_Z$  of  $U_Z^\square - Z^\square$  such that the natural map

$$C_Z \rightarrow \mathrm{Hom}_{\mathbf{Mon}}(\Lambda_Z^+/k^\times, (0, 1]), \quad x \mapsto (f \mapsto |f(x)|)$$

is a homeomorphism. Covering  $X$  by the open subschemes  $U_Z$ , we can glue the cones  $C_Z$  along common faces in  $\Omega(V)^{\mathrm{an}}$  to define a cone complex  $\mathfrak{S}(V)$ . This gluing is compatible with local retractions, so we get a retraction of  $\Omega(V)^{\mathrm{an}}$  onto  $\mathfrak{S}(V)$ .

The following property of the conical complex  $\mathfrak{S}(V)$  is specific to our situation and is the key point to prove Proposition 2.1. It may be interesting to look for other “natural” toroidal compactifications satisfying this condition.

LEMMA 2.2. *The map*

$$\iota : \mathfrak{S}(V) \rightarrow \mathrm{Hom}_{\mathbf{Ab}}\left(\mathcal{O}(\Omega(V))^\times/k^\times, \mathbf{R}_{>0}\right), \quad x \mapsto (f \mapsto |f(x)|)$$

*is a closed embedding inducing the integral affine structure on each cone. Moreover, (the images of) distinct cones span distinct linear spaces.*

*Proof.* Roughly speaking, this statement means that there are enough invertible functions on  $\Omega(V)$ . Consider a stratum  $Z$  of  $D$  corresponding to a flag  $\mathcal{F}$  of non-trivial linear subspaces of  $\mathbf{P}(V)$  and pick a basis  $(e_0, \dots, e_n)$  of  $V$  such that  $\mathcal{F}$  is a subflag of

$$Z(e_1, \dots, e_n) \subset Z(e_2, \dots, e_n) \subset \dots \subset Z(e_n).$$

The explicit description of  $X$  given at the end of the proof of Lemma 1.3 shows that the tuple  $(e_1/e_0, e_2/e_1, \dots, e_n/e_{n-1})$  of elements in  $\mathcal{O}_{X, \eta_Z}$  contains a regular system of parameters defining  $D$  at  $\eta_Z$ . Therefore, the map  $\iota$  induces an integral affine embedding of the cone  $C_Z$ .

Furthermore, we claim that the following fact is true: *given two distinct cones  $C, C'$ , there exists  $f \in \mathcal{O}(\Omega(V))^\times$  such that  $|f| = 1$  on one of them and  $|f| < 1$  on the interior of the other.* Injectivity of the map  $\iota$  and the last statement of the Lemma follow immediately.

We finish the proof by establishing the claim. Given two non-zero vectors  $v, v' \in V$  and a non-trivial linear subspace  $L \subset \mathbf{P}(V)$ , the function  $v/v'$  is either a unit, a uniformizer or the inverse of a uniformizer at the generic point of  $E_L$ , according to the position of  $Z(v)$  and  $Z(v')$  with respect to  $L$ . It follows that

- (a)  $|v/v'| < 1$  on  $\varepsilon_L(0, 1)$ , if  $L \subset Z(v)$  and  $L \not\subset Z(v')$
- (b)  $|v/v'| > 1$  on  $\varepsilon_L(0, 1)$ , if  $L \subset Z(v')$  and  $L \not\subset Z(v)$
- (c)  $|v/v'| = 1$  on  $\varepsilon_L(0, 1]$ , if the hyperplanes  $Z(v)$  and  $Z(v')$  are in the same position with respect to  $L$ .

Consider two distinct strata  $Z, Z'$  of  $D$ , corresponding to distinct flags  $\mathcal{F}, \mathcal{F}'$  of non-trivial linear subspaces. Pick a linear space  $L$  occurring in only one of them, say  $\mathcal{F}$ , and set  $i = \dim L$ . We embed  $\mathcal{F}'$  into a complete flag  $(L_0 \subset L_1 \subset \dots \subset L_{n-1})$  such that  $L_i \neq L$ .

We claim that this assumption guarantees the existence of two hyperplanes  $H, H'$  such that

- $L \subset H$  and  $L_i \not\subset H$
- $L_i \cap H = L_i \cap H'$  and  $L \not\subset H'$ .

In order to prove this claim, we argue with the corresponding linear quotient spaces of  $V$ . Let  $L = \mathbf{P}(V/W)$  and  $L_i = \mathbf{P}(V/W_i)$  where  $W$  and  $W_i$  are different linear subspaces of  $V$  of

dimension  $n - i$ . Choose a vector  $u \in W$  which is not contained in  $W_i$ , and a vector  $u_i \in W_i$  which is not contained in  $W$ . We denote by  $U$  the line in  $V$  generated by  $u$  and by  $U'$  the line generated by  $u' = u + u_i$ . The corresponding hyperplanes  $H = \mathbf{P}(V/U)$  and  $H' = \mathbf{P}(V/U')$  have the desired properties.

In particular,  $H$  and  $H'$  are in the same position with respect to  $L_0, \dots, L_{n-1}$ . Given any equations  $v, v' \in V$  of  $H$  and  $H'$  respectively, we thus obtain  $|v/v'| = 1$  on  $C_{Z'}$ . Let us now consider the flag  $\mathcal{F}$ . Any linear subspace  $M \in \mathcal{F}$  contained in  $H'$  is necessarily contained in  $L$ , hence in  $H$ , therefore  $|v/v'| \leq 1$  on the ray  $\varepsilon_M(0, 1]$ . Since  $|v/v'| < 1$  on the interior of the ray  $\varepsilon_L(0, 1]$ , we deduce that  $|v/v'| < 1$  on the interior of the cone  $C_Z$ .  $\square$

*Proof of proposition 2.1.* First, we observe that  $\mathfrak{S}(V)$  coincides with the set  $\Omega(V)_{\max}^{\text{an}}$  of maximal points of  $\Omega(V)^{\text{an}}$  for the following ordering:

$$x \preceq y \iff \forall f \in \mathcal{O}(\Omega(V)^{\text{an}}), |f(x)| \leq |f(y)|.$$

For any point  $x \in \Omega(V)^{\text{an}}$ , we have  $x \preceq \tau(x)$  because the fiber  $\tau^{-1}(\tau(x))$  is a  $k$ -affinoid domain with Shilov boundary  $\{\tau(x)\}$ . This implies the inclusion  $\Omega(V)_{\max}^{\text{an}} \subset \mathfrak{S}(V)$ .

We apply Lemma 2.2 to get the converse inclusion. If a point  $x \in \mathfrak{S}(V)$  is dominated by a point  $x' \in \Omega(V)^{\text{an}}$ , then it is also dominated by  $\tau(x')$ . However, for any two distinct points  $x, y$  in  $\mathfrak{S}(V)$ , there exists  $f \in \mathcal{O}(\Omega(V)^{\text{an}})^{\times}$  such that  $|f(x)| \neq |f(y)|$ , hence such that  $|f(x)| < |f(y)|$  and  $|(\frac{1}{f})(x)| > |(\frac{1}{f})(y)|$  or vice versa, and therefore  $x$  and  $y$  are incomparable. In particular, we get  $x = \tau(x')$  and thus  $x$  is maximal.

The above characterization of  $\mathfrak{S}(V)$  as a closed subset of  $\Omega(V)^{\text{an}}$  implies that it is preserved by any  $k$ -automorphism  $\varphi$  of  $\Omega(V)$ . It remains to check that the homeomorphism of  $\mathfrak{S}(V)$  induced by  $\varphi$  also preserves the conical structure. Let  $\Phi$  denote the linear automorphism of  $\text{Hom}_{\mathbf{Ab}}(\mathcal{O}(\Omega(V)^{\text{an}})^{\times}, \mathbf{R}_{>0})$  deduced from  $\varphi$ . Given an  $n$ -dimensional cone  $C \subset \mathfrak{S}(V)$ , the image of its interior is disjoint from the  $(n-1)$ -skeleton of  $\mathfrak{S}(V)$ ; otherwise it would meet the interiors of two distinct  $n$ -dimensional cones  $C', C''$ , hence  $\langle \iota C' \rangle = \Phi(\langle \iota C \rangle) = \langle \iota C'' \rangle$  contradicting Lemma 2.2. It follows that if  $\varphi(C)$  is contained in some  $n$ -dimensional cone  $C'$ , and thus  $\varphi(C) = C'$  by considering  $\varphi^{-1}$ . The assertion for lower dimension cones follows at once by considering faces since the automorphism  $\Phi$  is linear.

In particular, we see that  $\varphi$  preserves the 1-skeleton of  $\mathfrak{S}(V)$ , hence the set  $\Gamma(V)$  of discrete valuations on  $\kappa(V)$  associated with irreducible components of  $D = X - \Omega(V)$ , together with the simplicial structure reflecting the incidence relations between these components. By Proposition 1.4 (i), this implies that  $\varphi$  extends to a  $k$ -automorphism of  $X$ .  $\square$

REMARK 2.3.

- (i) Let  $D$  be a simple normal crossing divisor on a smooth and proper (connected) scheme  $X$  over  $k$ . Even if  $\Omega(V) = X - D$  is affine, Lemma 2.2 and its consequences may fail. For example, consider the case  $X = \mathbf{P}_k^n$ . If  $D$  is a hyperplane, then  $\mathfrak{S}_0(X, D)$  is a 1-dimensional cone whereas  $\Omega(V)_{\max}^{\text{an}}$  is empty. If  $D$  is the union of the coordinate hyperplanes, then  $\Omega(V) = \mathbf{G}_m^n$  and  $\mathfrak{S}_0(X, D) = \Omega(V)_{\max}^{\text{an}}$  is the toric fan, but the map  $\iota$  is bijective, hence all maximal cones span the same linear space. In fact, the inversion  $(t_1, \dots, t_n) \mapsto (t_1^{-1}, \dots, t_n^{-1})$  on  $\mathbf{G}_m^n$  transforms the fan  $\mathfrak{S}_0(X, D)$  into its opposite, hence does not preserve the conical structure. This reflects the fact that this automorphism of  $\mathbf{G}_m^n$  does not extend to  $\mathbf{P}^n$ .

- (ii) The conical complex  $\mathfrak{S}(V)$  is also the *vectorial building* of  $\mathrm{PGL}(V)$ , but this is somehow fortuitous and irrelevant from the viewpoint of automorphisms. In general, there exists for any connected and split semi-simple  $k$ -group  $G$  a canonical embedding of the vectorial building  $\mathcal{V}(G, k)$  of  $G(k)$  into the analytification of an open affine subscheme  $\Omega$  in any flag variety  $Y$  of  $G$  [Ber90, Section 5.5]. However, this observation does not lead to a generalization of Theorem 1.1, at least along the lines of the present proof. Indeed, while we made crucial use of the fact that  $\mathfrak{S}(V)$  is the fan of a normal crossing divisor, we doubt that  $\mathcal{V}(G, k)$  can be realized as the fan of a toroidal compactification of  $\Omega(V)$  if  $(G', Y) \neq (\mathrm{PGL}(V), \mathbf{P}(V)), (\mathrm{PGL}(V), \mathbf{P}(V^\vee))$ .
- (iii) It may be interesting to try to extend our method, based on the study of toroidal compactifications, to determine the automorphism groups of other period domains.
- (iv) Whether the above proposition can be proved without analytic geometry is not clear.

### 3. Step 2 – Geometry of the blow-up

The last step in the proof of the theorem relies on elementary intersection theory on  $X$ , which we review in this section. The standard reference is [Ful97].

The Chow ring  $\mathrm{CH}^*$  is a contravariant functor from the category of smooth  $k$ -schemes to the category of graded commutative rings. For any smooth  $k$ -scheme  $X$ , the abelian group underlying  $\mathrm{CH}^*(X)$  is the free abelian group on integral subschemes of  $X$  modulo rational equivalence, and it is graded by codimension. Multiplication comes from the intersection product. We write  $[Z]$  for the class of a closed subscheme  $Z$  of  $X$ .

We are going to use the following two basic facts.

- (a) Let  $Y$  be a regularly embedded closed subscheme of  $X$  of codimension  $\geq 2$  and let  $\pi : \tilde{X} \rightarrow X$  be the blow-up of  $X$  along  $Y$ , with exceptional divisor  $\tilde{Y}$ . The canonical map

$$\mathrm{CH}^1(X) \oplus \mathbf{Z}[\tilde{Y}] \rightarrow \mathrm{CH}^1(\tilde{X}), \quad (z, n[\tilde{Y}]) \mapsto \pi^*(z) + n[\tilde{Y}]$$

is an isomorphism [Ful97, Proposition 6.7].

- (b) In the situation of (a), let  $V$  be an integral subscheme of  $X$  with strict transform  $\tilde{V}$ . If  $\mathrm{codim}(Y, X) \leq \mathrm{codim}(V \cap Y, V)$ , then

$$\pi^*[V] = [\tilde{V}]$$

in  $\mathrm{CH}^*(\tilde{X})$  [Ful97, Corollary 6.7.2].

Now we focus on the particular case where  $\pi : X \rightarrow \mathbf{P}(V)$  is the blow-up along the full hyperplane arrangement, with exceptional divisor  $D$ .

LEMMA 3.1. *We have*

$$\mathrm{CH}^1(X) = \mathbf{Z}h \oplus \bigoplus_{L} \mathbf{Z}[E_L],$$

where  $h = \pi^*[H]$  denotes the pull-back of the hyperplane class  $[H]$  on  $\mathbf{P}(V)$  and  $L$  runs over the set of non-trivial linear subspaces of  $\mathbf{P}(V)$  of codimension at least 2.

*Proof.* For any non-trivial linear subspace  $L$  of  $\mathbf{P}(V)$  of dimension  $i \in \{0, \dots, n-1\}$ , let  $E_L^{(i)} \subset X_i$  denote the blow-up of its strict transform in  $X_{i-1}$ ; this is a smooth irreducible hypersurface. Recall that we have  $\pi = \pi_0 \circ \pi_1 \circ \dots \circ \pi_{n-1}$ , where  $\pi_{n-1}$  is an isomorphism. Applying (a) iteratively to

each blow-up  $\pi_0, \dots, \pi_{n-2}$ , we obtain that  $\text{CH}^1(X)$  is the free abelian group on  $h$  and the classes  $(\pi_{i+1} \circ \dots \circ \pi_{n-1})^* [E_L^{(i)}]$ , where  $i \in \{0, \dots, n-2\}$  and  $L$  runs over the set of  $i$ -dimensional linear subspaces of  $\mathbf{P}(V)$ .

The conclusion follows from the additional fact that we have an equality

$$(\pi_{i+1} \circ \dots \circ \pi_{n-1})^* [E_L^{(i)}] = [E_L]$$

in  $\text{CH}^1(X)$  for any linear subspace  $L$  of dimension  $i \in \{0, \dots, n-2\}$ . This is an immediate consequence of (b), since the center of each blow-up  $\pi_j$ , with  $j \in \{i+1, \dots, n-1\}$ , is transversal to the strict transform of  $E_L^{(i)}$  in  $X_{j-1}$ .  $\square$

By Proposition 2.1, every  $k$ -automorphism  $\varphi$  of  $\Omega(V)$  extends to a  $k$ -automorphism  $\tilde{\varphi}$  of  $X$ , hence it induces a permutation  $\hat{\varphi}$  of non-trivial linear subspaces of  $\mathbf{P}(V)$  defined by

$$\tilde{\varphi}(E_L) = E_{\hat{\varphi}(L)}.$$

Let us denote by  $K_Y$  the canonical divisor on a smooth  $k$ -scheme  $Y$ . For  $i \in \{0, \dots, n-1\}$ , we recall that  $\pi_i : X_i \rightarrow X_{i-1}$  is the blow-up along the strict transforms of all  $i$ -dimensional linear subspaces  $L \subset \mathbf{P}(V)$  in  $X_{i-1}$  (Definition 1.2). By the classical formula for the canonical divisor of a blow-up [Har77, Exercice II.8.5],

$$K_{X_i} = \pi_i^* K_{X_{i-1}} - (n-i-1) \sum_{L \in \mathcal{L}^i(V)} [E_L^{(i)}]$$

for every  $i \in \{0, \dots, n-1\}$ , where  $E_L^{(i)}$  denotes the irreducible component of the exceptional divisor of  $\pi_i$  coming from  $L$ . We deduce :

$$K_X = \pi^* K_{\mathbf{P}(V)} - \sum_{i=0}^{n-2} (n-i-1) \sum_{L \in \mathcal{L}^i(V)} (\pi_{i+1} \circ \dots \circ \pi_{n-1})^* [E_L^{(i)}] = -(n+1)h - \sum_{L \in \mathcal{L}} (\text{codim } L - 1) [E_L]$$

since  $\pi^* K_{\mathbf{P}(V)} = -(n+1)h$  and  $(\pi_{i+1} \circ \dots \circ \pi_{n-1})^* [E_L^{(i)}] = [E_L]$  (see the proof of Lemma 3.1). Then, we have the following identity in  $\text{CH}^1(X)$

$$(n+1)\tilde{\varphi}^* h + \sum_L (\text{codim } L - 1) \tilde{\varphi}^* [E_L] = (n+1)h + \sum_L (\text{codim } L - 1) [E_L]. \quad (1)$$

Let  $\Gamma$  denote the subgroup of  $\text{CH}^1(X)$  spanned by all  $[E_L]$  for  $L \in \mathcal{L}(V)$  with codimension at least 2. Since

$$h = [E_H] + \sum_{L \subsetneq H} [E_L]$$

for any hyperplane  $H$ , we deduce that

$$\tilde{\varphi}^* [E_L] \equiv \begin{cases} h & \text{if } \hat{\varphi}^{-1}(L) \text{ is a hyperplane} \\ 0 & \text{otherwise} \end{cases} \pmod{\Gamma}$$

This implies that

$$\tilde{\varphi}^* h \equiv \tilde{\varphi}^* [E_H] + \sum_{L \subsetneq H} \tilde{\varphi}^* [E_L] \equiv mh \pmod{\Gamma}$$

for some non-negative integer  $m$ .

Thanks to these observations, equation (1) leads to

$$\left( (n+1)m + \sum_{i=0}^{n-2} (n-i-1)m_i \right) h \equiv (n+1)h \pmod{\Gamma}$$

where  $m_i$  is a non-negative integer (the number of linear subspaces of dimension  $i$  mapped to hyperplanes under  $\widehat{\varphi}^{-1}$ ). Let us deduce from this identity that  $m = 1$  and  $m_i = 0$  for all  $i$ . If  $m = 0$ , then  $\widehat{\varphi}$  sends hyperplanes to lower dimensional linear subspaces and therefore each hyperplane  $H$  equals  $\widehat{\varphi}^{-1}L$  for some  $L \in \mathcal{L}^i(V)$  with  $i \leq n-2$ . This implies the following estimate:

$$\sum_{i=0}^{n-2} m_i = \#\mathbf{P}(V)(k) = \#\mathbf{P}^n(k) = (\#k)^n + \dots + \#k + 1 > n + 1$$

and

$$\sum_{i=0}^{n-2} (n-i-1)m_i \geq \sum_{i=0}^{n-2} m_i > n + 1.$$

Thus we must have  $m \geq 1$ , and then it is obvious that  $m = 1$  and  $m_i = 0$  for every  $i$ . Therefore  $\widehat{\varphi}$  has to preserve the set of hyperplanes.

The idea of considering the canonical bundle of  $X$  in order to show that hyperplanes are mapped to hyperplanes was suggested to us by Carlo Gasbarri. The first version of this paper contained the following alternative argument for this fact. One checks first that  $\text{rk CH}^1(E_L)$  as a function on  $\dim L$  is symmetric with respect to  $(n-1)/2$  and decreasing for  $0 \leq \dim L \leq \lfloor (n-1)/2 \rfloor$ . This follows from the formula

$$\text{rk CH}^1(E_L) = \lambda(d) + \lambda(n-1-d) + \varepsilon(d)$$

where  $d = \dim L$ ,  $\varepsilon(d) = 1$  if  $d \in \{0, 1\}$  and  $\varepsilon(d) = 2$  otherwise,  $\lambda(0) = 0$  and

$$\lambda(d) = \# \left\{ \begin{array}{l} \text{non-trivial linear subspaces} \\ \text{of codimension } \geq 2 \text{ in } \mathbf{P}_k^d \end{array} \right\} \quad \text{for } d \geq 1.$$

In particular, the permutation  $\widehat{\varphi}$  of  $\mathcal{L}(V)$  induced by  $\varphi$  can only map a hyperplane to a hyperplane or to a point. Then, our claim follows by induction on  $n = \dim V - 1 \geq 1$ . For  $n = 1$ , the result is obvious. For  $n = 2$ , it is enough to compute self-intersections of components of  $D$  to conclude: for a point  $p$  and a line  $\ell$ ,

$$\deg [E_p]^2 = -1 \quad \text{and} \quad \deg [E_\ell]^2 = \deg \left( h - \sum_{q \in \ell(k)} [E_q] \right)^2 = 1 - \# \ell(k) = -(\# k),$$

thus  $\widehat{\varphi}$  maps a line to a line.

Let us now assume that  $n$  is at least 3 and that the theorem has been proved in lower dimension. If  $H$  is a hyperplane and  $\widehat{\varphi}(H)$  is a rational point  $p$ , then  $\widehat{\varphi}$  induces a  $k$ -isomorphism  $\widetilde{\varphi}$  between  $E_H$  and  $E_p$  which maps the divisor  $D_H = \bigcup_{L \neq H} E_H \cap E_L$  onto the divisor  $D_p = \bigcup_{L \neq \{p\}} E_p \cap E_L$ . Since  $E_H$  (resp.  $E_p$ ) is the blow-up of  $H$  (resp.  $\mathbf{P}(T_p^\vee)$ ), where  $T_p$  denotes the tangent space of  $\mathbf{P}(V)$  at  $p$ ) along the full hyperplane arrangement, with exceptional divisor  $D_H$  (resp.  $D_p$ ), the theorem in dimension  $n-1$  implies that  $\widetilde{\varphi}$  is induced by a  $k$ -isomorphism between  $H$  and  $\mathbf{P}(T_p^\vee)$ , hence maps the components of  $D_H$  defined by rational points of  $H$  to components of  $D_p$  defined by rational points of  $\mathbf{P}(T_p^\vee)$ , which is to say by (rational) lines in  $\mathbf{P}(V)$  containing  $p$ .

Let  $q$  be a rational point of  $H$  and let  $\ell$  denote the line in  $\mathbf{P}(V)$  such that  $\tilde{\varphi}(E_H \cap E_q) = E_p \cap E_\ell$ . The two hypersurfaces  $E_\ell$  and  $\tilde{\varphi}(E_q)$  have the same non-empty intersection with  $\tilde{\varphi}(E_H) = E_p$ , so  $\tilde{\varphi}(E_q) = E_\ell$  since  $D$  is a normal crossing divisor. On the other hand, since  $n \geq 3$ , we have  $\text{rk } CH^1(E_q) > \text{rk } CH^1(E_\ell)$  and we thus get a contradiction. Therefore,  $\hat{\varphi}$  preserves the set of hyperplanes.

#### 4. Extension of the ground field

We now indicate how to prove the second part of Theorem 1.1. For every field extension  $K/k$ , the base change  $\Omega(V)_K$  of  $\Omega(V)$  coincides with the complement in  $\mathbf{P}(V)_K$  of all  $k$ -rational hyperplanes. Since blowing-up commutes with base change, the  $K$ -scheme  $X_K = X \otimes_k K$  can be obtained by blowing up  $\mathbf{P}(V)_K$  along the arrangement of all  $k$ -rational hyperplanes. Moreover, every irreducible component  $E_L$  of  $D$  is geometrically irreducible, and its base change  $(E_L)_K$  is the irreducible component of  $X_K - \Omega(V)_K$  corresponding to the  $k$ -rational linear subspace  $L_K$  of  $\mathbf{P}(V)_K$ .

Let us consider a  $K$ -automorphism  $\varphi$  of  $\Omega(V)_K$ . One proves exactly as in Proposition 1.4 that  $\varphi$  extends to a  $K$ -automorphism of  $X_K$  (resp. of  $\mathbf{P}(V)_K$ ) if and only if  $\varphi$  preserves the simplicial set  $\Gamma(V_K)$  of discrete valuations on  $\kappa(V_K)$  coming from irreducible components of  $D_K$  (resp. preserves the subset of  $\Gamma(V_K)$  corresponding to hyperplanes). Once again, this condition is established via analytic geometry over the field  $K$  endowed with the trivial absolute value. The key point is Lemma 2.2, which holds for the fan  $\mathfrak{S}(V_K)$  of the normal crossing divisor  $D_K$  on  $X_K$ . The proof works verbatim, but one could also argue that  $\mathfrak{S}(V_K)$  coincides with the inverse image of  $\mathfrak{S}(V)$  under the projection map  $p : X_K^{\text{an}} \rightarrow X^{\text{an}}$ , so the statement holds for  $\mathfrak{S}(V_K)$  since it holds for  $\mathfrak{S}(V)$ . We then prove as above that  $\varphi$  extends to a  $K$ -automorphism of  $X_K$ .

Lemma 3.1 and the argument with the canonical divisor (or with the rank of Chow groups) also apply to  $X \otimes_k K$ , when we replace "linear subspaces" by " $k$ -rational linear subspaces". It follows that the permutation of  $k$ -rational linear subspaces induced by  $\tilde{\varphi}$  preserves the hyperplanes, hence  $\varphi$  induces a  $K$ -automorphism of  $\mathbf{P}(V)_K$ . This automorphism preserves the set of  $k$ -rational hyperplanes. Pick a  $k$ -basis of  $V$  and consider the corresponding coordinate hyperplanes; since they are mapped to  $k$ -rational hyperplanes,  $\varphi$  is induced by a  $k$ -automorphism of  $\mathbf{P}(V)$ .

#### REFERENCES

- Alon06 Gil Alon, Automorphisms of products of Drinfeld half planes, unpublished paper (2006)
- Ber90 Vladimir Berkovich, *Spectral Theory and Analytic Geometry over non-Archimedean Fields*, volume **33** of *Mathematical Surveys and Monographs*, AMS (1990)
- Ber95 ———, The automorphism group of the Drinfeld half-plane, *C. R. Acad. Sci. Paris* **321** (1995), no. 9, 1127-1132
- Ber99 ———, Smooth  $p$ -adic analytic spaces are locally contractible, *Inventiones Math.* **137** (1999), no. 1, 1-84
- DOR10 Jean-François Dat, Sascha Orlik and Michael Rapoport, *Period domains over finite and  $p$ -adic fields*, Cambridge Tracts in Mathematics **183**, Cambridge University Press (2010)
- Drin74 Vladimir G. Drinfeld, Elliptic Modules, *Math. USSR Sbornik* **23** (1974) 561-592.
- EGA Alexander Grothendieck, *Éléments de géométrie algébrique*, chapitre IV, Publ. Math. IHÉS **20**, **24**, **28**, **32** 1964-1967.
- Ful97 William Fulton, *Intersection theory*, Second Edition, Springer (1997)

- Har77 Robin Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer (1977)
- Orl01 Sascha Orlik, The cohomology of period domains for reductive groups over finite fields, *Ann. Scient. Éc. Norm. Sup.* **34** 63-77 (2001).
- OR08 Sascha Orlik, Michael Rapoport, Deligne-Lusztig varieties and period domains over finite fields, *J. of Algebra* **320** (2008) 1220-1234.
- Rap97 Michael Rapoport, Period domains over finite and local fields, *Proc. Symp. Pure Math.* **62**, part 1, 361-381 (1997).
- RZ96 Michael Rapoport, Thomas Zink: *Period spaces for  $p$ -divisible groups*. Princeton University Press 1996.
- Thu07 Amaury Thuillier, Géométrie toroïdale et géométrie analytique non archimédienne. Application au type d'homotopie de certains schémas formels, *Manuscripta Math* **123**, 381-451 (2007)

Bertrand Rémy [remy@math.univ-lyon1.fr](mailto:remy@math.univ-lyon1.fr)

Université de Lyon, Université Lyon1-CNRS, Institut Camille Jordan - UMR5208, 43 bd. du 11 novembre 1918 , F-69622 Villeurbanne cedex

Amaury Thuillier [thuillier@math.univ-lyon1.fr](mailto:thuillier@math.univ-lyon1.fr)

Université de Lyon, Université Lyon1-CNRS, Institut Camille Jordan - UMR5208, 43 bd. du 11 novembre 1918 , F-69622 Villeurbanne cedex

Annette Werner [werner@mathematik.uni-frankfurt.de](mailto:werner@mathematik.uni-frankfurt.de)

Institut für Mathematik , Goethe-Universität Frankfurt, Robert-Maier-Str. 6-8, D-60325 Frankfurt a.M.