



**HAL**  
open science

# Multifractal analysis of some multiple ergodic averages for the systems with non-constant Lyapunov exponents

Lingmin Liao, Michal Rams

► **To cite this version:**

Lingmin Liao, Michal Rams. Multifractal analysis of some multiple ergodic averages for the systems with non-constant Lyapunov exponents. 2012. hal-00705104v1

**HAL Id: hal-00705104**

**<https://hal.science/hal-00705104v1>**

Preprint submitted on 6 Jun 2012 (v1), last revised 21 Jun 2012 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Multifractal analysis of some multiple ergodic averages for the systems with non-constant Lyapunov exponents

Lingmin Liao \*

LAMA UMR 8050, CNRS, Université Paris-Est Créteil,  
61 Avenue du Général de Gaulle, 94010 Créteil Cedex, France

e-mail: lingmin.liao@u-pec.fr

Michał Rams †

Institute of Mathematics, Polish Academy of Sciences  
ul. Śniadeckich 8, 00-956 Warszawa, Poland

e-mail: rams@impan.gov.pl

## Abstract

We study certain multiple ergodic averages of a two iterated functions system on the unit interval. By using the dynamical coding  $\{0, 1\}^{\mathbb{N}}$  of the attractor, we compute the Hausdorff dimension of the set of points with a given frequency of the pattern 11 in positions  $k, 2k$ .

## 1 Introduction and statement of results

Initiated by the paper of Fan Liao and Ma [FLM], the study of the multiple ergodic average from a point view of multifractal analysis have attracted much attention. The major achievements have been made by Fan, Kenyon, Peres, Schmeling, Seuret, Solomyak, Wu and etc. ([KPS11, KPS12, PS12b,

---

\*partially supported by 12R03191A - MUTADIS (France).

†supported by MNiSW grant N201 607640 (Poland). This paper was written during the visit of M.R. in Université Paris-Est Créteil.

2000 *Mathematics Subject Classification*: Primary 28A80, Secondary 37C45, 28A78

PS12a, FSW11, FSW12, PSSS12]). For a short history, we refer the readers the paper of Peres and Solomyak [PS12a].

In [PS] and [FSW], the authors considered the symbolic space  $\Sigma = \{0, 1\}^{\mathbb{N}}$  with the metric  $d(x, y) = 2^{-\min\{n: x_n \neq y_n\}}$ , and calculated the Hausdorff dimension spectrum of level sets of some special multiple ergodic averages. Among others, they obtained the Hausdorff dimension of the following sets

$$A_\alpha := \left\{ (x_k)_1^\infty \in \Sigma : \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n x_k x_{2k} = \alpha \right\}.$$

**Theorem 1.1** (Keyon-Peres-Solomyak, Fan-Schmeling-Wu). *For any  $\alpha \in [0, 1]$ , we have*

$$\dim_H A_\alpha = -\log(1-p) - \frac{\alpha}{2} \log \frac{q(1-p)}{p(1-q)},$$

where

$$p^2(1-q) = (1-p)^3, \quad \alpha = \frac{2pq}{2+p-q}.$$

However, since the Lyapunov exponent is constant for the shift transformation on the symbolic space, what is obtained is in fact the entropy spectrum.

Let  $\{f_0, f_1\}$  be an iterated function system on  $[0, 1]$  given by

$$f_0(x) = e^{-\lambda_0}x, \quad f_1(x) = e^{-\lambda_1}x + c,$$

satisfying the open set condition. It has the usual symbolic description by  $\Sigma = \{0, 1\}^{\mathbb{N}}$  with a natural projection

$$\pi(\omega) = \lim_{n \rightarrow \infty} f_{\omega_1} \circ f_{\omega_2} \circ \dots \circ f_{\omega_n}(0).$$

Similarly to [PS, FSW], we consider in  $\Sigma$  subsets

$$S = \{\omega \in \Sigma : \omega_k \omega_{2k} = 0 \ \forall k \in \mathbb{N}\}$$

and

$$S_\alpha = \left\{ \omega \in \Sigma : \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \omega_k \omega_{2k} = \alpha \right\}.$$

Denote

$$L = \pi(S), \quad L_\alpha = \pi(S_\alpha).$$

We remark that if  $\lambda_0 = \lambda_1 = \log 2$ , i.e. the Lyapunov exponent is constant and  $\pi(\Sigma) = [0, 1]$ , then the Hausdorff dimensions of  $L, L_\alpha$  are the same as

those of  $S, S_\alpha$ . Our goal is to calculate the Hausdorff dimension of sets  $L$  and  $L_\alpha$  for  $\lambda_0 \neq \lambda_1$ .

Our results are as follows:

**Theorem 1.2.** *We have*

$$\dim_H L = \dim_H L_0 = -\frac{\log(1-p)}{\lambda_0},$$

where

$$p^{2\lambda_0} = (1-p)^{2\lambda_1+\lambda_0}.$$

For any  $\alpha \in (0, 1]$ , we have

$$\dim_H L_\alpha = \frac{\alpha \log \frac{p(1-q)}{(1-p)q} - 2 \log(1-p)}{2\lambda_0},$$

where

$$\alpha(\lambda_1 - \lambda_0) \log \frac{p(1-q)}{(1-p)q} + \lambda_0 \log \frac{p^2(1-q)}{1-p} - 2\lambda_1 \log(1-p) = 0$$

and

$$\frac{2pq}{2+p-q} = \alpha.$$

The paper is strongly related to [PS12a], we mostly repeat the calculations there in a more complicated situation. For the lacking details, in particular for [PS12a, Lemma 2] we refer the reader there. In the following two sections we calculate the lower bound: in Section 2 we introduce a family of measures and then we find the measure in this family that is supported on the set  $L_\alpha$  and has maximal Hausdorff dimension, in Section 3 we find a formula for this dimension. In Section 4 we check that this formula is also the upper bound for the dimension of  $L_\alpha$ .

## 2 Telescopic product measures

The same measures that were used to calculate the entropy spectrum (see [PS12a]) will be useful for the Hausdorff spectrum as well.

Let us start from the multiplicative golden shift case. Given  $p \in [0, 1]$ , let  $\mu_p$  be a probability measure on  $S$  given by

- if  $k$  is odd then  $\omega_k = 1$  with probability  $p$ ,

- if  $k$  is even and  $\omega_{k/2} = 0$  then  $\omega_k = 1$  with probability  $p$ ,
- if  $k$  is even and  $\omega_{k/2} = 1$  then  $\omega_k = 0$ .

Let  $\nu_p = \pi_*\mu_p$ . The Hausdorff dimension of  $L$  will turn out to be the supremum of Hausdorff dimensions of  $\nu_p$ .

Similarly, to deal with the spectrum of the sets  $L_\alpha$  we will define two parameter family of probabilistic measures. Given  $p, q \in [0, 1]$  we define  $\mu_{p,q}$  as a measure on  $\Sigma$  by

- if  $k$  is odd then  $\omega_k = 1$  with probability  $p$ ,
- if  $k$  is even and  $\omega_{k/2} = 0$  then  $\omega_k = 1$  with probability  $p$ ,
- if  $k$  is even and  $\omega_{k/2} = 1$  then  $\omega_k = 1$  with probability  $q$ .

Once again,  $\nu_{p,q} = \pi_*\mu_{p,q}$ . Please note that this notation is a little bit different than one in [PS12a]. Note also that  $\mu_p = \mu_{p,0}$ .

**Lemma 2.1.** *We have*

$$\mu_{p,q}(S_\alpha) = 1$$

for

$$\alpha = \frac{2pq}{2 + p - q}.$$

*Proof.* This lemma is proven in [PS12a, Lemma 3]. However, we will need this proof as a starting point for the proof of Lemma 2.2.

Denote

$$x_n(\omega) = \frac{2}{n} \sum_{k=n/2+1}^n \omega_k.$$

For a  $\mu_{p,q}$ -typical  $\omega$  the Law of Large Numbers implies

$$x_{2n}(\omega) = \frac{1}{2}p + \frac{x_n(\omega)}{2}q + \frac{1 - x_n(\omega)}{2}p + o(1).$$

Hence, as  $k \rightarrow \infty$ ,

$$x_{2^k n}(\omega) \rightarrow \frac{2p}{2 + p - q}.$$

By [PS12a, Lemma 5], it implies that  $\mu_{p,q}$ -almost surely

$$\lim_{n \rightarrow \infty} x_n(\omega) = \frac{2p}{2 + p - q}. \tag{2.1}$$

Then, for  $\mu_{p,q}$ -a.e.  $\omega$ ,

$$\frac{2}{n} \sum_{k=n/2+1}^n \omega_k \omega_{2k} = x_n(\omega)(q + o(1)) \rightarrow \frac{2pq}{2+p-q}.$$

Thus the assertion follows.  $\square$

Let us denote

$$H(p) = -p \log p - (1-p) \log(1-p)$$

with convention  $H(0) = H(1) = 0$ .

**Lemma 2.2.** *We have*

$$\dim_H \nu_p = \frac{2H(p)}{2p\lambda_1 + (2-p)\lambda_1}.$$

and

$$\dim_H \nu_{p,q} = \frac{(2-q)H(p) + pH(q)}{2p\lambda_1 + (2-p-q)\lambda_0}.$$

*Proof.* As  $\nu_p = \nu_{p,0}$ , it is enough to prove the second part of the assertion. For  $\omega \in \Sigma$  denote

$$C_n(\omega) = \{\tau \in \Sigma; \tau_k = \omega_k \forall k \leq n\}.$$

Let

$$h_n(\omega) := \log \mu_{p,q}(C_{2n}(\omega)) - \log \mu_{p,q}(C_n(\omega))$$

and

$$\lambda_n(\omega) := \log \text{diam } \pi(C_{2n}(\omega)) - \log \text{diam } \pi(C_n(\omega)).$$

By the Law of Large Numbers, for  $\mu_{p,q}$ -typical  $\omega$  and for big enough  $n$  we have

$$\frac{2}{n} h_n(\omega) = (2-x_n(\omega))(p \log p + (1-p) \log p) + x_n(\omega)(q \log q + (1-q) \log(1-q)) + o(1)$$

and

$$\frac{2}{n} \lambda_n(\omega) = (2-x_n(\omega))(-p\lambda_1 - (1-p)\lambda_0) + x_n(\omega)(-q\lambda_1 - (1-q)\lambda_0) + o(1).$$

Hence, by (2.1)

$$\frac{h_n(\omega)}{\lambda_n(\omega)} \rightarrow \frac{(2-q)H(p) + pH(q)}{2p\lambda_1 + (2-p-q)\lambda_0} \quad \mu_{p,q} - a.e.$$

Hence, for  $\mu_{p,q}$ -a.e.  $\omega$  we have

$$\lim_{n \rightarrow \infty} \frac{\log \nu_{p,q}(\pi(C_n(\omega)))}{\log \text{diam } \pi(C_n(\omega))} = \frac{(2-q)H(p) + pH(q)}{2p\lambda_1 + (2-p-q)\lambda_0}.$$

□

We will denote

$$\gamma_\alpha = \left\{ (p, q) \in [0, 1]^2 : \alpha = \frac{2pq}{2+p-q} \right\}.$$

**Lemma 2.3.** *The maximal Hausdorff dimension among measures  $\nu_p$  is achieved for  $p$  satisfying*

$$p^{2\lambda_0} = (1-p)^{2\lambda_1 + \lambda_0}. \quad (2.2)$$

For  $\alpha \in (0, 1)$ , the maximal Hausdorff dimension among measures  $\{\nu_{p,q} : (p, q) \in \gamma_\alpha\}$  is achieved for  $(p, q)$  satisfying

$$\alpha(\lambda_1 - \lambda_0) \log \frac{p(1-q)}{(1-p)q} + \lambda_0 \log \frac{p^2(1-q)}{1-p} - 2\lambda_1 \log(1-p) = 0. \quad (2.3)$$

Such  $(p, q)$  is unique in  $\gamma_\alpha$  and is always in  $(0, 1)^2$ .

*Proof.* Let us start from the second part of assertion. We need to find the maximum of the function

$$D(p, q) = \frac{(2-q)H(p) + pH(q)}{2p\lambda_1 + (2-p-q)\lambda_0}$$

over the curve  $\gamma_\alpha$ . For  $\alpha > 0$  this curve's endpoints are  $(1, 3\alpha/(2+\alpha))$  and  $(\alpha/(2+\alpha), 1)$ . Moreover, we have

$$d\alpha = \frac{2}{(2+p-q)^2} (q(2-q)dp + p(2+p)dq).$$

Hence, we need to solve the equation

$$p(2+p) \frac{\partial D}{\partial p} - q(2-q) \frac{\partial D}{\partial q} = 0.$$

After expanding the left hand side and collecting the terms, it turns out that it is divisible by  $p(2-q)$ . We get

$$\begin{aligned}
& (2pq\lambda_1 + (4 + 2p - 2q - 2pq)\lambda_0) \cdot \log p \\
& + ((-4 - 2p + 2q - 2pq)\lambda_1 + (-2 - p + q + 2pq)\lambda_0) \cdot \log(1 - p) \\
& + (-2pq\lambda_1 + 2pq\lambda_0) \cdot \log q \\
& + (2pq\lambda_1 + (2 + p - q - 2pq)\lambda_0) \cdot \log(1 - q) = 0.
\end{aligned} \tag{2.4}$$

It will be convenient to use  $\beta = 2/\alpha$ . As  $(p, q) \in \gamma_\alpha$ , we have

$$2 + p - q = \beta pq.$$

Substituting this into (2.4), we get

$$\begin{aligned}
& (2\lambda_1 + (2\beta - 2)\lambda_0) \log p + ((-2\beta - 2)\lambda_1 + (-\beta + 2)\lambda_0) \log(1 - p) \\
& + (-2\lambda_1 + 2\lambda_0) \log q + (2\lambda_1 + (\beta - 2)\lambda_0) \log(1 - q) = 0
\end{aligned} \tag{2.5}$$

and (2.3) follows.

To get the first part of assertion it is enough to remove all terms with  $q$  and substitute  $\alpha = 0$  into (2.3).

What remains is the third part of the assertion. Denoting by  $F(p, q)$  the left hand side of (2.5), we have

$$F(1, 3\alpha/(2 + \alpha)) = \infty$$

and

$$F(\alpha/(2 + \alpha), 1) = -\infty.$$

We will check that  $F$  restricted to  $\gamma_\alpha$  is strictly monotone. We have

$$p(p + 2) \frac{\partial F}{\partial p} - q(2 - q) \frac{\partial F}{\partial q} = \lambda_0((2\beta - 2)(p + 2) - 2(2 - q)) + \text{spt},$$

where spt stands for some positive terms (in particular, all the terms with  $\lambda_1$  are positive). However, as

$$(2\beta - 2)(p + 2) - 2(2 - q) = 2p + 2q + 2(\beta - 2)(p + 2),$$

the coefficient for  $\lambda_0$  is also positive. Hence,  $F$  restricted to  $\gamma_\alpha$  indeed has no extrema, so it must have only one zero.  $\square$



*Remark.* The curves  $\gamma_\alpha$  cover whole  $(0, 1)^2$ . However, not all pairs  $(p, q) \in (0, 1)^2$  are solutions of (2.5) for any  $\lambda_1, \lambda_0$ . Indeed, we can write (2.5) in the form

$$\frac{\lambda_1}{\lambda_0} a_1 + a_2 = 0$$

with

$$a_1 = \alpha \log p + (-2 - \alpha) \log(1 - p) - \alpha \log q + \alpha \log(1 - q)$$

and

$$a_2 = (2 - \alpha) \log p + (\alpha - 1) \log(1 - p) + \alpha \log q + (1 - \alpha) \log(1 - q).$$

Both  $a_1$  and  $a_2$  converge to  $\infty$  as  $p \rightarrow 1$  and to  $-\infty$  as  $q \rightarrow 1$ . They are also both strictly monotone on  $\gamma_\alpha$ , which can be checked like in the third part of the proof of Lemma 2.3 (using  $(2 - \alpha)(p + 2) > \alpha(2 - q)$  in case of  $a_2$ ), so they both have unique zeros. As the equation

$$r a_1 + a_2 = 0$$

can have positive solution only if  $a_1$  and  $a_2$  have different signs, only those  $(p, q) \in \gamma_\alpha$  between zeros of  $a_1$  and  $a_2$ , or equivalently satisfying

$$\alpha \log \frac{p(1 - q)}{(1 - p)q} > \max \left( 2 \log(1 - p), \log \frac{p^2(1 - q)}{1 - p} \right),$$

are solutions of (2.5) for some choice of  $\lambda_1, \lambda_0$ .

*Remark.* The measures  $\mu_{p,q}$  for  $p = q$  are Bernoulli. Each  $\gamma_\alpha$  intersects the diagonal  $\{p = q\}$  in exactly one point  $(\alpha^{1/2}, \alpha^{1/2})$  and at this point  $a_1 > 0, a_2 < 0$ . So, (2.5) has a Bernoulli measure as a solution for each  $\alpha \in (0, 1)$ . It happens when

$$\lambda_0 \log p = \lambda_1 \log(1 - p),$$

that is, when  $\nu_{\alpha^{1/2}, \alpha^{1/2}}$  is the Hausdorff measure (in dimension  $\dim_H \pi(\Sigma)$ ) on  $\pi(\Sigma)$ .

### 3 Exact formulas

To be able to provide the upper bounds in the following section, we need to substitute the results of Lemma 2.3 to Lemma 2.2 and obtain simpler formulas for our lower bound. We start with the golden shift case. Given  $\lambda_1, \lambda_0$  let  $p$  be given by (2.2).

**Lemma 3.1.** *We have*

$$\dim_H \nu_p = -\frac{\log(1-p)}{\lambda_0}.$$

*Proof.* By Lemma 2.2,

$$\dim_H \nu_p = \frac{2H(p)}{2p\lambda_1 + (2-p)\lambda_0}.$$

Applying (2.2) it is easy to check that

$$(2p\lambda_1 + (2-p)\lambda_0) \log(1-p) = -2H(p)\lambda_0$$

and the assertion follows.  $\square$

The calculations for the multifractal case are a little bit more complicated. Given  $\lambda_1, \lambda_0$ , and  $\alpha$ , let  $p, q$  be given by (2.3).

**Lemma 3.2.** *We have*

$$\dim_H \nu_{p,q} = \frac{\alpha \log \frac{p(1-q)}{(1-p)q} - 2 \log(1-p)}{2\lambda_0}. \quad (3.1)$$

*If  $\lambda_1 \neq \lambda_0$  then we have another formula:*

$$\dim_H \nu_{p,q} = \frac{\log \frac{p^2(1-q)}{(1-p)^3}}{2(\lambda_0 - \lambda_1)}. \quad (3.2)$$

*Proof.* By Lemma 2.2,

$$\dim_H \nu_{p,q} = \frac{(2-q)H(p) + pH(q)}{2p\lambda_1 + (2-p-q)\lambda_0}.$$

Using (2.3) one can check that

$$(2p\lambda_1 + (2-p-q)\lambda_0) \left( \alpha \log \frac{p(1-q)}{(1-p)q} - 2 \log(1-p) \right) = 2\lambda_0((2-q)H(p) + pH(q)).$$

This gives (3.1). Applying (2.3) once again we get

$$\dim_H \nu_{p,q} = \frac{\alpha \log \frac{p(1-q)}{(1-p)q} + \log \frac{1-p}{p^2(1-q)}}{2\lambda_1}. \quad (3.3)$$

Together with (3.1) this gives (3.2).  $\square$

## 4 Upper bounds

The last part of the proof is the upper bound.

**Lemma 4.1.** *We have*

$$\dim_H L \leq \sup_p \dim_H \nu_p$$

and, for all  $\alpha \in [0, 1]$ ,

$$\dim_H L_\alpha \leq \sup_{(p,q) \in \gamma_\alpha} \dim_H \nu_{p,q}.$$

*Proof.* As  $L \subset L_0$ , it is enough to prove the second part of the assertion. Fix  $\alpha$  and let  $\omega \in S_\alpha$ . Let  $p, q$  be as in (2.3). We denote for all  $n \in \mathbb{N}$

$$X_1^n = \#\{k \in [1, n] : \omega_k = 1\}$$

and for all even  $n \in \mathbb{N}$

$$X_{11}^n = \#\{k \in [1, n/2] : \omega_k = \omega_{2k} = 1\}.$$

We also denote

$$\tilde{h}_n = -\log \mu_{p,q}(C_n(\omega))$$

and

$$\tilde{l}_n = -\log \text{diam } \pi(C_n(\omega)).$$

For any even  $n$  we have

$$-\tilde{h}_n = n \log(1-p) + X_1^{n/2} \log \frac{1-q}{1-p} + X_1^n \log \frac{p}{1-p} - X_{11}^n \log \frac{p(1-q)}{(1-p)q},$$

see [PS12a, Section 4]. We also have

$$\tilde{l}_n = (\lambda_1 - \lambda_0)X_1^n + n\lambda_0.$$

Substituting (3.1) and (3.2) we get

$$\tilde{l}_n \dim_H \nu_{p,q} = -\frac{1}{2}X_1^n \log \frac{p^2(1-q)}{(1-p)^3} + \frac{n}{2} \left( \alpha \log \frac{p(1-q)}{(1-p)q} - 2 \log(1-p) \right).$$

Hence,

$$\frac{1}{n}(\tilde{l}_n \dim_H \nu_{p,q} - \tilde{h}_n) = \left( \frac{\alpha}{2} - \frac{X_{11}^n}{n} \right) \log \frac{p(1-q)}{(1-p)q} + \frac{1}{2} \left( \frac{X_1^{n/2}}{n/2} - \frac{X_1^n}{n} \right) \log \frac{1-q}{1-p}.$$

As the first summand converges to 0 and the second telescopes,

$$\liminf_{n \rightarrow \infty} \frac{1}{n}(\tilde{l}_n \dim_H \nu_{p,q} - \tilde{h}_n) \leq 0$$

and we are done. □

## References

- [FLM] Ai-hua Fan, Lingmin Liao, and Ji-hua Ma. Level sets of multiple ergodic averages. , (5 november 2011, pp. 1-10, doi:10.1007/s00605-011-0358-5). *Monatshe. Math.*, 148, 2011 online.
- [FSW11] Aihua Fan, Jörg Schmeling, and Meng Wu. Multifractal analysis of multiple ergodic averages. *C. R. Math. Acad. Sci. Paris*, 349(17-18):961–964, 2011.
- [FSW12] Aihua Fan, Jörg Schmeling, and Meng Wu. Multifractal analysis of multiple ergodic averages. preprint, 2012.
- [KPS11] Richard Kenyon, Yuval Peres, and Boris Solomyak. Hausdorff dimension of the multiplicative golden mean shift. *C. R. Math. Acad. Sci. Paris*, 349(11-12):625–628, 2011.
- [KPS12] Richard Kenyon, Yuval Peres, and Boris Solomyak. Hausdorff dimension for fractals invariant under the multiplicative integers. *Ergodic Theory Dynamical Systems*, arXiv:1102.5136, 2012.
- [PS12a] Yuval Peres and Boris Solomyak. Dimension spectrum for a non-conventional ergodic average. *Real Anal. Ex.*, arXiv:1107.1749, 2012.
- [PS12b] Yuval Peres and Boris Solomyak. The multiplicative golden mean shift has infinite hausdorff measure. In *Proceedings of the Conference on Fractals and Related Fields II*, volume arXiv:1201.5842. Springer, Berlin, 2012.
- [PSSS12] Yuval Peres, Jörg Schmeling, Stéphane Seuret, and Boris Solomyak. Dimensions of some fractals defined via the semi-group generated by 2 and 3. preprint, 2012.