



HAL
open science

Asymptotic estimates for the parabolic-elliptic Keller-Segel model in the plane

Juan Campos Serrano, Jean Dolbeault

► **To cite this version:**

Juan Campos Serrano, Jean Dolbeault. Asymptotic estimates for the parabolic-elliptic Keller-Segel model in the plane. 2010, pp.533-542. hal-00706194v1

HAL Id: hal-00706194

<https://hal.science/hal-00706194v1>

Submitted on 9 Jun 2012 (v1), last revised 10 Oct 2013 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Asymptotic estimates for the parabolic-elliptic Keller-Segel model in the plane

JUAN F. CAMPOS AND JEAN DOLBEAULT

ABSTRACT. We investigate the large-time behavior of the solutions of the two-dimensional Keller-Segel system in self-similar variables, when the total mass is subcritical, that is less than 8π after a proper adimensionalization. It was known from previous works that all solutions converge to stationary solutions, with exponential rate when the mass is small. Here we remove this restriction and show that the rate of convergence measured in relative entropy is exponential for any mass in the subcritical range, and independent of the mass. The proof relies on symmetrization techniques, which are adapted from a paper of J.I. Diaz, T. Nagai, and J.-M. Rakotoson, and allow us to establish uniform estimates for L^p norms of the solution. Exponential convergence is obtained by the mean of a linearization in a space which is defined consistently with relative entropy estimates and in which the linearized evolution operator is self-adjoint. The core of proof relies on several new spectral gap estimates which are of independent interest.

1. INTRODUCTION

Consider the two-dimensional parabolic-elliptic Keller-Segel system

$$(1.1) \quad \begin{cases} \frac{\partial u}{\partial t} = \Delta u - \nabla \cdot (u \nabla v) & x \in \mathbb{R}^2, \quad t > 0, \\ v = G_2 * u & x \in \mathbb{R}^2, \quad t > 0, \\ u(0, x) = n_0 \geq 0 & x \in \mathbb{R}^2, \end{cases}$$

where G_2 denotes the Green function associated to $-\Delta$ on \mathbb{R}^2 :

$$G_2(x) := -\frac{1}{2\pi} \log |x|, \quad x \in \mathbb{R}^2.$$

The equation for the *mass density* u is parabolic, while the *chemo-attractant density* v solves an (elliptic) Poisson equation: $-\Delta v = u$. The drift term corresponds to an attractive mean-field nonlinearity, which has attracted lots of attention in mathematical biology in the recent years: see [24, 25, 26, 36, 38, 39] for some recent overviews. According to [27, 23, 10, 15], it is known that if

$$(1.2) \quad n_0 \in L^1_+(\mathbb{R}^2, (1 + |x|^2) dx), \quad n_0 |\log n_0| \in L^1(\mathbb{R}^2) \quad \text{and} \quad M := \int_{\mathbb{R}^2} n_0 dx < 8\pi,$$

then there exists a solution u , in the sense of distributions, that is global in time and such that $M = \int_{\mathbb{R}^2} u(t, x) dx$ is conserved along the evolution in the euclidean space \mathbb{R}^2 . There is no non-trivial stationary solution of (1.1) and any solution converges to zero locally as time gets large. In order to study the asymptotic behavior of u , it is convenient to work in self-similar variables. We define $R(t) := \sqrt{1 + 2t}$, $\tau(t) := \log R(t)$, and the rescaled functions n and c by

$$u(t, x) := R^{-2}(t) n(\tau(t), R^{-1}(t)x) \quad \text{and} \quad v(t, x) := c(\tau(t), R^{-1}(t)x).$$

Key words and phrases. Keller-Segel model; chemotaxis; large time asymptotics; subcritical mass; self-similar solutions; relative entropy; free energy; Lyapunov functional; spectral gap; logarithmic Hardy-Littlewood-Sobolev inequality.

2010 *Mathematics subject classification.* 35B40, 92C17

This time-dependent rescaling is the one of the heat equation. We observe that the non-linear term is also invariant under such a rescaling. The rescaled systems reads

$$(1.3) \quad \begin{cases} \frac{\partial n}{\partial t} = \Delta n + \nabla \cdot (n x) - \nabla \cdot (n \nabla c) & x \in \mathbb{R}^2, \quad t > 0, \\ c = G_2 * n & x \in \mathbb{R}^2, \quad t > 0, \\ n(0, x) = n_0 \geq 0 & x \in \mathbb{R}^2. \end{cases}$$

Under Assumptions (1.2), it has been shown in [10, Theorem 1.2] that

$$\lim_{t \rightarrow \infty} \|n(t, \cdot) - n_\infty\|_{L^1(\mathbb{R}^2)} = 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} \|\nabla c(t, \cdot) - \nabla c_\infty\|_{L^2(\mathbb{R}^2)} = 0,$$

where (n_∞, c_∞) solves

$$(1.4) \quad n_\infty = M \frac{e^{c_\infty - |x|^2/2}}{\int_{\mathbb{R}^2} e^{c_\infty - |x|^2/2} dx} \quad \text{with} \quad c_\infty = G_2 * n_\infty.$$

Moreover, n_∞ is smooth and radially symmetric. Existence of a solution to (1.4) has been established in [2] by ordinary differential equation techniques and in [34] by partial differential equation methods. The uniqueness has been shown in [5]. To recall the dependence of n_∞ in M , we will write it as $n_{\infty, M}$ whenever needed.

A simple computation of the second moment shows that smooth solutions with mass larger than 8π blow-up in finite time; see for instance [27]. The case $M = 8\pi$ has been extensively studied. We shall refer to [5, 6, 7] for some recent papers on this topic. The asymptotic regime is of a very different nature in such a critical case. In the present paper, we shall restrict our purpose to the sub-critical case $M < 8\pi$.

In [8] it has been proved that there exists a positive mass $M_\star \leq 8\pi$ such that for any initial data $n_0 \in L^2(n_\infty^{-1} dx)$ of mass $M < M_\star$ satisfying (1.2), System (1.3) has a unique solution n such that

$$\int_{\mathbb{R}^2} |n(t, x) - n_\infty(x)|^2 \frac{dx}{n_\infty(x)} \leq C e^{-\delta t} \quad \forall t \geq 0$$

for some positive constants C and δ . Moreover δ can be taken arbitrarily close to 1 as $M \rightarrow 0$. If $M < 8\pi$, we may notice that the condition $n_0 \in L^2(n_\infty^{-1} dx)$ is stronger than (1.2). Our main result is that $M_\star = 8\pi$ and $\delta \geq 1$, at least for a large subclass of solutions with initial datum n_0 satisfying the following technical assumption:

$$(1.5) \quad \exists \varepsilon \in (0, 8\pi - M) \quad \text{such that} \quad \int_0^s n_{0,*}(\sigma) d\sigma \leq \int_{B(0, \sqrt{s/\pi})} n_{\infty, M+\varepsilon}(x) dx \quad \forall s \geq 0.$$

Here $n_{0,*}(\sigma)$ stands for the symmetrized function associated to n_0 . Details will be given in Section 2.

Theorem 1. *Assume that n_0 satisfies (1.5),*

$$n_0 \in L^2_+(n_\infty^{-1} dx) \quad \text{and} \quad M := \int_{\mathbb{R}^2} n_0 dx < 8\pi.$$

Then any solution of (1.3) with initial datum n_0 is such that

$$\int_{\mathbb{R}^2} |n(t, x) - n_\infty(x)|^2 \frac{dx}{n_\infty(x)} \leq C e^{-2t} \quad \forall t \geq 0$$

for some positive constant C , where n_∞ is the unique stationary solution to (1.4) with mass M .

This result is consistent with the recent results of [11] for the two-dimensional radial model and its one-dimensional counterpart (see Proposition 16 for more comments). For completeness, let us mention that results of exponential convergence for problems with mean field have been

obtained earlier in [16, 17], but only for interaction potentials involving much smoother kernels than G_2 .

Our paper is organized as follows. In Section 2, we will apply symmetrization techniques as in [20, 21] to establish uniform estimates on $\|n\|_{L^p(\mathbb{R}^2)}$. Then we will prove the uniform convergence of n to n_∞ using Duhamel's formula: see Corollary 7 in Section 3. Section 4 is devoted to the linearization of the problem around n_∞ and to the study of the spectral gap of the linearized operator. A strict positivity result for the linearized entropy is also needed and will be proved in Section 5. The proof of Theorem 1 is completed in the last section. It is based on two estimates: Theorems 15 and 17 (also see Corollary 18) that are established in Sections 4 and 5 respectively. Some of the results of Sections 4 and 5 (see Theorem 15 and Corollary 18) have been announced without proof in [12] in connection with a new Onofri type inequality, whose linearized form is given in Inequality (5.2).

2. SYMMETRIZATION

In this section, we adapt the results of J.I. Diaz, T. Nagai, and J.-M. Rakotoson in [21] to the setting of self-similar variables. Several key estimates are based on an earlier work of J.I. Diaz and T. Nagai for the bounded domain case: see [20]. We shall therefore only sketch the main steps of the method and emphasize the necessary changes.

To any measurable function $u : \mathbb{R}^2 \mapsto [0, +\infty)$, we associate the distribution function defined by $\mu(t, \tau) := |\{u > \tau\}|$ and its decreasing rearrangement given by

$$u_* : [0, +\infty) \rightarrow [0, +\infty] , \quad s \mapsto u_*(s) = \inf\{\tau \geq 0 : \mu(t, \tau) \leq s\} .$$

We adopt the following convention: for any time-dependent function $u : (0, +\infty) \times \mathbb{R}^2 \rightarrow [0, +\infty)$, we will also denote by u_* the decreasing rearrangement of u with respect to the spatial variable, that is, $u_*(t, s) = u(t, \cdot)_*(s)$.

Rearrangement techniques are a standard tool in the study of partial differential equations: in the framework of chemotaxis, see for instance [1, 18, 33] in case of bounded domains, and [21] for unbounded domains. Let us briefly recall some properties of the decreasing rearrangement:

- (a) For every measurable function $F : \mathbb{R}^+ \mapsto \mathbb{R}^+$, we have

$$\int_{\mathbb{R}^2} F(u) \, dx = \int_0^\infty F(u_*) \, ds .$$

In particular, if $u \in L^p(\mathbb{R}^N)$ with $1 \leq p \leq \infty$, then $\|u\|_{L^p(\mathbb{R}^N)} = \|u_*\|_{L^p(\mathbb{R}^N)}$.

- (b) If $u \in W^{1,q}(0, T; L^p(\mathbb{R}^N))$ is a nonnegative function, with $1 \leq p < \infty$ and $1 \leq q \leq \infty$, then $u_* \in W^{1,q}(0, T; L^p(0, \infty))$ and the formula

$$\int_0^{\mu(t, \tau)} \frac{\partial u_*}{\partial t}(t, \sigma) \, d\sigma = \int_{\{u(t, \cdot) > \tau\}} \frac{\partial u}{\partial t}(t, x) \, dx$$

holds for almost every $t \in (0, T)$. Here $\mu(t, \tau)$ denotes $|\{u(t, \cdot) > \tau\}|$. See [21, Theorem 2.2, (ii), p. 167] for a statement and a proof.

As in [21], let us consider a solution (n, c) of (1.3) and define

$$k(t, s) := \int_0^s n_*(t, \sigma) \, d\sigma$$

The strategy consists in finding a differential inequality for $k(t, s)$. Then, using a comparison principle, we will obtain an upper bound on the L^p norm of n . In [21], the method was applied to (1.1). Here we adapt it to the solution in rescaled variables, that is (1.3).

Lemma 2. *If n is a solution of (1.3) with initial datum n_0 satisfying the assumptions of Theorem 1, then the function $k(t, s)$ satisfies*

$$k \in L^\infty([0, +\infty) \times (0, +\infty)) \cap H^1\left([0, +\infty); W_{\text{loc}}^{1,p}(0, +\infty)\right) \cap L^2\left([0, +\infty); W_{\text{loc}}^{2,p}(0, +\infty)\right)$$

and

$$\begin{cases} \frac{\partial k}{\partial t} - 4\pi s \frac{\partial^2 k}{\partial s^2} - (k + 2s) \frac{\partial k}{\partial s} \leq 0 & \text{a.e. in } (0, +\infty) \times (0, +\infty), \\ k(t, 0) = 0, \quad k(t, +\infty) = \int_{\mathbb{R}^2} n_0 dx & \text{for } t \in (0, +\infty), \\ k(0, s) = \int_0^s (n_0)_* d\sigma & \text{for } s \geq 0. \end{cases}$$

Proof. The proof follows the method of [21, Proposition 3.1]. We will therefore only sketch the main steps that are needed to adapt the results to the setting of self-similar variables and refer to [21] for all technical details.

For $\tau \in (0, n_*(t, 0))$ and $h > 0$, define the truncation function $T_{\tau, h}$ on $(-\infty, +\infty)$ by

$$T_{\tau, h} = \begin{cases} 0 & \text{if } s \leq \tau \\ s - \tau & \text{if } \tau < s \leq \tau + h \\ h & \text{if } \tau + h < s \end{cases}$$

and observe that $T_{\tau, h}(n(t, \cdot))$ belongs to $W^{1,p}(\mathbb{R}^2)$ since $n(t, \cdot) \in W^{1,p}(\mathbb{R}^2)$ and $T_{\tau, h}$ is Lipschitz continuous. Now we integrate (1.3) against $T_{\tau, h}(n)$ over \mathbb{R}^2 , and integrate by parts to obtain

$$\int_{\mathbb{R}^2} \frac{\partial n}{\partial t} T_{\tau, h}(n) dx + \int_{\mathbb{R}^2} \nabla n \nabla T_{\tau, h}(n) dx = \int_{\mathbb{R}^2} n (\nabla c - x) \nabla T_{\tau, h}(n) dx.$$

We have that $|\{n = \tau\}| = 0$ for almost every $\tau \geq 0$. Hence one can prove that

$$\lim_{h \rightarrow 0} \frac{1}{h} \int_{\mathbb{R}^2} \frac{\partial n}{\partial t} T_{\tau, h}(n) dx = \int_{\{n > \tau\}} \frac{\partial n}{\partial t}(t, x) dx = \frac{\partial k}{\partial t}(t, \mu(t, \tau)).$$

Next we observe that

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{1}{h} \int_{\mathbb{R}^2} \nabla n \nabla T_{\tau, h}(n) dx &= \lim_{h \rightarrow 0} \frac{1}{h} \left(\int_{\{n > \tau\}} |\nabla n|^2 dx - \int_{\{n > \tau + h\}} |\nabla n|^2 dx \right) \\ &= \frac{\partial}{\partial \tau} \int_{\{n > \tau\}} |\nabla n|^2 dx. \end{aligned}$$

Consider the function

$$\Phi_{\tau, h} = \int_0^s \sigma \frac{\partial T_{\tau, h}}{\partial \sigma}(\sigma) d\sigma = \begin{cases} 0 & \text{if } s \leq \tau, \\ \frac{1}{2}(s^2 - \tau^2) & \text{if } \tau < s \leq \tau + h, \\ h(\tau + \frac{h}{2}) & \text{if } \tau + h < s. \end{cases}$$

Integrating the Poisson equation for c against $\Phi_{\tau, h}(n)$, we get

$$\int_{\mathbb{R}^2} \nabla c \nabla \Phi_{\tau, h}(n) dx = \int_{\mathbb{R}^2} n \nabla c \nabla T_{\tau, h}(n) dx = \int_{\mathbb{R}^2} n \Phi_{\tau, h}(n) dx,$$

thus proving that

$$\begin{aligned} & \lim_{h \rightarrow 0^+} \frac{1}{h} \int_{\mathbb{R}^2} n \nabla c \nabla T_{\tau, h}(n) \, dx \\ &= \lim_{h \rightarrow 0^+} \left(\frac{1}{2h} \int_{\{\tau < n \leq \tau + h\}} n (n^2 - \tau^2) \, dx + \int_{\{n > \tau + h\}} n \left(\tau + \frac{h}{2} \right) \, dx \right) \\ &= \tau \int_{\{n > \tau\}} n \, dx = \frac{\partial k}{\partial s}(t, \mu(t, \tau)) k(t, \mu(t, \tau)) \end{aligned}$$

since $\tau = n_*(t, \mu(t, \tau)) = \frac{\partial k}{\partial s}(t, \mu(t, \tau))$ and $\int_{\{n > \tau\}} n \, dx = \int_0^{\mu(t, \tau)} n_*(t, s) \, ds = k(t, \mu(t, \tau))$. On the other hand,

$$\begin{aligned} \lim_{h \rightarrow 0^+} \frac{1}{h} \int_{\mathbb{R}^2} n(x) x \cdot \nabla T_{\tau, h}(x) \, dx \, dx &= \lim_{h \rightarrow 0^+} \frac{1}{h} \int_{\mathbb{R}^2} x \cdot \nabla \Phi_{\tau, h}(n) \, dx = - \lim_{h \rightarrow 0^+} \frac{2}{h} \int_{\mathbb{R}^2} \Phi_{\tau, h}(n) \, dx \\ &= -2\tau |\{n > \tau\}| = -2 \frac{\partial k}{\partial s}(t, \mu(t, \tau)) \mu(t, \tau) . \end{aligned}$$

Using the inequality

$$4\pi \mu(t, \tau) \leq \frac{\partial \mu}{\partial \tau}(t, \tau) \frac{\partial}{\partial \tau} \int_{\{n > \tau\}} |\nabla n|^2 \, dx ,$$

(see [20, Proof of Lemma 4, p. 669], and also [33, pp. 25-26] or [18, p. 20], and [37] for an earlier reference) we obtain

$$1 \leq - \frac{\frac{\partial \mu}{\partial \tau}(t, \tau)}{4\pi \mu(t, \tau)} \left(- \frac{\partial k}{\partial t}(t, \mu(t, \tau)) + \frac{\partial k}{\partial s}(t, \mu(t, \tau)) (k(t, \mu(t, \tau)) + 2\mu(t, \tau)) \right)$$

for almost every $\tau \in (0, n_*(t, 0))$. Integrating over $(\tau_1, \tau_2) \subset (0, n_*(t, 0))$, as in [19, Lemma 2], we get

$$\frac{1}{4\pi} \int_{\mu(t, \tau_1)}^{\mu(t, \tau_2)} \left(- \frac{\partial k}{\partial t}(t, s) + \frac{\partial k}{\partial s}(t, s) (k(t, s) + 2s) \right) \frac{ds}{s} \leq \tau_1 - \tau_2$$

where

$$\tau_1 - \tau_2 = \frac{\partial k}{\partial s}(t, \mu(t, \tau_1)) - \frac{\partial k}{\partial s}(t, \mu(t, \tau_2)) .$$

Hence dividing by $(\mu(t, \tau_2) - \mu(t, \tau_1))$ and then taking the limit completes the proof. \square

The next result is adapted from [20, Proposition A.1, p. 676] and [21, Proposition 3.2, p. 172]. Although it is unnecessarily general for our purpose, as the function g below is extremely well defined (and independent of t), we keep it as in J.I. Diaz *et al.* and give a sketch of the proof, for completeness.

Proposition 3. *Let f, g be two continuous functions on $Q = \mathbb{R}^+ \times (0, +\infty)$ such that*

- (i) $f, g \in L^\infty(Q) \cap L^2(0, +\infty; W_{\text{loc}}^{2,2}(0, +\infty))$, $\frac{\partial f}{\partial t}, \frac{\partial g}{\partial t} \in L^2(0, +\infty; L_{\text{loc}}^2(0, +\infty))$,
- (ii) $\left| \frac{\partial f}{\partial s}(t, s) \right| \leq C(t)$ and $\left| \frac{\partial g}{\partial s}(t, s) \right| \leq C(t) \max\{s^{-1/2}, 1\}$, for some continuous function $t \mapsto C(t)$ on \mathbb{R}^+ .

If f and g satisfy

$$\left\{ \begin{array}{l} \frac{\partial f}{\partial t} - 4\pi s \frac{\partial^2 f}{\partial s^2} - (f + 2s) \frac{\partial f}{\partial s} \leq \frac{\partial g}{\partial t} - 4\pi s \frac{\partial^2 g}{\partial s^2} - (g + 2s) \frac{\partial g}{\partial s} \text{ a.e. in } Q , \\ f(t, 0) = 0 = g(t, 0) \quad \text{and} \quad f(t, +\infty) \leq g(t, +\infty) \text{ for any } t \in (0, +\infty) , \\ f(0, s) \leq g(0, s) \text{ for } s \geq 0 , \text{ and } g(t, s) \geq 0 \text{ in } Q , \end{array} \right.$$

then $f \leq g$ on Q .

Proof. Take $w = f - g$. We have

$$\frac{\partial w}{\partial t} - 4\pi s \frac{\partial^2 w}{\partial s^2} - 2s \frac{\partial w}{\partial s} \leq w \frac{\partial f}{\partial s} + g \frac{\partial w}{\partial s}.$$

Multiplying by w_+/s , and integrating over (δ, L) with $0 < \delta < 1 < L$, we obtain

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} \int_{\delta}^L \frac{w_+^2}{s} ds + 4\pi \int_{\delta}^L \left(\frac{\partial w_+}{\partial s} \right)^2 ds - \left[4\pi \frac{\partial w_+}{\partial s}(t, s) w_+(t, s) \right]_{s=\delta}^{s=L} - \int_{\delta}^L \frac{\partial}{\partial s} (w_+^2) ds \\ \leq \int_{\delta}^L \left(w_+^2 \frac{\partial f}{\partial s} + w_+ \frac{\partial w}{\partial s} g \right) \frac{ds}{s} \end{aligned}$$

thus showing that

$$\frac{1}{2} \frac{\partial}{\partial t} \int_{\delta}^L \frac{w_+^2}{s} ds + 4\pi \int_{\delta}^L \left(\frac{\partial w_+}{\partial s} \right)^2 ds \leq C(t) \int_{\delta}^L \frac{w_+^2}{s} ds + \int_{\delta}^L \frac{w_+}{s} \frac{\partial w}{\partial s} g ds + G(t, \delta, L),$$

where $G(t, \delta, L)$ uniformly (with respect to $t \geq 0$) converges to 0 as $\delta \rightarrow 0$ and $L \rightarrow +\infty$. Now using the fact that $g(t, s)/\sqrt{s} \leq C(t)$ we obtain that, for some constant $K > 0$,

$$\int_{\delta}^L \frac{w_+}{s} \frac{\partial w}{\partial s} g ds \leq 4\pi \int_{\delta}^L \left(\frac{\partial w_+}{\partial s} \right)^2 ds + K C^2(t) \int_{\delta}^L \frac{w_+^2}{s} ds,$$

yielding

$$\frac{1}{2} \frac{\partial}{\partial t} \int_{\delta}^L \frac{w_+^2}{s} ds \leq (1 + K C(t)) C(t) \int_{\delta}^L \frac{w_+^2}{s} ds + G(t, \delta, L).$$

From Gronwall's lemma and $w_+(0, s) = 0$, with $R(t) := 2 \int_0^t (1 + K C(\tau)) C(\tau) d\tau$, it follows that

$$\int_{\delta}^L \frac{w_+^2}{s} ds \leq 2 e^{R(t)} \int_0^t e^{-R(\tau)} G(\tau, \delta, L) d\tau.$$

Taking the limit as $\delta \rightarrow 0$ and $L \rightarrow +\infty$, we obtain

$$\int_0^{\infty} \frac{w_+^2}{s} ds \leq 0,$$

which implies $f \leq g$ and concludes the proof. \square

Using Lemma 2 and Proposition 3, we can now establish uniform bounds on $\|n\|_{L^p(\mathbb{R}^2)}$ and $\|\nabla c\|_{L^\infty(\mathbb{R}^2)}$.

Theorem 4. *Assume that $n_0 \in L_+^2(n_\infty^{-1} dx)$ satisfies (1.5) and $M := \int_{\mathbb{R}^2} n_0 dx < 8\pi$. Then there exist positive constants $C_1 = C_1(M, p)$ and $C_2 = C_2(M, p)$ such that*

$$\|n\|_{L^p(\mathbb{R}^2)} \leq C_1 \quad \text{and} \quad \|\nabla c\|_{L^\infty(\mathbb{R}^2)} \leq C_2.$$

Proof. The function $M_\varepsilon(s) := \int_{B(0, \sqrt{s/\pi})} n_{\infty, M+\varepsilon} dx$ satisfies

$$4\pi s M_\varepsilon'' + 2s M_\varepsilon' + M_\varepsilon M_\varepsilon' = 0.$$

By direct application of Proposition 3, we obtain

$$k(t, s) \leq M_\varepsilon(s) \quad \forall (t, s) \in Q.$$

By [1, p. 74] or [18, Lemma 1.33], we deduce

$$\|n_*\|_{L^p(0, \infty)} \leq \|M_\varepsilon'\|_{L^p(0, \infty)},$$

which yields the result. More details on M_ε and *cumulated densities* will be given in Section 4.5. \square

3. UNIFORM CONVERGENCE

With the boundedness results of Section 2 in hands, we can now prove a result of uniform convergence for n and ∇c , if (n, c) is given as a solution of (1.3) satisfying the assumptions of Theorem 1.

Consider the kernel associated to the Fokker-Planck equation

$$K(t, x, y) := \frac{1}{2\pi(1 - e^{-2t})} e^{-\frac{1}{2} \frac{|x - e^{-t}y|^2}{1 - e^{-2t}}} \quad x \in \mathbb{R}^2, \quad y \in \mathbb{R}^2, \quad t > 0.$$

This definition deserves some explanations. If n is a solution of

$$\frac{\partial n}{\partial t} = \Delta n + \nabla \cdot (n x)$$

with initial datum n_0 , then $u(\tau, \xi) = R^{-2} n(\log R, R^{-1} \xi)$ with $R = R(\tau) = \sqrt{1 + 2\tau}$ is a solution of the heat equation

$$\frac{\partial u}{\partial \tau} = \Delta u, \quad u(\tau = 0, \cdot) = n_0,$$

whose solution is given by

$$u(\xi, \tau) = \frac{1}{4\pi\tau} \int_{\mathbb{R}^2} e^{-\frac{|\xi - y|^2}{4\tau}} n_0(y) dy.$$

By undoing the change of variables, we get that the solution of the Fokker-Planck equation is given by

$$n(t, x) = \int_{\mathbb{R}^2} K(t, x, y) n_0(y) dy.$$

Consider now a solution of (1.3). We have the following Duhamel formula.

Lemma 5. *Assume that n is a solution of (1.3) with initial data satisfying (1.2). Then for any $t > 0$, $x \in \mathbb{R}^2$, we have*

$$n(t, x) = \int_{\mathbb{R}^2} K(t, x, y) n_0(y) dy + \int_0^t \int_{\mathbb{R}^2} \nabla_x K(t - s, x, y) \cdot n(s, y) \nabla c(s, y) dy ds.$$

This is a standard fact whose proof relies on the fact that $(t, x) \mapsto K(t, x, y)$ is a solution of the Fokker-Planck equation with a δ -Dirac function initial value. Details are left to the reader.

Using the semi-group property, the expression for $n(2t, x)$ found in Lemma 5 can be written in terms of $n(t, x)$ as

$$n(2t, x) = \int_{\mathbb{R}^2} K(t, x, y) n(t, y) dy + \int_t^{2t} \int_{\mathbb{R}^2} \nabla_x K(2t - s, x, y) \cdot n(s, y) \nabla c(s, y) dy ds$$

for any $t \geq 0$. Since n_∞ is a stationary solution, we can also write that

$$n_\infty(x) = \int_{\mathbb{R}^2} K(t, x, y) n_\infty(y) dy + \int_t^{2t} \int_{\mathbb{R}^2} \nabla_x K(2t - s, x, y) \cdot n_\infty(y) \nabla c_\infty(y) dy ds$$

for any $t \geq 0$. By taking the difference of the two expressions written for $n(t, x)$ and n_∞ respectively, we get that

$$\begin{aligned} n(2t, x) - n_\infty(x) &= \int_{\mathbb{R}^2} K(t, x, y) (n(t, y) - n_\infty(y)) dy ds \\ &\quad + \int_t^{2t} \int_{\mathbb{R}^2} \nabla_x K(2t - s, x, y) \cdot (n(s, y) \nabla c(s, y) - n_\infty(y) \nabla c_\infty(y)) dy. \end{aligned}$$

This provides a straightforward estimate, which goes as follows:

$$\begin{aligned} \|n(2t, \cdot) - n_\infty\|_{L^\infty(\mathbb{R}^2)} &\leq \|K(t, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^2 \times \mathbb{R}^2)} \|n(t, \cdot) - n_\infty\|_{L^1(\mathbb{R}^2)} \\ &\quad + \int_0^t \|\nabla K(s, \cdot, \cdot)\|_{L^\infty(\mathbb{R}_y^2; L^r(\mathbb{R}_x^2))} ds + \mathcal{R}(t) \end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1$ with $p \in [1, \infty)$, $q \in [2, \infty)$ and $r \in (1, 2)$, and where

$$\mathcal{R}(t) := \sup_{s \in (t, 2t)} \left(\|n(s, \cdot)\|_{L^p(\mathbb{R}^2)} \|\nabla c(s, \cdot) - \nabla c_\infty\|_{L^q(\mathbb{R}^2)} + \|n(s, \cdot) - n_\infty\|_{L^p(\mathbb{R}^2)} \|\nabla c_\infty\|_{L^q(\mathbb{R}^2)} \right)$$

converges to 0 by Theorem 4 and the fact that

$$\lim_{t \rightarrow \infty} \|n(t, \cdot) - n_\infty\|_{L^1(\mathbb{R}^2)} = 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} \|\nabla c(t, \cdot) - \nabla c_\infty\|_{L^2(\mathbb{R}^2)} = 0$$

according to [10, Theorem 1.2]. Hence we have shown the *uniform convergence* of n towards n_∞ as $t \rightarrow \infty$ and

$$\lim_{t \rightarrow \infty} \|n(t, \cdot) - n_\infty\|_{L^p(\mathbb{R}^2)} = 0$$

for any $p \in [1, \infty]$, by Hölder's interpolation. As for the convergence of $\nabla c(t, \cdot)$ towards ∇c_∞ as $t \rightarrow \infty$ in $L^q(\mathbb{R}^2)$ for $q \in (2, \infty]$, we need one more interpolation inequality.

Lemma 6. *If $h = (-\Delta)^{-1}\rho$ for some function $\rho \in L^{2-\varepsilon} \cap L^{2+\varepsilon}(\mathbb{R}^2)$, with $\varepsilon \in (0, 1)$, then there exists an explicit positive constant $C = C(\varepsilon)$ such that*

$$\|\nabla h\|_{L^\infty(\mathbb{R}^2)} \leq C \left(\|\rho\|_{L^{2-\varepsilon}(\mathbb{R}^2)} + \|\rho\|_{L^{2+\varepsilon}(\mathbb{R}^2)} \right).$$

Proof. This follows by a direct computation. We can estimate $|\nabla h|$ by

$$|\nabla h(x)| = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{\rho(y)}{|x-y|} dy$$

for any $x \in \mathbb{R}^2$ and split the integral into two pieces corresponding to $|x-y| < 1$ and $|x-y| \geq 1$: by Hölder's inequality, we obtain that

$$\frac{1}{2\pi} \int_{|x-y| < 1} \frac{\rho(y)}{|x-y|} dy \leq C_1(\varepsilon) \|\rho\|_{L^{2+\varepsilon}(\mathbb{R}^2)}$$

with $C_1(\varepsilon) = \frac{1}{2\pi} (2\pi(1+\varepsilon)/\varepsilon)^{(1+\varepsilon)/(2+\varepsilon)}$ and

$$\frac{1}{2\pi} \int_{|x-y| \geq 1} \frac{\rho(y)}{|x-y|} dy \leq C_2(\varepsilon) \|\rho\|_{L^{2-\varepsilon}(\mathbb{R}^2)}$$

with $C_2(\varepsilon) = \frac{1}{2\pi} (2\pi(1-\varepsilon)/\varepsilon)^{(1-\varepsilon)/(2-\varepsilon)}$. The conclusion holds with $C = \max_{i=1,2} C_i$. \square

Hence we have also shown the *uniform convergence* of ∇c towards ∇c_∞ as $t \rightarrow \infty$. By Hölder's interpolation, the convergence holds in $L^q(\mathbb{R}^2)$ for any $q \in [2, \infty]$. Summarizing all results of this section, we have shown the following limits.

Corollary 7. *Assume that n is a solution of (1.3) with initial data satisfying the assumptions of Theorem 1. Then*

$$\lim_{t \rightarrow \infty} \|n(t, \cdot) - n_\infty\|_{L^p(\mathbb{R}^2)} = 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} \|\nabla c(t, \cdot) - \nabla c_\infty\|_{L^q(\mathbb{R}^2)} = 0$$

for any $p \in [1, \infty]$ and any $q \in [2, \infty]$.

4. SPECTRAL GAP OF THE LINEARIZED OPERATOR \mathcal{L}

Assume that n is a solution of (1.3) and consider f and g defined for any $(t, x) \in \mathbb{R}^+ \times \mathbb{R}^2$ by

$$n(t, x) = n_\infty(x) (1 + f(t, x)) \quad \text{and} \quad c(t, x) = c_\infty(x) (1 + g(t, x)) .$$

Then (f, g) is a solution of the nonlinear problem

$$(4.1) \quad \begin{cases} \frac{\partial f}{\partial t} - \mathcal{L} f = -\frac{1}{n_\infty} \nabla \cdot [f n_\infty \nabla (g c_\infty)] & x \in \mathbb{R}^2, \quad t > 0, \\ -\Delta (g c_\infty) = f n_\infty & x \in \mathbb{R}^2, \quad t > 0, \end{cases}$$

where \mathcal{L} is the linear operator

$$\mathcal{L} f = \frac{1}{n_\infty} \nabla \cdot [n_\infty \nabla (f - g c_\infty)] .$$

The goal of this section is to establish that \mathcal{L} has a spectral gap in an appropriate functional setting. To characterize the spectrum of \mathcal{L} , it is indeed necessary to specify the domain of the operator \mathcal{L} . Heuristically, it is simpler to identify the eigenfunctions corresponding to the lowest eigenvalues and define only afterwards the norm for which \mathcal{L} turns out to be self-adjoint. We will

- identify some eigenfunctions of the linearized Keller-Segel operator \mathcal{L} in Section 4.1,
- characterize the kernel of \mathcal{L} in Section 4.2,
- determine an adapted functional setting for \mathcal{L} and related operators in Section 4.3,
- show that the spectrum of \mathcal{L} is discrete in Section 4.4,
- and finally establish a spectral gap inequality in Section 4.5.

4.1. Some eigenfunctions of the linearized Keller-Segel operator \mathcal{L} . Using the fact that n_∞ depends on $x = (x_1, x_2) \in \mathbb{R}^2$ and on the mass parameter M , we observe that the functions

$$\begin{aligned} f_{0,0} &= \partial_M \log n_{\infty, M} , \\ f_{1,i} &= \partial_{x_i} \log n_{\infty, M} , \quad i = 1, 2 , \\ f_{0,1} &= x \cdot \nabla \log n_{\infty, M} , \end{aligned}$$

are eigenfunctions of \mathcal{L} . Here $\partial_M n_{\infty, M}$ denotes the derivative of the function $n_\infty = n_{\infty, M}$ with respect to the mass parameter M , while ∂_{x_i} stands for $\partial/\partial x_i$. We shall use two indices for the numbering of the eigenfunctions because of a spherical harmonics decomposition that will be studied in Section 4.5. A precise statements goes as follows.

Lemma 8. *With the above notations, we have*

$$\begin{aligned} \mathcal{L} f_{0,0} &= 0 , \\ \mathcal{L} f_{1,i} &= -f_{1,i} , \\ \mathcal{L} f_{0,1} &= -2 f_{0,1} . \end{aligned}$$

Proof. Assume that $M \in (0, 8\pi)$ and consider the unique solution n_∞ of (1.4), which is also the unique stationary solution of (1.1) such that (1.2) holds. For brevity, we shall omit to mention the dependence of $n_\infty = n_{\infty, M}$ in M .

Let us differentiate with respect to M each term of $\Delta n_\infty + \nabla \cdot (n_\infty x) - \nabla \cdot (n_\infty \nabla c_\infty) = 0$, where $c_\infty = G_2 * n_\infty$. It is straightforward to check that $g_{0,0} := \partial_M \log c_\infty$ is such that $g_{0,0} c_\infty = G_2 * (f_{0,0} n_\infty)$ and $\mathcal{L} f_{0,0} = 0$. Since

$$-\Delta c_\infty = M \frac{e^{c_\infty - \frac{1}{2}|x|^2}}{\int_{\mathbb{R}^2} e^{c_\infty - \frac{1}{2}|x|^2} dx} = n_\infty ,$$

it is clear that $g_{0,0}$ is non-trivial, and therefore $f_{0,0} = \partial_M \log n_\infty$ is a non-trivial solution to $\mathcal{L} f = 0$.

By computing

$$0 = \frac{\partial}{\partial x_1} \left(\Delta n_\infty + \nabla \cdot (x n_\infty) - \nabla \cdot (n_\infty \nabla c_\infty) \right) \quad \text{with} \quad -\Delta \left(\frac{\partial c_\infty}{\partial x_1} \right) = \frac{\partial n_\infty}{\partial x_1}$$

and observing that

$$\frac{\partial}{\partial x_1} \nabla \cdot (x n_\infty) = \frac{\partial n_\infty}{\partial x_1} + \nabla \cdot \left(x \frac{\partial n_\infty}{\partial x_1} \right),$$

we obtain that $f_{1,1} := \partial_{x_1} \log n_\infty$ associated with $g_{1,1} = \frac{1}{c_\infty} \partial_{x_1} c_\infty$ is an eigenfunction of \mathcal{L} , such that $-\mathcal{L} f_{1,1} = f_{1,1}$. The same observation holds if we differentiate with respect to x_i , $i = 2$.

Next consider the dilation operator $D := x \cdot \nabla$. If a is a vector valued function, an elementary computation shows that

$$D(\nabla \cdot a) = \nabla \cdot (D a) - \nabla \cdot a.$$

Since $a = \nabla n_\infty + x n_\infty - n_\infty \nabla c_\infty$ is such that $\nabla \cdot a = 0$, we get $D(\nabla \cdot a) = \nabla \cdot (D a)$ and hence

$$0 = D \left(\Delta n_\infty + \nabla \cdot (x n_\infty) - \nabla \cdot (n_\infty \nabla c_\infty) \right) = \nabla \cdot D \left(\nabla n_\infty + x n_\infty - n_\infty \nabla c_\infty \right).$$

Next, we observe that

$$D(\nabla n_\infty) = \nabla(D n_\infty) - \nabla n_\infty$$

so that

$$\nabla \cdot D(\nabla n_\infty) = \Delta(D n_\infty) - \Delta n_\infty.$$

It is also straightforward to observe that

$$D(x n_\infty) = x n_\infty + x D n_\infty \quad \text{and} \quad D(\nabla c_\infty) = \nabla(D c_\infty) - \nabla c_\infty.$$

Let $f_{0,1} = 1 + \frac{1}{2} D \log n_\infty = 1 + \frac{1}{2 n_\infty} D n_\infty$. By writing $D(\Delta c_\infty + n_\infty) = 0$, we get

$$-\Delta(D c_\infty) + 2 \Delta c_\infty = D n_\infty = 2(f_{0,1} - 1) n_\infty,$$

since

$$D(\Delta c) = \Delta(D c) - 2 \Delta c.$$

Hence, using the fact that $2 \Delta c_\infty = -2 n_\infty$, the function $g_{0,1} := \frac{1}{c_\infty} (-\Delta)^{-1}(n_\infty f_{0,1})$ is given by

$$c_\infty g_{0,1} = \frac{1}{2} D c_\infty.$$

Collecting these identities, we have found that

$$2 n_\infty \mathcal{L}(D \log n_\infty) - \nabla \cdot \left[\nabla n_\infty - x n_\infty - 2 n_\infty \left(\nabla(c_\infty g_{0,1}) - \nabla c_\infty \right) + n_\infty D(\nabla c_\infty) \right] = 0.$$

Using

$$2 \left(\nabla(c_\infty g_{0,1}) - \nabla c_\infty \right) = \nabla(D c_\infty) - 2 \nabla c_\infty = D(\nabla c_\infty) - \nabla c_\infty,$$

this gives

$$n_\infty \mathcal{L}(D \log n_\infty) - \Delta n_\infty + \nabla \cdot (x n_\infty - n_\infty \nabla c_\infty) = 0.$$

Hence, owing to the fact that $D \log n_\infty = 2(f_{0,1} - 1)$ and

$$n_\infty \mathcal{L}(D \log n_\infty) = 2 n_\infty \mathcal{L} f_{0,1} + 2 \nabla \cdot (n_\infty \nabla c_\infty),$$

we get

$$\begin{aligned} -2 n_\infty \mathcal{L} f_{0,1} &= -\Delta n_\infty + \nabla \cdot (x n_\infty + n_\infty \nabla c_\infty) \\ &= 2 \nabla \cdot (x n_\infty) = 4 n_\infty \left(1 + \frac{D n_\infty}{2 n_\infty} \right) = 4 n_\infty f_{0,1}. \end{aligned}$$

We have finally found that $-\mathcal{L} f_{0,1} = 2 f_{0,1}$, which completes the proof. \square

Remark 1. *The fact that 1 and 2 are eigenvalues of \mathcal{L} was known in the limit $M \rightarrow 0_+$: see [8]. It is remarkable that these two eigenvalues are independent of M but this can be explained by noticing that the corresponding eigenfunctions are associated with invariances of the problem before rescaling.*

The functions $\partial_{x_i} \log n_\infty$, $i = 1, 2$ correspond to the invariance under translation in the directions x_i . A decentered self-similar solution would converge in self-similar variables to the stationary solution, in relative entropy, exactly at a rate e^{-t} , thus showing that $\lambda_{1,1} = \lambda_{1,2} = 1$ are eigenvalues by considering the asymptotic regime.

The function $D \log n_\infty$ is associated with the scaling invariance. In original variables, a scaling factor corresponds to a translation in time at the level of the self-similar solution and it can easily be checked that, in self-similar variables, a solution corresponding to the stationary solution rescaled by a factor different from 1 converges, in relative entropy, exactly at a rate e^{-2t} , thus showing that $\lambda_{0,1} = 2$ is also an eigenvalue by considering the asymptotic regime.

4.2. The kernel of the linearized Keller-Segel operator \mathcal{L} . By definition of n_∞ , we know that $\log n_\infty = \mu_0(M) + c_\infty - \frac{1}{2}|x|^2$, so that $f_{0,0} = \mu'_0(M) + g_{0,0} c_\infty$ where $g_{0,0} = \partial_M \log c_\infty$ is such that $-\Delta(g_{0,0} c_\infty) = -\Delta f_{0,0} = f_{0,0} n_\infty$. The normalization constant μ_0 is determined by the condition that $M = \int_{\mathbb{R}^2} n_\infty dx$, that is $\mu_0 = \log M - \log \left(\int_{\mathbb{R}^2} e^{c_\infty - |x|^2/2} dx \right)$. By differentiating with respect to M , we also get that

$$\mu'_0(M) = \frac{1}{M} \left[1 - \int_{\mathbb{R}^2} g_{0,0} n_\infty c_\infty dx \right].$$

The function $f = f_{0,0}$ solves $\mathcal{L} f = 0$ and is such that the equation for $g = f/c_\infty$ reads

$$-\Delta f = n_\infty f.$$

It is not *a priori* granted that such an equation has at most one solution, up to a multiplication by a constant. The uniqueness issue is the purpose of our next result.

Proposition 9. *The kernel $Ker(\mathcal{L})$ is generated by $f_{0,0} = \partial_M \log n_\infty$, which is the unique solution in $L^2(\mathbb{R}^2, n_\infty dx)$, up to a multiplication by a constant, to*

$$-\Delta f_{0,0} = f_{0,0} n_\infty.$$

Proof. We have already seen that $f_{0,0} \in Ker(\mathcal{L})$. It remains to prove that $Ker(\mathcal{L})$ is one-dimensional. Let f be such that $\mathcal{L} f = 0$ and $g = c_\infty^{-1} G_2 * (f n_\infty)$. An elementary computation shows that

$$0 = \int_{\mathbb{R}^2} \mathcal{L} f (f - g c_\infty) n_\infty dx = \int_{\mathbb{R}^2} |\nabla(f - g c_\infty)|^2 n_\infty dx,$$

thus proving that $f = g c_\infty + \mu'_0$ for some real constant μ'_0 (depending eventually on M , with the same notations as above). Hence any solution of $\mathcal{L} f = 0$ has to solve

$$\mathcal{H} f = 0$$

where $\mathcal{H} := -\Delta - n_\infty$ is a Schrödinger operator with potential n_∞ , at least if one assumes that $\nabla(f - G_2 * (f n_\infty))$ belongs to $L^2(n_\infty dx)$. As we shall see later in the discussion of the domain of definition of \mathcal{L} , this is indeed a natural assumption. Altogether, we are interested in characterizing the ground state of the Schrödinger operator \mathcal{H} (with energy level 0) and prove that it is uniquely determined, up to a multiplication by a constant. It is clear that \mathcal{H} has no negative eigenvalue, otherwise the *free energy* functional

$$F[n] := \int_{\mathbb{R}^2} n \log \left(\frac{n}{n_\infty} \right) dx + \frac{1}{4\pi} \iint_{\mathbb{R}^2 \times \mathbb{R}^2} (n(x) - n_\infty(x)) \log |x - y| (n(y) - n_\infty(y)) dx dy$$

would not achieve its minimum for $n = n_\infty$ (see [10] for a proof).

Since n_∞ is radially symmetric (see for instance [10] for a summary of known results), Schwarz' symmetrization applied to \mathcal{H} shows that the ground state is radially symmetric. The function n_∞ seen as a potential, is smooth. By standard elliptic theory, the ground state is smooth as well. Hence, if $f \in H^1(\mathbb{R}^2)$ solves $\mathcal{H}f = 0$, it is uniquely determined as a solution of an ordinary differential equation by the Cauchy-Lipschitz theorem, up to a standard analysis at the origin. Indeed, by considering abusively n_∞ and f as functions of $r = |x|$, we find that f is given by

$$\begin{aligned} f'' + \frac{1}{r} f' + n_\infty f &= 0 \\ f(0) &= 1, \quad f'(0) = 0 \end{aligned}$$

(up to a multiplication by an arbitrary constant). This concludes the proof. \square

4.3. Functional setting and operators. In order to go further in the spectral analysis, to define correctly the domain of the operator \mathcal{L} , to justify the assumption that $\nabla(f - G_2 * (f n_\infty))$ belongs to $L^2(n_\infty dx)$ and to establish spectral gap estimates which are crucial for our analysis, some considerations on the functional setting are in order.

Lemma 10. *Assume that $M \in (0, 8\pi)$ and consider n_∞ defined by (1.4). Then*

$$(4.2) \quad -\frac{1}{2\pi} \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f(x) n_\infty(x) \log|x-y| f(y) n_\infty(y) dx dy = \int_{\mathbb{R}^2} f n_\infty g c_\infty dx \leq \int_{\mathbb{R}^2} |f|^2 n_\infty dx$$

for any $f \in L^2(\mathbb{R}^2, n_\infty dx)$, where $g c_\infty = G_2 * (f n_\infty)$. Moreover, if

$$(4.3) \quad \int_{\mathbb{R}^2} f f_{0,0} n_\infty dx = 0,$$

then equality holds in the above inequality if and only if $f = 0$.

Notice that, if $f \in L^2(\mathbb{R}^2, n_\infty dx)$ is such that

$$(4.4) \quad \int_{\mathbb{R}^2} f n_\infty dx = 0,$$

then (4.2) can be written as

$$(4.5) \quad \int_{\mathbb{R}^2} |\nabla(g c_\infty)|^2 dx \leq \int_{\mathbb{R}^2} |f|^2 n_\infty dx.$$

It is indeed well known that $\nabla(g c_\infty)$ is in $L^2(\mathbb{R}^2, dx)$ as a solution of $-\Delta(g c_\infty) = f n_\infty$ only if (4.4) holds. Lemma 10 will be improved in Section 5 (see Corollary 18); the proof of such a result is independent of the remainder of this section.

Proof. To prove the inequality, we recall that the free energy $n \mapsto F[n]$ achieves its minimum for $n = n_\infty$ according to the logarithmic Hardy-Littlewood-Sobolev inequality (see [22] for detailed considerations on this formulation of the inequality), and observe that

$$Q_1[f] := \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2} F[n_\infty(1 + \varepsilon f)] \geq 0$$

for any smooth function f with compact support satisfying (4.3). The inequality then holds for any $f \in L^2(\mathbb{R}^2, n_\infty dx)$ by density of smooth functions with compact support in $L^2(\mathbb{R}^2, n_\infty dx)$.

If equality holds in (4.2), then the Euler-Lagrange equation amounts to $-\Delta f = f n_\infty$, which characterizes the kernel $\text{Ker}(\mathcal{L})$ according to Proposition 9. \square

Consider the quadratic form Q_1 on $L^2(\mathbb{R}^2, n_\infty dx)$, which takes the form

$$Q_1[f] = \int_{\mathbb{R}^2} |f|^2 n_\infty dx + \frac{1}{2\pi} \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f(x) n_\infty(x) \log|x-y| f(y) n_\infty(y) dx dy.$$

By (4.2), it is nonnegative, and positive semi-definite on the orthogonal of the kernel of \mathcal{L} , for the natural scalar product on $L^2(n_\infty dx)$, *i.e.* for any $f \in L^2(n_\infty dx)$ such that (4.3) holds. Using previous notations, we may write

$$\mathbf{Q}_1[f] = \int_{\mathbb{R}^2} f(f - g c_\infty) n_\infty dx \quad \text{with} \quad g c_\infty = G_2 * (f n_\infty).$$

If (4.4) holds, we can also observe that

$$\mathbf{Q}_1[f] = \int_{\mathbb{R}^2} |f|^2 n_\infty dx - \int_{\mathbb{R}^2} |\nabla(g c_\infty)|^2 dx.$$

To \mathbf{Q}_1 we associate its polar form \mathbf{L}_1 defined on smooth functions with compact support such that (4.3) holds and define its Friedrich's extension on $L^2(n_\infty dx)$, that we still denote by \mathbf{L}_1 . By construction, \mathbf{L}_1 is a positive self-adjoint operator with domain $\mathcal{D}(\mathbf{L}_1) \subset L^2(n_\infty dx)$. On $\mathcal{D}(\mathbf{L}_1)$, we shall denote by $\langle \cdot, \cdot \rangle$ the scalar product induced by \mathbf{L}_1 . Explicitly, this means that

$$\langle f, \tilde{f} \rangle = \int_{\mathbb{R}^2} f \tilde{f} n_\infty dx + \frac{1}{2\pi} \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f(x) n_\infty(x) \log|x-y| \tilde{f}(y) n_\infty(y) dx dy.$$

The scalar product $\langle \cdot, \cdot \rangle$ induced by \mathbf{L}_1 is defined on the orthogonal of $f_{0,0}$, but can be extended as a bilinear form to $L^2(n_\infty dx)$. If $f \in L^2(n_\infty dx)$ is such that (4.4) holds, then we notice that

$$\langle f, f_{0,0} \rangle = \int_{\mathbb{R}^2} f(f_{0,0} - G_2 * f_{0,0}) n_\infty dx = 0$$

because $f_{0,0} = G_2 * (f_{0,0} n_\infty) + \mu'_0$. With these notations, notice that we have

$$\langle f, f \rangle = \mathbf{Q}_1[f] \geq 0$$

for any $f \in \mathcal{D}(\mathbf{L}_1)$, with equality if and only if $f = 0$.

We can also define the quadratic form \mathbf{Q}_2 as

$$\mathbf{Q}_2[f] := \int_{\mathbb{R}^2} |\nabla(f - g c_\infty)|^2 n_\infty dx \quad \text{with} \quad g = \frac{1}{c_\infty} G_2 * (f n_\infty).$$

As for \mathbf{Q}_1 , we define \mathbf{Q}_2 on the set of smooth functions such that (4.3) holds and extend it. The associated self-adjoint nonnegative operator is denoted by \mathbf{L}_2 and it is again a self-adjoint operator, with domain $\mathcal{D}(\mathbf{L}_2) \subset L^2(n_\infty dx)$.

Proposition 11. *With the above notations, the restriction of \mathcal{L} to $\mathcal{D}(\mathbf{L}_1)$ is a self-adjoint operator for the scalar product $\langle \cdot, \cdot \rangle$ with domain $\mathcal{D}(\mathbf{L}_2)$, such that*

$$\langle f, \mathcal{L} f \rangle = -\mathbf{Q}_2[f] \quad \forall f \in \mathcal{D}(\mathbf{L}_2)$$

and $\text{Ker}(\mathcal{L}) \cap \mathcal{D}(\mathbf{L}_2) = \{0\}$.

Remark 2. *The function $f_{0,0}$ is an eigenfunction of \mathcal{L} but this is not the case of $f \equiv 1$. With the notations of Section 4.1, the functions $f_{1,i}$ are orthogonal to $f \equiv 1$ in $L^2(\mathbb{R}^2, n_\infty dx)$ for $i = 1, 2$, but this is the case neither for $f_{0,0}$ nor for $f_{0,1}$.*

4.4. The spectrum of \mathcal{L} is discrete. We define

$$\Lambda_1 := \inf_{f \in \mathcal{D}(\mathbf{L}_2) \setminus \{0\}} \frac{\mathbf{Q}_2[f]}{\mathbf{Q}_1[f]} \quad \text{and} \quad \Lambda_\infty := \lim_{R \rightarrow \infty} \inf_{\substack{f \in \mathcal{D}(\mathbf{L}_2) \setminus \{0\} \\ \text{supp}(f) \subset \mathbb{R}^2 \setminus B(0, R)}} \frac{\mathbf{Q}_2[f]}{\mathbf{Q}_1[f]}.$$

First, let us give a heuristic approach of the problem. As an application of Persson's method (see [35]), the bottom of the essential spectrum of \mathcal{L} can be characterized as

$$\inf \sigma_{\text{ess}}(\mathcal{L}) = \Lambda_\infty.$$

To prove that \mathcal{L} has a spectral gap on $\mathcal{D}(L_2)$, it is enough to show that Λ_∞ is positive: either $\Lambda_1 = \Lambda_\infty$, or $\Lambda_1 < \Lambda_\infty$ is a nonnegative eigenvalue, which cannot be equal to 0. This is summarized in the following statement.

Proposition 12. *With the above notations, Λ_1 is positive and*

$$(4.6) \quad \Lambda_1 \mathbf{Q}_1[f] \leq \mathbf{Q}_2[f] \quad \forall f \in \mathcal{D}(L_2) .$$

For any $f \in \mathcal{D}(L_2)$, if (4.4) holds, then Inequality (4.6) can be reformulated as

$$\Lambda_1 \int_{\mathbb{R}^2} |f|^2 n_\infty dx \leq \int_{\mathbb{R}^2} |\nabla(f - g c_\infty)|^2 n_\infty dx + \Lambda_1 \int_{\mathbb{R}^2} |\nabla(g c_\infty)|^2 dx .$$

The proof of Proposition 12 can be done by considering \mathcal{L} as a perturbation of the operator $f \mapsto n_\infty^{-1} \nabla \cdot (n_\infty \nabla f)$ defined on $L^2(\mathbb{R}^2, n_\infty dx)$. This was the method of [8]. However on such a space \mathcal{L} is not self-adjoint and justifications are delicate because of the logarithmic kernel, away from the small mass regime.

In practice, Persson's method is not well designed either to handle convolution operators, although it can probably be adapted with little effort. This may even have been done, but we are not aware of such a result. Moreover, as we shall see below, we have: $\Lambda_\infty = \infty$, which further simplifies the proof. For these reasons, we will therefore give a direct proof, based on some of the tools of the concentration-compactness method (see [28, 29, 30, 31, 32]) and adapted to the case of a bounded measure, $n_\infty dx$, as in [9]. In that framework, Λ_∞ corresponds to the *problem at infinity*. For simplicity, let us split the proof into Lemmas 13 and 14.

Lemma 13. *With the above notations, $\Lambda_\infty = \infty$.*

Proof. Recall that

$$n_\infty = M \frac{e^{c_\infty - \frac{1}{2}|x|^2}}{\int_{\mathbb{R}^2} e^{c_\infty - \frac{1}{2}|x|^2} dx}$$

where $c_\infty = (-\Delta)^{-1} n_\infty$ is such that

$$\limsup_{|x| \rightarrow \infty} \left| c_\infty(x) + \frac{M}{2\pi} \log |x| \right| < \infty .$$

As a consequence, we know that

$$n_\infty(x) \sim |x|^{-\alpha} e^{-\frac{1}{2}|x|^2} \quad \text{as } |x| \rightarrow +\infty, \quad \text{with } \alpha = \frac{M}{2\pi} .$$

We can expand the square $|\nabla(f - g c_\infty)|^2$ and get

$$\begin{aligned} \mathbf{Q}_2[f] &= \int_{\mathbb{R}^2} |\nabla(f - g c_\infty)|^2 n_\infty dx \\ &= \int_{\mathbb{R}^2} |\nabla f|^2 n_\infty dx + \int_{\mathbb{R}^2} |\nabla(g c_\infty)|^2 n_\infty dx \\ &\quad + 2 \int_{\mathbb{R}^2} f \nabla(g c_\infty) \cdot \nabla n_\infty dx - 2 \int_{\mathbb{R}^2} f (-\Delta(g c_\infty)) n_\infty dx . \end{aligned}$$

Assume that f is supported in $\mathbb{R}^2 \setminus B(0, R)$, for $R > 0$, large. Then

$$\begin{aligned} \int_{\mathbb{R}^2} f (-\Delta(g c_\infty)) n_\infty dx &= \int_{\mathbb{R}^2} |f|^2 n_\infty^2 dx \\ &\leq \sup_{|x| > R} n_\infty(x) \int_{\mathbb{R}^2} |f|^2 n_\infty dx \sim R^{-\alpha} e^{-\frac{1}{2}R^2} \int_{\mathbb{R}^2} |f|^2 n_\infty dx \end{aligned}$$

on the one hand, and we know from Persson's method that

$$\lim_{R \rightarrow \infty} \inf_{\substack{f \in \mathcal{D}(\mathbf{L}_2) \setminus \{0\} \\ \text{supp}(f) \subset \mathbb{R}^2 \setminus B(0, R)}} \frac{\int_{\mathbb{R}^2} |\nabla f|^2 n_\infty dx}{\int_{\mathbb{R}^2} |f|^2 n_\infty dx} = +\infty$$

on the other hand, so that, for any $\varepsilon > 0$, there exists $R > 0$ large enough for which

$$\int_{\mathbb{R}^2} |f|^2 n_\infty dx \leq \varepsilon \int_{\mathbb{R}^2} |\nabla f|^2 n_\infty dx$$

for any function $f \in H^1(\mathbb{R}^2, n_\infty dx)$. Equivalently, we can write that there exists a positive function $R \mapsto \varepsilon(R)$ such that $\lim_{R \rightarrow +\infty} \varepsilon(R) = 0$ and

$$0 \leq \int_{\mathbb{R}^2} f (-\Delta(g c_\infty)) n_\infty dx \leq R^{-\alpha} e^{-\frac{1}{2} R^2} \varepsilon(R)$$

uniformly with respect to f as soon as it is supported in $\mathbb{R}^2 \setminus B(0, R)$.

Assume first that Condition (4.4) is satisfied. We notice that

$$\left| \int_{\mathbb{R}^2} f \nabla(g c_\infty) \cdot \nabla n_\infty dx \right| \leq 2 \int_{\mathbb{R}^2} |f \sqrt{n_\infty}| |\nabla(g c_\infty)| |(\nabla c_\infty - x) \sqrt{n_\infty}| dx$$

can be estimated by

$$\left| \int_{\mathbb{R}^2} f \nabla(g c_\infty) \cdot \nabla n_\infty dx \right| \leq 2 \sup_{|x| > R} |(\nabla c_\infty - x) \sqrt{n_\infty}| \left(\int_{\mathbb{R}^2} |f|^2 n_\infty dx \int_{\mathbb{R}^2} |\nabla(g c_\infty)|^2 dx \right)^{1/2}.$$

By Lemma 10, we know that $\mathbf{Q}_1[f] \geq 0$ is equivalent to (4.5) and find that

$$\begin{aligned} \left| \int_{\mathbb{R}^2} f \nabla(g c_\infty) \cdot \nabla n_\infty dx \right| &\leq 2 \sup_{|x| > R} |(\nabla c_\infty - x) \sqrt{n_\infty}| \int_{\mathbb{R}^2} |f|^2 n_\infty dx \\ &\leq 2 \sup_{|x| > R} |(\nabla c_\infty - x) \sqrt{n_\infty}| \varepsilon(R) \int_{\mathbb{R}^2} |\nabla f|^2 n_\infty dx. \end{aligned}$$

On the other hand, since Condition (4.4) is satisfied, we know for free that

$$\mathbf{Q}_1[f] = \int_{\mathbb{R}^2} |f|^2 n_\infty dx - \int_{\mathbb{R}^2} |\nabla(g c_\infty)|^2 dx \leq \int_{\mathbb{R}^2} |f|^2 n_\infty dx.$$

As a consequence, we have obtained that

$$\lim_{R \rightarrow \infty} \inf_{\substack{f \in \mathcal{D}(\mathbf{L}_2) \setminus \{0\} \\ \text{supp}(f) \subset \mathbb{R}^2 \setminus B(0, R)}} \frac{\mathbf{Q}_2[f]}{\mathbf{Q}_1[f]} = \lim_{R \rightarrow \infty} \inf_{\substack{f \in \mathcal{D}(\mathbf{L}_2) \setminus \{0\} \\ \text{supp}(f) \subset \mathbb{R}^2 \setminus B(0, R)}} \frac{\int_{\mathbb{R}^2} |\nabla f|^2 n_\infty dx}{\int_{\mathbb{R}^2} |f|^2 n_\infty dx} = +\infty,$$

which proves our claim.

If Condition (4.4) is not satisfied, the proof is more complicated. By homogeneity, there is no restriction to assume that $\frac{1}{M} \int_{\mathbb{R}^2} f^2 n_\infty dx = 1$. Let $\theta := \frac{1}{M} \int_{\mathbb{R}^2} f n_\infty dx$ and $\tilde{f} := f - \theta$, $\tilde{g} := g - \theta$. Then $\int_{\mathbb{R}^2} \tilde{f} n_\infty dx = 0$. Notice that, by the Cauchy-Schwarz inequality, we have: $\theta \in [-1, 1]$. Moreover, if $\theta \neq 0$, then $B(0, R)$ is contained in $\text{supp}(\tilde{f})$.

With these notations, we first have to estimate

$$\int_{\mathbb{R}^2} f \nabla(g c_\infty) \cdot \nabla n_\infty dx = 2\theta \int_{\mathbb{R}^2} f \sqrt{n_\infty} \nabla c_\infty \cdot \nabla \sqrt{n_\infty} dx + 2 \int_{\mathbb{R}^2} f \sqrt{n_\infty} \nabla(\tilde{g} c_\infty) \cdot \nabla \sqrt{n_\infty} dx.$$

By the Cauchy-Schwarz inequality, we have

$$\left| \int_{\mathbb{R}^2} f \nabla c_\infty \cdot \nabla \sqrt{n_\infty} \, dx \right|^2 \leq \int_{\mathbb{R}^2} f^2 n_\infty \, dx \int_{\mathbb{R}^2 \setminus B(0,R)} |\nabla c_\infty|^2 |\nabla \sqrt{n_\infty}|^2 \, dx ,$$

and it is simple to check that the last integral in the right hand side converges to 0 as $R \rightarrow \infty$. The second integral can be estimated as before by writing

$$\left| \int_{\mathbb{R}^2} f \sqrt{n_\infty} \nabla(\tilde{g} c_\infty) \cdot \nabla \sqrt{n_\infty} \, dx \right|^2 \leq \sup_{|x|>R} |(\nabla c_\infty - x) \sqrt{n_\infty}|^2 \int_{\mathbb{R}^2} |f|^2 n_\infty \, dx \int_{\mathbb{R}^2} |\nabla(\tilde{g} c_\infty)|^2 \, dx$$

and by recalling that $\int_{\mathbb{R}^2} |\tilde{f}|^2 n_\infty \, dx = M$. From these estimates, we conclude that

$$\lim_{R \rightarrow \infty} \inf_{\substack{f \in \mathcal{D}(\mathbf{L}_2) \setminus \{0\} \\ \text{supp}(f) \subset \mathbb{R}^2 \setminus B(0,R)}} \mathbf{Q}_2[f] = \infty .$$

We also need to estimate $\mathbf{Q}_1[f]$ and this can be done by showing that

$$\mathbf{Q}_1[f] = M + \frac{1}{2\pi} \iint_{\mathbb{R}^2 \times \mathbb{R}^2} (f n_\infty)(x) \log |x - y| (f n_\infty)(y) \, dx \, dy$$

is bounded from above if we still impose that $\int_{\mathbb{R}^2} |f|^2 n_\infty \, dx = M$. Using the crude estimate

$$2 \log |x - y| \leq |x - y|^2 \leq 2(|x|^2 + |y|^2) \quad \forall (x, y) \in \mathbb{R}^2 \times \mathbb{R}^2$$

and, as a consequence,

$$\iint_{\mathbb{R}^2 \times \mathbb{R}^2} (f n_\infty)(x) \log |x - y| (f n_\infty)(y) \, dx \, dy \leq 2 \int_{\mathbb{R}^2} f n_\infty \, dx \int_{\mathbb{R}^2} |y|^2 (f n_\infty)(y) \, dy ,$$

we conclude by observing that

$$\int_{\mathbb{R}^2} f n_\infty \, dx \leq \sqrt{M} \left(\int_{\mathbb{R}^2 \setminus B(0,R)} n_\infty \, dx \right)^{1/2}$$

$$\text{and } \int_{\mathbb{R}^2} |y|^2 (f n_\infty)(y) \, dy \leq \sqrt{M} \left(\int_{\mathbb{R}^2 \setminus B(0,R)} |y|^4 n_\infty \, dy \right)^{1/2}$$

both converge to 0 as $R \rightarrow \infty$. □

Lemma 14. *With the above notations, $\Lambda_1 > 0$.*

Proof. Tools for the proof of this lemma are to a large extent standard in concentration-compactness methods or when applied to models of quantum chemistry, so we shall only sketch the main steps and omit as much as possible the technicalities of such an approach. We will actually prove a result that is stronger than the one of Lemma 14: Λ_1 is achieved by some function $f \in \mathcal{D}(\mathbf{L}_2)$.

Consider a minimizing sequence $(f_n)_{n \in \mathbb{N}}$ for the functional $f \mapsto \mathbf{Q}_2[f]/\mathbf{Q}_1[f]$ defined on $\mathcal{D}(\mathbf{L}_2) \setminus \{0\}$. By homogeneity, we may assume that $\int_{\mathbb{R}^2} f^2 n_\infty \, dx = 1$ for any $n \in \mathbb{N}$, with no restriction, while $\mathbf{Q}_2[f_n]$ is bounded uniformly in $n \in \mathbb{N}$. Let $F_n := f_n \sqrt{n_\infty}$. In the framework of concentration-compactness methods, for any given $\varepsilon > 0$, it is a standard result that one can decompose F_n as

$$F_n = F_n^{(1)} + F_n^{(2)} + \tilde{F}_n$$

for any $n \in \mathbb{N}$, with

$$\int_{\mathbb{R}^2} |F_n^{(1)}|^2 \, dx + \int_{\mathbb{R}^2} |F_n^{(2)}|^2 \, dx + \int_{\mathbb{R}^2} |\tilde{F}_n|^2 \, dx = 1$$

where $F_n^{(1)}$, $F_n^{(2)}$ and \tilde{F}_n are supported respectively in $B(0, 2R)$, $\mathbb{R}^2 \setminus B(0, R_n)$ and $B(0, 2R_n) \setminus B(0, R)$, $R_n > R > 1$, $\lim_{n \rightarrow \infty} R_n = \infty$,

$$\int_{\mathbb{R}^2} |F_n^{(1)}|^2 dx \geq \theta - \varepsilon \quad \text{and} \quad \int_{\mathbb{R}^2} |F_n^{(2)}|^2 dx \geq 1 - \theta - \varepsilon$$

for some $\theta \in [0, 1]$. As a consequence, we also have that $\int_{\mathbb{R}^2} |\tilde{F}_n|^2 dx \leq 2\varepsilon$. A standard method to obtain such a decomposition is based on the IMS truncation method, which goes as follows. Take a smooth truncation function χ with the following properties: $0 \leq \chi \leq 1$, $\chi(x) = 1$ for any $x \in B(0, 1)$, $\chi(x) = 0$ for any $x \in \mathbb{R}^2 \setminus B(0, 2)$, and define $\chi_R(x) := \chi(x/R)$ for any $x \in \mathbb{R}^2$. Then for an appropriate choice of R and $(R_n)_{n \in \mathbb{N}}$, we can choose

$$F_n^{(1)} = \chi_R F_n \quad \text{and} \quad F_n^{(2)} = \sqrt{1 - \chi_{R_n}^2} F_n.$$

If $\theta = 1$, then $(F_n^{(1)})_{n \in \mathbb{N}}$ strongly converges in $L_{\text{loc}}^2(\mathbb{R}^2, dx)$ to some limit F and we have

$$\liminf_{n \rightarrow \infty} \int_{\mathbb{R}^2} |\nabla F_n|^2 dx \geq \int_{\mathbb{R}^2} |\nabla F|^2 dx \quad \text{and} \quad \int_{\mathbb{R}^2} |F|^2 dx \geq 1 - \varepsilon.$$

Now we repeat the argument as $\varepsilon = \varepsilon_n \rightarrow 0_+$, take a diagonal subsequence that we still denote by $(F_n)_{n \in \mathbb{N}}$, define

$$F_n^{(1)} = \chi_{R_n^{(1)}} F_n, \quad F_n^{(2)} = \sqrt{1 - \chi_{R_n^{(2)}}^2} F_n \quad \text{and} \quad \tilde{F}_n = F_n - F_n^{(1)} - F_n^{(2)},$$

where $\lim_{n \rightarrow \infty} R_n^{(1)} = +\infty$ and $R_n^{(2)} \geq 2R_n^{(1)}$ for any $n \in \mathbb{N}$. Since limits obtained above by taking a diagonal subsequence coincide on larger and larger centered balls (that is when R increases), we find a nontrivial minimizer $f = F/\sqrt{n_\infty}$ such that $\int_{\mathbb{R}^2} F^2 dx = \int_{\mathbb{R}^2} f^2 n_\infty dx = 1$, since all other terms are relatively compact. Notice that $\mathbf{Q}_1[f] > 0$ because the condition (4.3) is preserved by passing to the limit. Hence Λ_1 is achieved and we know that Λ_1 is positive because \mathbf{Q}_1 is positive semi-definite.

Assume now that $\theta < 1$. We know that $\int_{\mathbb{R}^2} |\tilde{F}_n|^2 dx \leq 2\varepsilon_n$ and hence

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} |\tilde{F}_n|^2 dx = 0.$$

It is not difficult to see that cross terms do not play any role in the integrals involving convolution kernels, as it is standard for Hartree type (or Schrödinger-Poisson) models. As a consequence, we can write that

$$\lim_{n \rightarrow \infty} \mathbf{Q}_1[f_n] = \lim_{n \rightarrow \infty} \mathbf{Q}_1[f_n^{(1)}] + \lim_{n \rightarrow \infty} \mathbf{Q}_1[f_n^{(2)}]$$

where $f_n^{(i)} := F_n^{(i)}/\sqrt{n_\infty}$, $i = 1, 2$. Proceeding as above, we may find a limit f of $(f_n^{(1)})_{n \in \mathbb{N}}$, in $L^2(n_\infty dx)$. It is then straightforward to observe that

$$\Lambda_1 = \lim_{n \rightarrow \infty} \frac{\mathbf{Q}_2[f_n]}{\mathbf{Q}_1[f_n]} \geq \lim_{n \rightarrow \infty} \frac{\mathbf{Q}_2[f] + \mathbf{Q}_2[f_n^{(2)}]}{\mathbf{Q}_1[f] + \mathbf{Q}_1[f_n^{(2)}]}.$$

If $\theta > 0$, we know that $\mathbf{Q}_2[f] \geq \Lambda_1 \mathbf{Q}_1[f]$ and $\lim_{n \rightarrow \infty} \mathbf{Q}_2[f_n^{(2)}]/\mathbf{Q}_1[f_n^{(2)}] > \Lambda_1$ by Lemma 13, so that $\lim_{n \rightarrow \infty} \mathbf{Q}_2[f_n^{(2)}] = 0$ and f is a nontrivial minimizer: we are back to the case $\theta = 1$, but with a different normalization of f . If $\theta = 0$, it is clear that $\mathbf{Q}_1[f] = 0$ and we get

$$\Lambda_1 \geq \lim_{n \rightarrow \infty} \frac{\mathbf{Q}_2[f_n^{(2)}]}{\mathbf{Q}_1[f_n^{(2)}]} = \infty,$$

again by Lemma 13, a contradiction with the fact that $\mathbf{Q}_2[f]/\mathbf{Q}_1[f]$ takes finite values for arbitrary test functions in $\mathcal{D}(\mathbf{L}_2) \setminus \{0\}$. This concludes our proof. \square

Remark 3. For any $k \in \mathbb{N}$, $k \geq 1$, define the Raleigh quotient

$$\Lambda_k := \inf_{\substack{f \in \mathcal{D}(\mathcal{L}_2) \setminus \{0\} \\ \langle f_j, f \rangle = 0, j = 0, 1, \dots, k-1}} \frac{\mathcal{Q}_2[f]}{\mathcal{Q}_1[f]}$$

where f_j denotes a critical point associated to Λ_j . Critical points are counted with multiplicity. Since the orthogonality condition $\langle f_j, f \rangle = 0$ is preserved by taking the limit along the weak topology of $L^2(n_\infty dx)$, building a minimizing sequence for $k \geq 1$ goes as in the case $k = 1$. It is easy to check that Λ_k is then an eigenvalue of \mathcal{L} considered as an operator on $\mathcal{D}(\mathcal{L}_2)$ with scalar product $\langle \cdot, \cdot \rangle$, for any $k \geq 1$ and $\lim_{k \rightarrow \infty} \Lambda_k = \infty$.

4.5. A spectral gap inequality. We are now going to prove that Ineq. (4.6) holds with $\Lambda_1 = \lambda_{1,i} = 1$, $i = 1, 2$. This is our first main estimate.

Theorem 15. For any function $f \in \mathcal{D}(\mathcal{L}_2)$, we have

$$\mathcal{Q}_1[f] \leq \mathcal{Q}_2[f].$$

Recall that, with the notations of Section 4, $\mathcal{Q}_1[f] = \langle f, f \rangle$ and $\mathcal{Q}_2[f] = \langle f, \mathcal{L}f \rangle$.

Proof. We have to compute the lowest positive eigenvalue of \mathcal{L} . After a reformulation in terms of cumulated densities for the solution of (1.3) and for the eigenvalue problem for \mathcal{L} , we will identify the lowest eigenvalue $\lambda_{0,1} = 2$ when \mathcal{L} is restricted to radial functions, and the lowest ones, $\lambda_{1,1} = \lambda_{1,2} = 1$, when \mathcal{L} is restricted to functions corresponding to the $k = 1$ component in the decomposition into spherical harmonics.

Step 1. Reformulation in terms of cumulated densities.

Among spherically symmetric functions, it is possible to reduce the problem to a single ordinary differential equation.

Consider first a stationary solution (n_∞, c_∞) of (1.3) and as in [4] or [3] (also see references therein), let us rewrite the system in terms of the *cumulated densities* Φ and Ψ defined by

$$\begin{aligned} \Phi(s) &:= \frac{1}{2\pi} \int_{B(0, \sqrt{s})} n_\infty(x) dx, \\ \Psi(s) &:= \frac{1}{2\pi} \int_{B(0, \sqrt{s})} c_\infty(x) dx. \end{aligned}$$

Notice that $\Phi(s) = \frac{1}{2\pi} M_\varepsilon(\pi s)$ for $\varepsilon = 0$, with the notations of the proof of Theorem 4. The motivation for such a reformulation is that the system can be rewritten in terms of a nonlinear, local, ordinary differential equation for Φ using the fact that n_∞ is radial. With a slight abuse of notations, we can consider n_∞ and c_∞ as functions of $r = |x|$. Elementary computations show that

$$\begin{aligned} n_\infty(\sqrt{s}) &= 2\Phi'(s) \quad \text{and} \quad n'_\infty(\sqrt{s}) = 4\sqrt{s}\Phi''(s), \\ c_\infty(\sqrt{s}) &= 2\Psi'(s) \quad \text{and} \quad c'_\infty(\sqrt{s}) = 4\sqrt{s}\Psi''(s). \end{aligned}$$

After one integration with respect to $r = \sqrt{s}$, the Poisson equation $-\Delta c_\infty = n_\infty$ can be rewritten as

$$-\sqrt{s}c'_\infty(\sqrt{s}) = \Phi(s)$$

while the equation for n_∞ , after an integration on $(0, r)$, is

$$n'_\infty(\sqrt{s}) + \sqrt{s}n_\infty(\sqrt{s}) - n_\infty(\sqrt{s})c'_\infty(\sqrt{s}) = 0.$$

These two equations written in terms of Φ and Ψ are

$$-4s\Psi'' = \Phi$$

and

$$\Phi'' + \frac{1}{2} \Phi' - 2 \Phi' \Psi'' .$$

After eliminating Ψ'' , we find that Φ is the solution of the ordinary differential equation

$$(4.7) \quad \Phi'' + \frac{1}{2} \Phi' + \frac{1}{2s} \Phi \Phi' = 0$$

with initial conditions $\Phi(0) = 0$ and $\Phi'(0) = \frac{1}{2} n(0) =: a$, so that all solutions can be parametrized in terms of $a > 0$.

Consider next the functions f and g involved in the linearized Keller-Segel system (1.3) and define the corresponding *cumulated densities* given by

$$\begin{aligned} \phi(s) &:= \frac{1}{2\pi} \int_{B(0, \sqrt{s})} (f n_\infty)(x) dx , \\ \psi(s) &:= \frac{1}{2\pi} \int_{B(0, \sqrt{s})} (g c_\infty)(x) dx . \end{aligned}$$

If $g c_\infty = (-\Delta)^{-1}(f n_\infty)$ and f is a solution of the eigenvalue problem

$$-\mathcal{L} f = \lambda f ,$$

then we can make a computation similar to the above one and get

$$(n_\infty f)(\sqrt{s}) = 2 \phi'(s) , \quad (n_\infty f')(\sqrt{s}) = 4 \sqrt{s} \phi''(s) - 2 \frac{n'_\infty}{n_\infty}(\sqrt{s}) \phi'(s) ,$$

$$(g c_\infty)(\sqrt{s}) = 2 \psi'(s) \quad \text{and} \quad (g c_\infty)'(\sqrt{s}) = 4 \sqrt{s} \psi''(s) .$$

The equations satisfied by f and g are

$$-\sqrt{s} (g c_\infty)'(\sqrt{s}) = \phi(s)$$

and

$$\sqrt{s} \left((n_\infty f')(\sqrt{s}) - n_\infty (g c_\infty)'(\sqrt{s}) \right) + \lambda \phi(s) = 0 .$$

These two equations written in terms of ϕ and ψ become

$$-4s \psi'' = \phi$$

and

$$4s \left(\phi'' - \frac{\Phi''}{\Phi'} \phi' - 2 \Phi' \psi'' \right) + \lambda \phi = 0 .$$

After eliminating ψ'' , we find that ϕ is the solution of the ordinary differential equation

$$\phi'' - \frac{\Phi''}{\Phi'} \phi' + \frac{\lambda + 2 \Phi'}{4s} \phi = 0 .$$

Taking into account the equation for Φ , that is

$$-\frac{\Phi''}{\Phi'} = \frac{1}{2} + \frac{\Phi}{2s} ,$$

we can also write that ϕ solves

$$(4.8) \quad \phi'' + \frac{s + \Phi}{2s} \phi' + \frac{\lambda + 2 \Phi'}{4s} \phi = 0 .$$

Recall that the set of solutions to (4.7) is parametrized by $a = \Phi'(0)$. It is straightforward to remark that $\phi = \frac{d}{da} \Phi$ solves (4.8) with $\lambda = 0$. The reader is invited to check that $s \mapsto s \Phi'(s)$ provides a nonnegative solution of (4.8) with $\lambda = 2$.

Step 2. Characterization of the radial ground state.

It is possible to rewrite (4.8) as

$$\frac{d}{ds} \left(e^{\alpha(s)} \frac{d\phi}{ds} \right) + \frac{\lambda + 2\Phi'}{4s} e^{\alpha(s)} \phi = 0 \quad \text{with} \quad \alpha(s) := \frac{s}{2} + \frac{1}{2} \int_0^s \frac{\phi(\sigma)}{\sigma} d\sigma .$$

The equation holds on $(0, \infty)$ and boundary conditions are $\phi(0) = 0$ and $\lim_{s \rightarrow \infty} \phi(s) = 0$. By the Sturm-Liouville theory, we know that $\lambda = 2 = \lambda_{0,1}$ is then the lowest positive eigenvalue such that ϕ is nonnegative and satisfies the above boundary conditions.

In other words, we have shown that the function $f_{0,1}$ found in Section 4.1 generates the eigenspace corresponding to the lowest positive eigenvalue of \mathcal{L} restricted to radial functions.

Step 3. Spherical harmonics decomposition.

We have to deal with non-radial modes of \mathcal{L} . Since n_∞ and c_∞ are both radial, we can use a spherical harmonics decomposition for that purpose. As in [13], the eigenvalue problem for the operator \mathcal{L} amounts to solve among radial functions f and g the system

$$\begin{aligned} -f'' - \frac{1}{r} f' + \frac{k^2}{r^2} f + (r - c'_\infty)(f' - (g c_\infty)') - n_\infty f &= \lambda f , \\ -(g c_\infty)'' - \frac{1}{r} (g c_\infty)' + \frac{k^2}{r^2} (g c_\infty) &= n_\infty f , \end{aligned}$$

for some $k \in \mathbb{N}$, $k \geq 1$. Here as above, we make the standard abuse of notations that amounts to write n_∞ and c_∞ as a function of $r = |x|$. It is straightforward to see that $k = 1$ realizes the infimum of the spectrum of \mathcal{L} among non-radial functions. The function $f = -n'_\infty$ provides a nonnegative solution for $k = 1$ and $\lambda = 1$. It is then possible to conclude using the following observation: f is a radial C^2 solution if and only if $r \mapsto r f =: \tilde{f}(r)$ solves $-\mathcal{L} \tilde{f} = (\lambda + 1) \tilde{f}$ among radial functions, and we are back to the problem studied in Step 2. The value we look for is therefore $\lambda = 1 = \lambda_{1,1} = \lambda_{1,2}$.




FIGURE 1. Using a shooting method, one can numerically compute the lowest eigenvalues of $-\mathcal{L}$ for $k = 0$ (radial functions) and for the $k = 1$ component of the spherical harmonics decomposition (dashed curve). The plot shows that 1 and 2 are the lowest eigenvalues, when mass varies between 0 and $8\pi \approx 25.1327$. See [13] for details and further numerical results.

In other words, we have shown that the functions $f_{1,1}$ and $f_{1,2}$ found in Section 4.1 generate the eigenspace corresponding to the lowest positive eigenvalue of \mathcal{L} corresponding to $k = 1$. We are now in position to conclude the proof of Theorem 15.

Either the spectral gap is achieved among radial functions and $\Lambda_1 = 2$, or it is achieved among functions in one of the non-radial components corresponding to the spherical harmonics decomposition: the one given by $k = 1$ minimizes the gap and hence we obtain $\Lambda_1 = 1$. See Fig. 1 for an illustration. \square

As a consequence of Step 2 in the proof of Theorem 15, we find that the following inequality holds.

Proposition 16. *For any radial function $f \in \mathcal{D}(L_2)$, we have*

$$2Q_1[f] \leq Q_2[f].$$

where, with the notations of Section 4, $Q_1[f] = \langle f, f \rangle$ and $Q_2[f] = \langle f, \mathcal{L}f \rangle$.

This observation has to be related with recent results of V. Calvez and J.A. Carrillo. As a consequence, the rate e^{-2t} in Theorem 1 can be replaced by e^{-4t} when solutions are radially symmetric, consistently with [11, Theorem 1.2]. The necessary adaptations (see Section 6) are straightforward.

5. A STRICT POSITIVITY RESULT FOR THE LINEARIZED ENTROPY

Lemma 10 can be improved and this is our second main estimate.

Theorem 17. *There exists $\Lambda > 1$ such that*

$$(5.1) \quad \Lambda \int_{\mathbb{R}^2} f n_\infty (-\Delta)^{-1} (f n_\infty) dx \leq \int_{\mathbb{R}^2} |f|^2 n_\infty dx$$

for any $f \in L^2(\mathbb{R}^2, n_\infty dx)$ such that (4.3) holds.

Proof. Let us give an elementary proof based on two main observations: the equivalence with a Poincaré type inequality using Legendre's transform, and the application of a concentration-compactness method for proving the Poincaré inequality. Recall that by Lemma 10 we already know that (5.1) holds with $\Lambda = 1$.

Step 1. We claim that Inequality (5.1) is equivalent to

$$(5.2) \quad \Lambda \int_{\mathbb{R}^2} |h|^2 n_\infty dx \leq \int_{\mathbb{R}^2} |\nabla h|^2 dx$$

for any $h \in L^2(\mathbb{R}^2, n_\infty dx)$ such that the condition $\int_{\mathbb{R}^2} h f_{0,0} n_\infty dx = 0$ holds, *i.e.* such that h satisfies (4.3). Let us prove this claim.

Assume first that (5.2) holds and take Legendre's transform of both sides with respect to the natural scalar product in $L^2(n_\infty dx)$: for any $f \in L^2(\mathbb{R}^2, n_\infty dx)$ such that (4.3) holds,

$$\sup_h \left(\int_{\mathbb{R}^2} f h n_\infty dx - \frac{1}{2} \int_{\mathbb{R}^2} h^2 n_\infty dx \right) \geq \sup_h \left(\int_{\mathbb{R}^2} f h n_\infty dx - \frac{1}{2\Lambda} \int_{\mathbb{R}^2} |\nabla h|^2 dx \right)$$

where the supremum is taken on both sides on all functions h in $L^2(\mathbb{R}^2, n_\infty dx)$ such that h satisfies (4.3). Since semi-definite positive quadratic forms are involved, the suprema are achieved by convexity. For the left hand side, we find that the optimal function satisfies

$$f = h + \mu f_{0,0}$$

for some Lagrange multiplier $\mu \in \mathbb{R}$. However, if we multiply by $f_{0,0} n_\infty$, we get that $\mu = 0$, so that the left hand side of the inequality is simply $\frac{1}{2} \int_{\mathbb{R}^2} f^2 n_\infty dx$. As for the right hand side, we find that the optimal function f is such that

$$f n_\infty = -\frac{1}{\Lambda} \Delta h + \mu f_{0,0} n_\infty$$

for some Lagrange multiplier $\mu \in \mathbb{R}$. In that case, if we multiply by $(-\Delta)^{-1}(f_{0,0} n_\infty) = f_{0,0}$, we get that

$$\mu \int_{\mathbb{R}^2} f_{0,0}^2 n_\infty dx = \frac{1}{\Lambda} \int_{\mathbb{R}^2} \Delta h (-\Delta)^{-1}(f_{0,0} n_\infty) dx = -\frac{1}{\Lambda} \int_{\mathbb{R}^2} h f_{0,0} n_\infty dx = 0$$

thus proving that $\mu = 0$ as well. Hence the right hand side of the inequality is simply $\frac{\Lambda}{2} \int_{\mathbb{R}^2} f n_\infty (-\Delta)^{-1}(f n_\infty) dx$, which establishes (5.1). It is left to the reader to check that Inequality (5.2) can also be deduced from (5.1) by a similar argument.

Step 2. Let us prove that (5.2) holds for some $\Lambda > 1$. Consider an optimizing sequence of functions $(h_n)_{n \geq 1}$ such that $\int_{\mathbb{R}^2} h_n^2 n_\infty dx = 1$ and $\int_{\mathbb{R}^2} h_n f_{0,0} n_\infty dx = 0$ for any $n \geq 1$, and $\lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} |\nabla h_n|^2 n_\infty dx = \Lambda$. As in the proof of Lemma 14, we are going to use the IMS truncation method. Consider a smooth function χ with the following properties: $0 \leq \chi \leq 1$, $\chi(x) = 1$ for any $x \in B(0, 1)$, $\chi(x) = 0$ for any $x \in \mathbb{R}^2 \setminus B(0, 2)$, and define $\chi_R(x) := \chi(x/R)$ for any $x \in \mathbb{R}^2$. It is standard in concentration-compactness methods that for any $\varepsilon > 0$, one can find a sequence of positive numbers $(R_n)_{n \geq 1}$ such that

$$h_n^{(1)} = \chi_{R_n} h_n \quad \text{and} \quad h_n^{(2)} = \sqrt{1 - \chi_{R_n}^2} h_n,$$

and, up to the extraction of a subsequence, there exists a function h such that

$$\int_{\mathbb{R}^2} |\nabla h_n^{(1)}|^2 n_\infty dx \geq \eta \Lambda - \varepsilon \quad \text{and} \quad \int_{\mathbb{R}^2} |\nabla h_n^{(2)}|^2 n_\infty dx \geq (1 - \eta) \Lambda - \varepsilon$$

for some $\eta \in [0, 1]$, where the sequence $(\nabla h_n^{(1)})_{n \geq 1}$ strongly converges to ∇h and

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} |h_n^{(1)}|^2 n_\infty dx = \int_{\mathbb{R}^2} h^2 n_\infty dx =: \theta$$

(this implies the strong convergence of $(h_n^{(1)})_{n \geq 1}$ towards h in $L^2(n_\infty dx)$) because

$$\int_{\mathbb{R}^2 \setminus B(0, R)} |h_n^{(1)}|^2 n_\infty dx \leq \left(\int_{\mathbb{R}^2} |h_n^{(1)}|^{\frac{2d}{d-2}} n_\infty dx \right)^{\frac{d-2}{d}} \left(\int_{\mathbb{R}^2 \setminus B(0, R)} n_\infty^{\frac{d}{2}} dx \right)^{\frac{2}{d}}$$

is uniformly small as $R \rightarrow \infty$ by Sobolev's inequality and because the last term of the right hand side is such that $\lim_{R \rightarrow \infty} \int_{\mathbb{R}^2 \setminus B(0, R)} n_\infty^{\frac{d}{2}} dx = 0$. Of course, we know that

$$\eta \Lambda \geq \int_{\mathbb{R}^2} |\nabla h|^2 dx \geq \Lambda \theta$$

by definition of Λ . The above estimate also guarantees that

$$\int_{\mathbb{R}^2 \setminus B(0, R)} |h_n^{(2)}|^2 n_\infty dx =: \varepsilon_n \rightarrow 0 \quad \text{as} \quad n \rightarrow \infty.$$

By construction of $(h_n^{(1)})_{n \geq 1}$ and $(h_n^{(2)})_{n \geq 1}$, we know that $|h_n^{(1)}|^2 + |h_n^{(2)}|^2 = |h_n|^2$ and hence $\theta = 1$. This also means that $\eta = 1$ and hence h is a minimizer, since the constraint passes to the limit:

$$\int_{\mathbb{R}^2} h f_{0,0} n_\infty dx = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} h_n f_{0,0} n_\infty dx = 0.$$

The function h is a solution of the Euler-Lagrange equation:

$$-\Delta h = \Lambda h n_\infty .$$

By Proposition 9, if $\Lambda = 1$, then h and $f_{0,0}$ are collinear, which is a contradiction with the constraint. This proves that $\Lambda > 1$. \square

Notice that the functions n_∞ and c_∞ being radial symmetric, we know that a decomposition into spherical harmonics allows to reduce the problem of computing all eigenvalues to radially symmetric eigenvalue problems. This provides a method to compute the explicit value of Λ , at least numerically.

Remark 4. *Inequality (5.2) is a Poincaré inequality, which has already been established in [14] as a linearized version of an Onofri type inequality. This Onofri inequality is dual of the logarithmic Hardy-Littlewood-Sobolev type inequality that has been established in [10] and according to which the free energy functional $F[n]$ is nonnegative.*

A straightforward consequence of Theorem 17 is that we can estimate $\int_{\mathbb{R}^2} f^2 n_\infty dx$ in terms of $\mathcal{Q}_1[f] = \int_{\mathbb{R}^2} f (f - g c_\infty) n_\infty dx$.

Corollary 18. *For the same value of $\Lambda > 1$ as in Theorem 17, we have*

$$\int_{\mathbb{R}^2} f^2 n_\infty dx \leq \frac{\Lambda}{\Lambda - 1} \mathcal{Q}_1[f]$$

for any $f \in L^2(\mathbb{R}^2, n_\infty dx)$ such that (4.3) holds.

Proof. We may indeed write

$$\mathcal{Q}_1[f] = \frac{\Lambda - 1}{\Lambda} \int_{\mathbb{R}^2} f^2 n_\infty dx + \frac{1}{\Lambda} \left(\int_{\mathbb{R}^2} |f|^2 n_\infty dx - \Lambda \int_{\mathbb{R}^2} f n_\infty (-\Delta)^{-1} (f n_\infty) dx \right)$$

and use the fact that the last term of the right hand side is nonnegative. \square

6. THE LARGE TIME BEHAVIOR

This section is devoted to the proof of Theorem 1. Our approach is guided by the analysis of the evolution equation corresponding to the linearization of the Keller-Segel system: see Section 6.1. The key estimates for the nonlinear evolution problem have been stated in Theorem 15, Theorem 17, and Corollary 18. Nonlinear terms are estimated using Corollary 7.

6.1. A linearized evolution problem. We recall that the restriction of \mathcal{L} to $\mathcal{D}(\mathcal{L}_1)$ is a self-adjoint operator with domain $\mathcal{D}(\mathcal{L}_2)$, such that

$$\langle f, \mathcal{L} f \rangle = -\mathcal{Q}_2[f] \quad \forall f \in \mathcal{D}(\mathcal{L}_2) .$$

and $\text{Ker}(\mathcal{L}) \cap \mathcal{D}(\mathcal{L}_2) = \{0\}$

By Proposition 11, any solution $(t, x) \mapsto f(t, x)$ of the linearized Keller-Segel model

$$\begin{cases} \frac{\partial f}{\partial t} = \mathcal{L} f & x \in \mathbb{R}^2, \quad t > 0, \\ -\Delta(g c_\infty) = f n_\infty & x \in \mathbb{R}^2, \quad t > 0, \end{cases}$$

has an exponential decay, since we know that

$$\frac{d}{dt} \langle f(t, \cdot), f(t, \cdot) \rangle = 2 \langle f(t, \cdot), \mathcal{L} f(t, \cdot) \rangle ,$$

that is

$$\frac{d}{dt} \mathcal{Q}_1[f(t, \cdot)] = -2 \mathcal{Q}_2[f(t, \cdot)] \leq -2 \mathcal{Q}_1[f(t, \cdot)]$$

by Theorem 15. Hence we obtain

$$\mathcal{Q}_1[f(t, \cdot)] \leq \mathcal{Q}_1[f(0, \cdot)] e^{-2t} \quad \forall t \in \mathbb{R}^+ .$$

Here we adopt the usual convention that $\mathbf{Q}_2[f] = +\infty$ for any $f \in \mathcal{D}(L_1) \setminus \mathcal{D}(L_2)$.

6.2. Proof of Theorem 1. As in [8], Eq. (1.3) can be rewritten in terms of $f = (n - n_\infty)/n_\infty$ and $g = (c - c_\infty)/c_\infty$ in the form of (4.1), that is

$$\frac{\partial f}{\partial t} = \mathcal{L} f - \frac{1}{n_\infty} \nabla [n_\infty f \nabla (g c_\infty)] .$$

The computation for the linearized problem established in Section 6.1 can be adapted to the nonlinear case and gives

$$\frac{d}{dt} \mathbf{Q}_1[f(t, \cdot)] = -2 \mathbf{Q}_2[f(t, \cdot)] + \int_{\mathbb{R}^2} \nabla (f - g c_\infty) f n_\infty \cdot \nabla (g c_\infty) dx ,$$

with $-2 \mathbf{Q}_2[f(t, \cdot)] \leq -2 \mathbf{Q}_1[f(t, \cdot)]$ according to Theorem 15. To get an estimate on the asymptotic behaviour, we have to establish an estimate of the last term of the right hand side, which is cubic in terms of f . For this purpose, we apply Hölder's inequality and Lemma 6 to get

$$\left(\int_{\mathbb{R}^2} \nabla (f - g c_\infty) f n_\infty \cdot \nabla (g c_\infty) dx \right)^2 \leq \mathbf{Q}_2[f] \int_{\mathbb{R}^2} f^2 n_\infty dx \|\nabla (g c_\infty)\|_{L^\infty(\mathbb{R}^2)}^2 .$$

Using Lemma 6 and Corollary 18, we find that the right hand side can be bounded by

$$C(\varepsilon) \frac{\Lambda}{\Lambda - 1} \mathbf{Q}_1[f] \mathbf{Q}_2[f] \left(\|f n_\infty\|_{L^{2-\varepsilon}(\mathbb{R}^2)} + \|f n_\infty\|_{L^{2+\varepsilon}(\mathbb{R}^2)} \right) .$$

From [10, Theorem 1.2] and Corollary 7 we know that $\lim_{t \rightarrow \infty} \|f(t, \cdot) n_\infty\|_{L^1(\mathbb{R}^2)} = 0$ and $\lim_{t \rightarrow \infty} \|f(t, \cdot) n_\infty\|_{L^\infty(\mathbb{R}^2)} = 0$ and therefore, for any given $\varepsilon \in (0, 1)$, there exists a continuous function $t \mapsto \delta(t, \varepsilon)$ with $\lim_{t \rightarrow \infty} \delta(t, \varepsilon) = 0$ such that

$$\|f(t, \cdot) n_\infty\|_{L^{2-\varepsilon}(\mathbb{R}^2)} + \|f(t, \cdot) n_\infty\|_{L^{2+\varepsilon}(\mathbb{R}^2)} \leq \delta(t, \varepsilon) .$$

As a consequence, we know that

$$\frac{d}{dt} \mathbf{Q}_1[f(t, \cdot)] \leq -2 \mathbf{Q}_2[f(t, \cdot)] + \delta(t, \varepsilon) \sqrt{\mathbf{Q}_1[f(t, \cdot)] \mathbf{Q}_2[f(t, \cdot)]} \leq (\delta(t, \varepsilon) - 2) \mathbf{Q}_2[f(t, \cdot)] ,$$

where the last inequality is a consequence of Theorem 15. This proves that $\mathbf{Q}_1[f(t, \cdot)]$ is uniformly bounded with respect to t as $t \rightarrow \infty$: there exists a constant $\mathcal{Q} > 0$ such that

$$\mathbf{Q}_1[f(t, \cdot)] \leq \mathcal{Q} \quad \forall t \geq 0 .$$

Now we can give a more detailed estimate. Using again Hölder's inequality, we find that

$$\int_{\mathbb{R}^2} (|f| n_\infty)^{2+\varepsilon} dx \leq \mathbf{Q}_1[f] \|f n_\infty\|_{L^\infty(\mathbb{R}^2)}^\varepsilon \|n_\infty\|_{L^\infty(\mathbb{R}^2)} ,$$

that is

$$\|f n_\infty\|_{L^{2+\varepsilon}(\mathbb{R}^2)} \leq (\mathbf{Q}_1[f])^{\frac{1}{2+\varepsilon}} \|f n_\infty\|_{L^\infty(\mathbb{R}^2)}^{\frac{\varepsilon}{2+\varepsilon}} \|n_\infty\|_{L^\infty(\mathbb{R}^2)}^{\frac{1}{2+\varepsilon}}$$

and

$$\|f n_\infty\|_{L^{2-\varepsilon}(\mathbb{R}^2)} \leq \|f n_\infty\|_{L^1(\mathbb{R}^2)}^{\frac{\varepsilon}{2-\varepsilon}} \|f n_\infty\|_{L^2(\mathbb{R}^2)}^{\frac{2(1-\varepsilon)}{2-\varepsilon}} \leq \|f n_\infty\|_{L^1(\mathbb{R}^2)}^{\frac{\varepsilon}{2-\varepsilon}} (\mathbf{Q}_1[f])^{\frac{1-\varepsilon}{2-\varepsilon}} \|n_\infty\|_{L^\infty(\mathbb{R}^2)}^{\frac{1-\varepsilon}{2-\varepsilon}} .$$

Recall that $\lim_{t \rightarrow \infty} \|f(t, \cdot) n_\infty\|_{L^1(\mathbb{R}^2)} = 0$ and $\lim_{t \rightarrow \infty} \|f(t, \cdot) n_\infty\|_{L^\infty(\mathbb{R}^2)} = 0$ according to [10, Theorem 1.2] and Corollary 7 respectively. For any given $\varepsilon \in (0, 1)$, there exists a continuous function (that we again denote by δ): $t \mapsto \delta(t, \varepsilon)$, with $\lim_{t \rightarrow \infty} \delta(t, \varepsilon) = 0$, such that

$$\frac{d}{dt} \mathbf{Q}_1[f] \leq -2 \mathbf{Q}_2[f] + \delta(t, \varepsilon) (\mathbf{Q}_2[f])^{\frac{1}{2}} (\mathbf{Q}_1[f])^{\frac{1}{2}} \left((\mathbf{Q}_1[f])^{\frac{1-\varepsilon}{2-\varepsilon}} + (\mathbf{Q}_1[f])^{\frac{1}{2+\varepsilon}} \right) .$$

Since $Q_1[f] \leq Q_2[f]$, this provides the estimate

$$\frac{d}{dt} Q_1[f] \leq -\sqrt{Q_2[f]} \left(2\sqrt{Q_2[f]} - \delta(t, \varepsilon) (Q_1[f])^{\frac{4-3\varepsilon}{2(2-\varepsilon)}} - \delta(t, \varepsilon) (Q_1[f])^{\frac{4+\varepsilon}{2(2+\varepsilon)}} \right).$$

Altogether, this proves that

$$\frac{d}{dt} Q_1[f] \leq -Q_2[f] \left[2 - \delta(t, \varepsilon) \left(Q^{\frac{1-\varepsilon}{2-\varepsilon}} + Q^{\frac{1}{2+\varepsilon}} \right) \right].$$

For $t > 0$ large enough, the right hand side becomes negative and we have found that

$$\frac{d}{dt} Q_1[f(t, \cdot)] \leq -Q_1[f(t, \cdot)] \left[2 - \delta(t, \varepsilon) \left(Q^{\frac{1-\varepsilon}{2-\varepsilon}} + Q^{\frac{1}{2+\varepsilon}} \right) \right],$$

thus showing that

$$\limsup_{t \rightarrow \infty} e^{(2-\eta)t} Q_1[f(t, \cdot)] < \infty$$

for any $\eta \in (0, 2)$. Actually we know from the above estimates that

$$2\sqrt{Q_2[f(t, \cdot)]} - \delta(t, \varepsilon) (Q_1[f(t, \cdot)])^{\frac{4-3\varepsilon}{2(2-\varepsilon)}} - \delta(t, \varepsilon) (Q_1[f(t, \cdot)])^{\frac{4+\varepsilon}{2(2+\varepsilon)}}$$

is positive for $t > 0$, large enough, thus showing that

$$\frac{d}{dt} Q_1[f(t, \cdot)] \leq -Q_1[f(t, \cdot)] \left[2 - \delta(t, \varepsilon) \left(Q_1[f(t, \cdot)]^{\frac{1-\varepsilon}{2-\varepsilon}} + Q_1[f(t, \cdot)]^{\frac{1}{2+\varepsilon}} \right) \right].$$

As a consequence, we finally get that

$$\limsup_{t \rightarrow \infty} e^{2t} Q_1[f(t, \cdot)] < \infty,$$

which completes the proof of Theorem 1. \square

Acknowledgments. The authors acknowledge support by the ANR projects *CBDif-Fr* and *EVOL* (JD), and by the MathAmSud project *NAPDE* (JC and JD).

© 2012 by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

REFERENCES

- [1] C. BANDLE, *Isoperimetric inequalities and applications*, vol. 7 of Monographs and Studies in Mathematics, Pitman (Advanced Publishing Program), Boston, Mass., 1980.
- [2] P. BILER, *Growth and accretion of mass in an astrophysical model*, Appl. Math. (Warsaw), 23 (1995), pp. 179–189.
- [3] P. BILER, L. CORRIAS, AND J. DOLBEAULT, *Large mass self-similar solutions of the parabolic–parabolic keller–segel model of chemotaxis*, Journal of Mathematical Biology, 63 (2011), pp. 1–32. 10.1007/s00285-010-0357-5.
- [4] P. BILER, J. DOLBEAULT, M. J. ESTEBAN, P. A. MARKOWICH, AND T. NADZIEJA, *Steady states for Streater’s energy-transport models of self-gravitating particles*, IMA Vol. Math. Appl., 135 (2004), pp. 37–56.
- [5] P. BILER, G. KARCH, P. LAURENÇOT, AND T. NADZIEJA, *The 8π -problem for radially symmetric solutions of a chemotaxis model in the plane*, Math. Methods Appl. Sci., 29 (2006), pp. 1563–1583.
- [6] A. BLANCHET, E. A. CARLEN, AND J. A. CARRILLO, *Functional inequalities, thick tails and asymptotics for the critical mass Patlak-Keller-Segel model*, Journal of Functional Analysis, 262 (2012), pp. 2142 – 2230.
- [7] A. BLANCHET, J. A. CARRILLO, AND N. MASMOUDI, *Infinite time aggregation for the critical Patlak-Keller-Segel model in \mathbb{R}^2* , Comm. Pure Appl. Math., 61 (2008), pp. 1449–1481.
- [8] A. BLANCHET, J. DOLBEAULT, M. ESCOBEDO, AND J. FERNÁNDEZ, *Asymptotic behaviour for small mass in the two-dimensional parabolic-elliptic Keller-Segel model*, Journal of Mathematical Analysis and Applications, 361 (2010), pp. 533 – 542.
- [9] A. BLANCHET, J. DOLBEAULT, AND M. KOWALCZYK, *Stochastic Stokes’ drift, homogenized functional inequalities, and large time behavior of brownian ratchets*, SIAM Journal on Mathematical Analysis, 41 (2009), pp. 46–76.
- [10] A. BLANCHET, J. DOLBEAULT, AND B. PERTHAME, *Two-dimensional Keller-Segel model: optimal critical mass and qualitative properties of the solutions*, Electron. J. Differential Equations, 44, 32 pages (2006).

- [11] V. CALVEZ AND J. A. CARRILLO, *Refined asymptotics for the subcritical Keller-Segel system and related functional inequalities*. Preprint ArXiv 1007.2837, to appear in Proc. AMS, Article electronically published on February 23, 2012.
- [12] J. CAMPOS AND J. DOLBEAULT, *A functional framework for the keller-segel system: logarithmic hardy-littlewood-sobolev and related spectral gap inequalities*. Preprint, 2012.
- [13] ———, *A numerical study of linearized Keller-Segel operator in self-similar variables*. Preprint, 2012.
- [14] J. CAMPOS SERRANO AND J. DOLBEAULT, *A functional framework for the Keller-Segel system: logarithmic Hardy-Littlewood-Sobolev and related spectral gap inequalities*. Preprint.
- [15] J. A. CARRILLO, L. CHEN, J.-G. LIU, AND J. WANG, *A note on the subcritical two dimensional Keller-Segel system*. Preprint.
- [16] J. A. CARRILLO, R. J. MCCANN, AND C. VILLANI, *Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates*, Rev. Mat. Iberoamericana, 19 (2003), pp. 971–1018.
- [17] ———, *Contractions in the 2-Wasserstein length space and thermalization of granular media*, Arch. Ration. Mech. Anal., 179 (2006), pp. 217–263.
- [18] J. I. DÍAZ, *Nonlinear partial differential equations and free boundaries. Vol. I*, vol. 106 of Research Notes in Mathematics, Pitman (Advanced Publishing Program), Boston, MA, 1985. Elliptic equations.
- [19] J. I. DIAZ AND J. MOSSINO, *Isoperimetric inequalities in the parabolic obstacle problems*, J. Math. Pures Appl. (9), 71 (1992), pp. 233–266.
- [20] J. I. DIAZ AND T. NAGAI, *Symmetrization in a parabolic-elliptic system related to chemotaxis*, Adv. Math. Sci. Appl., 5 (1995), pp. 659–680.
- [21] J. I. DIAZ, T. NAGAI, AND J.-M. RAKOTOSON, *Symmetrization techniques on unbounded domains: application to a chemotaxis system on \mathbb{R}^N* , J. Differential Equations, 145 (1998), pp. 156–183.
- [22] J. DOLBEAULT, *Sobolev and Hardy-Littlewood-Sobolev inequalities: duality and fast diffusion*, Math. Research Letters, (2012).
- [23] J. DOLBEAULT AND B. PERTHAME, *Optimal critical mass in the two-dimensional Keller-Segel model in \mathbb{R}^2* , C. R. Math. Acad. Sci. Paris, 339 (2004), pp. 611–616.
- [24] T. HILLEN AND K. J. PAINTER, *A user's guide to PDE models for chemotaxis*, J. Math. Biol., 58 (2009), pp. 183–217.
- [25] D. HORSTMANN, *From 1970 until present: the Keller-Segel model in chemotaxis and its consequences. I*, Jahresber. Deutsch. Math.-Verein., 105 (2003), pp. 103–165.
- [26] ———, *From 1970 until present: the Keller-Segel model in chemotaxis and its consequences. II*, Jahresber. Deutsch. Math.-Verein., 106 (2004), pp. 51–69.
- [27] W. JÄGER AND S. LUCKHAUS, *On explosions of solutions to a system of partial differential equations modelling chemotaxis*, Trans. Amer. Math. Soc., 329 (1992), pp. 819–824.
- [28] P.-L. LIONS, *Principe de concentration-compactité en calcul des variations*, C. R. Acad. Sci. Paris Sér. I Math., 294 (1982), pp. 261–264.
- [29] ———, *Applications de la méthode de concentration-compactité à l'existence de fonctions extrémales*, C. R. Acad. Sci. Paris Sér. I Math., 296 (1983), pp. 645–648.
- [30] P.-L. LIONS, *The concentration-compactness principle in the calculus of variations. The locally compact case. II*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 1 (1984), pp. 223–283.
- [31] ———, *The concentration-compactness principle in the calculus of variations. The limit case. I*, Rev. Mat. Iberoamericana, 1 (1985), pp. 145–201.
- [32] ———, *The concentration-compactness principle in the calculus of variations. The limit case. II*, Rev. Mat. Iberoamericana, 1 (1985), pp. 45–121.
- [33] J. MOSSINO, *Inégalités isopérimétriques et applications en physique*, Travaux en Cours. [Works in Progress], Hermann, Paris, 1984.
- [34] Y. NAITO AND T. SUZUKI, *Self-similar solutions to a nonlinear parabolic-elliptic system*, in Proceedings of Third East Asia Partial Differential Equation Conference, vol. 8, 2004, pp. 43–55.
- [35] A. PERSSON, *Bounds for the discrete part of the spectrum of a semi-bounded Schrödinger operator*, Math. Scand., 8 (1960), pp. 143–153.
- [36] B. PERTHAME, *Transport equations in biology*, Frontiers in Mathematics, Birkhäuser Verlag, Basel, 2007.
- [37] G. TALENTI, *Nonlinear elliptic equations, rearrangements of functions and Orlicz spaces*, Ann. Mat. Pura Appl. (4), 120 (1979), pp. 160–184.
- [38] M. J. TINDALL, P. K. MAINI, S. L. PORTER, AND J. P. ARMITAGE, *Overview of mathematical approaches used to model bacterial chemotaxis. II. Bacterial populations*, Bull. Math. Biol., 70 (2008), pp. 1570–1607.
- [39] M. J. TINDALL, S. L. PORTER, P. K. MAINI, G. GAGLIA, AND J. P. ARMITAGE, *Overview of mathematical approaches used to model bacterial chemotaxis. I. The single cell*, Bull. Math. Biol., 70 (2008), pp. 1525–1569.

(J. Dolbeault) CEREMADE (UMR CNRS no. 7534), UNIVERSITÉ PARIS-DAUPHINE, PLACE DE LATTRE DE TASSIGNY, F-75775 PARIS CÉDEX 16, FRANCE

E-mail address: `dolbeaul@ceremade.dauphine.fr`

(J.F. Campos) CEREMADE (UMR CNRS no. 7534), UNIVERSITÉ PARIS-DAUPHINE, PLACE DE LATTRE DE TASSIGNY, F-75775 PARIS CÉDEX 16, FRANCE & DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CMM, UNIVERSIDAD DE CHILE, CASILLA 170 CORREO 3, SANTIAGO, CHILE

E-mail address: `campos@ceremade.dauphine.fr`, `juanfcampos@gmail.com`